

Commutators on ℓ_1

Detelin Dosev ¹

Abstract

The main result is that the commutators on ℓ_1 are the operators not of the form $\lambda I + K$ with $\lambda \neq 0$ and K compact. We generalize Apostol's technique (1972, Rev. Roum. Math. Appl. 17, 1513 - 1534) to obtain this result and use this generalization to obtain partial results about the commutators on spaces \mathcal{X} which can be represented as $\mathcal{X} \simeq \left(\bigoplus_{i=0}^{\infty} \mathcal{X} \right)_p$ for some $1 \leq p < \infty$ or $p = 0$. In particular, it is shown that every compact operator on L_1 is a commutator. A characterization of the commutators on $\ell_{p_1} \oplus \ell_{p_2} \oplus \dots \oplus \ell_{p_n}$ is given. We also show that strictly singular operators on ℓ_∞ are commutators.

1 Introduction

The commutator of two elements A and B in a Banach algebra is given by

$$[A, B] = AB - BA.$$

A natural problem that arises in the study of derivations on a Banach algebra is to classify the commutators in the algebra. The first major contribution was due to Wintner([11]), who proved that the identity in a unital Banach algebra is not a commutator. This immediately implies that no non-zero multiple of the identity is a commutator and with no effort one can also obtain that no operator of the form $\lambda I + K$, where K is a compact operator and $\lambda \neq 0$, is a commutator in the Banach algebra $\mathcal{L}(\mathcal{X})$ of all bounded linear operators on the Banach space \mathcal{X} . The latter fact can be easily seen just by observing that the quotient algebra $\mathcal{L}(\mathcal{X})/K(\mathcal{X})$ ($K(\mathcal{X})$ is the space of compact operators on \mathcal{X}) also satisfies the conditions of the Wintner's theorem. Let us note also that instead of considering the ideal of compact operators one can consider any proper ideal in $\mathcal{L}(\mathcal{X})$. The observations we made are valid in this case as well.

For a Banach space \mathcal{X} for which there is a unique maximal proper ideal in $\mathcal{L}(\mathcal{X})$ (which is the case for the spaces L_p and ℓ_p for $1 \leq p \leq \infty$ and c_0) one can hope to obtain a complete classification of the commutators on the space. The natural conjecture is that the only operators on \mathcal{X} that are not commutators are the ones of the form $\lambda I + K$, where K belongs to the unique maximal ideal in $\mathcal{L}(\mathcal{X})$ and $\lambda \neq 0$. In 1965 Brown and Percy ([1]) made a breakthrough in this direction by proving that this is in fact a classification of the commutators on a Hilbert space. Note that if $\mathcal{X} = \ell_p$ ($1 \leq p < \infty$) or $\mathcal{X} = c_0$, the ideal of compact operators $K(\mathcal{X})$ is the largest proper ideal in $\mathcal{L}(\mathcal{X})$ ([12, Theorem 6.2]). Whitley's proof actually shows that the ideal of strictly singular operators is the largest ideal in the aforementioned spaces, but as he points out, a result of Feldman, Gohberg and Markus ([14]) shows that the compact operators are in fact the only closed proper ideal in $L(\mathcal{X})$ for $\mathcal{X} = \ell_p$ ($1 \leq p < \infty$) or $\mathcal{X} = c_0$. In 1972, Apostol ([2]) proved that every non-commutator on the space ℓ_p for $1 < p < \infty$ is of the form $\lambda I + K$, where K is compact and $\lambda \neq 0$. One year later he proved that the same classification holds in the case of $\mathcal{X} = c_0$ ([3]). While Apostol's approach in ([2]) gave some information about the commutators in $\mathcal{L}(\ell_1)$, he was unable to give a complete characterization. His proof uses the fact that the unit vector basis in ℓ_p for $1 < p < \infty$ is shrinking and this does not hold for ℓ_1 . We overcome this obstacle by using the structure of the infinite dimensional subspaces of ℓ_1 rather than just

¹Research supported in part by NSF grant DMS-0503688

the properties of the basis.

In the Section 2 we study spaces \mathcal{X} for which $\mathcal{X} \simeq \left(\bigoplus_{i=0}^{\infty} \mathcal{X} \right)_p$, $1 \leq p < \infty$ or $p = 0$, where for $p \geq 1$

$$\left(\bigoplus_{i=0}^{\infty} \mathcal{X} \right)_p := \{f = (f_1, f_2, \dots) : f_i \in \mathcal{X}, i = 1, 2, \dots, \|f\|^p = \sum_{i=0}^{\infty} \|f_i\|^p < \infty\}$$

(for $p = 0$, $\|f\| = \sup_{0 \leq i < \infty} \|f_i\|$), and we generally assume that $\mathcal{L}(\mathcal{X})$ has a largest proper ideal. The notion of a decomposition of \mathcal{X} will be introduced and it will be shown how it can be used to obtain results about commutators on these spaces. In Section 3 we show that the compact operators on \mathcal{X} , where \mathcal{X} admits the decomposition $\mathcal{X} \simeq \left(\bigoplus_{i=0}^{\infty} \mathcal{X} \right)_p$ (in the case $p = 1$ we will assume that $\mathcal{X} = L_1$ or $\mathcal{X} = \ell_1$), are commutators and as a corollary we will get that an operator that has a compact restriction to a sufficiently large subspace of \mathcal{X} is also a commutator. Section 3 contains our main result - Theorem 4.6 - which shows that the the only operators on ℓ_1 that are not commutators are the ones of the form $\lambda I + K$, where K is a compact operator and $\lambda \neq 0$. In the last section we give a characterization of the commutators on $\ell_{p_1} \oplus \ell_{p_2} \oplus \dots \oplus \ell_{p_n}$, where $1 \leq p_n < p_{n-1} < \dots < p_1 < \infty$ and we also show that the strictly singular operators on ℓ_{∞} are commutators.

The author thanks W.B. Johnson for useful conversations and suggestions on the subject of this paper.

2 Notation and basic results

We will follow the ideas in Apostol's paper [2], which in turn extend those of Brown & Pearcy [1] and earlier work referenced in [1], to develop a base for investigating the commutators on ℓ_1 and L_p ($1 \leq p < \infty$).

Consider a fixed decomposition $\mathcal{D} = \{X_i\}$ of \mathcal{X} , $1 \leq p < \infty$ or $p = 0$, meaning $\mathcal{X} \simeq \left(\bigoplus_{i=0}^{\infty} X_i \right)_p$, where X_i are complemented subspaces of \mathcal{X} which are also isomorphic to \mathcal{X} . Let $\psi_i : X_i \rightarrow \mathcal{X}$ be an isomorphism and let us also assume that $\|\psi_i^{-1}\| = 1$ for $i \in \mathbb{N}$ and $\lambda = \sup_{i \in \mathbb{N}} \|\psi_i\| < \infty$. Denote by P_i the projection from \mathcal{X} onto X_i . Let us also have a collection of uniformly bounded isomorphisms $\{\varphi_i\}_1^{\infty}$ as shown below

$$X_0 \xrightarrow{\varphi_0} X_1 \xrightarrow{\varphi_1} X_2 \xrightarrow{\varphi_2} X_3 \xrightarrow{\varphi_3} \dots$$

which make the following diagram commute

$$\begin{array}{ccc} X_i & \xrightarrow{\varphi_i} & X_{i+1} \\ & \searrow \psi_i & \downarrow \psi_{i+1} \\ & & \mathcal{X} \end{array}$$

for every $i \in \mathbb{N}$. Clearly $\varphi_i = \psi_{i+1}^{-1} \circ \psi_i$ and $\|\varphi_i\| \leq \|\psi_i\| \leq \lambda$, $\|\varphi_i^{-1}\| \leq \lambda$. Assume that for $p \geq 1$ we also have $\left\| \sum_{j=1}^n P_{i_j} x \right\|^p = \sum_{j=1}^n \|P_{i_j} x\|^p$ for every $n > 0$ and $i_1, i_2, \dots, i_n > 0$ with $i_l \neq i_k$ for $l \neq k$. Note that using the last condition in the case $p \geq 1$ we have $\|P_i\| \leq \|I - P_0\| \leq \|P_0\| + 1 = C_1$. The last inequality is clearly true for $p = 0$. For $x = (x_i) \in \mathcal{X}$, $x_i \in X_i$ define the following two operators :

$$R_{\mathcal{D}}(x) = \sum_{i=0}^{\infty} \varphi_i(x_i) \quad , \quad L_{\mathcal{D}}(x) = \sum_{i=0}^{\infty} \varphi_i^{-1}(x_{i+1}).$$

The operators $L_{\mathcal{D}}$ and $R_{\mathcal{D}}$ are, respectively, the left and the right shift associated with the decomposition \mathcal{D} . As one may observe, the operators $L_{\mathcal{D}}$ and $R_{\mathcal{D}}$ move the components of x one position to the left/right, respectively, via the isomorphisms φ_i . Throughout the paper we will simply use the letters L and R for the shifts when the decomposition \mathcal{D} associated with the shifts is clear from the context. Our first proposition shows some basic properties of the left and the right shift as well as the fact that all the powers of L and R are uniformly bounded, which will play an important role in the sequel.

Proposition 2.1. *Let \mathcal{D} be a decomposition of \mathcal{X} . Then we have*

$$\|L^n\| \leq 2\lambda C_1, \quad \|R^n\| \leq 2\lambda C_1 \quad \text{for every } n = 1, 2, \dots \quad (1)$$

$$LR = I, \quad RL = I - P_0, \quad RP_i = P_{i+1}R, \quad P_iL = LP_{i+1} \quad \text{for } i \geq 0. \quad (2)$$

$$\lim_{n \rightarrow \infty} \|L^n(x)\|_p = 0 \quad \text{for all } 1 \leq p < \infty \text{ and } p = 0. \quad (3)$$

Proof. The relations $LR = I$, $RL = I - P_0$ are clear from the definition of the left and the right shift. For $x = (x_i) \in \mathcal{X}$, $x_i \in X_i$ we have

$$\begin{aligned} P_{i+1}R(x) &= P_{i+1} \left(\sum_{i=0}^{\infty} \varphi_i(x_i) \right) = \varphi_i(x_i) = RP_i(x) \\ P_iL(x) &= P_i \left(\sum_{i=0}^{\infty} \varphi_i^{-1}(x_{i+1}) \right) = \varphi_i^{-1}(x_{i+1}) = LP_{i+1}(x). \end{aligned}$$

Now using the fact that

$$\begin{aligned} \varphi_{k+n} \circ \varphi_{k+n-1} \circ \dots \circ \varphi_k &= \psi_{k+n+1}^{-1} \circ \psi_k \\ \varphi_k^{-1} \circ \varphi_{k+1}^{-1} \circ \dots \circ \varphi_{k+n}^{-1} &= \psi_k^{-1} \circ \psi_{k+n+1} \end{aligned}$$

we deduce that

$$\begin{aligned} R^{n+1}(x) &= \sum_{i=0}^{\infty} \varphi_{i+n} \circ \varphi_{i+n-1} \circ \dots \circ \varphi_i(x_i) = \sum_{i=0}^{\infty} \psi_{i+n+1}^{-1} \circ \psi_i(x_i) \\ L^{n+1}(x) &= \sum_{i=0}^{\infty} \varphi_i^{-1} \circ \varphi_{i+1}^{-1} \circ \dots \circ \varphi_{i+n}^{-1}(x_{i+n+1}) = \sum_{i=0}^{\infty} \psi_i^{-1} \circ \psi_{i+n+1}(x_{i+n+1}) \end{aligned}$$

for $n = 0, 1, \dots$. Finally we estimate the norms of $\|R^n\|$ and $\|L^n\|$ for $p \geq 1$:

$$\begin{aligned} \|R^{n+1}(x)\|_p^p &= \left\| \sum_{i=0}^{\infty} \psi_{i+n+1}^{-1} \circ \psi_i(x_i) \right\|_p^p \leq \sum_{i=0}^{\infty} \|\psi_{i+n+1}^{-1} \circ \psi_i\|_p^p \|x_i\|_p^p \\ &\leq \sum_{i=0}^{\infty} \lambda^p \|x_i\|_p^p = \lambda^p \|P_0x\|_p^p + \lambda^p \left\| \sum_{i=1}^{\infty} P_i x \right\|_p^p \\ &= \lambda^p \|P_0x\|_p^p + \lambda^p \|(I - P_0)x\|_p^p \leq 2^p \lambda^p (\|P_0\| + 1)^p \|x\|_p^p \end{aligned}$$

$$\begin{aligned}
\|L^{n+1}(x)\|_p^p &= \|\psi_0^{-1} \circ \psi_{n+1}(x_{n+1}) + \sum_{i=1}^{\infty} \psi_i^{-1} \circ \psi_{i+n+1}(x_{i+n+1})\|^p \\
&\leq 2^p(\|\psi_0^{-1} \circ \psi_{n+1}(x_{n+1})\|^p + \|\sum_{i=1}^{\infty} \psi_i^{-1} \circ \psi_{i+n+1}(x_{i+n+1})\|^p) \\
&\leq 2^p(\lambda^p\|P_{n+1}x\|^p + \sum_{i=1}^{\infty} \lambda^p\|P_{i+n+1}x\|^p) \leq 2^p\lambda^p \sum_{i=1}^{\infty} \|P_i x\|^p \\
&= \|2\lambda \sum_{i=1}^{\infty} P_i x\|^p = 2^p\lambda^p\|(I - P_0)x\|^p \leq 2^p\lambda^p(\|P_0\| + 1)^p\|x\|^p.
\end{aligned}$$

Note that $\|L^m(x)\|_p^p \leq 2^p\lambda^p \sum_{i=0}^{\infty} P_{i+m}x\|^p \rightarrow 0$ shows (3) for $p \geq 1$. In the case $p = 0$ the computations are somewhat simpler and are shown below:

$$\begin{aligned}
\|R^{n+1}(x)\|_{\infty} &= \|\sum_{i=0}^{\infty} \psi_{i+n+1}^{-1} \circ \psi_i(x_i)\|_{\infty} = \max_{0 \leq i < \infty} \|\psi_{i+n+1}^{-1} \circ \psi_i(x_i)\| \\
&\leq \max_{0 \leq i < \infty} \|\psi_{i+n+1}^{-1} \circ \psi_i\| \|x_i\| \leq \lambda \max_{0 \leq i < \infty} \|x_i\| = \lambda \|x\|_{\infty} \\
\|L^{n+1}(x)\|_{\infty} &= \|\sum_{i=0}^{\infty} \psi_i^{-1} \circ \psi_{i+n+1}(x_{i+n+1})\|_{\infty} \\
&= \max_{0 \leq i < \infty} \|\psi_i^{-1} \circ \psi_{i+n+1}(x_{i+n+1})\| \\
&\leq \max_{0 \leq i < \infty} \|\psi_i^{-1} \circ \psi_{i+n+1}\| \|x_{i+n+1}\| \leq \lambda \max_{0 \leq i < \infty} \|x_{i+n+1}\| \\
&\leq \lambda \|x\|_{\infty}.
\end{aligned}$$

In this case $\|L^m(x)\|_0 \leq \lambda \max_{0 \leq i < \infty} \|x_{i+n+1}\| \rightarrow 0$ proves (3) for $p = 0$. □

Denote by D_S the inner derivation determined by S in $\mathcal{L}(\mathcal{X})$ i.e.

$$D_S T = ST - TS.$$

In the notation introduced above, an operator $T \in \mathcal{L}(\mathcal{X})$ is a commutator if and only if there exists $S \in \mathcal{L}(\mathcal{X})$ such that $T \in D_S \mathcal{L}(\mathcal{X})$. For a given decomposition \mathcal{D} of \mathcal{X} denote by $\mathcal{A}(\mathcal{D})$ the set

$$\mathcal{A}(\mathcal{D}) = \{T \in \mathcal{L}(\mathcal{X}) : \sum_{n=0}^{\infty} R^n T L^n \text{ is strongly convergent}\}. \quad (4)$$

For $T \in \mathcal{A}(\mathcal{D})$ we define

$$T_{\mathcal{D}} = \sum_{n=0}^{\infty} R^n T L^n.$$

Our next lemma shows that each operator $T \in \mathcal{A}(\mathcal{D})$ is a commutator and also gives an explicit expression for T as the commutator of two operators.

Lemma 2.2. *Let $T \in \mathcal{A}(\mathcal{D})$ for some decomposition $\mathcal{D} = \{X_i\}$ of \mathcal{X} . Then we have*

$$T = D_L(RT_{\mathcal{D}}) = -D_R(T_{\mathcal{D}}L). \quad (5)$$

Proof. We will show one of the equalities via direct computation. The proof of the other is similar.

$$\begin{aligned} D_L(RT_{\mathcal{D}}) &= LRT_{\mathcal{D}} - RT_{\mathcal{D}}L = T_{\mathcal{D}} - R\left(\sum_{n=0}^{\infty} R^n TL^n\right)L \\ &= T_{\mathcal{D}} - \sum_{n=1}^{\infty} R^n TL^n = T. \end{aligned}$$

In the computation above we used the convention $L^0 = R^0 = I$. □

Lemma 2.3. *For a decomposition $\mathcal{D} = \{X_i\}$ of \mathcal{X} we have the following relations*

$$\mathcal{A}(\mathcal{D}) = D_R(\mathcal{L}(\mathcal{X})RL) = D_L(RL\mathcal{L}(\mathcal{X})).$$

Proof. We will show the first of the relations. The proof of the second, as one may expect, is similar.

If $T \in \mathcal{A}(\mathcal{D})$, then $T_{\mathcal{D}}L = T_{\mathcal{D}}LRL = (T_{\mathcal{D}}L)RL \in \mathcal{L}(\mathcal{X})RL$. Then using $T = -D_R(T_{\mathcal{D}}L)$ from (5) we have $T \in D_R(\mathcal{L}(\mathcal{X})RL)$.

Now, to prove the other direction, assume that $T \in \mathcal{L}(\mathcal{X})RL$. Then $T = SRL$ for some operator S (hence $TR = SR$). Then

$$\begin{aligned} \sum_{n=0}^m R^n (D_R T) L^n &= \sum_{n=0}^m R^n (RT - TR) L^n = \sum_{n=0}^m R^{n+1} T L^n - \sum_{n=0}^m R^n T R L^n \\ &= \sum_{n=0}^m R^{n+1} S R L L^n - \sum_{n=0}^m R^n S R L R L^n \\ &= \sum_{n=0}^m R^{n+1} S R L^{n+1} - \sum_{n=0}^m R^n S R L^n \\ &= R^{m+1} S R L^{m+1} - S R = R^{m+1} T L^m - T R. \end{aligned}$$

Since $\lim_{m \rightarrow \infty} \|L^m(x)\| = 0$ for any $x \in \mathcal{X}$ from (3) and $\|R^m\| < 2\lambda C_1$ for every $m > 0$, we have

$\lim_{m \rightarrow \infty} \sum_{n=0}^m R^n (D_R T) L^n = -TR$. From the last equation we conclude that $D_R T \in \mathcal{A}(\mathcal{D})$ and $(D_R T)_{\mathcal{D}} = -TR$. Moreover, from $TR = SR$ we have $(D_R T)_{\mathcal{D}} = -SR$ and multiplying both sides by L we obtain $(D_R T)_{\mathcal{D}} L = -T$. □

We proved that for a given decomposition \mathcal{D} all operators $T \in \mathcal{A}(\mathcal{D})$ are commutators, but in general the condition in (4) is hard to check for a given operator T . We want to have a condition on T which is easy to check and which ensures the containment $T \in \mathcal{A}(\mathcal{D})$. To be more precise, given an operator T , we want to have a condition on T which will allow us to build a decomposition \mathcal{D} for which $T \in \mathcal{A}(\mathcal{D})$. Our next lemma gives us such a condition (as will become clear later) and it will be our main tool for constructing decompositions in the sequel.

Lemma 2.4. *Let $T \in \mathcal{L}(\mathcal{X})$ and $\mathcal{D} = \{X_i\}$ be a decomposition of \mathcal{X} . Fix $\varepsilon > 0$ and denote $\tilde{P}_n = \sum_{i=0}^n P_i$, where P_i is the projection onto X_i . Let us also assume that*

$$\lim_{n \rightarrow \infty} \|(I - \tilde{P}_n)T\| = \lim_{n \rightarrow \infty} \|T(I - \tilde{P}_n)\| = 0. \quad (6)$$

Then there exists an increasing sequence of integers $\{m_j\}_{j=0}^{\infty}$ such that

$$\sum_{j=0}^{\infty} \|(I - \tilde{P}_{m_j})T\| + \sum_{j=0}^{\infty} \|T(I - \tilde{P}_{m_j})\| + \sum_{i,j=0}^{\infty} \|(I - \tilde{P}_{m_i})T(I - \tilde{P}_{m_j})\| < \varepsilon.$$

Proof. Note first that $\|I - \tilde{P}_i\| \leq \|P_0\| + 1 = C_1$ for every $i \in \mathbb{N}$. Let $\{n_j\}_{j=0}^{\infty}$ be an increasing sequence of integers such that

$$\sum_{j=0}^{\infty} \|T(I - \tilde{P}_{n_j})\| < \frac{\varepsilon}{3C_1}, \quad \sum_{j=0}^{\infty} \|(I - \tilde{P}_{n_j})T\| < \frac{\varepsilon}{3C_1}.$$

Now we can use the inequality

$$\sum_{j=0}^{\infty} \|(I - \tilde{P}_i)T(I - \tilde{P}_{n_j})\| \leq \sum_{j=0}^m \|(I - \tilde{P}_i)T(I - \tilde{P}_{n_j})\| + C_1 \sum_{j=m+1}^{\infty} \|T(I - \tilde{P}_{n_j})\|$$

to deduce that

$$\lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} \|(I - \tilde{P}_i)T(I - \tilde{P}_{n_j})\| = 0.$$

Using the last equation we can find an increasing sequence of integers $\{m_j\}_{j=0}^{\infty}$, $m_j \geq n_j$ such that

$$\sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \|(I - \tilde{P}_{m_i})T(I - \tilde{P}_{n_j})\| < \frac{\varepsilon}{3C_1}.$$

Now it is easy to deduce that the sequence $\{m_j\}_{j=0}^{\infty}$ satisfies the condition of the lemma. In fact

$$\begin{aligned} \|T(I - \tilde{P}_{m_j})\| &= \|T(I - \tilde{P}_{n_j})(I - \tilde{P}_{m_j})\| \leq C_1 \|T(I - \tilde{P}_{n_j})\| \\ \|(I - \tilde{P}_{m_j})T\| &= \|(I - \tilde{P}_{m_j})(I - \tilde{P}_{n_j})T\| \leq C_1 \|(I - \tilde{P}_{n_j})T\| \\ \|(I - \tilde{P}_{m_i})T(I - \tilde{P}_{m_j})\| &= \|(I - \tilde{P}_{m_i})T(I - \tilde{P}_{n_j})(I - \tilde{P}_{m_j})\| \\ &\leq C_1 \|(I - \tilde{P}_{m_i})T(I - \tilde{P}_{n_j})\|. \end{aligned}$$

This finishes the proof. □

Lemma 2.5. *Let $\mathcal{D} = \{X_i\}$ be a decomposition of \mathcal{X} . Then for any $T \in \mathcal{L}(\mathcal{X})$ we have*

$$P_i T P_j \in \mathcal{A}(\mathcal{D}), \quad \|(P_i T P_j)_{\mathcal{D}}\| \leq C \|P_i T P_j\|$$

where C depends on \mathcal{D} only.

Proof. Let us consider the case $p \geq 1$ first. Note first that $\|L^n\| \leq 2\lambda C_1$ and $\|R^n\| \leq 2\lambda C_1$. For any $x \in \mathcal{X}$ we have (see Prop. (2.1))

$$\begin{aligned} \left\| \sum_{n=m}^{m+r} R^n P_i T P_j L^n x \right\|^p &= \left\| \sum_{n=m}^{m+r} R^n P_i T P_j L^n P_{j+n} x \right\|^p \\ &\leq 4\lambda^2 C_1^2 \|P_i T P_j\|^p \sum_{n=m}^{m+r} \|P_{j+n} x\|^p \end{aligned}$$

$$\begin{aligned}
&\leq 4\lambda^2 C_1^2 \|P_i T P_j\|^p \sum_{n=m}^{\infty} \|P_{j+n} x\|^p \\
&\leq 4\lambda^2 C_1^3 \|P_i T P_j\|^p \|x\|^p.
\end{aligned}$$

Since $\sum_{n=m}^{\infty} \|P_{j+n} x\|^p \rightarrow 0$ as $m \rightarrow \infty$ we have that $\sum_{n=0}^{\infty} R^n P_i T P_j L^n$ is strongly convergent and $P_i T P_j \in \mathcal{A}(\mathcal{D})$.

The inequality in the Theorem in this case follows from the inequality above with $C = 4\lambda^2 C_1^3$. For $p = 0$ a similar calculation shows

$$\begin{aligned}
\left\| \sum_{n=m}^{m+r} R^n P_i T P_j L^n x \right\|_{\infty} &= \left\| \sum_{n=m}^{m+r} R^n P_i T P_j L^n P_{j+n} x \right\|_{\infty} \\
&= \max_{m \leq n \leq m+r} \|R^n P_i T P_j L^n P_{j+n} x\| \\
&\leq 4\lambda^2 C_1^2 \|P_i T P_j\| \max_{m \leq n \leq m+r} \|P_{j+n} x\|.
\end{aligned}$$

Since $\max_{m \leq n \leq m+r} \|P_{j+n} x\| \rightarrow 0$ as $m \rightarrow \infty$ we apply the same argument as in the case $p \geq 1$ to obtain the conclusion of the theorem. \square

Corollary 2.6. *Let $T \in \mathcal{L}(\mathcal{X})$ and $\mathcal{D} = \{X_i\}$ be a decomposition of \mathcal{X} . Then we have*

$$\begin{aligned}
T P_0 &= D_R(L T P_0 - (P_0 T P_0)_{\mathcal{D}} L) \\
P_0 T &= D_L(-P_0 T R + R(P_0 T P_0)_{\mathcal{D}}).
\end{aligned}$$

Proof. We will prove the first equation. Note that from Lemma 2.2 we have $-D_R((P_0 T P_0)_{\mathcal{D}} L) = P_0 T P_0$ and $D_L(R(P_0 T P_0)_{\mathcal{D}}) = P_0 T P_0$. Now

$$\begin{aligned}
D_R(L T P_0 - (P_0 T P_0)_{\mathcal{D}} L) &= R L T P_0 - L T P_0 R + P_0 T P_0 \\
&= (I - P_0) T P_0 + P_0 T P_0 = T P_0 \\
D_L(-P_0 T R + R(P_0 T P_0)_{\mathcal{D}}) &= -L P_0 T R + P_0 T R L + P_0 T P_0 \\
&= P_0 T (I - P_0) + P_0 T P_0 = P_0 T.
\end{aligned}$$

Above we used the equality $P_0 R = 0 = L P_0$, which is clear from the definitions of R and L . \square

The following theorem shows the importance of the decompositions in determining whether an operator is a commutator.

Theorem 2.7. *Under the hypotheses of Lemma 2.4, there is a decomposition \mathcal{D} of \mathcal{X} for which*

$$T \in \mathcal{A}(\mathcal{D}), \quad \|T_{\mathcal{D}}\| \leq C \|T\| + \varepsilon$$

where C depends on \mathcal{D} only. In particular, using Lemma 2.2 we conclude that T is a commutator.

Proof. Using the sequence $\{m_j\}$ from Lemma 2.4, define a new decomposition where $\tilde{X}_0 = \bigoplus_{k=0}^{m_0} X_k$, $\tilde{X}_i =$

$\bigoplus_{k=m_{i-1}+1}^{m_i} X_k$. Note that the new decomposition also satisfies the condition about the projections. For simplicity of notation we will denote the new decomposition by $\{X_i\}$ and the projections onto X_i by P_i . In the new notation the conclusion from Lemma 2.4 can be written as

$$\sum_{j=0}^{\infty} \|(I - \tilde{P}_j)T\| + \sum_{j=0}^{\infty} \|T(I - \tilde{P}_j)\| + \sum_{i,j=0}^{\infty} \|(I - \tilde{P}_i)T(I - \tilde{P}_j)\| < \varepsilon.$$

Now using $P_i(I - \tilde{P}_{i-1}) = (I - \tilde{P}_{i-1})P_i = P_i$ we have

$$\begin{aligned} \sum_{i,j=0}^{\infty} \|P_i T P_j\| &\leq \|P_0 T P_0\| + C_1 \sum_{i=1}^{\infty} \|P_i T\| + C_1 \sum_{j=1}^{\infty} \|T P_j\| + \sum_{i,j=1}^{\infty} \|P_i T P_j\| \\ &\leq \|P_0 T P_0\| + C_1 \sum_{i=1}^{\infty} \|P_i (I - \tilde{P}_{i-1}) T\| \\ &\quad + C_1 \sum_{j=1}^{\infty} \|T (I - \tilde{P}_{j-1}) P_j\| \\ &\quad + \sum_{i,j=1}^{\infty} \|P_i (I - \tilde{P}_{i-1}) T (I - \tilde{P}_{j-1}) P_j\| \\ &\leq \|P_0 T P_0\| + C_1^2 \sum_{i=1}^{\infty} \|(I - \tilde{P}_{i-1}) T\| + C_1^2 \sum_{j=1}^{\infty} \|T (I - \tilde{P}_{j-1})\| \\ &\quad + C_1^2 \sum_{i,j=1}^{\infty} \|(I - \tilde{P}_{i-1}) T (I - \tilde{P}_{j-1})\| \\ &\leq \|P_0 T P_0\| + C_1^2 \varepsilon. \end{aligned}$$

Since the series $\sum_{i=0}^{\infty} P_i$ is strongly convergent to I , we have $T = \sum_{i,j=0}^{\infty} P_i T P_j$ in the norm topology of $\mathcal{L}(\mathcal{X})$. Using Lemma 2.5, the operator

$$S = \sum_{i,j=0}^{\infty} (P_i T P_j)_{\mathcal{D}}$$

is well defined and using Lemma 2.2 we have that $T = D_R(-SL) \in \mathcal{A}(\mathcal{D})$. Now $D_R(T_{\mathcal{D}}L - SL) = 0$ and by the proof of Lemma 2.3 we have

$$0 = -(D_R(T_{\mathcal{D}}L - SL))_{\mathcal{D}}L = (T_{\mathcal{D}} - S)L.$$

From the equation above we conclude that $T_{\mathcal{D}} = S$ and $\|T_{\mathcal{D}}\| \leq C\|T\| + \varepsilon$. \square

3 Compactness and commutators on ℓ_p and L_p ($1 \leq p < \infty$)

In order to prove the conjecture about the structure of the commutators on a given space we have to show that all the elements in the unique maximal ideal are commutators. We prove a lemma that takes care of

this in the case $\mathcal{X} = \ell_1$ and also shows that the ideal of compact operators consists of commutators only, provided the space \mathcal{X} has some additional structure. Before that we will show a lemma about the operators T on \mathcal{X} which do not preserve a copy of \mathcal{X} in the cases of $\mathcal{X} = \ell_1$ and $\mathcal{X} = L_1$, which we will use and is interesting on its own.

Lemma 3.1. *Let $\mathcal{X} = L_1$ or $\mathcal{X} = \ell_1$ and suppose that $T \in \mathcal{L}(\mathcal{X})$ does not preserve a copy of \mathcal{X} . Then, for every $\delta > 0$ and for every $\tilde{X} \subset \mathcal{X}$, $\tilde{X} \equiv \mathcal{X}$, there exists $Y \subset \tilde{X}$, such that Y is $(1 + \delta)$ isomorphic to \mathcal{X} , $(1 + \delta)$ complemented in \mathcal{X} , and $\|T|_Y\| < \delta$.*

Proof. Consider the case $\mathcal{X} = L_1$ first. By assumption T does not preserve a copy of L_1 which implies that T is not an E -operator (actually this can be taken as an equivalent definition for an operator not to be an E -operator [6, Theorem 4.1]) and hence it is not sign-preserving either ([9, Theorem 1.5]). Now [9, Lemma 3.1] gives us a subspace $Z \subset \tilde{X}$ such that $Z \simeq \tilde{X}$ and $\|T|_Z\| < \delta$. Using [9, Theorem 1.1] we find $Y \subset Z$, which is $(1 + \delta)$ isomorphic to $\tilde{X} \equiv L_1$, $(1 + \delta)$ complemented in \tilde{X} and Y clearly satisfies $\|T|_Y\| < \delta$. If Q is the norm one projection onto \tilde{X} , and $R : \tilde{X} \rightarrow Y$ is a projection of norm less than $1 + \delta$, then $P := RQ$ is a projection from L_1 onto Y and $\|P\| < 1 + \delta$.

For the case $\mathcal{X} = \ell_1$ we use the fact that if \tilde{X} is isometric to ℓ_1 , then $\tilde{X} = \overline{\text{span}}\{\psi_i : i = 0, 1, \dots\}$ for some vectors $\{\psi_i\}_{i=1}^\infty$ of norm one, such that

$$\psi_j = \sum_{i \in \sigma_j} \lambda_i e_i, \quad \text{with } \sigma_j \cap \sigma_k = \emptyset \text{ for } j \neq k$$

where $\{e_i\}_{i=1}^\infty$ is the standard unit vector basis of ℓ_1 . This follows trivially from the observation that Ue_i and Ue_j must have disjoint supports if $U : \tilde{X} \rightarrow \ell_1$ is an into isometry (cf. [13, Proposition 2.f.14]). Note also that since every infinite dimensional subspace of ℓ_1 contains an isomorphic copy of ℓ_1 ([13, Proposition 2.a.2]), then the operator T is automatically strictly singular and hence compact ([14]). Then, $\{T\psi_i\}_{i=0}^\infty$ is relatively compact in ℓ_1 and hence there exist $y \in \ell_1$ and a subsequence $\{\psi_{i_j}\}$ such that $T\psi_{i_j} \rightarrow y$. WLOG we may assume that $T\psi_i \rightarrow y$. Finally, define $\varphi_i = \frac{\psi_{2i} - \psi_{2i+1}}{2}$ for $i = 0, 1, \dots$. Clearly $\{\varphi_i\}_{i=0}^\infty$ is a normalized block basis of \tilde{X} such that $\|T\varphi_i\|_1 \rightarrow 0$. Assume WLOG that $\|T\varphi_i\|_1 < \varepsilon$ (this can be easily achieved by passing to a subsequence). Then for $Y = \overline{\text{span}}\{\varphi_i : i = 0, 1, \dots\}$ we have $\|T|_Y\| < \varepsilon$. Note also that $Y \subset \tilde{X}$ is 1-complemented in \tilde{X} as it is the closed span of a normalized block basis and clearly is isometric to \tilde{X} ([7, Lemma 1]). Finally, let $R : \tilde{X} \rightarrow Y$ be the norm one projection onto Y and $Q : \ell_1 \rightarrow \tilde{X}$ be the norm one projection onto \tilde{X} . Then clearly $P := RQ$ is a norm one projection onto Y . \square

Lemma 3.2. *Let \mathcal{X} be a Banach space for which $\mathcal{X} \simeq \left(\bigoplus_{i=0}^\infty \mathcal{X}\right)_p$ for some $1 \leq p < \infty$ or $p = 0$. In the case $p = 1$ we will assume that $\mathcal{X} = L_1$ or $\mathcal{X} = \ell_1$. Let $T \in \mathcal{L}(\mathcal{X})$ be a compact operator and $\varepsilon > 0$. Then there exists a decomposition \mathcal{D} of \mathcal{X} such that $T \in \mathcal{A}(\mathcal{D})$ and $\|T|_{\mathcal{D}}\| \leq C\|T\| + \varepsilon$ for some constant C depending on \mathcal{D} only. Consequently T is a commutator and $T = -D_R(T|_{\mathcal{D}}L)$.*

Proof. The result is known in the case of $\mathcal{X} = L_p$ and $\mathcal{X} = \ell_p$ for $1 < p < \infty$ (cf. [10] and [2]), and for $\mathcal{X} = c_0$ and $\mathcal{X} = C(K)$ ([3]). The proof presented here in these cases follows Apostol's ideas from [2] and our generalized context gives a shorter proof in the case of L_p for $1 < p < \infty$. Partial results were known in the case $\mathcal{X} = \ell_1$ ([2, Theorem 2.6]).

Case I. $p > 1$ or $p = 0$. In this case we proceed as in Theorem 2.4 in [2]. Consider an arbitrary decomposition \mathcal{D} of \mathcal{X} and denote $\tilde{P}_n = \sum_{i=0}^n P_i$. Now we have

$$\lim_{n \rightarrow \infty} \|(I - \tilde{P}_n)T\| = \lim_{n \rightarrow \infty} \|T(I - \tilde{P}_n)\| = 0.$$

In fact if we choose $\varphi_i, \psi_i \in \mathcal{X}$ such that

$$\|(I - \tilde{P}_n)T\varphi_n\|_{\mathcal{X}} > \|(I - \tilde{P}_n)T\| - \frac{1}{n+1}, \quad \|\varphi_n\| = 1$$

$$\|T(I - \tilde{P}_n)\psi_n\|_{\mathcal{X}} > \|T(I - \tilde{P}_n)\| - \frac{1}{n+1}, \quad \|\psi_n\| = 1, \quad (I - \tilde{P}_n)\psi_n = \psi_n.$$

Since the set $\{T\varphi_i\}_{i=0}^{\infty}$ is relatively compact in \mathcal{X} and the sequence $\{(I - \tilde{P}_i)\}_{i=0}^{\infty}$ converges strongly to 0 we have $\lim_{n \rightarrow \infty} \|(I - \tilde{P}_n)T\| = 0$. On the other hand, the sequence $\{\psi_i\}_{i=0}^{\infty}$ is weakly convergent to 0. Now using the fact that T is compact, it follows that the sequence $\{T\psi_i\}_{i=0}^{\infty}$ converges to 0 in norm and hence $\lim_{n \rightarrow \infty} \|T(I - \tilde{P}_n)\| = 0$. Now Theorem 2.7 gives the result.

Case II. $p = 1$. Fix $\varepsilon > 0$ and let $\mathcal{D} = \{X_i\}$ be the fixed decomposition of \mathcal{X} defined by $X_i = L_1[\frac{1}{2^{i+1}}, \frac{1}{2^i})$ in the case of $\mathcal{X} = L_1$ and by $X_i = P_{N_i}\ell_1$ (where $\mathbb{N} = \cup_{i=0}^{\infty} N_i$ such that $\text{card } N_i = \text{card } \mathbb{N}$ for all $i \in \mathbb{N}$ and $N_j \cap N_i = \emptyset$ for $i \neq j$) in the case of $\mathcal{X} = \ell_1$. Using Lemma 3.1 for each X_i with $\delta = \frac{\varepsilon}{2^i}$ will give us

$1 + \varepsilon$ complemented subspaces $\{Y_i\}$ of \mathcal{X} which are isomorphic to \mathcal{X} and $\|T|_{Y_i}\| < \frac{\varepsilon}{2^i}$. Set $Y_0 = (I - \sum_{i=1}^{\infty} P_i)\mathcal{X}$.

Note that $\mathcal{D} = \{Y_i\}$ is a decomposition for \mathcal{X} since all the spaces are complemented and isomorphic to \mathcal{X} . This is clear for Y_i for $i = 1, 2, \dots$ and it also holds for Y_0 , since $X_0 \subset Y_0$ is complemented in \mathcal{X} , isomorphic to \mathcal{X} , and using [6, Corollary 5.3] in the case $\mathcal{X} = L_1$, and [7, Proposition 4] in the case $\mathcal{X} = \ell_1$, it follows

that Y_0 is isomorphic to \mathcal{X} as well. Now, if $\tilde{P}_n = \sum_{i=0}^n P_i$, then clearly we have $\lim_{n \rightarrow \infty} \|T(I - \tilde{P}_n)\| = 0$. Since

T is compact operator, then we have $\lim_{n \rightarrow \infty} \|(I - \tilde{P}_n)T\| = 0$ as well (the argument provided in **Case I** above works in this case as well), so using Theorem 2.7 we conclude that T is a commutator. \square

Remark 3.3. *Using the previous lemma we immediately conclude that [10, Theorem 4.3] holds for $p = 1$. Namely, a multiplication operators M_ϕ on L_1 is a commutator if and only if the spectrum of M_ϕ contains more than one limit point or contains zero as the unique limit point.*

Corollary 3.4. *Let \mathcal{X} be a Banach space for which $\mathcal{X} \simeq \left(\bigoplus_{i=0}^{\infty} \mathcal{X}\right)_p$ for some $1 \leq p < \infty$ or $p = 0$. In the case $p = 1$ we will assume that $\mathcal{X} = L_1$ or $\mathcal{X} = \ell_1$. Let $T \in \mathcal{L}(\mathcal{X})$ and suppose that P is a projection on \mathcal{X} such that $P\mathcal{X} \simeq \mathcal{X} \simeq (I - P)\mathcal{X}$ and that either TP or PT is a compact operator. Then T is a commutator.*

Proof. First we treat the case when TP is compact operator. Let $\mathcal{D} = \{X_i\}_{i=0}^{\infty}$ be a decomposition for which $TP \in \mathcal{A}(\mathcal{D})$ and $\|(TP)_{\mathcal{D}}\|_{\mathcal{X}} \leq C\|TP\|_{\mathcal{X}} + \frac{\varepsilon}{2}$ for a fixed $\varepsilon > 0$ (by Lemma 3.2). We also want \mathcal{D} to be such that $(I - P)\mathcal{X} = X_0$ hence we may assume $(I - P) = P_0$, where P_0 is the projection onto X_0 . This can obviously be done for $1 < p < \infty$ (since the decomposition used in the proof was arbitrary). In the case of L_1 we consider the operator $\tilde{T} = GTG^{-1}$ where $G : P\mathcal{X} \oplus (I - P)\mathcal{X} \rightarrow (I - P_0)\mathcal{X} \oplus X_0$ is an isomorphism such that $GP\mathcal{X} = (I - P_0)\mathcal{X}$, $G(I - P)\mathcal{X} = X_0$. In this case $\tilde{T}GPG^{-1}$ is compact and clearly we can choose the decomposition as in Lemma 3.2 and apply the same argument. Now WLOG we will assume that $\tilde{T} = T$. In the case of ℓ_1 we can make a similarity as in the previous case and reduce to the case where TP_M is a compact operator for some $M \subset \mathbb{N}$. Define

$$S = LT(I - P) - (P_0T(I - P)P_0)_{\mathcal{D}}L - (TP)_{\mathcal{D}}L.$$

Use equation (5) applied to TP and $P_0T(I - P)P_0$ to get

$$-D_R((TP)_{\mathcal{D}}L) = TP \tag{7}$$

$$-D_R((P_0T(I - P)P_0)_{\mathcal{D}}L) = P_0T(I - P)P_0 = P_0T(I - P). \tag{8}$$

Now

$$D_R(LT(I - P)) = RLT(I - P) - LT(I - P)R = (I - P_0)T(I - P) \quad (9)$$

since $(I - P)R = 0$. Combining (7), (8) and (9) we conclude that $D_R S = T$. If PT is compact we consider $S = -(I - P)TR + R(P_0(I - P)TP_0)_{\mathcal{D}} + R(PT)_{\mathcal{D}}$ and a similar calculation shows that $T = D_L(S)$. \square

4 Commutators on ℓ_1

We already saw in the previous section that the compact operators on ℓ_1 are commutators and in order to prove the conjecture in the case of $\mathcal{X} = \ell_1$ we have to show that all operators not of the form $\lambda I + K$, where K is compact and $\lambda \neq 0$, are commutators. To do that we are going to show that if T is not of the form $\lambda I + K$, then there exist disjoint complemented subspaces X and Y of \mathcal{X} which are isomorphic to \mathcal{X} and such that $T|_X : X \rightarrow Y$ is an onto isomorphism. As we will see, this last property of T will be enough to

show that T is a commutator on any space \mathcal{X} for which $\mathcal{X} \simeq \left(\bigoplus_{i=0}^{\infty} \mathcal{X} \right)_p$.

Definition 4.1. *The left essential spectrum of $T \in \mathcal{L}(\mathcal{X})$ is the set ([4] Def 1.1)*

$$\sigma_{l.e.}(T) = \{ \lambda \in \Lambda : \inf_{\substack{x \in Y \\ \|x\| = 1}} \|(\lambda - T)x\| = 0, \text{codim}(Y) < \infty, Y - \text{closed} \}.$$

For any $T \in \mathcal{L}(\mathcal{X})$, $\sigma_{l.e.}(T)$ is a closed non-void set ([4, Theorem 1.4]). The following lemma is an analog of Lemma 4.1 from [2] and the proof follows the steps in the proof there.

Lemma 4.2. *Let $\mathcal{X} = \ell_1$ and let $T \in \mathcal{L}(\mathcal{X})$, $0 \in \sigma_{l.e.}(T)$. If T is not compact, then it is similar to an operator $T' \in \mathcal{L}(\mathcal{X})$ for which there exists a projection P_M such that $M \subset \mathbb{N}$, $\text{card } M = \text{card } (\mathbb{N} - M)$, and $P_{\mathbb{N}-M}T'P_M$ is not a compact operator.*

Proof. For simplicity of notation we will denote P_M simply by P . By Lemma 3.2 in [2] and using a similarity we can obtain a subset M of \mathbb{N} so that $\sum_{n \in \mathbb{N}-M} \|TP_{\{n\}}\| < \infty$, where $P_{\{n\}}$ is the projection onto the n -th coordinate. From this inequality we have that $T(I - P)$ is compact. If $(I - P)TP$ is not compact we are done, thus we may suppose that $(I - P)TP$ is compact. The equality

$$T = T(I - P) + (I - P)TP + PTP$$

gives us that PTP is not compact. Using $\mathcal{X} \equiv P\mathcal{X} \oplus_1 (I - P)\mathcal{X}$, let $\varphi : P\mathcal{X} \rightarrow (I - P)\mathcal{X}$ be an isometry and define the operators V and V' in the following way

$$V(x) = \varphi(Px), \quad V'x = \varphi^{-1}((I - P)x).$$

It is easy to see that $PV = V(I - P) = V'P = (I - P)V' = V^2 = (V')^2 = 0$ and $VV' + V'V = I$. Define

$$\sqrt{2}S = P - (I - P) + V + V'.$$

Now a simple check gives us

$$\begin{aligned}
2S^2 &= (P - (I - P) + V + V')(P - (I - P) + V + V') \\
&= P + PV' + (I - P) - (I - P)V + VP + VV' - V'(I - P) + V'V \\
&= 2 + PV' - (I - P)V + VP - V'(I - P) \\
&= 2 + PV' - V + VP - V' = 2,
\end{aligned}$$

hence $S = S^{-1}$. Now consider the operator $2(I - P)S^{-1}TSP$. Again a simple calculation shows that

$$\begin{aligned}
2(I - P)S^{-1}TSP &= (-(I - P) + (I - P)V)T(P + VP) \\
&= (-(I - P) + V)T(P + VP) \\
&= -(I - P)TP - (I - P)TVP + VTP + VTVP = \\
&= -(I - P)TP - (I - P)T(I - P)VP \\
&\quad + VPTP + VT(I - P)VP = VPTP + K
\end{aligned}$$

where $K = -(I - P)TP - (I - P)T(I - P)VP + VT(I - P)VP$ is a compact operator because $T(I - P)$ and $(I - P)TP$ are compact operators (the first one by construction, the second by assumption). Since $V|_{P\mathcal{X}}$ is an isometry, we conclude that $VPTP$ is not compact and hence $(I - P)S^{-1}TSP$ is not compact either. Taking $T' = S^{-1}TS$ we finish the proof. \square

Proposition 4.3. *Let $\mathcal{X} = \ell_1$ and $T \in \mathcal{L}(\mathcal{X})$ be such that there exists a projection P such that $P\mathcal{X} \simeq \mathcal{X}$, $(I - P)\mathcal{X} \simeq \mathcal{X}$ and the operator $(I - P)TP$ is not compact. Then there exists a complemented subspace $Y \subset P\mathcal{X}$ such that $Y \simeq \mathcal{X}$, $(I - P)TP|_Y$ is an isomorphism into and $(I - P)TP(Y)$ is complemented in \mathcal{X} .*

Proof. Clearly $(I - P)TP$ is not a strictly singular operator since in $\mathcal{L}(\mathcal{X})$ the ideal of compact operators and strictly singular operators coincide ([14]). In particular, this implies that there exists an infinite dimensional subspace $Z \subset P\mathcal{X}$ such that $(I - P)TP$ is an isomorphism on Z . Consider the infinite dimensional subspace $(I - P)TPZ$. Using [7, Lemma 2] (cf. also [13, Proposition 2.a.2]) we conclude that there exists $U \subset (I - P)TPZ$ which is complemented in \mathcal{X} and isomorphic to \mathcal{X} . Clearly $(I - P)TP$ is an isomorphism on $((I - P)TP)^{-1}U$ and since U is complemented in \mathcal{X} , we also have that $((I - P)TP)^{-1}U$ is complemented in \mathcal{X} as well. \square

Theorem 4.4. *Let $\mathcal{X} \simeq (\bigoplus_{i=0}^{\infty} \mathcal{X})_p$, $1 \leq p < \infty$ or $p = 0$, and let $T \in \mathcal{L}(\mathcal{X})$ be an operator for which there exists a projection P such that $P\mathcal{X} \simeq \mathcal{X}$, $(I - P)\mathcal{X} \simeq \mathcal{X}$ and there exists a complemented subspace $Y \subseteq P\mathcal{X}$ such that $Y \simeq \mathcal{X}$, $(I - P)TP|_Y$ is an isomorphism into and $X = (I - P)TP(Y)$ is also complemented in \mathcal{X} . Then there exists a decomposition \mathcal{D} such that T is similar to a matrix operator*

$$\begin{pmatrix} * & L \\ * & * \end{pmatrix}$$

on $\mathcal{X} \oplus \mathcal{X}$, where L is the left shift associated with \mathcal{D} .

Proof. Let P_X be a projection onto X . Note that $X = P_X\mathcal{X} \subset (I - P)\mathcal{X}$ and hence $Y \subset P\mathcal{X} \subset (I - P_X)\mathcal{X}$. The previous observation shows that the operator $P_X T(I - P_X)|_Y$ is an isomorphism from Y onto X . Also, $X = P_X\mathcal{X} \simeq \mathcal{X}$ by the assumption of the theorem, and $(I - P_X)\mathcal{X}$ contains Y - a complemented copy of \mathcal{X} and hence $(I - P_X)\mathcal{X} \simeq \mathcal{X}$ (using a result of Pelczynski [7, Proposition 4]). The observations we made imply that WLOG we may assume $P_X = I - P$. Consider two decompositions $\mathcal{D}_1 = \{X_i\}$, $\mathcal{D}_2 = \{Y_i\}$ of \mathcal{X} such that $X = Y_0 = X_1 \oplus X_2 \oplus \dots$,

$X_0 = Y_1 \oplus Y_2 \oplus \dots$ and $Y_1 = Y$. Define a map S

$$S\varphi = L_{\mathcal{D}_1}\varphi \oplus L_{\mathcal{D}_2}\varphi, \quad \varphi \in \mathcal{X}$$

from \mathcal{X} to $\mathcal{X} \oplus \mathcal{X}$. The map S is invertible ($S^{-1}(a, b) = R_{\mathcal{D}_1}a + R_{\mathcal{D}_2}b$). Just using the definition of S and the formula for S^{-1} it is easy to see that

$$\begin{aligned} STS^{-1}(a, b) &= ST(R_{\mathcal{D}_1}a + R_{\mathcal{D}_2}b) = S(TR_{\mathcal{D}_1}a + TR_{\mathcal{D}_2}b) \\ &= (L_{\mathcal{D}_1}TR_{\mathcal{D}_1}a + L_{\mathcal{D}_1}TR_{\mathcal{D}_2}b) \oplus (L_{\mathcal{D}_2}TR_{\mathcal{D}_1}a + L_{\mathcal{D}_2}TR_{\mathcal{D}_2}b), \end{aligned}$$

hence

$$STS^{-1} = \begin{pmatrix} * & L_{\mathcal{D}_1}TR_{\mathcal{D}_2} \\ * & * \end{pmatrix}.$$

Let

$$A = P_{Y_0}TR_{\mathcal{D}_2} = (I - P)TR_{\mathcal{D}_2} \quad (10)$$

and note that $A|_{P_{Y_0}\mathcal{X}} \equiv A|_{(I-P)\mathcal{X}} : (I-P)\mathcal{X} \rightarrow (I-P)\mathcal{X}$ is onto and invertible since $R_{\mathcal{D}_2}$ is an isomorphism on $P_{Y_0}\mathcal{X}$ and $R_{\mathcal{D}_2}(P_{Y_0}\mathcal{X}) = Y_1 = Y$. Here we used the fact that $P_{Y_0}T$ is an isomorphism on Y ($PY = Y$). Denote by T_0 the inverse of $A|_{P_{Y_0}\mathcal{X}}$ (note that T_0 is an automorphism on $(I-P)\mathcal{X}$) and consider

$$G = I + T_0(I - P) - T_0A.$$

We will show that $G^{-1} = A + P$. In fact, from the definitions of A and T_0 it is clear that

$$AT_0(I - P) = T_0A(I - P) = I - P, PT_0 = PA = 0, (I - P)A = A \quad (11)$$

and since A maps onto $(I - P)\mathcal{X}$ and $AT_0|_{(I-P)\mathcal{X}} = I|_{(I-P)\mathcal{X}}$ we also have

$$A - AT_0A = 0. \quad (12)$$

Now using (11) and (12) it is easy to see that

$$\begin{aligned} (A + P)G &= (A + P)(I + T_0(I - P) - T_0A) \\ &= A + AT_0(I - P) - AT_0A + P = I - P + P = I \\ G(A + P) &= (I + T_0(I - P) - T_0A)(A + P) \\ &= A + P + T_0(I - P)A + T_0(I - P)P - T_0AA - T_0AP \\ &= A + P + T_0A - T_0AA - T_0AP \\ &= P + (I - T_0A)A + T_0A(I - P) \\ &= P + (I - T_0A)(I - P)A + (I - P) \\ &= I + ((I - P) - T_0A(I - P))A \\ &= I + (I - P - (I - P))A = I. \end{aligned}$$

Using a similarity we obtain

$$\begin{pmatrix} I & 0 \\ 0 & G^{-1} \end{pmatrix} \begin{pmatrix} * & L_{\mathcal{D}_1}TR_{\mathcal{D}_2} \\ * & * \end{pmatrix} \begin{pmatrix} I & 0 \\ 0 & G \end{pmatrix} = \begin{pmatrix} * & L_{\mathcal{D}_1}TR_{\mathcal{D}_2}G \\ * & * \end{pmatrix}.$$

It is clear that we will be done if we show that $L_{\mathcal{D}_1} = L_{\mathcal{D}_1}TR_{\mathcal{D}_2}G$. In order to do this consider the equation $(A + P)G = I \Leftrightarrow AG + PG = I$. Multiplying both sides of the last equation on the left by $L_{\mathcal{D}_1}$ gives us $L_{\mathcal{D}_1}AG + L_{\mathcal{D}_1}PG = L_{\mathcal{D}_1}$. Using $L_{\mathcal{D}_1}P \equiv L_{\mathcal{D}_1}P_{X_0} = 0$ we obtain $L_{\mathcal{D}_1}AG = L_{\mathcal{D}_1}$. Finally, substituting A from (10) in the last equation yields

$$L_{\mathcal{D}_1} = L_{\mathcal{D}_1}AG = L_{\mathcal{D}_1}P_{Y_0}TR_{\mathcal{D}_2}G = L_{\mathcal{D}_1}(I - P_{X_0})TR_{\mathcal{D}_2}G = L_{\mathcal{D}_1}TR_{\mathcal{D}_2}G$$

which finishes the proof. \square

The following theorem was proved in [2] for $\mathcal{X} = \ell_p$, but inessential modifications give the result in the general case.

Theorem 4.5. *Let \mathcal{D} be a decomposition of \mathcal{X} and let L be the left shift associated with it. Then the matrix operator*

$$\begin{pmatrix} T_1 & L \\ T_2 & T_3 \end{pmatrix}$$

acting on $\mathcal{X} \oplus \mathcal{X}$ is a commutator.

Proof. Let $\mathcal{D} = \{X_i\}$ be the given decomposition. Consider a decomposition $\mathcal{D}_1 = \{Y_i\}$ such that $Y_0 = \bigoplus_{i=1}^{\infty} X_i$ and $X_0 = \bigoplus_{i=1}^{\infty} Y_i$. Now there exists an operator G such that $D_{L_{\mathcal{D}}}G = R_{\mathcal{D}_1}L_{\mathcal{D}_1}(T_1 + T_3)$. This can be done using Corollary 2.6, since $R_{\mathcal{D}_1}L_{\mathcal{D}_1} = I - P_{Y_0} = P_{X_0}$. By making the similarity

$$\tilde{T} := \begin{pmatrix} I & 0 \\ G & I \end{pmatrix} \begin{pmatrix} T_1 & L \\ T_2 & T_3 \end{pmatrix} \begin{pmatrix} I & 0 \\ -G & I \end{pmatrix} = \begin{pmatrix} T_1 - LG & L \\ * & T_3 + GL \end{pmatrix}$$

we have $T_1 + T_3 - LG + GL = T_1 + T_3 - D_LG = T_1 + T_3 - R_{\mathcal{D}_1}L_{\mathcal{D}_1}(T_1 + T_3) = P_{Y_0}(T_1 + T_3)$. Using again Corollary 2.6 we deduce that $T_1 + T_3 - LG + GL$ is a commutator. Thus by replacing T by \tilde{T} we can assume that $T_1 + T_3$ is a commutator, say $T_1 + T_3 = AB - BA$ and $\|A\| < \frac{1}{2}$ (this can be done by scaling). Denote by M_T left multiplication by the operator T . Then $\|M_R D_A\| < 1$ where R is the right shift associated with \mathcal{D} . The operator $T_0 = (M_I - M_R D_A)^{-1} M_R (T_3 B - T_2)$ is well defined and it is easy to see that

$$\begin{pmatrix} A & 0 \\ T_3 & A - L \end{pmatrix} \begin{pmatrix} B & I \\ T_0 & 0 \end{pmatrix} - \begin{pmatrix} B & I \\ T_0 & 0 \end{pmatrix} \begin{pmatrix} A & 0 \\ T_3 & A - L \end{pmatrix} = \begin{pmatrix} T_1 & L \\ T_2 & T_3 \end{pmatrix}.$$

This finishes the proof. □

Theorem 4.6. *Let $\mathcal{X} = \ell_1$. An operator $T \in \mathcal{L}(\mathcal{X})$ is a commutator if and only if $T - \lambda$ is not compact for any $\lambda \neq 0$.*

Proof. Note first that if T is a commutator, from the remarks we made in the introduction it follows that $T - \lambda$ cannot be compact for any $\lambda \neq 0$. For proving the other direction we have to consider two cases:

Case I. If T is compact operator ($\lambda = 0$), the statement of the theorem follows from Lemma 3.2.

Case II. If $T - \lambda$ is not compact for any λ , then we consider $\sigma_{l.e.}(T)$. Since $\sigma_{l.e.}(T)$ is a non-empty set, there exists $\lambda \in \sigma_{l.e.}(T)$ such that $T - \lambda$ is not compact and we are in a position to apply Lemma 4.2 for the operator $T - \lambda$. Note that the conclusion of Lemma 4.2 for $T - \lambda$ implies that the same claim is true for T as well. Now we are in position to apply Theorem 4.4 (which we can because of Proposition 4.3) and obtain that T is similar to an operator of the form $\begin{pmatrix} * & L \\ * & * \end{pmatrix}$. Finally, we apply Theorem 4.5 to complete the proof. □

5 Commutators on $\ell_{p_1} \oplus \ell_{p_2} \oplus \dots \oplus \ell_{p_n}$ and ℓ_{∞}

Lemma 5.1. *Let X and Y be Banach spaces and $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ an operator from $X \oplus Y$ into $X \oplus Y$. If A and D are commutators on the corresponding spaces then T is a commutator on $X \oplus Y$.*

Proof. Let $A = [A_1, A_2]$ and $D = [D_1, D_2]$. Assume without loss of generality that $\max(\|A_2\|, \|D_2\|) < \frac{1}{4}$. We need to find operators E_1 and E_2 such that

$$T = \begin{pmatrix} A_1 & E_1 \\ E_2 & D_1 \end{pmatrix} \begin{pmatrix} A_2 + I & 0 \\ 0 & D_2 \end{pmatrix} - \begin{pmatrix} A_2 + I & 0 \\ 0 & D_2 \end{pmatrix} \begin{pmatrix} A_1 & E_1 \\ E_2 & D_1 \end{pmatrix},$$

or equivalently, we have to solve the equations

$$B = E_1 D_2 - (A_2 + I) E_1 \quad (13)$$

$$C = E_2 (A_2 + I) - D_2 E_2 \quad (14)$$

for E_1 and E_2 . Let $G : L(X, Y) \rightarrow L(X, Y)$ be defined by $G(S) = -SA_2 + D_2S$. Clearly $\|G\| < 1$ by our choice of A_2 and D_2 and hence $I - G$ is invertible. Now it is enough to observe that (14) is equivalent to $C = (I - G)(E_2)$ which will give us $E_2 = (I - G)^{-1}C$. Analogously we define $F : L(Y, X) \rightarrow L(Y, X)$ by $F(S) = -A_2S + SD_2$ and then (13) will be equivalent to $-B = (I - F)(E_1)$. Applying the same argument as above we get that $I - F$ is invertible and hence $E_1 = (I - F)^{-1}(-B)$. \square

Theorem 5.2. *Let $\mathcal{X} = \ell_p \oplus \ell_q$ where $1 \leq q < p < \infty$ and $T \in \mathcal{L}(\mathcal{X})$. Let P_{ℓ_p} and P_{ℓ_q} be the natural projections from \mathcal{X} onto ℓ_p and ℓ_q respectively. Then T is a commutator if and only if $P_{\ell_p}TP_{\ell_p}$ and $P_{\ell_q}TP_{\ell_q}$ are commutators as operators acting on ℓ_p and ℓ_q respectively.*

Proof. Throughout the proof we will work with the matrix representation of T as an operator acting on \mathcal{X} . Let $T = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ where $A : \ell_p \rightarrow \ell_p, D : \ell_q \rightarrow \ell_q, B : \ell_q \rightarrow \ell_p, C : \ell_p \rightarrow \ell_q$. The well known fact that the operator C is compact ([13, Proposition 2.c.3]) will play an important role in the proof. If T is a commutator, then $T = [T_1, T_2]$ for some $T_1, T_2 \in \mathcal{L}(\mathcal{X})$. Write $T_i = \begin{pmatrix} A_i & B_i \\ C_i & D_i \end{pmatrix}$ for $i = 1, 2$. A simple computation shows that

$$T = \begin{pmatrix} [A_1, A_2] + B_1C_2 - B_2C_1 & A_1B_2 + B_1D_2 - A_2B_1 - B_2D_1 \\ C_1A_2 + D_1C_2 - C_2A_1 - D_2C_1 & [D_1, D_2] + C_1B_2 - C_2B_1 \end{pmatrix}.$$

From the classification of the commutators on ℓ_p for $1 \leq p < \infty$ and the fact that the C_i 's are compact we immediately deduce that the diagonal entries in the last representation of T are commutators. For the other direction we apply Lemma 5.1 which concludes the proof. \square

The classification given in the theorem can be immediately generalized to a space which is finite sum of ℓ_p spaces, namely, we have the following

Corollary 5.3. *Let $\mathcal{X} = \ell_{p_1} \oplus \ell_{p_2} \oplus \dots \oplus \ell_{p_n}$ where $1 \leq p_n < p_{n-1} < \dots < p_1 < \infty$ and $T \in \mathcal{L}(\mathcal{X})$. Let $P_{\ell_{p_i}}$ be the natural projections from \mathcal{X} onto ℓ_{p_i} for $i = 1, 2, \dots, n$. Then T is a commutator if and only if for each $1 \leq i \leq n$, $P_{\ell_{p_i}}TP_{\ell_{p_i}}$ is a commutator as an operator acting on ℓ_{p_i} .*

Proof. We will proceed by induction on n and clearly Theorem 5.2 gives us the result for $n = 2$. If the statement is true for some n , then to show it for $n + 1$, denote $Y = \ell_{p_2} \oplus \ell_{p_3} \oplus \dots \oplus \ell_{p_n}$. Now $\mathcal{X} = \ell_{p_1} \oplus Y$ and using the same argument as in Theorem 5.2 we can see that if T is a commutator, then both $P_{\ell_{p_1}}TP_{\ell_{p_1}}$ and $P_{\ell_Y}TP_{\ell_Y}$ are commutators on ℓ_{p_1} and Y respectively. Here we use the induction step to show that compact perturbation of a commutator on Y is still a commutator. The other direction is exactly as in Theorem 5.2. It is worthwhile noticing that for this direction we do not need any assumption on the spaces in the sum. \square

Our last result shows that every strictly singular operator in $L(\ell_\infty)$ is a commutator. Clearly this is an essential step in proving the conjecture about the classification of the commutators on ℓ_∞ , namely, that an operator $T \in L(\ell_\infty)$ is not a commutator if and only if $T = \lambda I + S$ for some strictly singular operator S and some $\lambda \neq 0$, but because of the structure of ℓ_∞ we cannot apply the method developed in this paper. Note also that the ideal of the strictly singular operators is the largest ideal in $L(\ell_\infty)$ (follows from [12, Theorem 1.2] and [8, Corollary 1.4]), the proof of which we include for completeness. In order to develop (if at all possible) a similar approach, one may have to find a suitable substitution for the set $\mathcal{A}(\mathcal{D})$ defined in (4) and an analog of the left essential spectrum (Definition 4.1). Also, a couple of times in this paper we have used the fact that every infinite dimensional subspace of ℓ_p ($1 \leq p < \infty$) contains a further subspace isomorphic to ℓ_p and complemented in ℓ_p , which does not hold for ℓ_∞ . This additional obstacle should be overcome as well. First we will prove

Lemma 5.4. *The ideal of strictly singular operators is the largest ideal in $L(\ell_\infty)$.*

Proof. Assume that T is not a strictly singular operator. Our goal will be to prove that any ideal that contains T must coincide with $L(\ell_\infty)$. Note first that on ℓ_∞ the ideals of the weakly compact and the strictly singular operators coincide ([12, Theorem 1.2]). Then we use the fact that any non-weakly compact operator is an isomorphism on some subspace Y of ℓ_∞ isomorphic to ℓ_∞ ([8, Corollary 1.4]). The subspaces Y and TY will be automatically complemented in ℓ_∞ because ℓ_∞ is an injective space. This automatically yields that I_{ℓ_∞} factors through T and hence any ideal containing T coincides with $L(\ell_\infty)$. \square

Theorem 5.5. *Let $T \in L(\ell_\infty)$ be a strictly singular operator. Then T is a commutator.*

Proof. Since T is a strictly singular operator, T is weakly compact ([8, Corollary 1.4]). Thus it follows that $T\ell_\infty$ is separable (since any weakly compact subset of the dual to any separable space is metrizable) and let $Y = \overline{T\ell_\infty}$. The space ℓ_∞/Y must be non-reflexive since assuming otherwise gives us that Y has a subspace isomorphic to ℓ_∞ ([5, Theorem 4]). Now consider the quotient map $Q : \ell_\infty \rightarrow \ell_\infty/Y$. Q is not weakly compact and hence (using again [8, Corollary 1.4]) there exists $X \simeq \ell_\infty$, $X \subset \ell_\infty$ such that $Q|_X$ is an isomorphism. Let P' be a projection onto QX and set $P = (Q|_X)^{-1}P'Q$. P is a projection in ℓ_∞ , $PY = \{0\}$ and by the construction, $P\ell_\infty$ is isomorphic to ℓ_∞ . Thus it follows that $PT = 0$ and we obtain that T is similar to an operator T' for which there exists a $M \subset \mathbb{N}$ such that $P_M T' = 0$. Using [2, Theorem 2.9] we conclude that T' is commutator and hence T is commutator. \square

References

- [1] A. Brown, C. Pearcy *Structure of commutators of operators*, Ann. of Math. **82** (1965), 112–127.
- [2] C. Apostol, *Commutators on ℓ_p spaces*, Rev. Roum. Math. Appl. **17** (1972), 1513–1534.
- [3] C. Apostol, *Commutators on c_0 -spaces and on ℓ_∞ -spaces*. Rev. Roum. Math. Pures Appl. **18** (1973), 1025–1032.
- [4] C. Apostol, *On the left essential spectrum and non-cyclic operators in Banach spaces*, Rev. Roum. Math. Appl. **17** (1972), 1141–1148.
- [5] J. Lindenstrauss, H.P. Rosenthal *Automorphisms in c_0, ℓ_1 and m* , Israel J. Math. **7** (1969), 227–239.
- [6] P. Enflo, T.W. Starbird *Subspaces of L_1 containing L_1* , Studia Math **65** (1979), 203–225.
- [7] A. Pelczynski, *Projections in certain Banach spaces*, Studia Math **19** (1960), 209–228.
- [8] H.P. Rosenthal, *On relatively disjoint families of measures, with some applications to Banach space theory*, Studia Math **37** (1970), 13–36.
- [9] H.P. Rosenthal, *Embeddings of L_1 in L_1* , Contemp. Math. **26** (1984), 335–349.
- [10] C. Schneeberger, *Commutators on a separable L_p space*, Proceedings of the American Mathematical Society **28**, Number **2** (1971), 464–472.
- [11] A. Wintner, *The unboundedness of quantum-mechanical matrices*, Phys. Rev **71** (1947), 738–739.
- [12] R.J. Whitley, *Strictly singular operators and their conjugates*. Trans. Amer. Math. Soc. **113** (1964), 252–261
- [13] J. Lindenstrauss, L. Tzafriri, *Classical Banach spaces. I*. Ergebnisse der Mathematik und ihrer Grenzgebiete, **Vol. 92** Springer-Verlag, Berlin-New York, (1977) xiii+188 pp
- [14] I.A. Feldman, I.C. Gohberg, A.S. Markus *Izv. Moldav. Filial AN SSSR* **10(76)** 51-69. (Russian)

Department of Mathematics, Texas A&M University, College Station, TX 77840
E-mail address : dossev@math.tamu.edu