

On rigid Hirzebruch genera

Oleg R. Musin *

Abstract

The classical multiplicative (Hirzebruch) genera of manifolds have the wonderful property which is called rigidity. Rigidity of a genus h means that if a compact connected Lie group G acts on a manifold X , then the equivariant genus $h^G(X)$ is independent on G , i.e. $h^G(X) = h(X)$.

In this paper we are considering the rigidity problem for complex manifolds. In particular, we are proving that a genus is rigid if and only if it is a generalized Todd genus.

1 Introduction

Let U_* be the complex bordism ring with coefficients in $R = \mathbb{Q}, \mathbb{R}$, or \mathbb{C} . For a closed complex manifold X , Hirzebruch [5] defined a multiplicative genus $h(X)$ by a homomorphism $h : U_* \otimes R \rightarrow R$.

Recall that according to Milnor and Novikov, two complex manifolds are complex cobordant if and only if they have the same Chern numbers. Therefore, for any multiplicative genus h there exists a multiplicative sequence of polynomials $\{K_i(c_1, \dots, c_i)\}$ such that $h(X) = K_n(c_1, \dots, c_n)$, where the c_k are the Chern classes of X and $n = \dim_{\mathbb{C}}(X)$ (see [5, 7, 3] and Section 2).

Let U_*^G be the ring of complex bordisms of manifolds with actions of a compact Lie group G . Then for any homomorphism $h : U_* \otimes R \rightarrow R$ we can define the *equivariant genus* h^G , i.e. a homomorphism

$$h^G : U_*^G \otimes R \rightarrow K(BG) \otimes R$$

(see details in [9]).

A multiplicative genus h is called *rigid* if for any connected compact group G the equivariant genus $h^G(X) = h(X)$. For the complex case rigidity means that

$$h^G : U_*^G \otimes R \rightarrow R \subset K(BG) \otimes R,$$

i.e. $h^G([X, G])$ belongs to the ring of constants. From the functorial properties of h^G it follows that it is sufficient to prove rigidity for the case $G = \mathbf{S}^1$.

For $G = \mathbf{S}^1$, the universal classifying space BG is $\mathbb{C}P^\infty$, and the ring $K(BG) \otimes R$ is isomorphic to the ring of formal power series $R[[t]]$. Then for any

*The author partially supported by NSF and NSA grants.

\mathbf{S}^1 -manifold X and a Hirzebruch genus h we have $h^{S^1}([X, \mathbf{S}^1])$ in $R[[t]]$. For instance, in the case of an \mathbf{S}^1 -action on an almost complex manifold X^n with isolated points p_1, \dots, p_m with weights w_{i1}, \dots, w_{in} , $i = 1, \dots, m$, h^{S^1} can be found explicitly (see [17, 14, 8, 9]):

$$h^{S^1}([X, \mathbf{S}^1]) = S_h(\{w_{ij}\}, t) := \sum_{i=1}^m \prod_{j \neq i} \frac{H(w_{ij} t)}{w_{ij} t}, \quad (1.1)$$

where H is the characteristic series of h (see Section 2). If h is rigid, then from (1.1) it follows that

$$h(X) = S_h(\{w_{ij}\}, t) \text{ for any } t. \quad (1.2)$$

Atiyah and Hirzebruch based on the Atiyah-Zinger index theorem proved that T_y -genus is rigid [1]. (Note that T_0 is the famous Todd genus.) Krichever [9] gives a proof of rigidity of the $T_{x,y}$ -genus using global analytic properties of $S_h(\{w_{ij}\}, t)$. It is not hard to see that (1.2) yields the Atiyah-Hirzebruch formula [1, 9]:

$$T_{x,y}(X) = \sum_{i=1}^m x^{s_i^+} (-y)^{s_i^-},$$

where s_i^+, s_i^- are numbers of positive and negative weights $\{w_{ij}\}$. Moreover, a similar formula holds for any stable (or quasi) complex \mathbf{S}^1 -manifolds with connected components of the set of fixed points F_1, \dots, F_m :

$$T_{x,y}(X) = \sum_{i=1}^m \sigma_i x^{s_i^+} (-y)^{s_i^-} T_{x,y}(F_i).$$

Here $\sigma_i = \pm 1$ is the ‘‘sign’’ of F_i . For almost complex manifolds all $\sigma_i = 1$ (see details in [4]).

Let h be a Hirzebruch genus for oriented manifolds, i.e. $h : \Omega_*^{\text{SO}} \otimes R \rightarrow R$ is a homomorphism of rings. Since the class of oriented manifolds is greater than the class of (stable) complex manifolds, the family of rigid genera for oriented manifolds is less than the family of complex rigid genera. Note that $T_{1,1}$ coincides with the famous L -genus (or Signature) [5]. Atiyah and Hirzebruch proved that the L -genus is rigid [1]. They also proved that the \hat{A} -genus vanishes (i.e. is rigid) for spin manifolds.

Krichever [10] extended the Atiyah-Hirzebruch theorem on the \hat{A} -genus to complex manifolds. Namely he proved that if for an integer $k > 1$ we have $c_1(X) \equiv 0 \pmod{k}$, then $A_k(X) = 0$.

The theory of elliptic genera and elliptic cohomologies which arose in the end of 1980s in the papers of Witten, Ochanine, Landweber, and Stong [20, 21, 18, 12] was stimulated by Witten’s conjecture concerning the rigidity of the character-index of ‘‘twisted’’ Dirac operators, or equivalently the rigidity of the equivariant elliptic genera [11, 12, 13]. Bott, Taubes, Hirzebruch, and Krichever [2, 19, 6, 11] have studied Witten’s conjecture and its extensions and proved

several rigidity theorems (see Liu [13] for references, historical overview, and rigidity theorems). Actually, Krichever [11] generalized elliptic genera studied in [2, 6, 12, 18, 19, 20, 21] and proved that they are rigid for SU-manifolds, i.e. for almost complex manifolds whose first Chern class is zero.

Note that among genera that were considered above only the family of $T_{x,y}$ genera is rigid for all complex manifolds. For all other rigid cases we need additional assumptions, for instance $c_1(X) = 0$ for Krichever's genera. In this paper we show that a multiplicative genus is rigid for all complex manifolds if and only if it is a generalized Todd genus (Sections 3 and 4).

2 Hirzebruch genera and algebraic rigidity

2.1 The genus of a formal power series

A sequence of polynomials $K_1(c_1), K_2(c_1, c_2), \dots$ in variables c_1, c_2, \dots with coefficients in R , where $R = \mathbb{Q}, \mathbb{R}$, or \mathbb{C} , is called *multiplicative* if

$$1 + c_1 t + c_2 t^2 + c_3 t^3 + \dots = (1 + a_1 t + a_2 t^2 + \dots)(1 + b_1 t + b_2 t^2 + \dots)$$

implies that

$$\sum_{j=0}^{\infty} K_j(c_1, \dots, c_j) t^j = \sum_{i=0}^{\infty} K_i(a_1, \dots, a_i) t^i \sum_{\ell=0}^{\infty} K_\ell(b_1, \dots, b_\ell) t^\ell.$$

Let

$$H(z) = 1 + r_1 t + r_2 t^2 + r_3 t^3 + \dots$$

be a formal power series in $R[[t]]$ with $r_i = K_i(1, 0, \dots, 0)$. The power series $H(t)$ is called the *characteristic (or Hirzebruch's) power series* of the multiplicative sequence $\{K_j\}$.

Note that a multiplicative sequence $\{K_j\}$ is uniquely defined by its characteristic power series $H(t)$ [5, Lemma 1.2.1]. Moreover, for any formal power series $H(t) = 1 + r_1 t + r_2 t^2 + \dots$ there exists a multiplicative sequence $\{K_j\}$ with Hirzebruch's power series $H(t)$ [5, Lemma 1.2.2].

The genus h of complex manifolds corresponding to H is given by

$$h(X) = K_n(c_1, \dots, c_n), \quad n = \dim_{\mathbb{C}}(X),$$

where the c_k are the Chern classes of X . Therefore, H defines the homomorphism $h : U_* \otimes R \rightarrow R$ from the complex cobordism ring $U_* \otimes R$ into R (see details in [3, 5, 7, 17]).

Novikov [17] proved that $t/H(t)$ is equal to $g_h^{-1}(t)$ in $R[[t]]$, where

$$g_h(t) = \sum_{n=0}^{\infty} \frac{h_n}{n+1} t^{n+1}$$

and

$$h_n := [H(t)^{n+1}]_n = h(\mathbb{C}P^n).$$

2.2 Linear circle actions on $\mathbb{C}P^n$

Consider the following action of \mathbf{S}^1 on $\mathbb{C}P^n$:

$$[z_0 : z_1 : \dots : z_n] \rightarrow [e^{iw_0\varphi}z_0 : e^{iw_1\varphi}z_1 : \dots : e^{iw_n\varphi}z_n]. \quad (2.1)$$

Denote $\mathbb{C}P^n$ with the circle action (2.1) by $\mathbb{C}P^n[w_0, \dots, w_n]$.

If w_0, \dots, w_n are distinct integers, then this action has $n + 1$ isolated fixed points p_0, \dots, p_n and weights of the representation of \mathbf{S}^1 in the tangent plane to p_i are $w_{0,i}, \dots, w_{i-1,i}, w_{i+1,i}, \dots, w_{n,i}$, where $w_{k,i} = w_k - w_i$.

Let $H = 1 + r_1t + r_2t^2 + \dots$ be a formal power series. Denote

$$F_H(t) := \frac{H(t)}{t} = \frac{1}{t} + r_1 + r_2t + r_3t^2 + \dots$$

and

$$S_H(w_0, \dots, w_n; t) := \sum_{i=0}^n \prod_{j \neq i} F_H([w_j - w_i]t).$$

Proposition 2.1. *Let $H(t) = 1 + r_1t + \dots \in R[[t]]$ and let w_0, \dots, w_m be distinct integers. Then $S_H(w_0, \dots, w_n; t) \in R[[t]]$ and $[S_H(w_0, \dots, w_n; t)]_0 = h_n$, i.e.*

$$S_H(w_0, \dots, w_n; t) = h_n + s_1t + s_2t^2 + \dots$$

Proof. It is known (see [17, 14, 8, 9]) that

$$h^{S^1}(\mathbb{C}P^n[w_0, \dots, w_n]) = S_H(w_0, \dots, w_n; t).$$

On the other hand, $h^{S^1}(\mathbb{C}P^n[w_0, \dots, w_n]) = h(\mathbb{C}P^n) + s_1t + s_2t^2 + \dots \in R[[t]]$. \square

2.3 Algebraically rigid series

Definition 2.1. *We say that a formal power series H is n -algebraically rigid and write $H \in \text{AR}^n$ if for any distinct integers w_0, \dots, w_m and for all $m \in [1, n]$ the formal series $S_H(w_0, \dots, w_m; t)$ is constant in $R[[t]]$, i.e.*

$$S_H(w_0, \dots, w_m; t) \in R \subset R[[t]].$$

If $H \in \text{AR}^n$ for all n , then we say that H is strong algebraically rigid and write $H \in \text{AR}^\infty$.

Proposition 2.2. *Rigidity yields strong algebraic rigidity, i.e. for any rigid Hirzebruch genus $h : U_* \otimes R \rightarrow R$ its characteristic power series $H \in \text{AR}^\infty$.*

Proof. Since

$$h^{S^1}(\mathbb{C}P^n[w_0, \dots, w_n]) = S_H(w_0, \dots, w_n; t),$$

the rigidity of h implies $S_H(w_0, \dots, w_n; t) = h_n \in R \subset R[[t]]$ for all n . \square

3 GT rigid series

Let $a \in R$. Consider the Euler characteristic, i.e. the Euler genus $a^n c_n(M^n)$. It is easy to see that for the Euler genus the multiplicative sequence of polynomials is

$$K_0 = 1, K_1 = ac_1, K_2 = a^2c_2, K_3 = a^3c_3, \dots$$

Then

$$H(t) = E_a(t) := 1 + at$$

and

$$h_n = (n + 1) a^n.$$

Let

$$H_0(t) = \frac{t}{1 - e^{-t}}.$$

Then h coincides with the Todd genus and all $h_n = 1$ [5, Lemma 1.7.1].

Hirzebruch [5, Lemma 1.8.1] considers the T_y -genus with

$$H_y(t) = \frac{t(e^{t(1+y)} + y)}{e^{t(1+y)} - 1}.$$

Note that T_0 is the Todd genus and

$$T_y(\mathbb{C}P^n) = \frac{1 - (-y)^{n+1}}{1 + y}.$$

T_y -genus was extended by Krichever [9]. He considered the $T_{x,y}$ -genus with

$$H_{x,y}(t) = \frac{t(xe^{t(x+y)} + y)}{e^{t(x+y)} - 1}$$

and

$$T_{x,y}(\mathbb{C}P^n) = \frac{x^{n+1} - (-y)^{n+1}}{x + y}. \quad (3.1)$$

Let $D_{a,b}(t) := H_{a+b,a-b}(t)$. Clearly,

$$D_{a,b}(t) = t(a \coth(at) + b).$$

Note that $D_{a,b}(t)$ is well defined for $a = 0$ and

$$D_{0,b}(t) = E_b(t).$$

Here we introduce one more family of genera. Let

$$G_{a,b}(t) := t(a \cot(at) + b).$$

Since $\cot(x) = i \coth(ix)$, we have

$$G_{a,b}(t) = D_{ia,b}(t). \quad (3.2)$$

Therefore, for $R = \mathbb{C}$, families of series D and G are the same.

Using (3.1) and (3.2) we get

$$g_{a,b}(\mathbb{C}P^n) = \frac{(b+ia)^{n+1} - (b-ia)^{n+1}}{2ia},$$

where $g_{a,b}$ is a genus with the characteristic series $G_{a,b}$.

Definition 3.1. *We say that a formal power series $H \in R[[t]]$ is the GT (Generalized Todd) series if there are $a, b \in R$ such that $H(t) = D_{a,b}(t)$ or $H(t) = G_{a,b}(t)$. (For the case $R = \mathbb{C}$ we may assume that $H(t) = D_{a,b}(t)$ only.)*

Theorem 3.1. *If $H \in R[[t]]$ is the GT series, then $h : U_* \otimes R \rightarrow R$ is rigid.*

Proof. Atiyah and Hirzebruch [1], Krichever [9] proved this theorem for the cases T_y and $T_{x,y}$ genera with $R = \mathbb{Q}$. We can repeat the proof in [9] (practically without changes) for the case $R = \mathbb{C}$. Therefore, the theorem holds for $T_{x,y}$, $x, y \in \mathbb{C}$, genera, i.e. for $D_{a,b}(t)$, $a, b \in \mathbb{C}$, series. From this it follows that the theorem holds also for $H = D_{a,b}$, $a, b \in R$, where $R = \mathbb{Q}$ or \mathbb{R} . Moreover, (3.2) implies a proof for the case $H = G_{a,b}$.

We also can propose a proof which is based on multiplicative generators of $U_*^{S^1}$ [15], where $U_*^{S^1}$ is the bordisms ring of complex manifolds with circle actions. The idea of this proof is very natural: we just verify the theorem for generators that are given in [15] explicitly. It is clear that rigidity of h for generators implies rigidity for all manifolds. Note that [15] gives a proof of the Atiyah - Hirzebruch theorem for the T_y -genus. \square

4 Converse theorem on rigid genera

In this section we prove the main theorem which implies that a Hirzebruch genus is rigid if and only if its characteristic is the GT series. In our paper [16] we consider the case of a rigid genus h such that $F_H(\log z)$ is a rational function in z . It is proved that in this case we have $H = H_{x,y}$. Surprisingly, the main line of a proof in [16] can be used also for the general case.

Lemma 4.1. *If $H = 1 + r_1t + r_2t^2 + \dots \in \text{AR}^2$, then*

$$F_H^2(-t) + h_1F_H(t) + F_H'(t) = h_2.$$

Proof. If $H \in \text{AR}^2$, then for any $a, b, c \in R$ we have

$$f((b-a)s)f((c-a)s) + f((a-b)s)f((c-b)s) + f((a-c)s)f((b-c)s) = h_2,$$

where $f(t) = F_H(t)$. Let $t = as$, $bs = \varepsilon$, and $cs = -\varepsilon$. Then

$$f(\varepsilon - t)f(-\varepsilon - t) + f(t + \varepsilon)f(2\varepsilon) + f(t - \varepsilon)f(-2\varepsilon) = h_2.$$

Let us denote by $O(\varepsilon)$ any series $\varphi(t, \varepsilon) = \varepsilon \psi(t, \varepsilon)$ with $\psi(t, \varepsilon) \in R[[t, \varepsilon]]$. We have

$$f(t + \varepsilon)f(2\varepsilon) + f(t - \varepsilon)f(-2\varepsilon)$$

$$\begin{aligned}
&= f(t + \varepsilon) \left(\frac{1}{2\varepsilon} + r_1 \right) + f(t - \varepsilon) \left(\frac{-1}{2\varepsilon} + r_1 \right) + O(\varepsilon) \\
&= 2r_1 f(t) + \frac{f(t + \varepsilon) - f(t - \varepsilon)}{2\varepsilon} + O(\varepsilon) = 2r_1 f(t) + f'(t) + O(\varepsilon).
\end{aligned}$$

(Here $g'(t) = \sum_k k g_k t^{k-1}$, where $g(t) = \sum_k g_k t^k$). Since $h_1 = 2r_1$, we obtain

$$f^2(-t) + h_1 f(t) + f'(t) + O(\varepsilon) = h_2,$$

and

$$F_H^2(-t) + h_1 F_H(t) + F_H'(t) = h_2.$$

□

Remark 4.1. *In fact, we have the following equality:*

$$h^{S^1}(\mathbb{C}P^2[1, 0, 0]) = F_H^2(-t) + h_1 F_H(t) + F_H'(t)$$

(see [9, 16]). So Lemma 4.1 shows that rigidity of h for $\mathbb{C}P^2[w_0, w_1, w_2]$ with distinct w_i yields that h is rigid for $\mathbb{C}P^2[w_0, w_1, w_1]$ also. It is clear that the same holds for all n , i.e. the rigidity of h for $\mathbb{C}P^n[w_0, \dots, w_n]$ with distinct w_i yields the rigidity of h for $\mathbb{C}P^n[w_0, \dots, w_n]$ with any w_i .

Theorem 4.1. *$H \in \text{AR}^2$ if and only if H is the GT series.*

Proof. Theorem 3.1 implies that if H is the GT series, then $H \in \text{AR}^\infty \subset \text{AR}^2$. (It is easy to prove this fact directly using elementary algebraic computations.)

Consider $H \in \text{AR}^2$. Then $H \in \text{AR}^1$ and

$$S_H(1, 0; t) = f(-t) + f(t) = h_1, \quad \text{where } f(t) := F_H(t).$$

This yields

$$f(-t) = h_1 - f(t). \tag{4.1}$$

Lemma 4.1 implies

$$(f(-t))^2 + h_1 f(t) + f'(t) = h_2.$$

Using (4.1), we get

$$(h_1 - f(t))^2 + h_1 f(t) + f'(t) = h_2.$$

Then

$$f' = -f^2 + h_1 f + h_2 - h_1^2.$$

From this follows that

$$t = \int \frac{df}{-f^2 + h_1 f + h_2 - h_1^2}. \tag{4.2}$$

Let

$$d = h_2 - 3h_1^2/4 = h_2 - 3r_1^2.$$

Consider two cases: (a) $d \neq 0$ and (b) $d = 0$

(a) $d \neq 0$. Let $q_1 = r_1 - \sqrt{d}$, $q_2 = r_1 + \sqrt{d}$. Then

$$\frac{q_2 - q_1}{-f^2 + h_1 f + h_2 - h_1^2} = \frac{1}{f - q_1} - \frac{1}{f - q_2}.$$

It follows from (4.2) that

$$(q_2 - q_1)t = \log\left(\frac{f - q_1}{f - q_2}\right) + c.$$

Note that $f(t) - 1/t \in R[[t]]$ implies $c = 0$. Thus,

$$H(t) = tf(t) = \frac{t(q_2 e^{(q_2 - q_1)t} - q_1)}{e^{(q_2 - q_1)t} - 1} = H_{q_2, -q_1}(t) = D_{\sqrt{d}, r_1}(t). \quad (4.3)$$

For the case $R = \mathbb{C}$, and for the cases $R = \mathbb{Q}$ or \mathbb{R} and $d > 0$ the equality (4.3) shows that H is the GT series.

In the cases $R = \mathbb{Q}$ or \mathbb{R} and $d < 0$ we have

$$H(t) = H_{q_2, -q_1}(t) = D_{ia, r_1}(t) = G_{a, r_1}(t), \quad \text{where } a = \sqrt{-d}.$$

(b) $d = 0$. We have

$$\frac{1}{-f^2 + h_1 f + h_2 - h_1^2} = \frac{-1}{(f - r_1)^2}.$$

It follows from (4.2) that

$$H(t) = tf(t) = 1 + r_1 t = E_{r_1}(t).$$

□

Corollary 4.1. *Let $R = \mathbb{Q}, \mathbb{R},$ or \mathbb{C} . A Hirzebruch genus $h : U_* \otimes R \rightarrow R$ is rigid if and only if H is the GT series.*

Corollary 4.2. *If H is 2-algebraically rigid, then H is strong algebraically rigid. In other words, $\text{AR}^2 = \text{AR}^3 = \dots = \text{AR}^\infty$.*

Consider the case of oriented manifolds. In this case the characteristic series H has to be even, i.e. $H(-t) = H(t)$ [5]. If we apply Theorem 4.1 for even series, then we get the following theorem.

Theorem 4.2. *Let $R = \mathbb{Q}, \mathbb{R},$ or \mathbb{C} . A Hirzebruch genus $h : \Omega_*^{\text{SO}} \otimes R \rightarrow R$ is rigid if and only if $H(t) = at \coth(at)$ or $H(t) = at \cot(at)$, where $a \in R$.*

Acknowledgments

The author thanks V. M. Buchstaber, I. M. Krichever, and S. P. Novikov for useful discussions and comments.

References

- [1] M. F. Atiyah and F. Hirzebruch, Spin manifolds and group actions, in Essays in Topology and Related Subjects, Springer-Verlag, Berlin, 1970, pp. 18-28.
- [2] R. Bott and C. Taubes, On the rigidity theorems of Witten, *J. Amer. Math. Soc.* **2** (1989) 137-186.
- [3] V. M. Buchstaber and S. P. Novikov, Formal groups, power systems and Adams operators, *Math. USSR-Sb.* **13** (1971), 80-116.
- [4] V. M. Buchstaber and N. Ray, The universal equivariant genus and Krichever's formula, *Russian Math. Surveys*, 62:1 (2007), 178 - 180.
- [5] F. Hirzebruch, Topological Methods in Algebraic Geometry, 3rd edition, Grundlehren der mathematischen Wissenschaften, no. 131, Springer, Berlin-Heidelberg 1966.
- [6] F. Hirzebruch, Elliptic genera of level N for complex manifolds, *Differential Geometric Methods in Theoretical Physics*, Kluwer, Dordrecht, 1988, 37-63.
- [7] F. Hirzebruch, T. Berger, and R. Jung, Manifolds and Modular Forms, Max-Planck-Institut für Math., Bonn, 1992.
- [8] G. G. Kasparov, Invariants of classical lens spaces in bordism theory, *Izv. Akad. Nauk SSSR Ser. Mat.*, **33** (1969), 735-747.
- [9] I. M. Krichever, Formal groups and the Atiyah - Hirzebruch formula, *Izv. Akad. Nauk SSSR Ser. Mat.* **38** (1974), 1289-1304.
- [10] I. M. Krichever, Obstructions to the existence of \mathbf{S}^1 -actions. Bordisms of ramified covering spaces, *Izv. Akad. Nauk SSSR*, **40** (1976), no. 4, 828-844.
- [11] I. M. Krichever, Generalized elliptic genera and Baker-Akhiezer functions, *Math. Notes* **47** (1990), no. 2, 132-142.
- [12] P. S. Landweber, Elliptic cohomology and modular forms, In Landweber, P. S., Elliptic curves and modular forms in algebraic topology, Lecture Notes in Math., Vol. **1326**, Springer, Berlin, 1988, 55-68.
- [13] K. Liu, On modular invariance and rigidity theorems, *J. Differential Geom.*, **41** (1995), 343-396.
- [14] A. S. Mischenko, Manifolds with actions and fixed points, *Math. Notes*, **4** (1968), 381-386.
- [15] O. R. Musin, Generators of \mathbf{S}^1 -bordism, *Math. USSR Sb.* **44** (1983), 325-334
- [16] O. R. Musin, Converse theorem on equivariant genera, to appear in *Russian Math. Surveys*.

- [17] S. P. Novikov, Adams operators and fixed points, *Math. USSR-Izv.* **2** (1968), 1193-1211.
- [18] S. Ochanine, Genres elliptiques equivariants, In Landweber, P.S., Elliptic curves and modular forms in algebraic topology, Lecture Notes in Math., Vol. **1326**, Springer, Berlin, 1988, 107-122.
- [19] C. Taubes, S^1 -actions and elliptic genera, *Comm. Math. Phys.* **122** (1989) 455-526.
- [20] E. Witten, Elliptic genera and quantum field theory, *Comm. Math. Phys.* **109** (1987) 525-536.
- [21] E. Witten, The index of the Dirac operator in loop space, In Landweber, P. S., Elliptic curves and modular forms in algebraic topology, Lecture Notes in Math., Vol. **1326**, Springer, Berlin, 1988, 161-186.

O. R. Musin, Department of Mathematics, University of Texas at Brownsville,
80 Fort Brown, Brownsville, TX, 78520.
E-mail address: oleg.musin@utb.edu