Flavor Symmetry Breaking and Vacuum Alignment on Orbifolds

Tatsuo Kobayashi, Yuji Omura, Koichi Yoshioka

Department of Physics, Kyoto University, Kyoto 606-8502, Japan

(September, 2008)

Abstract

Flavor symmetry has been widely studied for figuring out the masses and mixing angles of standard-model fermions. In this paper we present a framework for handling flavor symmetry breaking where the symmetry breaking is triggered by boundary conditions of scalar fields in extra-dimensional space. The alignment of scalar expectation values is achieved without referring to any details of scalar potential and its minimization procedure. As applications to non-abelian discrete flavor symmetries, illustrative lepton mass models are constructed where the S_3 and A⁴ flavor symmetries are broken down to the directions leading to the tribimaximal form of lepton mixing and realistic mass patterns.

1 Introduction

The masses and mixing angles of quarks and leptons have been long-standing and inspiring problems in particle physics. Their observed patterns, e.g. the recent experimental result for neutrino generation mixing [\[1\]](#page-18-0), suggest underlying principles such as grand unification and flavor symmetry lying behind the data.

Flavor symmetry has been widely investigated as a solution to the fermion mass problems. Among various flavor groups, a particular attention has recently been paid for non-abelian discrete groups such as S_3 and A_4 . These non-abelian discrete symmetries have several advantages that they provide a definite meaning for the generation and also link up different generations. However flavor symmetry should be broken in a nontrivial way while preserving the trace of symmetry structure. Flavor symmetry breaking is caused, e.g., by non-vanishing vacuum expectation values of scalar fields, which are determined by the minimization of scalar potential. For a non-abelian discrete symmetry to leave some imprints in low-energy symmetry-broken theory, the expectation values must take specific directions, namely, the vacuum alignment is required. That is usually achieved by performing the detailed analysis of scalar potential invariant under flavor symmetries [\[2,](#page-18-1) [3\]](#page-18-2). In particular, the vacuum alignment has recently been discussed in numbers of lepton mass models which account for the tri-bimaximal form of generation mixing [\[4\]](#page-18-3) as a good approximation of the experimental data. The tri-bimaximal mixing implies a specific form of mass matrix for light Majorana neutrinos: $M_L = \sum_i m_i v_i v_i^{\dagger}$ with $v_1 = (2, -1, -1), v_2 = (1, 1, 1)$ and $v_3 = (0, 1, -1)$. These inter-family related values of matrix elements seem to indicate non-abelian symmetry, vacuum alignment, and so hidden structure of nature beyond the standard model.

In this paper, we present a framework to realize aligned flavor symmetry breaking without any elaborated potential analysis. A central ingredient is the global property of bulk scalar fields in higher-dimensional space. At the boundaries of extra space, dynamical conditions for bulk fields are specified to fix the model. That reduces the number of degrees of freedom of symmetry groups [\[5\]](#page-19-0). This type of symmetry breaking with extra dimensions has been applied to, for example, the electroweak gauge symmetry [\[6\]](#page-19-1) and grand unified theory [\[7\]](#page-19-2). As for flavor symmetry, a viable breaking mechanism is provided by flavor-twisted boundary conditions for bulk fermion fields [\[8\]](#page-19-3). In this paper we discuss bulk scalar fields charged under the flavor symmetry, whose expectation values are aligned to cause discrete patterns of symmetry breaking. The vacuum alignment is archived in our approach by adopting the boundary conditions such that only one light

mode survives and it governs all vacuum expectation values in low-energy theory. Our method is also applicable to the quark sector and grand unified theory.

This paper is organized as follows. In Section 2, we first present the general procedure to have aligned scalar expectation values and classify the flavor-group matrix elements in the five-dimensional orbifold theory. The next two parts (Sections 3 and 4) are devoted to the applications of S_3 and A_4 flavor symmetries. We also present illustrative lepton mass models with flavor symmetries and vacuum alignments which are implemented on higher-dimensional orbifolds. Section 5 summarizes our results.

2 General recipe

In this section, we consider as the simplest example a five-dimensional orbifold theory on the flat gravitational background. The generalization to higher dimensions is straightforward. The five-dimensional coordinates are denoted by (x^{μ}, x^5) where x^5 corresponds to the fifth dimension. There are two types of operations on this space; the reflection $\hat{Z}: x^5 \to -x^5$ and the translation $\hat{T}: x^5 \to x^5 + 2\pi L$, where L is a constant. They trivially satisfy $\hat{T}\hat{Z} = \hat{Z}\hat{T}^{-1}$. On field variables living in the bulk, these operations are expressed in terms of matrices in the field space:

$$
\hat{Z}\phi(x^5) = Z^{-1}\phi(-x^5), \qquad \hat{T}\phi(x^5) = T^{-1}\phi(x^5 + 2\pi L). \tag{2.1}
$$

Let us suppose the extra space is compactified on the S^1/Z_2 orbifold with the radius L. This is achieved by the identifications $\hat{Z}(x^5) = x^5$ and $\hat{T}(x^5) = x^5$. On this orbifold, there are two fixed points (boundaries) at $x^5 = 0$ and $x^5 = \pi L$. The boundary conditions of bulk fields in compactifying the extra dimension are then defined by the identifications $\hat{Z}\phi(x^5) = \phi(x^5)$ and $\hat{T}\phi(x^5) = \phi(x^5)$.

It is noted that possible boundary conditions for bulk fields are limited by several consistency relations. First, since \hat{Z} behaves as parity, available choices are

$$
Z^2 = 1.\tag{2.2}
$$

In addition, for the translation, the matrix T must satisfy the relation

$$
ZT = T^{-1}Z.\t\t(2.3)
$$

Clearly not all types of boundary conditions are allowed. For example, in the case that Z is trivial (the unit matrix), the translation matrix T is limited to $T^2 = 1$. That is

more easily seen by defining another parity operation $\hat{Z}' = \hat{T}\hat{Z}$, which is interpreted as the reflection about $x^5 = \pi L$. The consistency relation [\(2.3\)](#page-2-0) just implies $(Z')^2 = 1$, where Z' is the matrix representation of \hat{Z}' on the field space.

With a set of boundary conditions, bulk fields are expanded by Kaluza-Klein modes. After integrating over the fifth dimension, we obtain a four-dimensional effective theory below the compactification scale $1/L$. In the low-energy theory, the physics is described by zero modes. In this paper, we focus on bulk scalar fields. They are assumed to be non-trivially charged under the flavor symmetry. The boundary condition determines which components of bulk scalars have (massless) zero modes. These zero modes are expected to develop their vacuum expectation values in low-energy effective theory and then contribute to Yukawa operators for quarks and leptons. The expectation values are determined, for example, à la Coleman-Weinberg [\[9\]](#page-19-4) or by use of boundary terms whose existence is however irrelevant to the following discussion, and further details are not considered in this paper. On the other hand, other excited modes are superheavy and stabilized at the origin with vanishing expectation values.

An idea is that a specific boundary condition is adopted under which only one zero mode remains in low-energy theory. All non-vanishing expectation values are controlled by this single zero mode and therefore related to each other, i.e., the vacuum alignment is established. If there are more than one zero modes, several directions of the vacua are degenerate and generally depend on free model parameters. It is noted that this procedure makes sense in phenomenology only for non-abelian and discrete symmetries. For an abelian or continuous symmetry, the vacuum described by multi field values is generally determined by continuous parameters and therefore the alignment does not work.

Here we give a general recipe for non-abelian discrete symmetry breaking in the fifth dimension S^1/Z_2 such that only a single zero mode survives. The bulk theory has a flavor symmetry acting on the three-generation quarks and leptons. We introduce bulk scalar fields ϕ_i (i = 1, 2, 3) which transform as a triplet under the flavor symmetry. The triplet representation can be reducible. The boundary conditions for the scalars ϕ_i are given by

$$
\hat{Z}: \qquad \phi(-x^5) = Z\phi(x^5), \tag{2.4}
$$

$$
\hat{T}: \phi(x^5 + 2\pi L) = T\phi(x^5). \tag{2.5}
$$

Here Z and T are the unitary representation matrices of flavor symmetry group and act on the generation space. The consistency relations require $Z^2 = (ZT)^2 = 1$. This

implies that for each choice of the matrix Z , it can be rotated to the diagonal form; $Z = \text{diag}(p_1, p_2, p_3)$ with $p_i = \pm 1$. For a positive (negative) parity element $p_i = +1$ $(p_i = -1)$, a corresponding bulk field has an even (odd) wavefunction about the extradimensional coordinate. Therefore the number of zero modes is given by that of positive parity elements when one takes into account the \hat{Z} boundary condition only. A fast conclusion at this stage is that at least one positive parity elements are needed to have one zero mode. Therefore we fix $p_1 = +1$ in the following discussion in this section. To include another boundary condition taken into account, it is useful to express the translation matrix as $T = e^{\pi i W}$ where W is the hermite 3×3 matrix written in the basis where Z is diagonal. From the boundary condition (2.5) , the mode expansion is found to be

$$
\phi_i(x^{\mu}, x^5) = \sum_n \left(\exp \left[i \left(\frac{W}{2} + n \right) \frac{x^5}{L} \right] \right)_{ij} \phi_j^{(n)}(x^{\mu}). \tag{2.6}
$$

The even (cosine) or odd (sine) function part is chosen depending on the parity assignment for ϕ_i . From this mode expansion, we find that massless zero modes are obtained when W has zero eigenvalues (mod 2). There are two types of boundary conditions for realizing a single zero mode:

I. The number of massless mode is reduced to 1 with either Z or T only. This is the case that the matrix Z has one positive parity, i.e., $p_1 = +1$ and $p_2 = p_3 = -1$. In this basis, W should take the following form:

$$
W = \begin{pmatrix} 0 & 0 & 0 \\ 0 & * & * \\ 0 & * & * \end{pmatrix} . \tag{2.7}
$$

The right-bottom 2×2 sub-matrix (denoted hereafter as w) is hermite and has integer eigenvalues so that the consistency relation $(2T)^2 = 1$ is satisfied. The translation \hat{T} is commutative with \hat{Z} and becomes a parity operation, but not necessarily flavor diagonal. The relative rotation between Z and W is undetermined at this stage, but such a freedom is irrelevant to the low-energy phenomenology of zero modes. We thus find that a single zero mode appears in ϕ_1 . For the flat extra dimension, the roles of two boundaries at $x^5 = 0$ and $x^5 = \pi L$ are interchangeable. This exchange gives another solution, namely $Z' = diag(+1, -1, -1)$ and w with integer eigenvalues. For any flavor symmetry group, we have such group elements of the form [\(2.7\)](#page-4-0). For example, $T = 1$ for $w = 0$ and $T = Z$ for $w = diag(\pm 1, \pm 1)$.

II. Another type of boundary conditions is that Z and T are in conspiracy to reduce the number of zero modes. This is the case that the matrix Z has two positive parities, i.e., $p_1 = p_2 = +1$ and $p_3 = -1$. The corresponding W has either of the following two forms.

The first case is that, like the Type I boundary condition, the translation matrix T becomes a parity and commutes with Z. It takes the form

$$
W = \begin{pmatrix} * & * & 0 \\ * & * & 0 \\ 0 & 0 & * \end{pmatrix} . \tag{2.8}
$$

The 3-3 element in the matrix [\(2.8\)](#page-5-0) is an integer, but the even integer case is equivalent to that in Type I if suitable exchanges of the generation indices and two boundaries are performed. The consistency relation $(ZT)^2 = 1$ and the requirement of a surviving zero mode restrict the left-top sub-matrix has one even and one odd eigenvalues. The zero mode is found in a definite linear combination of ϕ_1 and ϕ_2 . If the matrix W is expressed in a basis where the first component has a zero mode, it has the form $W = diag(0, 1, 1)$.

The second case is that W is non-commutative with Z . We find the most general expression to satisfy $(ZT)^2 = 1$ and to have a single zero mode:

$$
W = \begin{pmatrix} \beta^* \\ \alpha^* \\ \beta & \alpha \end{pmatrix}, \tag{2.9}
$$

with α and β being arbitrary complex numbers. The exception is that $|\alpha|^2 + |\beta|^2 =$ (even integer)² in which case there appear two zero modes. The translation matrix T obtained from [\(2.9\)](#page-5-1) coincides with [\(2.8\)](#page-5-0) just on the point $\alpha = 1$ and $\beta = 0$ (or vice versa), at which we have a zero mode in ϕ_1 .

Several comments are in order. The interchange of Z and Z' also gives a solution. Unlike the Type I boundary condition, there is no relative freedom between Z and T. This fact indicates the conspiracy of flavor symmetry breaking. Note also that the Type II boundary condition is viable only when the flavor symmetry group has a matrix representation X with det $X = -1$.

To summarize the result, the general recipe for obtaining a single zero mode is the following. First, we diagonalize all representation matrices for a triplet and list up the parity matrices with one positive and two negative eigenvalues. The boundary conditions

with this class of elements directly lead to one zero mode and realize the Type I boundary condition. The parity matrices with two positive and one negative eigenvalues are also useful if combined with the specific forms of translation, i.e. [\(2.8\)](#page-5-0) or [\(2.9\)](#page-5-1). The matrix [\(2.8\)](#page-5-0) has one even and one odd integer eigenvalues in the left-top sub-matrix. The Z and T operations conspire to lead to a single zero mode, called the Type II boundary condition. Through these procedures, one can list up all possible boundary conditions. Finally, we write down such boundary conditions in a unified basis.

This recipe is easily extended to higher-dimensional theories, for example, the sixdimensional theory on the orbifold T^2/Z_N for $N = 2, 3, 4, 6$.^{[1](#page-6-0)} The only difference from the present case is the form of consistency relations and mode expansion.

3 S_3 flavor symmetry

We first consider the permutation group S_3 as an example of flavor symmetry, which is the simplest non-abelian discrete group and has attractive features for flavor phe-nomenology [\[11\]](#page-19-5). The S_3 group is composed of six elements which are the identity I, two types of cyclic rotations R, R^2 , and three types of odd permutations P, PR and RP . They satisfy the characteristic relations for the S_3 group:

$$
P^2 = (PR)^2 = R^3 = 1.
$$
\n(3.1)

In addition to the trivial singlet, the S_3 group has two non-trivial representations, the doublet and pseudo singlet, the latter of which changes the sign under a permutation. Consequently, S_3 has two types of (reducible) triplet representations: the triplet 3_S and the pseudo triplet 3_A . On the field space, the circulation matrices are commonly represented on both triplets but the permutation matrices are different. They are explicitly written in the three-generation space by

$$
P_A = -\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, \quad P_S = \frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2 & 2 & -1 \\ 2 & -1 & 2 \end{pmatrix}, \quad R = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, \quad (3.2)
$$

and their derivatives R^2 , P_xR and RP_x ($x = A, S$). The parity operations \hat{Z}, \hat{Z}' are therefore given by P_x , P_xR , and RP_x . Their representation matrices have the determinants $+1$ and -1 for 3_A and 3_S , respectively.

¹See for various types of orbifolds in higher dimensions, e.g. [\[10\]](#page-19-6).

3.1 Vacuum alignment

For a scalar ϕ_A in the 3_A representation, only Type I boundary condition is possible since the parity matrices on 3_A have two negative eigenvalues. Let us consider the Z boundary condition associated with the twisting by P_A , i.e.,

$$
\phi_A(-x^5) = P_A \phi_A(x^5). \tag{3.3}
$$

The boundary condition with $P_A R$ or RP_A is equivalent to that with P_A just by label exchange. The matrix P_A is diagonalized by the unitary transformation $\phi_A = U_A \overline{\phi}_A$:

$$
U_A^{\dagger} P_A U_A = \begin{pmatrix} 1 & 0 & 0 & \frac{1}{\sqrt{3}} & \frac{-2}{\sqrt{6}} \\ -1 & -1 & 0 & 0 \end{pmatrix}, \qquad U_A = \begin{pmatrix} 0 & \frac{1}{\sqrt{3}} & \frac{-2}{\sqrt{6}} \\ \frac{-1}{\sqrt{2}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{6}} \end{pmatrix}.
$$
 (3.4)

The diagonalizing matrix U_A has a freedom in that its left-bottom 2×2 sub-matrix is undetermined due to the degenerate eigenvalues. However that is irrelevant to the following discussion of zero mode physics. From [\(2.7\)](#page-4-0) (i.e. the consistency relation), the boundary condition with the translation \hat{T} is found to be given by the matrix $T = I$ or P_A . Either choice for T leads to the same zero-mode physics. It is found from the parity sign in [\(3.4\)](#page-7-0) that only $\bar{\phi}^1_A$ has the massless zero mode with a constant wave function. It survives in four-dimensional low-energy effective theory and is expected to develop a vacuum expectation value. In terms of the original fields, the vacuum expectation values become

$$
\langle \phi_A \rangle = \langle U_A \bar{\phi}_A \rangle \propto (0, -1, 1). \tag{3.5}
$$

Thus the vacuum described by ϕ_A is determined discretely, that is, the vacuum alignment is achieved.

For a scalar ϕ_S in the 3_S representation, the Type II boundary condition is viable. Similar to the ϕ_A case, the \hat{Z} boundary condition is given by

$$
\phi_S(-x^5) = P_S \phi_S(x^5). \tag{3.6}
$$

The matrix P_S is diagonalized as follows $(\phi_S = U_S \overline{\phi}_S)$:

$$
U_S^{\dagger} P_S U_S = \begin{pmatrix} 1 & & \\ & 1 & \\ & & -1 \end{pmatrix}, \qquad U_S = \begin{pmatrix} \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & \frac{-1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & 0 & \frac{-2}{\sqrt{6}} \end{pmatrix}.
$$
 (3.7)

The parity signs in [\(3.7\)](#page-7-1) imply $\bar{\phi}_S^1$ and $\bar{\phi}_S^2$ could have massless zero modes. The Type II boundary condition further needs non-trivial \hat{T} twisting [\(2.8\)](#page-5-0) or [\(2.9\)](#page-5-1), satisfying the consistency relation. It is easily found that, in the basis (3.7) where P_S is diagonal, no S_3 group element satisfies [\(2.8\)](#page-5-0), but the matrix R (and R^2) has the form [\(2.9\)](#page-5-1) with $\alpha = -2i/3$ ($\alpha = +2i/3$). Therefore, imposing the \hat{T} boundary condition

$$
\phi_S(x^5 + 2\pi L) = R \phi_S(x^5), \tag{3.8}
$$

we find the mode expansion (2.6) with the translation matrix of the form (2.9) , and then obtain a single zero mode in $\bar{\phi}_S^1$. The vacuum expectation values of ϕ_S become

$$
\langle \phi_S \rangle = \langle U_S \bar{\phi}_S \rangle \propto (1, 1, 1). \tag{3.9}
$$

Thus the vacuum described by ϕ_S is aligned to the specific direction. Notice that this direction is independent and physically different from that of ϕ_A . They are not converted to each other by label exchanges.

3.2 Illustrative model

With the aligned vacuum at hand, it is an interesting task to construct realistic flavor symmetric theory for fermion masses and mixing in higher dimensions. In this section, we present an illustrative neutrino model based on the S_3 flavor symmetry.

Let us consider a six-dimensional model on the orbifold $T^2/(Z_2)^2$. While a straightforward application of the previous result leads to a viable five-dimensional model with the vacuum alignment, we here take a six-dimensional extension for Yukawa operators and flavor symmetry invariance (see the last part of this subsection). All the standard-model fields including the Higgs doublet h and three-generation lepton doublets ℓ_i (i = 1, 2, 3) as well as the right-handed neutrinos ν_i are confined on the fourdimensional fixed point at $x^5 = x^6 = 0$. Besides the gravity, only standard-model gauge singlet fields can propagate in the extra space not to violate the charge conservations. We thus introduce gauge singlet five-dimensional scalars ϕ_A and ϕ_S , which induce effective neutrino Yukawa operators. The five-dimensional fields ϕ_A and ϕ_S are extended to the fifth and sixth dimension, respectively.

The bulk theory has the flavor S_3 and parity symmetries (other than the parity

operation \hat{Z} in the extra dimensions) under which the fields have the following charges:^{[2](#page-9-0)}

$$
\begin{array}{c|cccc}\n\ell & \nu_1 & \nu_2 & \nu_3 & h & \phi_S & \phi_A \\
\hline\nS_3 & 3_S & 1_A & 1_S & 1_S & 1_S & 3_S & 3_A \\
Z & + & + & - & + & + & - & + \\
Z' & + & + & - & + & + & -\n\end{array}
$$
\n(3.10)

In addition to the kinetic and potential terms for each field, we have boundary interactions between the bulk and boundary fields, which give the effective Yukawa operators invariant under the flavor symmetry:

$$
\mathcal{L}_Y = \bar{\ell}_i \big(y_S \delta_{ij} + y_S' D_{ij} \big) \phi_{S_j} \nu_2 h + \bar{\ell}_i \big(y_A \delta_{ij} + y_A' D_{ij} \big) \phi_{A_j} \nu_3 h + \cdots \tag{3.11}
$$

where the ellipsis denotes higher-dimensional operators including ν_1 and the bulk field wavefunctions are evaluated at the fixed point. These terms are described by two types of symmetry-invariant tensors: the unit matrix δ_{ij} and the democratic matrix D_{ij} where all elements are equal to 1. It is noted that the parameters y_A and y'_A obey the relation $y_A = -3y'_A$, which reflects the fact that the tensor product $3_A \times 3_S$ contains only one trivial singlet.

In compactifying the extra dimensions, the boundary conditions on the bulk scalar fields must be specified. Along the line discussed above, we impose the following conditions at the two boundaries of the extra dimension:

$$
\phi_A(-x^5) = P_A \phi_A(x^5), \qquad \phi_A(x^5 + 2\pi L_5) = \phi_A(x^5), \tag{3.12}
$$

$$
\phi_S(-x^6) = P_S \phi_S(x^6), \qquad \phi_S(x^6 + 2\pi L_6) = R \phi_S(x^6), \tag{3.13}
$$

and all the other boundary conditions are trivial (i.e., non-twisted). According to the general recipe, they result in a single zero mode in each ϕ_A and ϕ_S , and the aligned vacuum is found to be

$$
\langle \phi_A \rangle = a(0, 1, -1), \quad \langle \phi_S \rangle = s(1, 1, 1).
$$
 (3.14)

The coefficients a and s involve the zero-mode expectation values and wavefunction factors evaluated at the fixed point where the standard-model fields reside. Substituting the scalar expectation values and integrating out heavy modes, we obtain the low-energy effective model below the compactification scale, which include the neutrino Yukawa couplings $\mathcal{L} = \overline{\ell_i} Y_{ij} \nu_j h + \text{h.c.}$

$$
Y = \begin{pmatrix} 0 & s(y_S + 3\bar{y}_S) & 0 \\ 0 & s(y_S + 3\bar{y}_S) & ay_A \\ 0 & s(y_S + 3\bar{y}_S) & -ay_A \end{pmatrix}.
$$
 (3.15)

² The right-handed neutrino ν_1 may belong to 1_S if it has a large Majorana mass and is decoupled.

Further, the flavor symmetry induces the generation-diagonal form of right-handed neutrino Majorana masses: $\mathcal{L}_{\nu} = \overline{\nu^c} M_R \nu$ with $M_R = \text{diag}(M_1, M_2, M_3)$. Implementing the seesaw mechanism, we obtain the Majorana mass matrix of light neutrinos

$$
M_L = m \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} + m' \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}, \tag{3.16}
$$

$$
m = \frac{s^2(y_S + 3\bar{y}_S)^2}{M_2}, \qquad m' = \frac{a^2 y_A^2}{M_3}.
$$
 (3.17)

This form of mass matrix is well consistent with the current experimental results. That is, it leads to the tri-bimaximal generation mixing [\[4\]](#page-18-3) and the mass squared differences $\Delta m_{21}^2 = 9|m|^2$ and $\Delta m_{32}^2 = 4|m'|^2$, which indicate the normal hierarchy spectrum. A non-vanishing mass eigenvalue for the first-generation light neutrino and the deviation from the tri-bimaximal mixing are both generated by the corrections from higherdimensional operators involving ν_1 . The flavor symmetry implies these operators of the forms $\ell \phi_A^2 \nu_1 h$ and $\ell \phi_S^2 \nu_1 h$. Compared with the leading terms above, they are suppressed by $\langle \phi_{A,S} \rangle / \Lambda$ where Λ is the cutoff scale of the theory. Other higher-dimensional operators suppressed by the cutoff scale are reasonably made small in a similar way. The corrections to low-energy physics from Kaluza-Klein excited modes could also be insignificant due to some extra-dimensional property [\[12\]](#page-19-7). While we do not explicitly discuss the charged-lepton sector here, an example of charged-lepton mass matrix is presented in [\[8\]](#page-19-3) where the right-handed charged leptons and a related scalar field are introduced in the four-dimensional boundary, and the induced hierarchy of charged-lepton mass eigenvalues realizes that the total lepton mixing is dominated by the neutrino tribimaximal mixing obtained above. Similar or more reasonable models might be available in a variety of scenarios which have been discussed in four or higher-dimensional theory.

The twisting boundary conditions of scalar fields generally induce flavor-breaking Lagrangian terms on the boundary of orbifold. For ϕ_A , it is easily found that the twisting (3.12) does not affect the induced Yukawa couplings Y (3.15) due to the aligned vacuum. On the other hand, for $\phi_{\mathcal{S}}$, the flavor symmetry breaking could split the matrix elements in the second column of Y. That is ameliorated, for example, by treating ϕ_S as a six-dimensional field (and by exchanging the two boundary conditions at $x^5 = 0$ and πR). In this case, the fixed point where the standard-model fields live respects the whole S_3 flavor symmetry and so does the induced Yukawa operators.

4 A_4 flavor symmetry

The next example is A_4 which consists of the even permutations of four objects. The A_4 flavor symmetry has been widely studied in the literature in view of the recent neutrino experimental data [\[13\]](#page-20-0). The A_4 group is composed of twelve elements generated by two fundamental elements P , R and their derivatives: PR , RP , R^2 , PR^2 , PRP , RPR , R^2P , $RPR²$, and $R²PR$. They satisfy the characteristic relations for the $A₄$ group:

$$
P^2 = (PR)^3 = R^3 = 1.
$$
\n(4.1)

In addition to the trivial singlet, the A_4 group has three non-trivial representations, the triplet and two pseudo singlets (denoted as 1′ and 1′′), the latters of which are respectively multiplied by the complex numbers χ and χ^2 under the R operation (χ = $e^{2\pi i/3}$). The representation matrices for the triplet are built up from

$$
P = \begin{pmatrix} 1 & & \\ & -1 & \\ & & -1 \end{pmatrix}, \qquad R = \begin{pmatrix} 1 & & \\ & 1 & \\ 1 & & \end{pmatrix}. \tag{4.2}
$$

The parity operations \hat{Z} , \hat{Z}' are written by P, RPR^{-1} and $R^{-1}PR$. All their representation matrices have the determinants $+1$. The tensor product of two triplets $\phi = (\phi_1, \phi_2, \phi_3)$ and $\varphi = (\varphi_1, \varphi_2, \varphi_3)$ are given by

$$
\phi \times \varphi = (\phi_2 \varphi_3, \phi_3 \varphi_1, \phi_1 \varphi_2)_3 + (\phi_3 \varphi_2, \phi_1 \varphi_3, \phi_2 \varphi_1)_3 + (\phi_1 \varphi_1 + \phi_2 \varphi_2 + \phi_3 \varphi_3)_1 + (\phi_1 \varphi_1 + \chi^2 \phi_2 \varphi_2 + \chi \phi_3 \varphi_3)_1 + (\phi_1 \varphi_1 + \chi \phi_2 \varphi_2 + \chi^2 \phi_3 \varphi_3)_1,
$$
\n(4.3)

where the suffices denote the A_4 representations.

4.1 Vacuum alignment

Let us consider a bulk scalar ϕ in the triplet representation of A_4 symmetry and examine the boundary conditions which leave a single zero mode in four-dimensional effective theory.

It is first noticed that, if the extra space is one dimensional, only one type of such boundary conditions is found to be allowed for A_4 symmetry. This is the Type I boundary condition with $T = I$ and $Z \neq I$. In the basis that the parity representation matrices are flavor diagonal, the other group matrices involve the rotation R and do not take the form [\(2.7\)](#page-4-0), i.e, they do not satisfy the consistency relation among the fifth-dimensional operations. There is no possibility for the Type II boundary condition since the A_4 group matrix elements have the determinants $+1$ which implies an even number of negative parity signs in each matrix. In this way, we find only possibility to have a single zero mode is $T = I$ and $Z \neq I$. However, with one type of boundary conditions, we obtain only one type of alignments, which is not compatible with rich flavor structure of quarks and leptons. This fact leads us to introduce two spatial extra dimensions for the A_4 flavor symmetry.^{[3](#page-12-0)}

Unlike the five-dimensional theory, there are several different ways to compactify the extra two-dimensional space on an orbifold. In this section, we consider the $T^2/(Z_2 \times Z_3)$ $(\simeq T^2/Z_6)$ orbifold. It is however stressed that the result obtained below is also applicable to T^2/Z_2 and T^2/Z_3 orbifolds because the Z_2 and Z_3 orbifolding are independent as will be seen. It is useful to denote the extra-dimensional coordinates (x^5, x^6) by a complex one $z = x^5 + ix^6$. The extra-dimensional space has four types of operations; the translations \hat{T} and \hat{T}' , the reflection \hat{Z}_2 , and the rotation \hat{Z}_3 . They act on z as

$$
\hat{T} : z \to z + 2\pi L, \qquad \hat{Z}_2 : z \to -z, \n\hat{T}' : z \to z + 2\pi L \chi, \qquad \hat{Z}_3 : z \to \chi z,
$$
\n(4.4)

with χ being the cubic root of the unity $(\chi = e^{2\pi i/3})$. The T^2/Z_6 orbifold is obtained by the identifications with these operations. The \hat{Z}_2 parity has the four fixed points $z=0$, πL , $\pi L \chi$ and $\pi L(1+\chi)$. The \hat{Z}_3 action leaves the three points $z = 0$, $(1+2\chi)/3$ and $(2 + \chi)/3$ unchanged. Therefore the T^2/Z_6 orbifold has one fixed point at the origin $z = 0$. We will later consider that the standard model fields live on this four-dimensional fixed point. On the field variables $\phi(x^{\mu}, z)$, the compactification is performed with the identification;

$$
\phi(z + 2\pi L) = T \phi(z), \qquad \phi(-z) = Z_2 \phi(z),
$$

$$
\phi(z + 2\pi L \chi) = T' \phi(z), \qquad \phi(\chi z) = Z_3 \phi(z), \qquad (4.5)
$$

where T, T' and $Z_{2,3}$ are the triplet representation matrices of A_4 group.

It is important that these matrices should satisfy several consistency relations associated with the property of orbifold. From the transformation rules [\(4.4\)](#page-12-1), we find the

³An alternative is to consider additional flavor symmetry acting on bulk scalar ϕ . In particular, if ϕ is a negative parity under an additional bulk symmetry, the odd number of negative parity eigenvalues is possible with boundary conditions involving this parity twisting. In this case, Type II boundary condition is also viable within five-dimensional A_4 theory.

following independent relations for the T^2/Z_6 orbifold:

$$
Z_2 Z_3 = Z_3 Z_2, \t\t(4.6)
$$

$$
(Z_2)^2 = (Z_2T)^2 = (Z_2T')^2 = 1,
$$
\n(4.7)

$$
TT' = T'T,\t\t(4.8)
$$

$$
(Z_3)^3 = (Z_3T)^3 = 1, \qquad TZ_3 = Z_3T'.
$$
 (4.9)

The first relation means that Z_2 and Z_3 are commutative and their representation matrices are simultaneously diagonalized. The second line implies that, like the S^1/Z_2 orbifold, other forms of parity operations are constructed which are the reflections about the Z_2 fixed points. The last two types of relations are characteristic for six-dimensional theory. It is interesting that these consistency relations rather reduce the number of possible boundary conditions. We find that only two types of non-trivial boundary conditions are viable in the A_4 theory on T^2/Z_6 :

(i)
$$
Z_2 = P
$$
, $Z_3 = T = T' = I$ (4.10)

(ii)
$$
Z_3 = R
$$
, $Z_2 = T = T' = I$ (4.11)

where the matrices P and R are given in (4.2) and I is the unit matrix. The other sets of matrices with exchanging generation labels also become the solutions. It is noticed that any non-trivial boundary condition associated with the translations is not allowed. The above classification of boundary conditions is general and valid for any flavor symmetry unless it has non-trivial commutative matrices A and B which satisfy $A^2 = B^3 = 1$.

For the boundary condition (i), i.e., $\phi(-z) = P\phi(z)$, it is easily found from the matrix form [\(4.2\)](#page-11-0) that we have a single zero mode in ϕ_1 due to a positive parity. The zero mode has a constant wavefunction profile. Then the vacuum expectation value becomes

$$
\langle \phi \rangle \propto (1, 0, 0). \tag{4.12}
$$

On the other hand, the boundary condition (ii) is explicitly written by

$$
\phi(\chi z) = R \phi(z),\tag{4.13}
$$

and the other conditions are trivial and do not affect the physics. It is convenient to write the matrix R in the diagonal form $(\phi = U_R \bar{\phi})$

$$
U_R^{\dagger} R U_R = \begin{pmatrix} 1 & & \\ & \chi & \\ & & \chi^2 \end{pmatrix}, \qquad U_R = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & \chi & \chi^2 \\ 1 & \chi^2 & \chi \\ 1 & 1 & 1 \end{pmatrix} . \tag{4.14}
$$

The eigenvalues obtained in [\(4.14\)](#page-13-0) indicate that only $\bar{\phi}_1$ has a zero mode with a flat wavefunction. The vacuum expectation value of the original field ϕ thus becomes

$$
\langle \phi \rangle \propto (1, 1, 1). \tag{4.15}
$$

Therefore the alignment of vacuum is established, similarly to the S_3 flavor theory in five dimensions.

We also have the vacuum alignments on other types of orbifolds T^2/Z_N with $N = 2, 3$ and 4. For T^2/Z_2 , the consistency conditions are a fewer sets than for T^2/Z_6 , i.e. (4.7) – (4.8) . Therefore in addition to (4.10) we generally have another solutions for the consistency conditions and one surviving zero mode. The most general solutions for T^2/Z_2 are given by $Z_2, T, T' = 1$ or P (and those with label exchanges). It is noticed that the aligned vacuum is not affected by this wider possibility of twisting and is given by [\(4.12\)](#page-13-4). The same is true for the T^2/Z_3 orbifold. The consistency conditions are [\(4.8\)](#page-13-2)–[\(4.9\)](#page-13-5) and the general solutions for these are given by $Z_3, T, T' = 1$ or R (and those with changing labels and phases). The light-mode physics is not altered with these choices of boundary conditions and the aligned vacuum becomes [\(4.15\)](#page-14-0). Finally, for the T^2/Z_4 orbifold, the independent consistency conditions are $(Z_4)^4 = (Z_4^2 T)^2 = 1$ and $TZ_4 = Z_4T'$. For the A_4 symmetry, the general solutions are given by $Z_4, T, T' = 1$ or P with $T = T'$ (and those with label exchanges) which leave only one zero mode in lowenergy theory. The vacuum alignment is the same as that on T^2/Z_2 .

4.2 Illustrative model

We apply the above result for the A_4 flavor symmetry to constructing of an explicit orbifold model for lepton masses and generation mixing. Similarly to the S_3 model in the previous section, a higher-dimensional extra space is introduced for flavor symmetry invariance, while the vacuum alignment itself is viable in lower dimensions.

Let us consider an eight-dimensional theory on the orbifold $T^2/Z_2 \times T^2/Z_3$. The standard-model fields are assumed to live on a fixed point of the orbifold, which we here choose the origin of extra-dimensional space. The three families of left-handed lepton doublets ℓ_i transform as a triplet of A_4 while the right-handed charged leptons e_1, e_2, e_3 are assigned to three different singlets, 1, 1', 1'', respectively. We introduce two gaugesinglet bulk scalars ϕ and ϕ' of the triplet representation, which give effective Yukawa and mass operators for leptons in low-energy theory. The fifth and sixth dimensions are compactified on T^2/Z_2 and the seventh and eighth ones on T^2/Z_3 . The scalars ϕ

and ϕ' live on T^2/Z_2 and T^2/Z_3 , respectively.^{[4](#page-15-0)} The basic building block is the same as the models presented in [\[3\]](#page-18-2). The bulk theory has the flavor A_4 and Z'_3 symmetries (other than the operation \hat{Z}_3). The field content and the flavor-symmetry charges are summarized below:

$$
\begin{array}{c|ccccccccc}\n & \ell & e_1 & e_2 & e_3 & h & \eta & \phi & \phi' \\
\hline\nA_4 & 3 & 1 & 1' & 1'' & 1 & 1 & 3 & 3 \\
Z & \chi & \chi & \chi & \chi & 1 & \chi & \chi & 1\n\end{array}
$$
\n(4.16)

Here h and η are the electroweak doublet and singlet Higgs fields, respectively. We have boundary interactions between the bulk and boundary fields which are invariant under the flavor symmetry:

$$
\mathcal{L}_Y = y_1 \bar{e}_1 \ell \phi h + y_2 \bar{e}_2 \phi' \ell h + y_3 \bar{e}_3 \ell \phi' h + w_1 \phi \bar{\ell}^c \ell h^2 + w_2 \eta \bar{\ell}^c \ell h^2 + \cdots, \qquad (4.17)
$$

where $w_{1,2}$ and $y_{1,2,3}$ are the coupling constants and the wavefunctions of bulk scalar fields are evaluated at the fixed point on which the standard-model fields reside.^{[5](#page-15-1)}

To fix the model, the boundary conditions for bulk fields must be specified. According to the result obtained in the above, we have the following non-trivial boundary conditions on the scalar fields:

$$
\phi(-z) = P\,\phi(z),\tag{4.18}
$$

$$
\phi'(\chi z') = R \phi'(z'), \qquad (4.19)
$$

where $z = x^5 + ix^6$ and $z' = x^7 + ix^8$, and the other boundary conditions are trivial. Under these conditions, the previous analysis says that the zero modes are found in ϕ_1 and $\phi'_1 + \phi'_2 + \phi'_3$ which develop the expectation values of the form

$$
\langle \phi \rangle = a(1, 0, 0), \qquad \langle \phi' \rangle = a'(1, 1, 1), \tag{4.20}
$$

in low-energy effective theory below the compactification scale. It is noticed that these aligned forms of expectation values were required in the models of Ref. [\[3\]](#page-18-2). Inserting the scalar expectation values in [\(4.17\)](#page-15-2), we obtain the operators for charged-lepton Dirac masses and neutrino Majorana masses; $\mathcal{L}_M = \bar{e}_i(M_e)_{ij} \ell_j + \overline{\ell_i^c}(M_\nu)_{ij} \ell_j$:

$$
M_e = a'v \begin{pmatrix} y_1 & y_1 & y_1 \\ y_2 & \chi^2 y_2 & \chi y_2 \\ y_3 & \chi y_3 & \chi^2 y_3 \end{pmatrix}, \qquad M_L = v^2 \begin{pmatrix} w_2 b & w_2 b & w_1 a \\ w_2 b & w_1 a & w_2 b \end{pmatrix}, \qquad (4.21)
$$

⁴The T^2/Z_2 orbifold can be replaced with S^1/Z_2 . In this case, we consider a seven-dimensional theory compactified on $S^1/Z_2 \times T^2/Z_3$.

⁵For the mode expansion on the T^2/Z_3 orbifold, see for example Ref. [\[14\]](#page-20-1).

where $v = \langle h \rangle$ and $b = \langle \eta \rangle$. These forms of mass matrices are known [\[3\]](#page-18-2) to be well fitted to the present experimental values. The lepton generation mixing is exactly given by the tri-bimaximal mixing. The neutrino mass eigenvalues are predicted as $\Delta m_{21}^2 =$ $-\text{Re}[w_1a(w_1a + 2w_2b)]v^2/2$ and $\Delta m_{31}^2 = -4\text{Re}(w_1w_2ab)v^2$. The hierarchy of chargedlepton masses is proportional to that of y_i . It may be implemented by, for example, the Froggatt-Nielsen mechanism [\[15\]](#page-20-2) or extra-dimensional scheme [\[16\]](#page-20-3), where the three right-handed charged leptons have different properties.

Finally, we comment on the boundary conditions (4.18) – (4.19) and the symmetry invariance of the boundary Lagrangian \mathcal{L}_Y . As mentioned before, the consistent boundary conditions for T^2/Z_2 and T^2/Z_3 have wider possibilities than [\(4.10\)](#page-13-3) and [\(4.11\)](#page-13-6). An important point is that the vacuum alignment itself is not affected by such choices of twisting as long as the light-mode physics is concerned. This fact can, in turn, be utilized to protect boundary terms by the flavor symmetry. Namely, flavor symmetry breaking occurs within the hidden sector which is separate in the extra-dimensional space from the visible sector where the standard-model fields reside. Such boundary conditions are given, for example, by $Z_2 = 1$ and $T = T' = P$ for T^2/Z_2 , and $Z_3 = 1$ and $T = T' = R$ for T^2/Z_3 . Bulk scalar fields interact both of these sectors and induce effective mass operators by connecting up the symmetry-invariant terms on the visible boundary and the aligned vacuum expectation values on the hidden boundary.

5 Summary

In this paper we have presented the scenario for breaking flavor symmetry with Scherk-Schwarz twisted boundary conditions on bulk scalar fields, where only one zero mode from multiple scalar fields survives in low-energy effective theory. For non-abelian discrete flavor symmetry, the vacuum alignment is shown to be achieved, that is, the symmetry-breaking expectation values of these scalars are aligned to definite directions. This implies that flavor symmetry in the high-energy regime can leave some imprints to the low-energy regime and provides a viable explanation for the fermion mass problems. Our scheme of vacuum alignment needs no elaboration of analyzing complicated scalar potential which generally involves multiple scalar fields and requires extra symmetries to realize the vacuum alignment.

As applications of our approach, the S_3 and A_4 flavor symmetries are considered in five and six dimensions compactified on the S^1/Z_2 and T^2/Z_N $(N = 2, 3, 4, 6)$ orbifolds, and possible types of boundary conditions are classified. Further we have constructed explicit models for lepton masses and mixing, and shown that simpler setups lead to realistic phenomenology such as the tri-bimaximal generation mixing of neutrinos. Other possibilities including the quarks sector/grand unification, curved gravitational backgrounds, geometrical origins of flavor symmetries in extra dimensions [\[17\]](#page-20-4) and so on will be investigated in future study.

Acknowledgments

This work is supported by the scientific grants from the ministry of education, science, sports, and culture of Japan No. 20540266 and 20740135) and also by the grant-in-aid for the global COE program "The next generation of physics, spun from universality and emergence" and the grant-in-aid for the scientific research on priority area $(\#441)$ "Progress in elementary particle physics of the 21st century through discoveries of Higgs boson and supersymmetry" (No. 16081209). The work of Y.O. is supported by the Japan society of promotion of science (No. $20 \cdot 324$).

References

- [1] M. Maltoni, T. Schwetz, M.A. Tortola and J.W.F. Valle, New J. Phys. 6 (2004) 122; R.N. Mohapatra *et al.*, Rept. Prog. Phys. **70** (2007) 1757; A. Strumia and F. Vissani, [hep-ph/0606054.](http://arxiv.org/abs/hep-ph/0606054)
- [2] For example, D. Wyler, Phys. Rev. D19 (1979) 3369; E. Derman and H.S. Tsao, Phys. Rev. D20 (1979) 1207; R. Barbieri, L.J. Hall, G.L. Kane and G.G. Ross, [hep-ph/9901228;](http://arxiv.org/abs/hep-ph/9901228) I. de Medeiros Varzielas and G.G. Ross, Nucl. Phys. B733 (2006) 31; W. Grimus and L. Lavoura, JHEP 0601 (2006) 018; C. Hagedorn, M. Lindner and F. Plentinger, Phys. Rev. D74 (2006) 025007; S.F. King and M. Malinsky, Phys. Lett. **B645** (2007) 351; F. Feruglio, C. Hagedorn, Y. Lin and L. Merlo, Nucl. Phys. B775 (2007) 120; S. Kaneko, et al, Prog. Theor. Phys. 117 (2007) 161; [hep-ph/0703250;](http://arxiv.org/abs/hep-ph/0703250) M.C. Chen and K.T. Mahanthappa, Phys. Lett. B652 (2007) 34; C. Luhn, S. Nasri and P. Ramond, Phys. Lett. B652 (2007) 27; F. Bazzocchi, S. Kaneko and S. Morisi, JHEP 0803 (2008) 063; B. Brahmachari, S. Choubey and M. Mitra, Phys. Rev. D77 (2008) 073008.
- [3] G. Altarelli and F. Feruglio, Nucl. Phys. B720 (2005) 64; G. Altarelli, F. Feruglio and C. Hagedorn, *JHEP* 0803 (2008) 052.
- [4] P.F. Harrison, D.H. Perkins and W.G. Scott, Phys. Lett. B530 (2002) 167; P.F. Harrison and W.G. Scott, Phys. Lett. B535 (2002) 163; Z.z. Xing, Phys. Lett. B533 (2002) 85; X.G. He and A. Zee, Phys. Lett. B560 (2003) 87; R.N. Mohapatra, S. Nasri and H.B. Yu, Phys. Lett. B639 (2006) 318; I. de Medeiros Varzielas, S.F. King and G.G. Ross, Phys. Lett. B648 (2007) 201; H. Zhang, Phys. Lett. B655 (2007) 132; P.D. Carr and P.H. Frampton, [hep-ph/0701034;](http://arxiv.org/abs/hep-ph/0701034) A. Dighe, S. Goswami and P. Roy, Phys. Rev. D76 (2007) 096005; Y. Koide, [arXiv:0707.0899;](http://arxiv.org/abs/0707.0899) N. Nimai Singh, H. Zeen Devi and M. Patgiri, [arXiv:0707.2713;](http://arxiv.org/abs/0707.2713) S. Pakvasa, W. Rodejohann and T.J. Weiler, *Phys. Rev. Lett.* **100** (2008) 111801; S. Antusch, S.F. King and M. Malinsky, JHEP 0805 (2008) 066; N. Haba, R. Takahashi, M. Tanimoto and K. Yoshioka, [arXiv:0804.4055;](http://arxiv.org/abs/0804.4055) F. Plentinger, G. Seidl and W. Winter, JHEP 0804 (2008) 077; C.S. Lam, [arXiv:0809.1185;](http://arxiv.org/abs/0809.1185) see for early work, L. Wolfenstein, Phys. Rev. D18 (1978) 958; Y. Yamanaka, H. Sugawara and S. Pakvasa, Phys. Rev. D25 (1982) 1895.
- [5] J. Scherk and J.H. Schwarz, Phys. Lett. B82 (1979) 60; Nucl. Phys. B153 (1979) 61. R. Barbieri, L.J. Hall and Y. Nomura, Nucl. Phys. B624 (2002) 63; J.A. Bagger, F. Feruglio and F. Zwirner, Phys. Rev. Lett. 88 (2002) 101601.
- [6] A. Delgado, A. Pomarol and M. Quiros, Phys. Rev. D60 (1999) 095008; C. Csaki, C. Grojean, H. Murayama, L. Pilo and J. Terning, Phys. Rev. D69 (2004) 055006; R. Barbieri, A. Pomarol and R. Rattazzi, Phys. Lett. B591 (2004) 141; Y. Hosotani, S. Noda and K. Takenaga, Phys. Lett. B607 (2005) 276; C.A. Scrucca, M. Serone, L. Silvestrini and A. Wulzer, JHEP 0402 (2004) 049.
- [7] Y. Kawamura, Prog. Theor. Phys. 105 (2001) 999; G. Altarelli and F. Feruglio, Phys. Lett. **B511** (2001) 257; L.J. Hall and Y. Nomura, Phys. Rev. **D64** (2001) 055003; A. Hebecker and J. March-Russell, Nucl. Phys. B625 (2002) 128.
- [8] N. Haba, A. Watanabe and K. Yoshioka, Phys. Rev. Lett. 97 (2006) 041601.
- [9] S.R. Coleman and E. Weinberg, Phys. Rev. D7 (1973) 1888.
- [10] L.J. Dixon, J.A. Harvey, C. Vafa and E. Witten, Nucl. Phys. B261 (1985) 678; ibid. B274 (1986) 285; Y. Katsuki, Y. Kawamura, T. Kobayashi, N. Ohtsubo, Y. Ono and K. Tanioka, *ibid.* **B341** (1990) 611; T. Kobayashi and N. Ohtsubo, Int. J. Mod. Phys. A9 (1994) 87.
- [11] S. Pakvasa and H. Sugawara, Phys. Lett. B73 (1978) 61; H. Harari, H. Haut and J. Weyers, Phys. Lett. B78 (1978) 459; N.G. Deshpande, M. Gupta and P.B. Pal, Phys. Rev. D45 (1992) 953; L.J. Hall and H. Murayama, Phys. Rev. Lett. 75 (1995) 3985; M. Fukugita, M. Tanimoto and T. Yanagida, Phys. Rev. D57 (1998) 4429; R.N. Mohapatra and S. Nussinov, Phys. Lett. B441 (1998) 299; Y. Koide, Phys. Rev. D60 (1999) 077301; R. Dermisek and S. Raby, Phys. Rev. D62 (2000) 015007; J. Kubo, A. Mondragon, M. Mondragon and E. Rodriguez-Jauregui, *Prog. Theor. Phys.* **109** (2003) 795; T. Kobayashi, J. Kubo and H. Terao, Phys. Lett. **B568** (2003) 83; S.L. Chen, M. Frigerio and E. Ma, Phys. Rev. $D70$ (2004) 073008; N. Haba and K. Yoshioka, Nucl. Phys. B739 (2006) 254; C.Y. Chen and L. Wolfenstein, Phys. Rev. D77 (2008) 093009.
- [12] I. Antoniadis and K. Benakli, Phys. Lett. B326 (1994) 69; M. Bando, T. Kugo, T. Noguchi and K. Yoshioka, Phys. Rev. Lett. 83 (1999) 3601; T. Kugo and K. Yoshioka, Nucl. Phys. B594 (2001) 301; J. Hisano and N. Okada, Phys. Rev. D61 (2000)

106003; S. Cullen, M. Perelstein and M.E. Peskin, Phys. Rev. D62 (2000) 055012; A. De Rujula, A. Donini, M.B. Gavela and S. Rigolin, Phys. Lett. B482 (2000) 195.

- [13] K.S. Babu, E. Ma and J.W.F. Valle, Phys. Lett. B552 (2003) 207; M. Hirsch, J.C. Romao, S. Skadhauge, J.W.F. Valle and A. Villanova del Moral, Phys. Rev. D69 (2004) 093006; A. Zee, Phys. Lett. B630 (2005) 58; X.G. He, Y.Y. Keum and R.R. Volkas, JHEP 0604 (2006) 039; E. Ma, H. Sawanaka and M. Tanimoto, Phys. Lett. $B641$ (2006) 301; L. Lavoura and H. Kuhbock, Mod. Phys. Lett. $A22$ (2007) 181; B. Adhikary, B. Brahmachari, A. Ghosal, E. Ma and M. K. Parida, Phys. Lett. B638 (2006) 345; F. Bazzocchi, S. Morisi, M. Picariello and E. Torrente-Lujan, [arXiv:0802.1693.](http://arxiv.org/abs/0802.1693)
- [14] Y. Kawamura, T. Kinami and T. Miura, [arXiv:0808.2333.](http://arxiv.org/abs/0808.2333)
- [15] C.D. Froggatt and H.B. Nielsen, Nucl. Phys. B147 (1979) 277.
- [16] N. Arkani-Hamed and M. Schmaltz, Phys. Rev. D61 (2000) 033005; K. Yoshioka, Mod. Phys. Lett. A15 (2000) 29; L.J. Hall, J. March-Russell, T. Okui and D. Tucker-Smith, JHEP 0409 (2004) 026; M. Bando, T. Kobayashi, T. Noguchi and K. Yoshioka, Phys. Lett. B480 (2000) 187; Phys. Rev. D63 (2001) 113017; S.J. Huber and Q. Shafi, Phys. Lett. B498 (2001) 256.
- [17] T. Kobayashi, S. Raby and R.J. Zhang, Nucl. Phys. B704 (2005) 3; G. Altarelli, F. Feruglio and Y. Lin, Nucl. Phys. B775 (2007) 31; T. Kobayashi, H.P. Nilles, F. Ploger, S. Raby and M. Ratz, Nucl. Phys. B768 (2007) 135.