INCOMPRESSIBILITY AND LEAST-AREA SURFACES

SIDDHARTHA GADGIL

ABSTRACT. We show that if F is a smooth, closed, orientable surface embedded in a closed, orientable 3-manifold M such that for each Riemannian metric g on M, F is isotopic to a least-area surface F(g), then F is incompressible.

1. INTRODUCTION

We assume throughout that all manifolds (and surfaces) we consider are orientable. Let M be a closed, smooth, 3-manifold and let F be a smoothly embedded surface in M.

For a Riemannian metric g on M, we can seek to minimise the area of embedded surfaces in the homotopy class of F. Here, we say two embedded surfaces F and F' in M are homotopic if there is a homeomorphism $\varphi : F \to F'$, so that if $i_F : F \to M$ and $i_{F'} : F' \to M$ denote the inclusion maps, then i_F is homotopic to $i'_F \circ \varphi : F \to M$. We consider the functional

 $A(q, F) = \inf\{Area_q(F') : F' \text{ embedded surface homotopic to } F\}$

Definition 1.1. A surface F is said to be *least area* with respect to the metric g if $Area_g(F) = A(g, F)$

We recall the concept of *incompressibility* of the surface F.

Definition 1.2. A closed, smoothly embedded surface $F \subset M$ in a 3-manifold M is said to be incompressible if the following conditions hold:

- If $D \subset M$ is a smoothly embedded 2-disc with $\partial D \subset F$ and $int(D) \cap F = \phi$, then there is a disc $E \subset F$ with $\partial E = \partial D$.
- If F is a 2-sphere, then F does not bound a 3-ball.

A disc $D \subset M$ such that $\partial D \subset F$ and $int(D) \cap F = \phi$ so that D is transversal to F is called a *compressing disc* if there is no disc $E \subset F$ with $\partial E = \partial D$.

If F is *incompressible* and M is *irreducible*, then a fundamental result ([11], see also [4]) is that for each metric g, the above infimum is attained for some smooth, embedded surface F(g), i.e., there is a least area surface. Recall that M is said to be irreducible if every embedded 2-sphere $S \subset M$ bounds a 3-ball in M. Least area surfaces enjoys several very useful properties [2] – for instance, leading to the equivariant Dehn's lemma of Meeks and Yau [12].

We show here that, conversely, the property of having least area representatives for each Riemannian metric characterises incompressibility. Fix henceforth a closed, smooth, orientable surface F embedded in a closed, orientable, 3-manifold M.

Date: October 31, 2018.

¹⁹⁹¹ Mathematics Subject Classification. Primary 57N10; Secondary 53A10.

SIDDHARTHA GADGIL

Theorem 1.3. Suppose for each Riemannian metric g on M, there is a smooth, embedded surface F(g) homotopic to F such that $Area_g(F(g)) = A(g, F)$, then F is incompressible.

Our result gives a geometric characterisation of incompressibility, which may be useful in proving incompressibility for surfaces constructed as limits. In particular, this result was motivated by Tao Li's proof [10] of the Waldhausen conjecture in the non-Haken case, where an incompressible surface was constructed as a limit of strongly irreducible Heegaard splittings.

Henceforth assume, without loss of generality, that F is connected. If F is a 2-sphere, either F is incompressible or F bounds a 3-ball in M. In the first case, there is nothing to prove. In the second case, F is homotopically trivial and hence homotopic to the boundary of any 3-ball in M. By considering the boundaries of arbitrarily small balls, we see that A(g, F) = 0 for any metric g. Thus there is no embedded surface F(g) with $Area_g(F(g)) = A(g, F)$. Thus, we can assume henceforth that F is not a 2-sphere.

Suppose F is not incompressible (and F is not a 2-sphere), then there is a compressing disc D for F. A regular neighbourhood of $F \cup D$ has two boundary components, one of which is parallel to F. Denote the other by F'. We call F' the result of compressing F along D. Observe that F is obtained from F' by adding a 1-handle. If F is a 2-sphere that bounds a 3-ball, we declare the empty set to be the result of compressing F.

Given any surface F, we can inductively define a sequence of compressions. Namely, if F is not incompressible, then we compress F along some compressing disc D to get F'. We repeat this process for each component of F' which is not compressible. As the maximum of the genus of the components of the surface F'obtained by compressing F is less than the genus of F, this process terminates after finitely many steps. The result is a (possibly empty) surface \hat{F} , each component of which is incompressible.

Suppose the result of the compressions is empty, then F is homotopic to the boundary of a handlebody, i.e., the boundary of a regular neighbourhood of a graph $\Gamma \subset M$. By considering arbitrarily small neighbourhoods of Γ , we see that for any metric g, A(g, F) = 0. Thus there is no embedded surface F(g) with $Area_g(F(g)) = A(g, F)$, i.e., the hypothesis of the theorem cannot be satisfied.

Thus, we can, and do, assume henceforth that F is not homotopic to the boundary of a handlebody and \hat{F} is not empty. Then F is obtained from \hat{F} by addition of 1-handles. Given $\epsilon > 0$, the 1-handles can be attached to \hat{F} so that the area of the resulting surface, which is homotopic to F, is at most $Area_g(\hat{F}) + \epsilon$. It follows that

$$A(g, F) \le Area_q(\hat{F})$$

We construct a metric g which is a warped product in a neighbourhood $N(\hat{F})$ of \hat{F} so that any least-area surface not contained in $N(\hat{F})$ has area greater than the area of \hat{F} . Thus, if a least-area surface homotopic to F exists, it must be contained in $N(\hat{F})$. The structure of the metric on $N(\hat{F})$ together with some topological arguments show that this cannot happen unless $F = \hat{F}$, i.e., F is incompressible.

2. Construction of the metric

In this section we construct the desired metric for which F has no least-area representative unless $F = \hat{F}$.

As \hat{F} and M are orientable, a regular neighbourhood $N(\hat{F})$ of \hat{F} is a product. We shall identify this with $\hat{F} \times [-T, T]$, with T to be specified later. We shall also consider the regular neighbourhood $n(\hat{F}) = \hat{F} \times [-1, 1] \subset N(\hat{F})$.

Choose and fix a metric of constant curvature 1, 0 or -1 on each component of \hat{F} and denote this g_0 . Let the area of \hat{F} with respect to g_0 be A_0 .

We shall use the *monotonicity lemma* of Geometric measure theory. We state this below in the form we need. For a stronger result in the Riemannian case, see [3].

Lemma 2.1 (Monotonicity lemma). There exist constants $\epsilon > 0$ and R > 0 such that if g is a Riemannian metric on M and x is a point so that the sectional curvature of g on the ball $B_g(x, R)$ of radius R around x (with respect to g) has sectional curvature satisfying $|K| \leq \epsilon$ and F is a least-area surface with $x \in F$, then $Area_g(F \cap B(x, R)) > A_0$.

We shall construct the desired metric in the following lemma.

Lemma 2.2. There is a Riemannian metric g on M satisfying the following properties.

- On $N(\hat{F})$, g is of the form $g = f(t)g_0 \oplus dt^2$, with f a smooth function with f(0) = 1 and f(t) > 1 for $t \neq 0$.
- For $x \in M int(N(\hat{F}))$, the sectional curvature of g on M satisfies $|K| \leq \epsilon$.
- For $x \in M int(N\hat{F})$, the injectivity radius at x is greater than R.

Proof. Observe that $N(\hat{F}) - int(n(\hat{F}))$ has two components for each component \hat{F}_0 of \hat{F} , each of which can be identified with $\hat{F}_0 \times [0,1]$ with $\hat{F} \times \{1\}$ a component of $\partial N(\hat{F})$ and $\hat{F} \times \{0\}$ a component of $\partial n(\hat{F})$. On each such component consider the product Riemannian metric $g_0 \oplus dt^2$. Extend this smoothly to a metric on the complement $M - int(n(\hat{F}))$ of the interior of $n(\hat{F})$. Rescale the metric by a constant s > 1 to ensure that it has sectional curvature satisfying $|K| \leq \epsilon$ and the injectivity radius at each point outside $int(N(\hat{F}))$ is at least R. We choose the constant s to be greater than 1 even if this is not necessary to ensure the bounds on curvature and the injectivity radius. We denote the rescaled metric, defined on $M - int(n(\hat{F}))$, by g. The restriction of g to each component of N(T) can be identified with the product metric $sg_0 + dt^2$.

Let T = 1 + s. Then there is a natural identification of $N(\hat{F})$ with $\hat{F} \times [-T, T]$, with $n(\hat{F})$ identified with $\hat{F} \times [-1, 1]$ and with the restriction of the metric g to $N(\hat{F}) - int(n(\hat{F}))$ given by $sg_0 \oplus dt^2$.

We extend the constant function f(t) = s on $[-T, -1] \cup [1, T]$ to a smooth function on [-T, T] with f(0) = 1 and f(t) > 1 if $t \neq 0$. The Riemannian metric on the complement of $n(\hat{F})$ extends smoothly to one given by $g = f(t)g_0 \oplus dt^2$ on $N(\hat{F})$. This satisfies all the conditions of the lemma.

Note that by construction $\hat{F} \times \{0\}$ is isometric to \hat{F} with the metric g_0 . Further the projection map $p: F \times [-T, T] \to F$ is (weakly) distance decreasing, and strictly distance decreasing outside $\hat{F} \times \{0\}$.

SIDDHARTHA GADGIL

3. Proof of incompressibility

Suppose now that there is a surface F(g) homotopic to F with area A(g, F).

Lemma 3.1. We have $F(g) \subset \hat{F}_0 \times (-T,T)$ for some component \hat{F}_0 of \hat{F} .

Proof. If there is a point $x \in F(g) - \hat{F} \times (-T, T)$, the monotonicity lemma applied to $F(g) \cap B(x, R)$ shows that the area of F(g) is greater than A_0 , a contradiction. Further, as F(g) is connected, for some component \hat{F}_0 of \hat{F} , $F(g) \subset \hat{F}_0 \times [-T, T]$. \Box

To simplify notation, we henceforth denote the surface F(g) by F. We consider the restriction of the projection map from $F_0 \times [-T,T]$ to F, which we denote, by abuse of notation, by $p: F \to \hat{F}_0$. We have seen that this is strictly distance decreasing unless $F = \hat{F}_0$.

We have two cases, depending on whether the embedded surface $F \subset \hat{F}_0 \times (-T,T)$ separates the boundary components of $\hat{F}_0 \times [-T,T]$.

In case F does not separate the boundary components of $\hat{F}_0 \times [-T, T]$, there is a curve γ joining the boundary components disjoint from F. By considering cup products, it follows that $p: F \to \hat{F}_0$ has degree zero. On the other hand, if F does separate the boundary components of $\hat{F}_0 \times [-T, T]$, as F is connected there is a curve γ joining the boundary components intersecting F transversely in one point. It follows that we can choose an orientation on F so that $p: F \to \hat{F}_0$ has degree one.

Recall that F is a connected, orientable surface that is not a 2-sphere. As $\hat{F}_0 \times (-T, T)$ deformation retracts to \hat{F}_0 , the homotopy class of the inclusion map is determined by the homotopy class of p. If \hat{F}_0 is not a 2-sphere, then the homotopy class of p is determined by the induced map on fundamental groups. If \hat{F}_0 is a 2-sphere, then the homotopy class of p is determined by the degree of p.

We show first that the case where F does not separate the boundary components of $\hat{F}_0 \times [-T, T]$ cannot occur.

Lemma 3.2. The surface F must separate the boundary components of $\hat{F}_0 \times [-T,T]$.

Proof. We have seen that $p: F \to \hat{F}_0$ has degree zero. Suppose first that \hat{F}_0 is not a 2-sphere. By a theorem of Hopf and Knesser [5][6][8][9], it follows that p is homotopic to a map whose image does not contain some point $p \in F_0$. It follows that $G = p_*(\pi_1(F))$ (which is finitely-generated) is conjugate to the subgroup of a free group and hence is a finitely generated free group. The projection p: $\hat{F}_0 \times [-T,T] \to \hat{F}_0$ induces an isomorphism of fundamental groups. This gives an identification of G with the image of $\pi_1(F)$ under the homomorphism induced by the inclusion $i: F \to \hat{F}_0 \times [-T,T]$.

By a theorem of Jaco [7], there is a finite graph Γ and $\pi_1(\Gamma)$ isomorphic to G and a map $f: F \to \Gamma$ so that the mapping cylinder M(f) of f is a handlebody. Moreover, we have an identification of $\pi_1(\Gamma)$ with G with respect to which the homomorphism from $\pi_1(F)$ to $G \subset \pi_1(\hat{F}_0 \times [-T, T])$ induced by the inclusion corresponds to the map induced by inclusion from $\pi_1(F)$ to $\pi_1(M(f)) = \pi_1(\Gamma)$.

Choose an embedding of Γ in $\hat{F}_0 \times [-T, T]$ with induced map on fundamental groups $\pi_1(\Gamma) \to G \subset \pi_1(\hat{F}_0 \times [-T, T])$ corresponding to the above identification of $\pi_1(\Gamma)$ with G. Then the surface F is homotopic to the boundary of a regular neighbourhood of the image of Γ , which is a handlebody in $\hat{F}_0 \times [-T, T]$. We have seen that in this case the hypothesis of the theorem cannot be satisfied.

Finally consider the case when \hat{F}_0 is a 2-sphere. As p has degree zero, p is homotopic to a constant map. Hence the inclusion map $i: F \to \hat{F}_0 \times (-T, T)$ is also homotopic to a constant map. It follows that F is homotopic to the boundary of a handlebody. As above, the hypothesis of the theorem cannot be satisfied in this case.

It thus suffices to consider the case when F does separate the boundary components of $\hat{F}_0 \times [-T, T]$.

Lemma 3.3. Suppose F does separate the boundary components of $\hat{F}_0 \times [-T, T]$, then $F = \hat{F}_0$.

Proof. We use a theorem of Edmonds [1] regarding degree-one maps $\varphi : F \to F'$ between closed surfaces. Namely, there is a map ψ homotopic to φ , a compact, connected subsurface Σ in F and a disc $D \subset F'$ such that $\psi(\Sigma) \subset D$ and ψ maps $F - int(\Sigma)$ homeomorphically onto F' - int(D). We can regard F as obtained from F' by attaching 1-handles to D. Further, all the cores and co-cores of these 1-handles are mapped to homotopically trivial curves by ψ .

It follows that an embedded surface homotopic to F is obtained from \hat{F}_0 by adding 1-handles (corresponding to those required to obtain F from \hat{F}_0). Thus, if A_1 is the area of $\hat{F}_0 \times \{0\}$, then $A(g, F) \leq A_1$, hence $Area_g(F) \leq A_1$.

Let $d\omega$ and dA denote the area forms on $\hat{F}_0 = \hat{F}_0 \times \{0\}$ and F respectively. Then for a smooth function h on F, $p^*(d\omega) = hdA$. As the projection map is distance-decreasing, $h(p) \leq 1$ for all p, with equality at all points only in the case where $F = \hat{F}_0$.

Observe that $A_1 = \int_{\hat{F}_0} d\omega = \int_F p^*(d\omega) = \int_F h dA$, where the second equality holds as p has degree one. Further, if $F \neq \hat{F}_0$, then $\int_F h dA < \int_F dA = Area(F)$. Thus, $A_1 < Area_q(F)$, a contradiction.

It follows that $F = \hat{F}_0$.

Thus, the surface F must be homotopic to \hat{F}_0 , which is incompressible. Recall that we can assume that F is not a 2-sphere. It follows that F is incompressible by the characterisation of incompressible surfaces as those for which the induced map on fundamental groups is injective (if F is not the 2-sphere). This contradicts our assumption that F is compressible, completing the proof of the theorem. \Box

Acknowledgements. I thank the referee for helpful comments and for pointing out that the hypothesis of irreducibility in an earlier version was unnecessary. I thank Harish Seshadri for helpful discussions.

References

- A. Edmonds, Deformation of maps to branched coverings in dimension two. Ann. Math. 110 (1979), 113–125.
- M. H. Freedman, J. Hass, G. P. Scott, Least area incompressible surfaces in 3-manifolds. Invent. Math. 71 (1983), no. 3, 609–642.
- K. R. Frensel, Stable complete surfaces with constant mean curvature. Bol. Soc. Brasil. Mat. (N.S.) 27 (1996), no. 2, 129–144.

SIDDHARTHA GADGIL

- J. Hass, G. P. Scott The existence of least area surfaces in 3-manifolds. Trans. Amer. Math. Soc. 310 (1988), no. 1, 87–114.
- H. Hopf, Zur Topologie der Abbildungen von Mannigfaltigkeiten. Math. Ann. 100, 579–608 (1928)
- H. Hopf, Zur Topologie der Abbildungen von Mannigfaltigkeiten. II. Math. Ann.102, 562–623 (1929)
- 7. W. Jaco, Heegaard splittings and splitting homeomorphisms. Trans. AMS 144 (1970), 365–379.
- 8. H. Kneser, Glttung von Flchenabbildungen. Math. Ann.100, 609-616 (1928)
- H. Kneser, Die kleinste Bedeckungszahl innerhalb einer Klasse von Flchenabbildungen. Math. Ann.103, 347–358 (1930)
- T. Li, Heegaard surfaces and measured laminations. II. Non-Haken 3-manifolds. J. Amer. Math. Soc. 19 (2006), no. 3, 625–657
- 11. W. H. Meeks III, S. T. Yau, Topology of three-dimensional manifolds and the embedding problems in minimal surface theory. Ann. of Math. (2) 112 (1980), no. 3, 441–484.
- W. H. Meeks III, S. T. Yau, The equivariant Dehn's lemma and loop theorem. Comment. Math. Helv. 56 (1981), no. 2, 225–239.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BANGALORE 560003, INDIA *E-mail address*: gadgil@math.iisc.ernet.in

 $\mathbf{6}$