

Magnetism, Russell-FeS 'bubbles', and a Quantum Coherent Ancestor

Gargi Mitra–Delmotte¹ and A.N. Mitra²

Abstract

The alkaline seepage site hydrothermal mound scenario of Russell et al is expanded to include rock-magnetism for enabling self-assembly of super-paramagnetic greigite clusters within FeS gel-membranes via thermomagnetic convection. Suggested nucleation of membranes by framboidal sacs (Russell et al 1989); possibility of nano-ramboids (Sawlowicz 1983); observed nested icosahedral forms of framboidal greigite; internal structure formation in non-ideal ferrofluids; field-assisted nano-crystal assembly; and convection aided phyllotactic patterns naturally lead to the idea of membrane-embedded, dynamically ordered ramboid-like magnetic assemblies. A predecessor scenario of molecular motors; pre-RNA world a la Cairns-Smith; and optical polarity follow logically from the plausibility of thermal-gradient driven propulsion of tiny clusters through cluster layers. Further possibilities via magnetism – primordial multicellularity to adaptive learning in nested structures – uncover a role for magnetism in the origin of life. The potential of ferromagnetism plus possible ferroelectricity in multi-faceted iron-sulphide phases are explored for enabling quantum coherent searches.

Key words: nested dynamic-assembly; pre-RNA world; phyllotaxis ; ferromagnetic-ferroelectric ; quantum search; FeS gel-membranes

¹ Corresponding author, present address: 39 Cite de l'Océan, Montgaillard, St.Denis 97400, REUNION; e.mail: gargijj@orange.fr ; Tel. and Fax. no.: 00-262-262307972

² Formerly INSA Einstein Professor, Department of Physics, Delhi University; address : 244 Tagore Park, Delhi -110009. INDIA ; e.mail: ganmitra@nde.vsnl.net.in

1 Introduction and background

Iron-sulphide (FeS) clusters in early evolved enzymes (and across species in a range of crucial roles, e.g. catalytic, electron transfer, structural), have been linked with exhalates on the Hadean ocean floor, based on the close resemblance of these clusters with greigite (Fe_5NiS_8) (Russell et al 1994; Huber and Wachtershauser 1997). Russell et al (1994) envisage Life as having emerged in moderate temperature hydrothermal systems, such as mild alkaline seepage springs. Water percolating down through cracks in the hot ocean crusts reacted exothermically with ferrous iron minerals, and returned in convective updrafts infused with H_2 , NH_3 , HCOO^- , HS^- , CH_3^- ; this fluid ($\text{pH} \sim 10 \leq 120^\circ \text{C}$) exhaled into CO_2 , Fe^{2+} bearing ocean waters ($\text{pH} \sim 5.5 \leq 20^\circ \text{C}$) (Russell and Arndt 2005). The interface, comprising (mainly) precipitating barriers of FeS gel membranes controlled this meeting, as they enclosed bubbles entrapping the alkaline exhalate : an aggregate growing by hydrodynamic inflation. At the growing front of the mound, the production of daughter bubbles by budding, was sustained by a constant supply of newly precipitated FeS. Note that fluids in alkaline hydrothermal environments contain very little hydrogen sulphide. So the entry of sulphide, likely to have been carried in alkaline solution on occasions where the solution met sulphide at depth (Russell and Hall, in press), was controlled; this was perhaps important for a gradual build-up of scale-free clusters leading to the envisaged gel-environment. (As pointed out by Sawlowicz (see below) colloids often form more readily in dilute solutions – suspension as a sol– than in concentrated ones where heavy precipitates are likely to form). The forced entry of buoyant seeps may have led to chimney-like protrusions. Sure enough, membrane structures with remarkable similarities to those in retrieved Irish orebodies were generated in experimental simulations of the mound conditions (Russell and Hall, 1997). Like cells, these mini FeS compartments protected and concentrated the spectrum of energy-rich molecules (Rickard and Luther 2007), borne out by harnessing important gradients across the mound (a true far-from equilibrium system, driven by energy released from geodynamic sources) : redox, pH and thermal gradients for electron transfers, primitive metabolism, and directed diffusion, respectively.

1.1 Greigite formation in FeS membranes

Fig.1 shows laboratory simulated FeS compartments; the chambers and walls are ~ 20 and 5μ , respectively. The amorphous precipitate turned out to be weakly magnetic and acid soluble (Russell et al 1989; Russell 1986). According to Russell et al. (2005), the permeable membranes likely comprise (ferredoxin-like) greigite and mackinawite, and whose metal and sulphide layers work for and against e^- conduction, respectively. An insight into this calls for a brief

outline of iron sulphide transformations under wet and moderate temperature conditions. Amorphous mackinawite is the first FeS phase formed from aqueous S(-II) and Fe(II) at ambient temperatures, apparently via two competing pathways governing the relative proportions of the two end-member phase mixture. The long-range ordered phase with bigger crystalline domain size and more compact lattice, increases at the cost of sheet-like precipitated aqueous FeS clusters (Wolthers et al. 2003). Rickard and Morse (2005) have reviewed the substantial presence of aqueous FeS clusters in water environments. They suggested the enhanced stability of some stoichiometries—stable magic number clusters— from among the apparent continuum of stoichiometries of aqueous FeS clusters. This ranges from Fe_2S_2 to $\text{Fe}_{150}\text{S}_{150}$, where the first condensed phase (FeS_m , mackinawite) appears, with a size and volume of 2 nm and 10 nm^3 , respectively. The accompanying density increase over the smallest aqueous FeS cluster is $\geq 10^6$. The structure of hydrated clusters is believed to determine that of the first condensed phase (formed by H_2O expulsion). X-ray diffraction of the first nano-precipitate shows a lattice expanded tetragonal mackinawite structure. That the data fit well with other independent estimates is ascribed to the plate-like form of FeS_m . The quick transformation of disordered mackinawite to the ordered form is followed by solid state transformation to the more stable but structurally congruent greigite, with a *12 percent decrease in volume*, involving a rearrangement of Fe atoms in a close-packed, cubic array of S atoms. (In bio-mineralized crystals too, selected-area electron diffraction patterns show the two minerals oriented relative to one another with $(011)_m // (222)_g$ and $[110]_m // [100]_g$, embedded in a continuous sulphur-substructure across the interfaces (Posfai et al. 1998)). Further, trace amounts of aldehydes are believed to bind to the mackinawite surface, initiating Fe(II) oxidation; they also prevent the formation of dissolved FeS aqueous clusters, crucial for pyrite formation, thus assisting in greigite formation at the cost of pyrite (perhaps as in bacteria) (Rickard et al. 2001). The possibility of organic ligands stabilizing aqueous (greigite-like) Fe(III)-bearing sulphide clusters, leaves open the question of mackinawite-greigite transformations being limited only to the solid-state (Rickard and Luther 2007). Importantly, FeS membranes formed in the laboratory show a 20-40 fold increased durability on adding a mixture of plausible abiogenic organics. Diffusion controlled reactions would slow down with thickening of aging membranes (Russell et al. 1994).

1.2 *The FeS Gel environment and framboids*

The possibility of gel-sol phase transitions underlying the dichotomy of the cell led to the suggestion of Life as having emerged in a hydrogel environment (Trevors and Pollack 2005). Now as noted by Russell et al. (1994), citing Kopelman (1989), gels lie between liquid and solid states with self-similar




Fig. 2. Framboids in chimneys. (a) Small pyrite vent structure: Reflected ore microscopy of transverse section shows a central area of empty black spaces plus (grey) fine framboidal pyrite, and a fine euhedral authigenic rim surrounded by baryte, with minor pyrite; (b) Sheaf system, formed from coalescing rods of anastomosing microcrystalline pyrite. Black areas are empty spaces; central regions are framboidal pyrite with an exterior of crystalline pyrite. (Labelled pictures given by Dr Adrian Boyce are reproduced with his kind permission ; Source : Boyce et al. 1983; Boyce 1990 : Exhalation, sedimentation and sulphur isotope geochemistry of the Silvermines Zn + Pb + Ba deposits, County Tipperary, Ireland ; Boyce, Unpublished Ph.D. thesis, University of Strathclyde, Glasgow).

The formation of microframboids was considered on the basis of (1) occurrence of granule-sized framboids building standard-sized ones; (2) particle aggregates $\leq 0.1\mu m$ in bacteria; (3) hollow microgranules like annular framboids; (4) the transformation of spherules into euhedral crystals, coexisting with corresponding framboids of all sizes; (5) similarities between grain forms building framboids and polyframboids. And since spheroidal microframboids are formed of equant nanocrystals, Sawlowicz (1993, 2000) suggested the formation of nano-framboids, comprising microcluster aggregations ($\sim 100atoms$), by analogy with the 3-scale framboidal hierarchy. From these observations, Sawlowicz proposed a formation mechanism by which the original super-saturated gel-droplet would undergo subsequent divisions into immiscible smaller droplets, where further subdivisions would depend on its initial size, plus concentration of iron, stabilization of gel (eg. by organic matter) and effect on its viscosity, activity of sulphur species, etc. (Note the key role of the colloid-gel phase leading to the fractal forms). The narrow distribution of sizes and uniform growth of thousands of crystals in framboids within a short time interval was attributed to a regulated balance between rates of nucleation and of crystal growth (par-

ticulation mechanism), as in the La Mer and Dinegar model (1950). Increasing solution concentration provokes condensation of small polyanions leading to macromolecules that by aggregation and further condensation lead to sol particles. Significantly, the formation of perfectly spherical clusters was found to be dependent on the extent of motional freedom of primary particles. Sawlowicz further observed (see also Wang and Morse (1996)) that with increasing supersaturation of the parent solution pyrite crystal morphology changed from cube, to octahedron, to spherule. Also the D/d ratio (D and d being diameter of framboid and constituent microcrystal, respectively) varies inversely with the size of primary droplets and their iron concentration (see Farrand (1970) and Kizilshtein and Minaeva (1972)), between 5:1 and 30:1. Furthermore, the nucleation of a supersaturated solution by the first-formed crystal triggers the separation of many crystals of the same size. This liquid-solid-like phase transition is dependent on packing considerations of hard-sphere-like microcrystals, whose ordering is an outcome of the interplay of close-packing and repulsive forces (see Oxtoby (1990) and Taylor (1982)).

A characteristic pattern of icosahedral frambooids (see Appendix I) - octahedral microcrystals, large D/d ratio - has been attributed to a high initial nucleation rate and low growth rate of microcrystals (Ohfuji and Akai 2002; Ohfuji and Rickard 2005). According to Sawlowicz, the interplay of surface-minimizing forces (e.g. surface-tension, limited space, magnetic, etc.) with repulsive interactions (e.g. steric) lead to close-packed frambooids (tending to polyhedrons) – a ramification of anastrophic supramolecular organization with its far-from-equilibrium conditions (see Nicolis and Prigogine (1977); Kauffman (1993)). Sure enough, the framboid morphology is strongly reminiscent of the ubiquitous phenomena of Phyllotaxis, from subnano to cosmological scales (see Appendix II), and regarded as a dynamical self-organizing process. Very significantly, the role of *convection* in generating phyllotactic patterns has been pointed out (Selvam 1998; Lofthouse (arXiv:physics/0411169v1)), showing the importance of directed forces on assembling elements (that enhance their motional energy). These patterns are produced when the sequential accretion/deposition or appearance/growth of elements, is governed by an energy-minimized optimization of the main opposing forces : largest available space vs repulsive interactions (see Appendix II).

1.4 Magnetic interactions

Magnetic interactions turned out to have an overwhelming influence when Wilkin and Barnes (1997) included them in the standard DLVO treatment for interacting colloidal particles that considers attractive van der Waals and double-layer repulsive interactions, for modeling framboidal pyrite formation. This is based on the alignment of precursor greigite, under the influence of the

weak geo-magnetic field that would help overcome the thermal energy of particles above a critical size. Ferrimagnetic greigite has a maximum or saturation magnetization value M_{sat} at 298K ranging between 110 and 130 kA/m . On the basis of microscopic observations by Hoffmann of natural greigite crystals, $< \mu$ meter - sized greigite can be roughly taken as single-domain particles. Assuming a spherical geometry, the critical grain diameter of constituent crystallites comprising the framboid interior $d_c = 2a$, where $a > 1$, is given by

$$d_c = (6k_B T / \mu_0 \pi M_{sat} |H|)^{1/3} \quad (1.1)$$

Here k_B is Boltzmann's constant and μ_0 the permeability of vacuum. When aligned parallel to weak fields (geomagnetic field ~ 70 microT), $d_c = 0.1$ micrometer. Although framboids can form in varied environments (see Sawlowicz), this magnetic greigite-precursor based mechanism is seen as operative only upto temperatures of 200 ° C (Wilkin and Barnes, 1997), eg. sediments in natural waters.

1.5 Summary of background and outline of paper

To recapitulate, we studied two essential stages of framboid formation: multiple nuclei followed by growth of each into a crystal. The dynamically ordered patterns in framboids are brought about by an interplay of forces. The observation of scale-free forms and suggestion of nanoframboids (Sawlowicz), implies the possibility of such patterns at even lower scales of hydrated clusters ordered in nuclei before the stage of crystal growth –this would depend on the availability of *opposing* forces at the scale of clusters plus an effective *directed* force for enabling this interplay. Next, recall the significance of the colloid environment and suggestion that framboids could form in the alkaline seepage-site scenario of Russell et al (1989). Note here that the greigite component of the newly forming colloidal barrier is super-paramagnetic. In this paper, it's resemblance to a ferrofluid, make us consider two properties of ferrofluids: i) super-paramagnetic particles in response to moderate fields and in the presence of thermal gradient can undergo thermomagnetic convection; ii) structure formation due to inter-particle interactions in non-ideal ferrofluids manifest in *dense* phases (a milder phase transition than to the solid-crystalline one). So we shall check for the feasibility of TMC (in the mound), as a directed force that could effectively aid the interplay of opposing forces: induced magnetic dipole interactions plus Van der Waals vs steric plus negative charge on nucleating clusters (under mound conditions). Here the non-ideal ferrofluid nature of the polydisperse, greigite clusters would manifest in attractive inter-particle interactions. Further, as material properties are likely to determine structure formation manifest in dense phases in ferrofluids, we consider the possibility of framboid-like assemblies based on observations of scale-free framboidal forms

of greigite plus Sawlowicz's fractal framboids. Now, structure formation owing to magnetic ordered assemblies of hydrated clusters, would increase the viscosity of the medium and reduce the possibility of TMC. Nevertheless, these assemblies would have been installed (c.f., nucleation of chimneys by Russell et al (1989)) in the barriers, and the opposing gradients of magnetic susceptibility and temperature would continue to fight each other: due to the latter (Baaske et al 2007) clusters would be pushed through the assemblies (probability inversely proportional to size) even as the former force would attract the assemblies and maintain them together in the face of the disruptive forces. This would lead to interesting possibilities, some of which are considered in this paper. Otherwise what can be expected in the absence of magnetic effects? The presence of a thermal gradient alone would lead to heat transfer across the distribution of hydrated clusters and assist their alignment by breaking bonds, permitting motional freedom and re-associations. Recall that elimination of interfacial water and crystallographic-coherent alignment of coalescing clusters (steps towards nucleation and growth) underlie the close matching between structure and bonding in clusters, and their crystalline counterparts (Navrotsky 2004). Now, owing to the aging process : 1) Increased intra-cluster bonds with water loss, would increase molecular size, and approach the solid state (rigid order); 2) Water loss means increased density and loss of gel-like, structured water property, culminating in a rigid network.

Could magnetic interactions have pre-empted this "dull" scenario and driven *life-like dynamics* by holding non-bonded greigite clusters together, in the face of pumped in heat, and harnessed a continuing inflow of heat energy? It's feasibility follows from the demonstration of controlled assemblies of superparamagnetic molecules attuned to external magnetic fields. And since moderate magnetic fields with a strength \sim ten milli-Tesla are required to exert control on such suspensions (Odenbach 2006), it is highly plausible that the thermal gradient across the mound was harnessed by the greigite clusters aided by a *local* magnetic field— allowing a 'bottom-top' assembly at nano-scales, viz. the formation stage of the nascent greigite clusters— leading to dynamical accretion of soft nuclei in the supersaturated aqueous phase. In this paper

1) We examine the chances of a local magnetic field as well as directed motional forces in the Russell mound which would have enabled the installation of dynamical nano-scale assemblies of greigite clusters, thus nucleating the FeS membranes (Sect.2). 2) Directed diffusion of small clusters through intra-cluster layers (I-CLs) of the dynamic assemblies are seen vis-a-vis biophenomena like budding, pre-molecular motors, pre-RNA world, and optical polarity (Sect.3). 3) Relevance of ferromagnetism in biology, especially quantum processing in the context of possibilities for a magnetically ordered Ancestor, reveal a possible role for magnetism in the origin of life (Sect.4). In the light of ferroelectric effects in present day biology (Tuszynski et al 1995; Mavromatos et al 1998), we also briefly check the status of corresponding

ferroelectric effects at the Ancestor level.

2 Controlled assembly tuned to external fields

We check for the feasibility of two ingredients: local moderate magnetic field in the mound (Sect. 2.1) and thermomagnetic convection as a directed force (Sect. 2.2). We check its plausibility in the mound scenario for propelling newly forming colloidal greigite clusters (Sect. 2.3). In Sect. 2.4 we briefly review correlation phenomena in ferrofluids (deviations from ideal monodisperse ferrofluid behaviour), with possible implications for the nucleation stage in framboid formation—the basis, apart from observations, for the proposed framboid-like assemblies in Sect. 2.5.

2.1 Remanent magnetism of mound constituents

Now, subsurface magnetic rocks are known to create sufficiently intense magnetic anomalies (relative to predictions of the global model of the geomagnetic field) used to track their location. As an example consider the rich iron ore province in the Pilbara region of Western Australia with a background ambient magnetic of about $55 \mu T$, where a helicopter survey recorded high anomaly amplitudes of up to $120 \mu T$, indicating the high percentage of iron ore composition (<http://www.foxresources.com.au/documents/318.pdf>). Since field strength decreases rapidly with distance ($\sim r^{-3}$) from the magnetic medium, the corresponding value on the rock surface is expected to be higher by few-fold orders of magnitude, thus overwhelming the contribution of the ambient geomagnetic field. Such high local field strength due to magnetic rocks, justify the magnetism of mound rocks as an important ingredient in the initial conditions leading to Life. Now thermo-remanent magnetization (TRM) of components would result from solidification of molten magmatic ‘compass-like’ elements, freezing in the geo-magnetic field direction, right during primary crust formation itself. For, using laser-heated ancient crystals of feldspar and quartz, Tarduno et al. (2007) estimated that the Earth’s magnetic field was about half as strong 3.2 billion years ago as it is today, implying the formation of a rotating and convecting iron inner core, more than 3 billion years ago. Sediments may also have contributed, e.g. greigite itself (Posfai and Dunin-Borkowski 2006), retaining remanence upto $300\text{-}350^\circ \text{C}$ (Hu et al. 2002). But considering levels of magnetization as offered by possibilities like TRM alone would be limited by the dependence of the initial magnetization upon the then effective geomagnetic field strength. For instance, the present geomagnetic field strength is too weak to explain the high NRM (natural remanent magnetization) to saturation isothermal remanent magnetization (SIRM) ra-

tios of lodestones, natural magnets with magnetic field strengths varying upto 0.1 Tesla (see Magnetism and magnetic surveys, AG Education Services Ltd., www.sta.ie) that eventually led to lightning remnant magnetism as a plausible mechanism (Wasilewski and Kletetschka 1999). Indeed, the most important contribution to the crust was quite possibly thanks to the presence of accreted highly magnetized meteoritic matter, with acquired isothermal remanent magnetism. Recently, Tunyi et al. (2003) have examined the possibility of nebular lightnings as a source of impulse magnetic fields in the accretion of Earth and other planets, that is seen as rendering the gravitational accretion process more efficient, by magnetizing the ferromagnetic dust grains to their saturation levels. Thus the possible presence of ferromagnetic matter (metallic Fe, Ni, Co) due to vestiges of iron and iron-alloy containing meteorite bodies in the primitive Hadean crust, could have even provided a local magnetic field strength of the order of a Tesla. For in contrast to today's picture, conditions in the primitive crust were highly reducing with the redox state depicted at Fe-FeO (Wustite) (Russell et al. 2003; Righter et al. 1997).

2.2 Directed force : thermo-magnetic convection (TMC)

Phyllotactic patterns are believed to be facilitated by convective forces (see Sect. 1.3; Appendix-II), as a search for the energy minimized location for contact would obviously be more effective than by a simple translational motion; the energy optimization is further aided by the disruptive effects of convection on the one hand and accretion on another. The importance of motional freedom for admitting long range order of dynamically organized elements can also be seen in synthesized nano-scale assemblies: Non-directional forces or insufficient mobility during assembly were identified as reasons leading to amorphous or short-range-ordered materials (Redl et al 2003). Now greigite particles ≤ 30 -50nm are superparamagnetic (Hoffmann 1992); negatively charged colloidal greigite clusters (since under mound conditions pH is well above 3 (see Wilkins and Barnes 1997)), closely resemble aqueous based ferrofluids (suspensions of single domain magnetic nanoparticles (~ 10 nm)) with negative charges to stabilize the dispersion. Imposing an external magnetic field on these isotropic media (in zero field) can line up the constituent randomly oriented ferrofluid particles (magnetic domains), acting as tiny magnets. Hence we investigate the feasibility of thermomagnetic convection in this scenario. The magnetization vector M (magnetic moment per unit volume, A/m) of the bulk ferrofluid obeys the Langevin equation, namely,

$$|M| = M_{sat}\phi(\coth\alpha - 1/\alpha); \quad \alpha = (4\pi a^3/3)(\mu_0 M_{sat}|H|/k_B T) \quad (2.1)$$

where ϕ is the volume fraction of magnetic particles in the ferrofluid; M_{sat} is the bulk saturation magnetization of the particles; α is the Langevin pa-

parameter; a is the nanoparticle radius, H is the imposed magnetic field, k_B the Boltzmann constant, and T the absolute temperature. The net magnetization of ferrofluids in an external magnetic (H) field that can help in convective heat transfer, depends upon its temperature and H. The Kelvin body force (KBF) per unit volume, viz., $\mathbf{f} = (\mathbf{M} \cdot \nabla) \mathbf{B}$, is the force that a magnetic fluid experiences in a spatially nonuniform magnetic field, propelling it towards stronger H-fields. Its variation with a temperature-gradient results in a non uniform magnetic force acting upon the ferrofluid, giving rise to TMC (Finlayson 1970), which is a function of particle size, shape and structure (Ganguly and Puri 2006). Ganguly et al (2004) have studied this form of convection using a square cavity (enclosure height h) with a temperature gradient ΔT due to the left-hand side vertical wall being maintained at a temperature of T_h and the other vertical wall being at an isothermal heat sink at T_c . The upper and lower walls were taken as adiabatic. The cavity was considered as infinity in the third dimension such that the flow developing inside it was two-dimensional – an effective two-dimensional external magnetic field was provided by a line dipole placed adjacent to the lower adiabatic wall halfway along the channel length at depth $h/4$ below its inner surface. The latter was used to approximate the field produced by a permanent magnet /electromagnet, held parallel to the channel wall. In Appendix IV, we have carried out a simple derivation of the average range of magnetic field over such a cavity of height h due to a symmetrically placed line dipole, on the basis of its $1/r^2$ dependence, leading to the following result (see Eq A. 2) :

$$h^2 < 1/r^2 > = \pi \ln 7 = 6.113 \quad (2.2)$$

Hence to calculate the average H - field strength for a given cavity height, it is necessary to replace the factor $< m/r^2 >$ by $6.113 \times m/h^2$.

In a narrow temperature range, the magnetization vector \mathbf{M} is approximately linear in T and H , when measured from its equilibrium value M^* corresponding to $T = T^*$ and $H = H^*$ respectively. Now since $M = \chi_m H$, where only χ_m varies slowly with T , substitution of this variation in the above formula for \mathbf{f} , and some routine simplification, leads finally to a result of the form $KBF = KBF_I + KBF_{II}$ where

$$\begin{aligned} KBF_I &= \frac{1}{2} \mu_0 (\chi_m + \chi_m^2) \nabla H^2; \\ KBF_2 &= -\frac{\mu_0 \chi_m^2 \beta_\rho}{1 + \beta_\rho (T - T^*)} \mathbf{H} (\mathbf{H} \cdot \nabla T) \end{aligned} \quad (2.3)$$

In the absence of KBF_2 as in isothermal system, $KBF = KBF_I$ creates a static pressure field symmetric about the line dipole (here rock source of magnetism). In a non isothermal system the temperature dependence of the

magnetic susceptibility alters the symmetry of the KBF. A ripple in temperature dependent KBF_2 term leads to imbalance in the main background term KBF_1 (magnetic pressure term even if temperature independent) that leads to thermomagnetic convection in a ferrofluid contained in such a cavity, under the influence of an H-field of a line-dipole. This has been correlated (Ganguly et al. 2004; Mukhopadhyay et al. 2005) with a dimensionless magnetic Rayleigh number Ra_m :

$$Ra_m = \mu_0 \rho_0 \beta_\rho \chi_{m,0} (Pr) m^2 \Delta T / [\eta^2 h^2]; \quad \beta_\rho = -\rho^{-1} \delta \rho / \delta t \quad (2.4)$$

Here m the magnetic moment (in units of Am) per unit length of line dipole; ΔT the difference in temperature between left (hot) and right (cold) walls of cavity ; and h is the enclosure height and the rest of the factors are defined and their corresponding values listed in the Table. (Here effects of magnetocrystalline anisotropy, magnetoviscosity and magnetodissipation were neglected). TMC was simulated for the aqueous ferrofluid described in the Table, using Eq. 2.4 (an m/D^2 scaling was also seen; see Mukhopadhyaya et al 2005). Two typical examples are : a) for the ferrofluid enclosed in a cavity height of 4cm with a gradient of 50K across, and a line dipole strength of 2.96 Am (giving $\langle H \rangle \sim 14$ mT ; see Appendix IV), a Ra_m value of 2.14×10^4 was obtained; b) for the same temperature difference but cavity height of 4mm, a line dipole strength of 2.96 Am ($\langle H \rangle \sim 1.4$ T), an Ra_m value of 2.14×10^6 was obtained. Further, the ratio of KBF_1 and KBF_2 was 3.8×10^2 for both these cases (also for all cases considered in their study, ratios were about 100 fold). Significantly, TMC in the range $Ra_m = 2 \times 10^4 - 2 \times 10^7$ was found to be as effective as the range $10^3 - 10^6$ for the conventional Rayleigh number (buoyant convection).

2.3 Feasibility of TMC in mound scenario

In the absence of data on greigite based ferrofluids, we look at magnetite-based ones for rough estimates. Now the saturation magnetization of magnetite ($M_s = 4.46 \times 10^5$ A/m) is about 3.5 times greater than that of greigite; from this one expects proportionate values for the fluid susceptibility of a corresponding greigite preparation. Next, the volume content of the aqueous dispersions of magnetite (10nm) is usually 2-3% (Rosensweig (1997); Ferrofluid database from German Ferrofluid Community Information Server). In the mound, a slow build up in local concentration of greigite across 1 cm colloidal barrier, upto $\sim 2\%$ (0.05M taking molecular weight of $Fe_3S_4 = 396$), is within conservative limits, if one considers that apart from it's formation there would be inter-particle dipole-dipole interactions plus screening effect due to ionic strength of natural waters (see also Phenrat et al 2007). The rest of the parameters are taken from the Table since in view of the dilute and aqueous nature of

Table 1

Representative aqueous ferrofluid (Ganguly et al (2004); Rosensweig (1997))

<i>Property (Symbol)</i>	<i>Value</i>
Permeability of vacuum (μ_0)	$4\pi \times 10^{-7} Hm$
Density (ρ_0)	$1180 kgm^{-3}$
Fluid compressibility/Thermal expansion coefficient (β_ρ)	$5.6 \times 10^{-4} K^{-1}$
Thermal conductivity (κ)	$0.59 Wm^{-1} K^{-1}$
Specific heat (C_p)	$4180 J/kg$
Viscosity (η)	$0.007 Nsm^{-2}$
Kinematic viscosity = η/ρ ($\nu_{k.v.}$)	$5.93 \times 10^{-6} m^2/s$
Thermal diffusivity = $\kappa/C_p\rho$ ($\alpha_{t.d.}$)	$1.19 \times 10^{-7} m^2/s$
Prandtl number = $\nu_{k.v.}/\alpha_{t.d.}$ or $C_p\rho/\kappa$ (Pr)	49.6
Fluid susceptibility at reference temperature 300K ($\chi_{m,0}$)	0.1

the suspensions, these values would not change drastically for greigite based suspensions.

The newly forming colloidal barrier containing super-paramagnetic greigite $\sim 1cm$, would have acted as the corresponding ‘cavity’ (as described in the Ganguly et al set up) in the presence of an external magnetic field owing to magnetic rocks in the mound base, it’s boundaries demarcated by the zero magnetic susceptibility of the immediately adjacent aqueous medium lacking the magnetically responsive particles. The temperature gradient across the thickness of this newly forming front (from hot mound interior to cool ocean outwards) corresponds to that in the set up of Ganguly et al (2004) (Sect. 2.2), where it forms an active dimension of the square cavity. Note that the essentially 2D nature of the geometry (third dimension linear in Ganguly et al; circular in the present case), allows this correspondence. As the thickness of the mixing front could be in the range $\sim mm$ to $\sim cm$, we shall consider two cases, that of cavity height of 0.04m and 0.004m, individually. Extrapolating the above results (Sect. 2.2) of case a) to an $\langle H \rangle$ of almost double the value ($\sim 30mT$ which requires $m = 6.246 Am$), keeping the other values same, Ra_m works out to 1.0252×10^5 , a five-fold increase over the Ganguly et al value. Again, a corresponding extrapolation from case b) (using $\langle H \rangle \sim 300mT$) also gives an almost identical value of Ra_m at 1.0265×10^5 . So even with a 3.5 fold reduction in susceptibility, TMC would still be in the efficient range for H-fields of 30mT and 300mT for a thickness of 4cm and 4mm, respectively, of the mixing front. Further, a temperature gradient of about 100K (c.f. 50K

for Ganguly et al) could exist across the interface of the colloidal barrier; this factor would provide an extra two-fold enhancement in the Ra_m values for both these extrapolated cases. An increased temperature gradient of 2-fold, means that the corresponding KBF_1/KBF_2 ratios would be 1.79×10^2 , as the values of KBF_1 and KBF_2 are of the order of $\mu_0\chi_{m,0}m^2/h^5$ and $\mu_0\chi_{m,0}^2\beta_\rho m^2\Delta T/h^5$, respectively. Thus it would still be ~ 100 -fold, like in all cases considered by Ganguly et al (2004). This approach treats the suspension as a single-component (magnetic) fluid. For when regarded as a binary mixture (Ryskin et al 2003), thermal convection sets in at Rayleigh numbers well below the critical number for a magnetic fluid, treated as pure. Here we have ignored temperature variations in H- field as in Ganguly et al. (2004); advective effects due to Lorentz force on ions (Uechi et al 2004; Stone and Goldstein 2004); and boundary effects like Rosenweig instabilities (Rannacher and Engel (2004)).

2.4 Correlations in non-ideal ferrofluids

While magnetoviscous effects may be ignored for non-interacting particles, studies on real ferrofluids reveal marked rheological changes even in moderate fields $\sim 38\text{mT}$, that cannot be explained on the basis of rotational viscosity due to the Brownian relaxation of large sized (hard) particle fraction, alone. Microscopic approaches assume internal structure formation due to interparticle interactions (non-ideal behaviour) leading to dense phases. The structure of soft ('molten') heterogenous aggregates (e.g. chain-like, drop-like) would depend on factors like the strength of the applied field, the nature of the ferrofluid (molecular shape, susceptibility) etc. (Odenbach 2004; Zubarev et al 2005; Zubarev and Iskakova 2004). An increase of chain size beyond a critical length, compactification due to interparticle magnetic interactions, formation of globules as nuclei for new dense phase are seen in the scheme of phase transitions leading to formation of bulk drop-like aggregates. Also, the action of hydrodynamical shear forces are seen to lead to a microemulsion of discrete drops (of a threshold size) of the dense phase at the final stage of phase separation (c.f., microemulsion formation stage of fractal fimbroids (see Sect. 1.3)). Further, Wang and Holm (2003) studied the role of polydispersity and concluded that the fraction of large particles, with larger relative dipole moments in proportion to their volume, would be responsive to weaker fields more easily and therefore dictate magnetization properties, even in case of dilute fluids. Their larger dipole-dipole interaction energy makes them overcome thermal forces more easily, and consequent structures formed influence properties like initial susceptibility. Importantly, Taketomi et al (1991) observed field-induced macrocluster formation in water and paraffin-based ferrofluids but not in an alkyl-naphthalene based one (even at 0.2 T). In contrast, macroclusters formed in the water-based fluid at very low fields and remained even after removing the field. (They also correlated these with ordering phenomena dictated by

surface energy like Nisoli et al (see Appendix II).

These aspects were studied by Li et al (2007) who pointed out that the field-induced formation of aggregates that break up in response to thermal effects upon removal of field, indicate their *dissipative* nature. Thus the total magnetic energy of ferrofluids obtained from an applied field: $W_T = W_M + W_S$; where $W_M = \mu_0 M_L H V$ and $W_S = -T \Delta S$ are the magnetized and the structured energies, respectively, V is the volume of the ferrofluid sample and ΔS is the entropic change due to the microstructure transition of the ferrofluid. The equivalence of W_T (zero interparticle interactions), with the Langevin magnetized energy $W_L = \mu_0 M_L H V$ leads to the correction in the magnetization, in terms of the entropy change. They also obtained an expression for ϕ_H or the particle volume fraction of the field-induced aggregates (particles located in ‘aggregate-space’ on the lines of a compressed gas) by considering its independent relations with the intensity of the applied field (via the Langevin parameter) and the particle volume fraction, plus boundary conditions. The proportionality coefficient γ (compression parameter) that goes to zero in the limit of a well-behaved ferrofluid (obeying Langevin function) turns out to be an intrinsic parameter of ferrofluids. It is related to the aggregation characteristics and appears to be dependent not only on the dipolar coupling constant $\lambda = m^2/4\pi d^3 \mu_0 kT$ but also molecular properties preventing aggregation.

2.5 Framboid-like assemblies : Quasiperiodicity

In the mound scenario, although the external field strength would be constant, the likely polydisperse nature of greigite in the evolving suspension would similarly dictate the rheological properties of the barrier. Further a number of reports demonstrate the self-assembly of super-paramagnetic nano-crystallites, controlled by external magnetic fields ($\sim 0.5T$), composed of different materials of varying sizes : nickel ferrite nanospheres (250nm) (Wang et al 2005); nickel zinc ferrite nanoparticles 30-40nm (Zhang et al 2005); magnetic superlattices using maghemite nanocubes $\sim 10nm$ (Ahniyaz et al 2007), FePt nanoparticles 3-10nm (Sun et al 2000), binary systems Co and Fe_3O_4 nanoparticles of 7.5 and 18.0nm, respectively (Cheon et al 2006). Note that coatings help prevent attractive interactions other than magnetic dipolar ones and in tuning inter-particle contact distances (Lalatonne et al 2004). Clustering patterns can be controlled by an external field and although strong anisotropy of dipolar forces and preference for head-tail arrangement lead to tendency for chain-like clusters, other superstructures are possible (Ahniyaz et al 2007 + ref). Despite a different scenario of an evaporation-mediated self-assembly in most of these studies, one can appreciate that apart from field- strength, the properties of nanocrystallites (geometry, susceptibility, coatings, etc) are important criteria for clustering patterns. In fact, in a study more relevant to

the mound scenario, a magnetic field assisted assembly under hydrothermal conditions has been demonstrated : chain-like structures > 10 magnetite nanoparticles, average size 80nm under hydrothermal conditions, 200 ° C, formed under a 0.18T external magnetic field (Wu et al 2005). Thus in addition to the investigations on internal structure formation in polydisperse ferrofluids (see Sect. 2.4) , concerning soft macro-ordered clusters, these experiments show the control of magnetic fields on the other end of the spectrum (nano-crystallites). Extrapolating from these observations, one can expect that an increase in size of soft-ordered nuclei would result in stronger induced dipolar interactions that would aid their assembly into larger superstructures. Now, as structure formation is related to molecular characteristics, we searched naturally occurring ordered structures of greigite for clues as to the possible nature of the nano-assemblies. This led to the smallest observed form of icosahedral framboidal greigite: about 2.1 μ diameter, (see Appendix-I for details), nested structures of which lead to larger ones (Preisinger and Aslanian 2004). In view of this *natural* propensity for framboid-formation ; the possibility of TMC-aided interplay of opposing forces (repulsive charge (as clusters carry negative charge at pH > 3 in mound conditions) plus steric vs attractive magnetic dipole plus Van der Waal’s interactions) generating phyllotactic patterns; and Sawlowicz’s suggestion of nano-framboids, we propose the formation of magnetic clusters–soft-nuclei as building blocks–ordered in framboid-like assemblies.

This is unlikely to have led to the conventional crystalline phase, where the nucleus surface acts as a template for copying a unit cell via local interactions. Rather, it naturally fits in with the aperiodic, long range order of systems known to form *quasicrystals* (see Appendix-III) whose growth occurs by accretion of preformed clusters in the liquid state by the growing nucleus. A recent simulation by Keys and Glotzer (2007) show that quasicrystal growth is facilitated by structurally persistent atoms (icosahedral clusters in this case) that become kinetically trapped in the supercooled liquid; these are incorporated into the growing solid nucleus in a way that minimizes expensive rearrangements and eases propagation. In icosahedral materials, atomic clusters seen as recurring structural motifs, usually comprise nested polyhedra of several tens of metal atoms; the localized properties of these clusters do contribute towards some physical properties in unusual combinations. Some of these are given in Appendix-III, where we also note their strong analogies in bioprocesses. Recall that a quasicrystalline arrangement of biomolecules is believed to enable quantum coherent processes (Jibu et al 1994). While the importance of order and crystal growth have been invoked earlier to understand a speculated transition from mineral forms of pre-life to life (Cairns-Smith 1985) , we note that a quasiperiodic order is softer, with flexibility in it’s ordering as compared to a rigid crystalline array. Aperiodic order (Schroedinger 1944) comes with extra degrees of freedom, and softer packing (due to defects, vacancies etc.) is better placed to accomodate collective moves, relative to close packed rigid layers (Roth and Gahler 1998); this could enhance the potential for quasiperi-

odicty vs crystalline order for generating ('computing') life-dynamics, as a dynamic order might have been an important pre-requisite leading to life (see also Davies 2003). Besides, the known links between quasiperiodic phases and phyllotactic patterns (see above) might be highly relevant for insights into the observed icosahedral domain organization (reviewed in Ohfuji and Akai 2002, see Appendix I) in closed packed icosahedral greigite frambooids, with a high degree of phyllotactic frambooidal texture. These reveal a predominance of edge to edge contacts (Roberts and Turner 1993; Ohfuji and Akai 2002) with a lattice-like space filling arrangement (solely) in the connecting edges of the five trapezoidal domains; this puts a limit to possible pathways for conduction in these structures.

The possibility of an icosahedral shape of the frambooid-like assemblies arises out of packing considerations on a spherical surface that brings in geometrical frustrations. And geometric constraints do not rule out long-range magnetic order (see studies on icosahedral magnetic quasicrystals (Lifshitz 1998)). The scale-free icosahedral forms of frambooidal greigite could be understood from a mathematical study where tessellation of spheres (number N ; radius a) packed on the surface of a large sphere (radius R), showed that energy minimization would lead to buckling into icosahedral forms for smaller R/a as $N \sim (R/a)^2$; for higher R/a values other factors become significant, and this correspondence cannot be assumed (arXiv:cond-mat/0311413v1). Also the nature of ordering would affect initial susceptibility characteristics, e.g. susceptibility of antiferromagnetic arrays would initially increase with temperature. Now, the viscosity changes in the barrier due to installation of dissipatively ordered (via interplay of thermal and magnetic forces), magnetic assemblies of soft nuclei – in agreement with the proposed nucleation of membranes by FeS by Russell et al (see Sect. 1.2) – would be expected to decrease the possibility of TMC. In any case, magnetically ordered intra-cluster layers (I-CLs) would be in place as templates for hosting oriented diffusion of small clusters as we shall next study.

3 Oriented transport of tiny clusters through I-CLs

The envisaged dynamical assemblies are embedded in the FeS-gel barrier, which as a first approximation demarcates a natural surface-limit for the ensuing processes. Embedded-ness offers protection, apart from benefits of the gradients, in the locally enriched (due to entrapped, synthesized 'metabolites' in the aqueous interior of 'Russell chambers') and shielded environment of the hydrothermal mound (Martin and Russell 2003). With formation of magnetically ordered frambooid-like assemblies, the increased viscosity of the medium could slow down TMC, but the system would be maintained away-from-equilibrium due to the fight between the two forces: The magnetic -field

due to mound rocks would try to harness the thermal flux by herding the super-paramagnetic clusters together. The temperature gradient on the other hand would propel molecules (Baaske et al 2007) across the gradient, in inverse proportion to size, and try to disrupt the assemblies by pushing tiny clusters through them. These in turn would be migrating through the I-CLs in an oriented manner due to magnetic dipole-dipole interactions : smaller clusters would thus become bigger– ‘cluster propagation’ a la Cairns-Smith’s (1985) ‘crystal propagation’ . As magnetic susceptibility of the clusters would be directly proportional to their size, the entire assembly would have a tendency to be propelled in the direction of the mound interior; here translational motion would be impeded by the viscosity of the medium but rotational effects could be possible. So in this section we look at the possibility of oriented-diffusion of tiny clusters through I-CLs of the dynamic assemblies in the context of biologically relevant phenomena. Note that besides being catalytic, the I-CLs *satisfy Schroedinger’s vision (60 years ago) of aperiodic surfaces as efficient information-holders*. This also implies that resistance to migration through I-CLs would have an asymmetric profile, with one direction offering (even marginally) greater ease of passage.

3.1 *Self-replication a la Cairns-Smith; pre- molecular motors*

In contrast to the growth of conventional crystals restricted to a growing surface, field-assisted alignment of diffusing tiny clusters, would occur both on cluster surfaces and *I-CLs*, thereby allowing access to topologically equivalent templates offered by smaller scale nested structures. Their consequent inflation into larger clusters resembles the ‘self-pattern’ conserving replication of biosystems. (This also matches the bigger scale phenomenon of growing FeS ‘bubbles’ (Russell et al. 1994)). This could perhaps answer the subsequent diversity of modes biosystems use for copying information: nucleic acids follow a template basis for assembly, membranes grow by extension of existing ones, entire structures can duplicate (e.g. spindle pole body or the dividing cell as a whole), etc. (Harold 2005). Next consider newly synthesized molecules in the ‘Russell chambers’ that would be ligated to tiny greigite clusters (Milner-White and Russell 2005, 2008). Now, diffusing tiny greigite units through I-CLs have a striking parallel to the directed movement of bio-molecular motors (in the translational, transcriptional, cytoskeletal assemblies) on aperiodic intracellular surfaces that indicate an invariant topological theme for a ratchet mechanism: Movement of a cargo loaded element on a template (representing a varying potential) that harvests thermal fluctuations for dissociating its bound state and spends energy for conformationally controlled directed binding, or an ionic gradient for direction [Astumian (1997)]. In the ‘Russell mound’ the direction of movement of tiny clusters gets dictated by the thermal gradient; the consequent interplay of thermal fluctuations with weak

binding forces (Van der Waal's, magnetic (Wilkins and Barnes 1997)) lead to the propulsion of the loaded tiny clusters, aligned on the I-CL template (that would have presented a locally varying magnetic potential to the aligning and diffusing clusters) thereby generating primitive molecular motors. Later on in evolution, the aperiodic surfaces of the I-CLs, this time formed by consecutive non-magnetic replacements, could have used features like structural anisotropy features for biasing the motion. Recall that a 'thermal gradient' was proposed by Feynman to circumvent the idea of 'biased' Brownian motion (based on structural anisotropy alone) which, despite a right magnitude for driving nano-sized particles [Phillips and Quake (2006)] is otherwise forbidden by the Second Law of Thermodynamics.

3.2 *Pre-RNA world; transfer reactions; optical polarity*

In the proposed RNA world [see Orgel (2004); Koonin and Martin (2005)], RNA played the roles of *both* DNA and protein – call them RNA-sequential and RNA-structural, respectively. Evidently, nature designed DNA for packaging information efficiently, satisfying Shannon's maximum entropy requirement (no correlations across sequences). This leads to the 'chicken-egg' conundrum, as the largely random sequential information encoded in DNA (neglecting structural signatures, e.g. regulation) is correlated via RNA with the high degree of stereo-chemical information in proteins. We suggest that RNA-sequential emerged to replace aperiodically varying I-CL templates hosting directed diffusion of loaded greigite clusters (see above). These very templates are envisaged as a primitive translational machinery, where Wachtershauser's (1988) 'bucket brigade-like' transfer reactions were carried out by oriented greigite units playing the key adaptor roles *a la* transfer RNAs - the directed diffusion of the greigite end on an aperiodic surface with no correlations (RNA-sequential-like), with the other, ligated to compounds rich in structural information (synthesized in the mound or imported). This 'magnetic letters-like' scenario bears a striking resemblance to the tRNA's bringing the amino acids together for stringing them up on the basis of the sequential information inscribed in the mRNA template. Further, in the transfer reactions, the directional asymmetry of transport (the directed passage of the tiny clusters through the asymmetric I-CLs would be guided by the pathway showing less resistance to their transit; see Sect. 3) may well have pushed the balance in favour of bond formation between juxtaposed activated units having the same chirality close to the ligand-binding site, further aided by the space constraints of such intra-layer activity, in tune with McBride and Tully proposal (2008) of a 'grinding start to life'. Thus, the optical activity of the first-bound unit – the symmetry-breaking choice – might have set the preferences for those of the subsequently selected units from the soup of chemicals waiting in the aqueous interiors of the 'Russell compartments' - leading to concentration of

one optical isomer over its counterpart.

3.3 'Mound'-decoupling; origin of gradients, multicellularity ?

The cytoskeleton, at least in eukaryotes, is organized via transmitted internal or external spatial cues, reflecting the polar organization of the cell (Drubin 2000). Now, the directed diffusion of synthesized products would lead to their gradients formed within the membrane-embedded, surface-limited structures. These gradients could have been coupled to nonequilibrium fluctuations via suitable far-from-equilibrium chemical reactions, e.g. magnetically sensitive reactions (Weaver et al. 2000) wherein orientation of reactants could influence reaction rates. Thus crucial pathways of the gradient-rich mound could have been replaced by such strategies, aiding the replicator's exit. Apart from amphipathic barriers for navigating outside the protective 'mound', energy rich molecules would need to take over the charge of pumping energy into the range of dissipative processes, outlined above, e.g. sustaining directed transport in the absence of a thermal gradient using energy-rich molecules would have generate primitive molecular motors. Independence from the mound in turn would require a progressively decreasing functional dependence on FeS (see Sect. 4). (Nevertheless, the continued presence of magnetic elements in the liberated, replicator (e.g. structural roles) would offer a magnetic basis for their association. A proposal on these lines, citing the multicellular life displayed by magnetotactic bacteria as a pathway for the evolution of a multicellular prokaryote, was also made recently (by Davila et al, Lunar Planet Sci XXXVIII (2007), p1495)). Martin and Russell (2003) envisage an initially confined universal ancestor that diverged into replicating systems, located separately on a single submarine seepage site, en route to proto-branches of life, e.g. different initial conditions in differently situated 'LUCA organic concentrates' (last universal common ancestor) would lead to independent schemes for lipid biosynthesis (Koonin and Martin 2005).

4 'The Importance of Being Magnetic': 'The Giant'

The strongest argument for a role of magnetism in the origin of life, via multifaceted iron sulphide phases on the Hadean floor, comes from ubiquitous magnetic influences across kingdoms: navigation sensing in bacteria, algae, protists, bees, ants, fishes, dolphins, turtles and birds (Kirschvink and Gould 1981; Winkelhofer 2005); field effects on growth patterns, differentiation, orientation of plants and fungi (Galland and Pazur 2005); ferromagnetic elements in brain tissues (Kirschvink et al. 1992; Brem et al. 2006), etc. New insights into biophysical mechanisms underlying magnetoreception (arXiv:0804.2646) reveal

the importance of magnetism in biological systems of today. Indeed, magnetism has myriad manifestations at different scales – quantum to cosmological (Skomski 2008). The repeated appearance of fractal themes is compelling – from magnetic critical phenomena to finer length scales where quasiparticle behaviour in a magnetic field can be explained by fractional quantum numbers (Jain 2007; Goerbig et al. 2004); Farey series elements, F_n ; Hausdorff dimension h (da Cruz 2005). Then again, similarities of biological phenomena with magnetic ones in other disciplines have led to shared theoretical approaches. A prominent example is the close mapping between the impact of frustration in magnetically disordered systems and bioprocesses, such as protein folding (Hollander and Toninelli 2004; Stein 1996); this includes a variant of the Ising spin glass as a model for the origin of life (Anderson 1983). Another association between the cytoskeletal network and percolation systems (Traverso 2005), recalls the long-range connectivity of magnetism (e.g., magnetic percolation clusters forming fractal networks (Itoh et al. 2006)). Again, the diamagnetic anisotropy of planar peptide bonds permits their oriented self-assembly in a magnetic field, seen for fibrous biostructures (Torbet and Ronziere 1984 plus ref). These magnetic-like bio-phenomena are seen here as ‘distant cousins’ of their ‘non-living’ counterparts, having evolved by replacing a magnetic Ancestor. Thus even quantum processing is viewed as a legacy and not a product of adaptive evolution (Doll and Finke 2003). For, magnetic ordering may stem from unpaired p - electron systems (Ohldag et al. 2007) (not just 3d, 4f!). The ‘substitutes’, despite increasing complexity, would need to pass-on the legacy of multi-dimensional properties of the Ancestor possessed, especially phase information. Indeed, biomolecules have many facets at different levels, e.g. DNA has positional information, with possible phase signatures in its helical structure (Kwon 2007). Magnetic effects are invasive, altering the expected course of phenomena, but they can also act in an orthogonal non-interfering manner. Its control on spin states gives magnetism a natural handle for manipulation in biology – from chemical reactivity (due to spin-selectivity of reactions; see Buchachenko 2000) to quantum coherent events (arXiv:0804.3503) as in the following.

4.1 Biological searches: quantum search algorithms

Biological systems are intrinsically adaptive: A ‘crisis’ (e.g. a depleted nutrient) could push a system into a search, leading to a new pathway (node) for succour. Among prominent biological-search examples are biological evolution itself, with divergences symbolized by tree nodes; the clonal Darwinian-like phase in the adaptive immune system; brain connections and protein folding (linked to heat-shock proteins occurring in all domains of life). In a classical search, the effect of conflicting forces on different components, plus contributions from environmental noise, can lead to frustrations in local minima. A

quantum search on the other hand, requires quantum coherence in the set of elements on the active front (e.g., a molecule and/or concerned local/non-local neighbour parts); their wave-property enables them to ‘step in phase’ with one another to form a superposition of states and give a ‘holistic’ decision, leading to a more efficient search programme. Thus in the face of crises, halted networked interactions in a subsystem would prompt the formation of a ‘quantum decision front’ with the rest of the system plus environment, waiting as a ‘composite environment’ – its fruitful interaction (a ‘measurement’ of the quantum front) with one chosen path would mean a simultaneous collapse of the quantum superposition of alternative paths (McFadden and Al-Khalili 1999). The newly created path would join up for propagating with other similarly born networked interactions, thus keeping the macroscopic classical system, away-from-equilibrium; due to feedback, this interactive phase would continue till the next challenge forced the system to seek help from the co-existing quantum domain. Now the two main questions are : 1) Does Nature employ quantum coherent processes ? 2) If so, how does it maintain coherence in its interactive milieu ? Today, apart from decoherence evasion mechanisms, e.g. ”dynamical embededness” (Bieberich 2000; Hagan et al. 2002), clear signatures of quantum processing in biology are coming in ((Engel et al. 2007), aided by femtosecond laser-based 2D spectroscopy and coherent control approaches, showing how phase relationships in nano-structures modulate the course of bio-reactions (Nagya et al 2006). It seems Nature has quietly been using these strategies all along. Using Grover’s quantum search method for a marked item in an unsorted database, Patel (2001) hit upon the base-pairing logic of nucleic acids in transcription and translation as an excellent quantum search algorithm – a directed walk through a superposition of all possibilities - resulting in a 2-fold increase in sampling efficacy over its classical counterpart (which at best permits a random walk). Similarly, in a database of dimension d , a quantum search gives a square root speed-up over its classical counterpart – also valid for the respective nested versions (Cerf et al. 2000). Quantum scaling laws (Davies 2004) seem relevant with nested searches for structured problems in fractal processes, e.g. protein folding (via different length scales).

4.2 Coherence: ferromagnetic-ferroelectric effects

Frohlich (1968, 1975) proposed the emergence of a long range coherent state via alignment of dipoles in cell membranes. Ordering of electric dipoles via interactions between structured water and the interior of microtubular cavities brings in a dynamic role of *ferroelectricity* as a frequency-dependent dielectric-constant $\epsilon(\omega)$, which gives a big dispersive (non-dissipative) interaction (robust against thermal losses) for small values of ω (since the factor $\epsilon(\omega)$ occurs in the denominator of the corresponding interaction) (Mavromatos et al 1998). Apart from the importance of such coherent electric dipole ordering

alignment of actin monomers prior to ATP-activation, Hatori et al (quant-ph/0104042) report the coherent alignment of magnetic dipoles induced along the filament, by the flow of protons released from ATP molecules during their hydrolysis (basically a Maxwell displacement current-like dynamical effect). Extrapolating backwards from these interesting present-day biological observations, it is natural to ask if *both* ferroelectric and ferromagnetic effects had co-operatively existed at the Ancestor level. But in contrast to the similar nature of magnetic ordering mechanisms conferring ferromagnetism via exchange interactions of predominantly localized magnetic moments, a variety of ferroelectric ordering mechanisms exist for different types of ferroelectrics, not all of which are well understood. In fact, in materials their co-existence can range from being mutually exclusive, such as due to incompatibility of d-electron criterion for magnetism with off-centering second-order Jahn-Teller effect, all the way to strongly coupled giant magneto-resistance effects (includes non-oxidic ferrimagnetic semi-conductor thiospinels $FeCr_2S_4$ and $Fe_{0.5}Cu_{0.5}Cr_2S_4$, that are Fe^{2+} and Fe^{3+} end members of solid solution $Fe_{1-x}C_xCr_2S_4$ ($0 < x < 0.5$) (Palmer and Greaves 1999)). While lattice distortions with lowered symmetry reduce competing interactions (Chern et al 2006; see also arXiv:cond-mat/0702362v1 ; arXiv:cond-mat/0402495v1), an insight into the loss of inversion symmetry comes via the spin-orbit coupling mechanism which gives the electric polarization \mathbf{P} ($\sim \mathbf{e} \times \mathbf{Q}$), where \mathbf{e} is the spin rotation axis and \mathbf{Q} is the wave vector of a spiral) induced upon transition to a spiral spin-density-wave state triggered by magnetic frustrations (Mostovoy 2006). Apart from the spin-orbit coupling factor, a reduction of crystal symmetry (Fd3m to non-centrosymmetric $F\bar{4}3m$) in several spinel compounds, including $FeCr_2S_4$ was attributed to a displacement of cations (Mertinat et al 2005, Charnock et al 1990). Similar off-centering was also found in oxide spinels (Charnock et al 1990), e.g. magnetite Fe_3O_4 (the inverse spinel greigite is it's sulphur analogue). Additionally, a combination of site-centred (extra holes or electrons on metal sublattice, e.g. Fe^{2+} and Fe^{3+} , where anions don't play a role) and bond-centred (the alternation of short and long bonds, in otherwise equivalent sites, lead to a bond-centered charge density wave) charge-ordering was suggested for explaining the multiferroic behavior of Fe_3O_4 below the Verwey transition at 120K (Khomskii 2004).

Note that the composition of iron sulphide clusters found in enzymes, Fe_5NiS_8 , lie between $FeNi_2S_4$ and Fe_3S_4 . Although a solid solution in this range has not been observed in synthetic dry condition, high temperature experiments, it has been observed in natural violarite (iron-nickel thiospinel) phases (Vaughan and Craig 1985). More recently, the supergene oxidation of pentlandite ($(Fe, Ni)_9S_8$) to violarite (includes extensions from $FeNi_2S_4$ towards both Fe_3S_4 and Ni_3S_4), was experimentally reproduced under mild hydrothermal conditions (Tenailleau et al 2006). Electron backscatter diffraction and backscattered electron imaging clearly showed a dissolution-precipitation transformation mechanism that was dependent upon a large number of solu-

tion parameters, including pH, oxidation/reduction potential, and speciation and concentration of sulphur, iron and nickel in solution. The reaction is promoted by acidic pH and high $H_2S(aq)$ concentrations. Further, assuming the oxidant as Fe^{3+} in reactions where it was added, they observed that the reaction was speeded up by higher Fe^{3+} and H_2S concentrations; here acetate chelated cation was added which shows the similarity to the Russell mound scenario (see Sect. 1.1). These results show the feasibility of high iron/nickel ratios in violarite forming under reducing mound conditions, despite the suggested metastability of these compositions from bonding models. Iron is believed to occur as low spin Fe^{2+} in $FeNi_2S_4$ that exhibits metallic, Pauli paramagnetic behaviour. In contrast, the Mossbauer spectrum of Fe_3S_4 is attributed to high-spin Fe^{3+} in tetrahedral A and Octahedral B sites and its electronic structure from molecular orbital calculations (Vaughan and Tossell 1981) reveal localized 3d electrons with unpaired spins, coupled anti-ferromagnetically at lower temperatures. According to Vaughan and Craig (1985), the greater ionic character and larger number of electrons in antibonding orbitals in Fe_3S_4 relative to $FeNi_2S_4$, could contribute to the instability of intermediate compositions, despite their natural occurrence. They expect a transition from (metallic) delocalized to localized d-electron behaviour at some point in the series, noting such a transition in $CuCr_{2-x}V_xS_4$. Now, the co-operative co-existence of ferroelectric and ferro-magnetic properties in the above structural relatives of greigite –due to a subtle interplay between charge, spin, orbital and lattice degrees of freedom (Hemberger et al 2006) –raise the possibility of a similar profile for Fe_3S_4 or close relatives found in enzymes, e.g. Fe_5NiS_8 , for which no direct evidence is so far available. Note also the possibility of a (effective) lower local temperature due to magnetic ordering (see Sect. 4.3). In any case, either scenario—an independent co-existence in locally differing subspaces or co-existence in the same phase itself—in the Ancestor would only reinforce the expected coherent effects of ferromagnetism, due to their 'dispersive' (non-dissipative) nature.

4.3 Quantum processing in dynamical assemblies

Significantly, Matsuno and Paton (2000) have shown that the gradual release of energy stored in ATP by actomyosin ATPase, in a sequence of quanta E_m over time intervals ($\approx 4.5 \times 10^{-9}$ sec), underlies the huge ($\sim 2 \times 10^6$) discrepancy between the observed time interval of hydrolysis of 1 molecule of $ATP \sim 10^{-2}$ sec, and that calculated via a singly emitted quantum, viz., $\sim 2 \times 10^{-15}$ sec, thus indicating about 2×10^6 number of coherent energy quanta released during one cycle, instead of a single one ! In Kelvin scale, each such energy quantum E_m amounts to 1.6×10^{-3} K associated with the actomyosin complex. This very *low local effective temperature* thus enables these molecular machines to act as heat engines for exploiting the thermodynamic gradient,

taking average physiological temperature as 310K. This inspires us to propose the following mechanism in the Ancestor that would also explain the above as one arising from the evolutionary process.

Consider the oriented transport of a small cluster across an I-CL, as depicted in Figure 3, due to the temperature gradient from the hot mound interior to the cooler ocean front, dictating the direction of transport in this away-from-equilibrium system. To the migrating particle (a dipole), the template (comprising aligned dipoles of the nano-assembly) appears locally as a fluctuating potential. Here energy barriers correspond to the potential faced by the migrating cluster: when it's correctly aligned to the template-dipole it sees an energy minimum (system at Well 1, 2 etc) as the magnetic interaction energy is greatest ('barrier off'); when it is not aligned then the interaction energy can vary, depending upon it's relative orientation, upto a maximum system energy (Hump 1, 2 etc—corresponding to 'barrier on'). At the basin of Well 1, we consider that the particle is completely aligned and therefore effective temperature of system (particle plus template) is lower than that of the surrounding heat bath (local aqueous environment). So the system absorbs energy and with that the particle is positioned on the Hump 1, preceding basin of Well 2 (direction of work done in transport defined by the gradient). At this point the system is at the same temperature as the surrounding bath but now the particle has reduced or lost orientation. At this point the random brownian motion due to molecules of the aqueous medium in which the particle is suspended, will help it overcome this potential barrier as the random hits will rotate the particle in all directions till the correct one allows it to align the system to the basin of Well 2. Due to the temperature gradient the particle does work by migrating to consecutive positions, pumping heat from one basin as it dissipates the rest of the heat (remaining from work done), into the basin of the next one. Superimposed on this activity is the harvesting of Brownian fluctuations (thermal noise is a fluctuating forces averaging to zero over time) by this away-from-equilibrium system which lowers the effective temperature by orienting it to the template dipole. [Absorption of heat from Well 1 that is dissipated in going to Well 2 plus change of position (work done), keeps account of the total entropy which does not decrease. The low effective temperature due to reduction in entropy (on the lines of Li et al (2007), see Sect. 2.4) is at the cost of the surroundings, so that the Second Law of Thermodynamics is not violated]. See the correspondence of this with present day motors where a gradual energy release by ATP serves the function of reducing effective temperature close to 0 ° K (above). Now in contrast to an electron-like particle with two spin states (up and down), the dipole (thanks to help from brownian random hits) aligning to a template, needs to search a continuous set of orientations for achieving the right orientation w.r.t. the template. This offers the possibility of a 'continuous measurement' by the 'waiting partner' on the subsequent position on the template. For example, the template element corresponding to system at Well 2 would be observing




Fig. 3. Oriented transport of tiny cluster (dipole) on I-CL : The x-axis represents a template comprising aligned elements 1,2,3,4 etc. of an intra-cluster layer of the dynamic assembly. The temperature gradient from the hot mound interior (T_1) to the cooler ocean front (T_2) dictates the direction of movement of the particle, for whom the template appears as a locally varying magnetic potential; the y-axis represents the potential of the combined system, i.e. particle plus template, where Wells and Humps represent energy minima and maxima respectively; see text

the aligning dipole corresponding to system at Hump 1. And the relative phase correlations of aligned template elements in turn are due to maintainance of long range order in the magnetic assemblies. Thus, the migrating dipole is like a quantum system from the microscopic realm being observed by an ensemble of ordered elements (template) acting as an observer from the macroscopic realm. [This Quantum Zeno scenario is rather naturally satisfied for a small to moderate magnetic field vis-a-vis a realistic (e.g. due to viscosity effects etc.) interval of 'continuous' observations (see Gagen, Ph.D.Thesis, Quantum measurement theory and the Quantum Zeno effect, University of Queensland, 1993)]. We strongly suspect that the small increments (contributions to total energy for the transition e.g. from Hump 1 to Well 2) from these small Brownian hits (individual bits of quanta close to zero energy) are the analogue of the gradual energy release from ATP in present day evolved molecular motors. Quantum effects would have been correspondingly facilitated due to the effective low temperature brought about, thanks to help from Brownian forces in the same way.

4.4 *A quantum coherent Ancestor*

The characteristics of dynamically self-assembled nano-structures with bottom-up complexity, formed by dissipating energy, depend on the constituent particle size, shape, hardness, composition (Min et al 2008), apart from their sensitivity to (control by) external fields; this approach was used in generating systems with hierarchial complexity via an interplay of magnetic and hydrodynamic interactions (Grzybowski and Campbell 2004). Here, we have addressed the possibility of magnetic-field-controlled dissipative-self-assembly generating nested frambooid-like dynamic order; note that hierarchial architecture is a key feature of living systems. Further, a gel state sustaining ordering of dipoles and a quasicrystalline packing are believed to be important ingredients for accessing quantum coherence in biological systems (Jibu et al 1994). Here firstly, the envisaged frambooid-like order of the gel-embedded assemblies should logically lead to limited pathways for electronic conduction, i.e. only along the connecting edges perpendicular to the five-fold axes (see Sect. 2.5), thus providing zones locally shielded from dissipative effects. Secondly, apart from ordering of dipoles, the mechanism outlined above for the hypothesized clusters would enable coherent connections between the microscopic and macroscopic domains. The co-evolution of both realms, right from the origin of life, could underlie the coherent response of biosystems to fields, e.g. by chemical gradients – critical players in cell events (Harold 2005). Indeed, Selvam (1998) proposed a coherence preservation mechanism via self-similar structures with quasicrystalline order as iterative principles – the main tools for handling non-linear dynamics of perturbations for evolving nested order that connect the microscopic and macroscopic realms with scale-free structures arising out of deterministic chaos. Such long-range spatio-temporal correlations (via a non-dimensional scale factor) are the hallmark of self-similarity, manifest as self-organized criticality in natural dynamical systems (Bak and Chen 1991). Here, the quantum mechanical ‘spin’ endows magnetism with a unique potential for coherently connecting the microscopic and macroscopic domains in the envisaged nested structures. The possible long range order (depending on frustration) in the dynamical assemblies would permit adaptive feedback within the network of interacting clusters, for harnessing the influx of thermal energy. Such a ‘feedback-permitting’ magnetic ancestor seems in line with Ling’s (2001) description of biological structures as *nested organizations based on coherent feedback through a lattice of self and non-self interactions of spatially oriented units*. (For even intracellular water is structured: ‘Site-dipoles’ have been proposed for resolving the apparent contradiction between random molecular movements in and out of sites, and correlated orientations in assemblies, manifesting in coherent patterns (Higo et al. 2001)). Recall some facets of magnetism in common with those of self-organizing systems : emergence of global order from local interactions, organizational closure, hierarchy, downward causation, distributed control underlying robustness, bifurcations

via boundary conditions, non-linearity due to feedback, etc. (Heylighen 2001).

5 Conclusions and scope

Russell et al have argued that life's hatchery could have been busy by 3.8 Gyr, evolving fast enough for a branch to have reached the ocean surfaces by 3.5 Gyr, as evidenced by photosynthetic signatures. The gestation period of life had to have been less than the umbilical mound's delivery of the formative hydrothermal solution, i.e., certainly less than 3 million years, and probably less than 30,000 years (Frueh-Green et al. 2003). This brings us to profound questions linked to the nature of life :

(A) Is life only a classical self-organizing phenomenon? (B) Is it's complexity linked with the quantum domain, enabling efficient *quantum* searches ?

The search for such a connection in the 'mound' scenario (Russell et al 1994) leads to the possibility of TMC forces, arising from an interplay of thermal gradient (50K) and magnetic field (~ 120 mT) acting on a colloidal dispersion (~ 1 cm across) of super-paramagnetic clusters. The resulting dynamical ordering ('bottom-up assembly) of greigite clusters in turn comprises an interplay of attractive and repulsive interactions, making for a spectrum of possibilities – from ratchet effects to heralding in of quantum coherence – which can be explained via the migration of tiny clusters through I-CLs of framboid-like assemblies ; these ingredients can give important insights into molecular motors that seem to have evolved to effectively harness the fluctuations, in their noisy environment. For at the scale of 1°A , the chemical, mechanical, and electrostatic energies converge with that of thermal fluctuations at an average value of $\sim 10^{-18}$ J (see Figure 2 of Phillips and Quake 2006). Especially noteworthy is the significance of the slow release in energy stored in ATP leading to proposed lowering of local temperature down close to 0°K (Matsuno and Paton 2000) ; this scenario matches with the cumulative addition of energy bits from random hits of Brownian motion, till the 'right' orientation is achieved for aligning to the waiting template partner, with lowering of entropy.

Again, 'bucket brigade-like' transfer reactions (Wachtershauser 1988) between tiny greigite clusters ligated to structurally rich compounds, as they migrate through the I-CLs in an oriented manner, could explain the evolution of the translational machinery. And optical polarity would arise by symmetry breaking due to the directional preference of migrating clusters, thanks to the 'grinding effect' (McBride and Tully 2008) brought about via space constraints of these transfer reactions occurring between the layers of the assemblies. In a gel-environment, magnetic interactions would give a semblance of 'bonds' and 'order' associated with the solid state, despite the inflow of thermal energy: in magnetically 'herded' arrays of small diffusing clusters, 'replacements' would continue to maintain the relative orientations of 'predecessors'. The Ancestor's

exit from the mound into a weak (geo-) magnetic field would have required reduced dependence on multi-faceted iron sulphide phases; in addition to ferromagnetic properties, a possible role of ferroelectricity has been discussed (Sect. 4.2). The adaptive nature of biological systems and their fractal organization, press for a coherent connection between the apparently co-evolving classical and quantum domains - a physical basis for linking them at the very origins of life is given in this paper. To that end, this rudimentary study has freely drawn on the findings of a spectrum of scientists (see bibliography), necessitating more interdisciplinary efforts.

Acknowledgements: We thank Prof.M.J.Russell for inspiration and constant support; Prof.Z. Sawlowicz for key references; Dr. A.Boyce for active help with his labelled frambooid-in-chimney pictures; Prof.K.Matsuno for urging a quantitative description and the (anonymous) Referee for pointing out a co-existing possibility of ferroelectricity. This work was entirely financed, with full infrastructural support, by Dr.Jean-Jacques Delmotte; Mr.Guy Delmotte gave computer assistance and Dr.Vineet Ghildyal helped with manuscript processing. GMD acknowledges the driving force of Dr.Anand K.Bachhawat in this project; it's course was transformed with Dr.Bani M.Sodermark's key suggestion "look for a fractal lattice" !

Appendix I : Ordered frambooids

Frambooids have been defined as microscopic spheroidal to sub-spheroidal clusters of equidimensional and equimorphic microcrystals which suggests a homogenous nucleation of the initial microcrystals. Other than the spherical frambooids, a highly ordered icosahedral type has been reported where this packing is maintained in its internal structure. The formational environment is evidently critical for the architecture of these structures that can vary from disordered to remarkably ordered arrangements of constituent crystals. Correlations between frambooid diameter (D) and microcrystal diameter (d) from modern sediments were investigated by Wilkin et al (1996). As pointed out by Ohfuji and Akai (2002), D/d ratios of frambooids dominated by irregular or loosely packed cubic-cuboidal microcrystals are low compared to high corresponding values observed for those composed of densely packed octahedral microcrystals. Interestingly, the latter variety were found in sedimentary rocks more than 11,000 years old [Roberts and Turner (1993)], where the central parts of the weakly magnetized frambooids were found to have greigite microcrystals. Sections from these show that the pentagonal arrangement comprise a central pentagonal domain with its sides connected to five rectangular/trapezoid-like regions which are in turn connected via fanshaped domains. The arrangement pattern of these densely packed octahedral micro-

crystals linked edge to edge is 'lattice-like' (space filled) in the rectangular domains, whereas in the triangular domains the triangles are formed by the (111) faces of the octahedral microcrystals and the voids between them. Thus within these domains the individual faces of the microcrystals do not make any contact. The icosahedral form is seen as generated by stacking twenty tetrahedral sectors packed on three faces out of four, and connected by their apexes at the centre. Generally acknowledged as dynamically stable, this form is known to have six 5-fold axes at each apex, and ten 3-fold axes at each face, as can be seen in a number of naturally occurring structures from microclusters like fullerene to some viruses (Ohfuji and Akai 2002). In an investigation of apparent biologically induced mineralization by symbiotically associating bacterial and archaeal species, framboidal greigites have been obtained from Black Sea sediments that are ordered clusters of octahedral crystals comprising Fe_3S_4 -spinel (Essentially cubic where sulphur forms a fcc lattice with 32 atoms in the unit cell, and Fe occupies 1/8 of the tetrahedral and 1/2 of the octahedral sites). Their size is restrained by their icosahedral symmetry and under greater pressures at depths of 200m, the diameters are mostly \sim (2.1, 4.2, 6.3 or 8.4) μ , with the two intermediate ones predominating. The smallest of these are formed from 20 octahedral crystals (0.35 μ) positioned at the apexes of an icosahedron and surrounding a 0.5 μ diameter vacancy that give rise to 12 pentagonal depressions on the outside. Nested structures building up from this smallest one lead to the higher sized clusters (Preisinger and Aslanian 2004). Sub-spheroidal pyrite-framboids, due to curved polyhedron-like outer facets, probably reflect an internal icosahedral microcrystal organisation (Ohfuji and Akai 2002), which are classically forbidden crystallographic symmetries (Ohfuji and Rickard 2005).

Appendix II : Phyllotaxis

Phyllotaxis was first identified in plants: the numerical regularity with which rows of nearest neighbours in lattices of plant elements (leaves, scales, etc.) form two families of intersecting spirals, with right and left-handed threads called parastichies (Levitov 1991), with their numbers as consecutive Fibonacci numbers (1, 1, 2, 3, 5, 8, 13, 21, ...). The divergence angle between consecutive primordia, is generally close to the 'golden angle' $\alpha = 360(5 - \sqrt{5})/2$ degrees. As pointed out by Hotton et al (2006), two simple geometrical rules: 1) equivalent or nearly equivalent units are added in succession; 2) position of new units is determined by interactions with those already in place, underlie the crystal-like periodicity of myriad natural patterns. The self-organizing system of Douady and Couder (1996) captures some of these features, with the radial motion of ferrofluid drops mimicking phyllotactic growth – an optimized interplay of repulsive forces between the drops and their periodic

introduction— demonstrating its success in applying the underlying physics as iterative principles. In fact, the repulsive interactions between sequentially arising/deposited elements in an energy minimized manner, and manifesting in mathematically precise phyllotaxis patterns are generic to countless phenomena from subnano to cosmological scales (Levitov 1991; Adler et al. 1997) : Repulsive magnetic dipoles, galactic structures, biostructures, from the molecular (proteins, DNA) to macroscopic levels (myriad forms such as marine-life), proportions in morphological and branching patterns (Dunlap 1997), Bernard convection cells, stress-driven self-assembly, bunched crystalline ion beams, atmospheric flows, and flux lattices in layered superconductors. Stressing on the underlying manifold, Nisoli et al (see arXiv:cond-mat/0702335v1) discuss the effect of repulsive interaction of magnetic dipoles on a cylinder, instead of on a flat space.

Appendix III : Quasiperiodicity

The discovery of quasicrystals (see Appendix III) with pentagonal symmetry had rekindled interest in phyllotactic phenomena by providing an illuminating link between observed periodicities in animate and inanimate forms (Adler et al 1997). The connection of the classically forbidden icosahedral form to quasiperiodic phases was sparked off by Shechtman et al. (1984). Further, the correspondence between the diffraction spectra of quasicrystalline solids and model spectra of substitution tilings, was realized from the models based on self-similar tilings (e.g. Penrose) providing the necessary aperiodicity and long range order (Frank in arXiv:0705.1142v1). For although aperiodic, quasicrystals give sharp diffraction spots like crystals, yet no unit cell can describe their atomic packing. Unlike classical crystals (30 symmetry groups), no corresponding group of isometric transformations leave quasicrystals invariant. But compensating self-similarities confer invariance under affine homothetic transformations. In fact, inflation rules determine hierarchical selfsimilar packing of atomic clusters in perfect icosahedral quasicrystals, whose structural stability demands bonding electrons to be recurrently localized to selfsimilar isomorphic subsets; the corresponding eigenstates generate a hopping mechanism for conductivity (Janot 1997). Further, unlike conventional crystals whose surfaces are defined by bulk planes with maximum atomic density, surface identification in quasicrystalline materials is compounded by the quasiperiodic stacking of infinite non-identical atomic layers. According to the inter-layer spacing rule (Sharma et al 2004), stable quasicrystal surfaces are a consequence of bulk truncations at positions where blocks of atomic layers are separated by larger interlayer spacings. Though so far studies have been mostly carried out on metallic alloy phases, more recently, studies on soft matter (Lifshitz and Diamant 2007) indicate that interwoven patterns characteristic of nested,

self-similar, quasiperiodic structures can form on increasing scales of tiling edges: metallic alloy (0.5nm), chalcogenide (2nm), dendron (10nm), and star block copolymer (80nm) (Henley et al. 2005). We note that properties of quasicrystalline order (based on faceted-phases of metal alloys) have analogies in bio-processes: allow diffusion and storage due to low friction coefficients; facile cleavage due to brittle nature; presence of metals in catalytic amounts; resistance against oxidation and wetting; low thermal conductivity; atypical electron transport, etc. (Roche and Fujiwara 1998; Suck et al. (Eds.) 2004; Thiel 2004; Trebin (Ed.) 2003) (also Zhong et al. arXiv:cond-mat/0011118v1).

Appendix IV : Derivation of average range of H-field

In this appendix, we outline a simple derivation of the average range of the magnetic field over a square cavity of height h due to a dipole placed symmetrically below the cavity at a distance of $h/4$, a la Fig 1 of Ganguli et al (2004). This quantity which is best expressed via the average H - field (proportional to $1/r^2$) in accordance with the double integral

$$h^2 \langle 1/r^2 \rangle = \int \int dx dy [x^2 + (y + h/4)^2]^{-1} \quad (\text{A.1})$$

where (x, y) are defined as in Fig 1 of Ganguli et al(2004) , over the domain $-h/2 \leq (x, y) \leq +h/2$; (note the 'shift ' of the ordinate y by an amount $h/4$ due to the position of the dipole $h/4$ below the X-axis). The result of integration is

$$h^2 \langle 1/r^2 \rangle = \pi \ln 7 = 6.113 \quad (\text{A.2})$$

To see how this comes about, change to polar coordinates as per the convention of fig. 1 in Ganguli et al(2004) , viz.,

$$x = \rho \sin \phi; \quad y = \rho \cos \phi$$

This gives $r^2 = \rho^2 + h\rho(\cos\phi)/2 + h^2/16$, whose substitution in (1) leads to

$$h^2 \langle 1/r^2 \rangle = \int \rho d\rho \int d\phi [\rho^2 + h\rho(\cos\phi)/2 + h^2/16]^{-1} \quad (\text{A.3})$$

where now $0 \leq \rho \leq h/\sqrt{2}$, and $0 \leq \phi \leq 2\pi$. The ϕ integration can be done by the standard method of contour integral, according to

$$\int_0^{2\pi} d\phi / (1 + a \cos \phi) = 2\pi / \sqrt{1 - a^2}; \quad a^2 < 1$$

Substitution of this result in (A3) gives

$$h^2 < 1/r^2 > = \int_0^{h/\sqrt{2}} 2\pi\rho d\rho/(\rho^2 - h^2/16) \quad (\text{A.4})$$

leading finally to the result (A2).

References

- [1] Adler I, Barabe D, Jean RV (1997) A history of the study of phyllotaxis. *Annals of Botany* 80, 231-244
- [2] Ahniyaz A, Sakamoto Y, Bergstrom L (2007) Magnetic field-induced assembly of oriented superlattices from maghemite nanocubes. *Proc Natl Acad Sci* 104(45), 17570-17574
- [3] Astumian RD (1997) Thermodynamics and kinetics of a Brownian motor. *Science* 276 (5314), 917 - 922
- [4] Anderson PW (1983) Suggested model for prebiotic evolution: The use of chaos. *Proc Natl Acad Sci USA* 80, 3386-3390
- [5] Baaske P, Weinert F, Duhr S et al (2007) Extreme accumulation of nucleotides in simulated hydrothermal pore systems. *Proc Natl Acad Sci USA* 104, 9346-9351
- [6] Bieberich E (2000) Probing quantum coherence in a biological system by means of DNA amplification. *Biosystems* 57(2), 109-24
- [7] Boyce AJ, Coleman ML, Russell MJ (1983) Formation of fossil hydrothermal chimneys and mounds from Silvermines, Ireland. *Nature*, 306, 545-550
- [8] Brem F, Hirt AM, Winklhofer M et al (2006) Magnetic compounds in the human brain : a comparison of tumor and hippocampal tissue. *J R Soc Interface*, 3(11), 833-841
- [9] Cairns-Smith AG (1985) Seven clues to the origin of life. Cambridge University Press, New York
- [10] Cerf NJ, Grover LK, Williams CP (2000) Nested quantum search and NP-complete problems. *Phys Rev A* 61, 032303
- [11] Charnock J, Garner CD, Pattick RAD et al (1990) An EXAFS study of thiospinel minerals. *Amer Mineral* 75, 247-255
- [12] Cheon J, Park J-I, Choi J-S et al (2006) Magnetic superlattices and their nanoscale phase transition effects. *Proc Natl Acad Sci* 103(9), 3023-3027

- [13] Chern G-W, Fennie CJ, Tchernyshyov O (2006) Broken parity and a chiral ground state in the frustrated magnet $CdCr_2O_4$. Phys Rev B 74, 060405(R) (4pages)
- [14] da Cruz W (2005) A fractal-like structure for the fractional quantum Hall effect. Chaos Solitons and Fractals 23, 373-378
- [15] Davies PCW (2003) How bio-friendly is the universe. Int J Astrobiol 2, 115
- [16] Davies PCW (2004) Does quantum mechanics play a non-trivial role in life? Biosystems 78, 69-79
- [17] Doll KM, Finke RG (2003) A compelling experimental test of the hypothesis that enzymes have evolved to enhance quantum mechanical tunneling in hydrogen transfer reactions: The β -neopentylcobalamin system combined with prior adocobalamin data. Inorg Chem 42 (16), 4849-4856
- [18] Douady S, Couder Y (1996) Phyllotaxis as a dynamical self-organizing process. Part I: The spiral modes resulting from time-periodic iterations. J theor Biol 178, 255-274
- [19] Drubin DG (2000) Cell polarity. Oxford University Press, Oxford
- [20] Dunlap RA (1997) The Golden Ratio and Fibonacci Numbers. World Scientific, New Jersey.
- [21] Engel GS, Calhoun TR, Read EL et al (2007) Evidence for wavelike energy transfer through quantum coherence in photosynthetic systems. Nature 446, 782-786
- [22] Finlayson BA (1970) Convective instability of ferromagnetic fluids. J Fluid Mech 40, 753-767
- [23] Frohlich H (1968) Longrange coherence and energy storage in biological systems. Int J Quantum Chem 2, 6419
- [24] Frohlich H (1975) The extraordinary dielectric properties of biological materials and the action of enzymes. Proc Natl Acad Sci USA 72, 4211-4215
- [25] Frueh-Green GL, Kelley DD, Bernasconi SM et al (2003) 30,000 years of hydrothermal activity at the Lost City vent field. Science 301, 495-498
- [26] Galland P, Pazur A (2005) Magnetoreception in plants. J Plant Res 118(6), 371-389
- [27] Ganguly R, Puri IK (2006) Field assisted self-assembly of superparamagnetic nanoparticles for biomedical, MEMS, and bioMEMS applications. Adv Appl Mechanics 41, 293-335
- [28] Ganguly R, Sen S, Puri IK (2004) Thermomagnetic convection in a square enclosure using a line dipole. Phys Fluids 16, 2228
- [29] Goerbig MO, Lederer P, Morais Smith C (2004) On the self-similarity

- [30] Goerbig MO, Lederer P, Morais Smith C (2004) On the self-similarity in quantum Hall systems. *Europhys Lett* 68, 72-78
- [31] Hagan S, Hameroff S, Tuszynski J (2002) Quantum computation in brain microtubules? Decoherence and biological feasibility. *Phys Rev E* 65, 061901
- [32] Harold FM (2005) Molecules into cells: Specifying spatial architecture. *Microbiol Mol Biol Rev* 69(4), 544-564
- [33] Hemberger J, Lunkenheimer P, Ficht R et al (2006) Multiferroicity and colossal magneto-capacitance in Cr-thiospinels. *Phase Trans* 79, 1065
- [34] Henley CL, de Boissieu M, Steurer W (2006) Discussion on clusters, phasons, and quasicrystal stabilisation. *Phil Mag* 86 (6-8), 1131-1151
- [35] Heylighen F (2001) The Science of Self-organization and Adaptivity. In: L. D. Kiel, (ed.) *Knowledge Management, Organizational Intelligence and Learning, and Complexity*, In: *The Encyclopedia of Life Support Systems ((EOLSS))*, Eolss Publishers, Oxford.
- [36] Higo J, Sasai M, Shirai H et al (2001) Large vortex-like structures of dipole field in computer models of liquid water and dipole-bridge between biomolecules. *Proc Natl Acad Sci USA* 98, 5961-5964
- [37] Hollander WThF den, Toninelli FL (2004) Spin glasses : A mystery about to be solved. *Nieuw Archief voor Wiskunde*, 5/5(4), 274-278
- [38] Hotton S, Johnson V, Wilbarger J et al (2006) The possible and the actual in phyllotaxis: Bridging the gap between empirical observations and iterative models. *J Plant Growth Regul* 25, 313-323
- [39] Hu S, Deng C, Appel E et al (2002) Environmental magnetic studies of lacustrine sediments. *Chin Sci Bull* 47(7), 613-616
- [40] Huber C, Wachtershauser G (1997) Activated acetic acid by carbon fixation on (Fe,Ni)S under primordial conditions. *Science* 276, 245-247
- [41] Itoh S, Kajimoto R, Iwasa K et al (2006) Fractal structure and critical scattering in the three-dimensional percolating antiferromagnet, RbMn_{0.31}Mg_{0.69}F₃. *Physica B* 385-386 (1), 441-443
- [42] Jain JK (2007) *Composite fermions*. Cambridge Univ Press, Cambridge
- [43] Janot C (1997) Atomic clusters, local isomorphism, and recurrently localized states in quasicrystals. *J Phys Sect Condens Matter* 9, 1493-1508
- [44] Jibu M, Hagan S, Hameroff SR et al (1994) Quantum optical coherence in cytoskeletal microtubules: implications for brain function. *Biosystems* 32(3), 195-209
- [45] Keys AS, Glotzer SC (2007) How do Quasicrystals Grow? *Phys Rev Lett* 99, 235503

- [46] Kirschvink JL, Gould JL (1981) Biogenic magnetite as a basis for magnetic field detection in animals. *Biosystems* 13(3), 181-201
- [47] Kirschvink JL, Kobayashi-Kirschvink A, Diaz-Ricci JC et al (1992) Magnetite in human tissues: a mechanism for the biological effects of weak ELF magnetic fields. *Bioelectromagnetics Suppl* 1, 101-13
- [48] Khomskii DI (2004) Multiferroics: different ways to combine magnetism and ferroelectricity. *J Magn Magn Mat* 306, 1
- [49] Koonin EV, Martin W (2005) On the origin of genomes and cells within inorganic compartments. *Trends Genet* 21, 647-654
- [50] Kopelman R (1989) Diffusion-controlled reaction kinetics. In: Avnir D (ed) *The fractal approach to heterogenous chemistry*. John Wiley, New York, pp 295-309
- [51] Kwon Y (2007) Quantum mechanism of biological search. *Chaos, Solitons and Fractals* 34 (4), 1037-1038
- [52] Lalatonne Y, Richardi J, Pileni MP (2004) Van der Waals versus dipolar forces controlling mesoscopic organizations of magnetic nanocrystals. *Nat Mater* 3(2), 121-5
- [53] Larter RCL, Boyce AJ, Russell MJ (1981) Hydrothermal pyrite chimneys from the Ballynoe baryte deposit, Silvermines, County Tipperary, Ireland. *Mineralium Deposita* 16, 309-317
- [54] Levitov LS (1991) Fibonacci numbers in botany and physics: Phyllotaxis. *Pis'ma Zh Eksp Teor Fiz* 54(9), 542-545
- [55] Li J, Huang Y, Liu X et al (2007) Effect of aggregates on the magnetization property of ferrofluids: A model of gaslike compression. *Sci Tech Adv Mater* 8, 448-454
- [56] Lifshitz R (1998) Symmetry of magnetically ordered quasicrystals. *Phys Rev Letts* 80(12), 2717-2720
- [57] Lifshitz R, Diamant H (2007) Soft quasicrystals - Why are they stable? *Phil Mag* 87, 3021-3030
- [58] Ling GN (2001) *Life at the cell and below-cell level: The hidden history of a functional revolution in Biology*. Pacific Press, New York
- [59] Martin W, Russell MJ (2003) On the origins of cells: a hypothesis for the evolutionary transitions from abiotic geochemistry to chemoautotrophic prokaryotes, and from prokaryotes to nucleated cells. *Philos Trans R Soc Lond* 358(1429), 59-83, discussion 83-5.
- [60] Matsuno K, Paton RC (2000) Is there a biology of quantum information ? *Biosystems* 55(1-3), 39-46
- [61] Mavromatos NE, Nanopoulos DV, Samaras I et al (1998) Ferroelectrics and their possible involvement in biology. *Adv Struct Biol* 5, 127 - 134

- [62] McBride JM, Tully JC (2008) Did life grind to a start? *Nature* 452, 161-162
- [63] McFadden J, Al-Khalili J (1999) A quantum mechanical model of adaptive mutations. *Biosystems* 50, 203-211
- [64] Mertinat M, Tsurkan V, Samusi D et al (2005) Low-temperature structural transition in $FeCr_2S_4$. *Phys Rev B* 71, 100408(R) (4 pages)
- [65] Milner-White EJ, Russell MJ (2005) Nests as sites for phosphates and iron-sulfur thiolates in the first membranes: 3 to 6 residue anion-binding motifs. *Origins Life Evol Biosphere* 35, 19-27
- [66] Milner-White EJ, Russell MJ (2008) Predicting the conformations of peptides and proteins in early evolution. *Biol Direct*, doi:10.1186/1745-6150-3-3
- [67] Min Y, Akbulut M, Kristiansen et al (2008) The role of interparticle and external forces in nanoparticle assembly. *Nature Materials* 7, 527-538
- [68] Mostovoy M (2006) Ferroelectricity in spiral magnets. *Phys Rev Lett* 96, 067601 (4 pages)
- [69] Mukhopadhyaya A, Ganguly R, Sena S, Puri IK (2005) A scaling analysis to characterize thermomagnetic convection. *Int J Heat Mass Transfer* 48 (17), 3485-3492
- [70] Nagya A, Prokhorenkoa V, Dwayne Millera RJ (2006) Do we live in a quantum world ? Advances in multidimensional coherent spectroscopies refine our understanding of quantum coherences and structural dynamics of biological systems. *Curr Opin Struct Biol* 16(5), 654-663
- [71] Navrotsky A (2004) Energetic clues to pathways to bio-mineralization: Precursors, clusters and nanoparticles. *Proc Natl Acad Sci, USA* 101 (33), 12096-12101
- [72] Odenbach S (2004) Recent progress in magnetic fluid research. *J Phys Condens Matter* 16, R1135-R1150
- [73] Ohfuji H, Akai J (2002) Icosahedral domain structure of framboidal pyrite. *Amer Mineral* 87, 176-180
- [74] Ohfuji H, Rickard D (2005) Experimental synthesis of framboids—a review *Earth-Sci Rev* 71, 147-170
- [75] Ohldag H, Tylliszczak T, Hohne R et al (2007) Pi-Electron ferromagnetism in metal-free carbon probed by soft X-ray dichroism. *Phys Rev Lett* 98, 187204
- [76] Orgel LE (2004) Prebiotic chemistry and the origin of the RNA world. *Crit Rev Biochem Mol Biol* 39(2), 99-123
- [77] Palmer HM, Greaves C (1999) Structural, magnetic and electronic properties of $Fe_{0.5}Cu_{0.5}Cr_2S_4$. *J Mater Chem* 9, 637-640
- [78] Patel AD (2001) Quantum algorithms and the genetic code. *Pramana* 56(2), 367

- [79] Phenrat T, Saleh N, Sirk K et al (2007) Aggregation and sedimentation of aqueous nanoscale zerovalent iron dispersions. *Environ Sci Technol* 41, 284-290
- [80] Phillips R, Quake SR (2006) The biological frontier of physics. *Physics Today* 59(5), 38-43
- [81] Posfai M, Buseck PR, Bazylinski DA et al (1998) Reaction sequence of iron sulphide minerals in bacteria and their use as biomarkers. *Science* 280, 880-883
- [82] Posfai M, Dunin-Borkowski RE (2006) Sulfides in biosystems. *Reviews in Mineralogy and Geochemistry* 61, 679-714
- [83] Preisinger A, Aslanian S (2004) The formation of framboidal greigites in the Black Sea. *Geophys Res Abstr* 6, 02702 (SRef-ID: 1607-7962/gra/EGU04-A-02702)
- [84] Rannacher D, Engel A (2004) Double Rosensweig instability in a ferrofluid sandwich structure. *Phys Rev E* 69, 66306
- [85] Redl FX, Cho K-S, Murray CB et al (2003) Three-dimensional binary superlattices of magnetic nanocrystals and semiconductor quantum dots. *Nature* 423, 968971
- [86] Rickard D, Butler IB, Oldroyd A (2001) A novel iron sulphide mineral switch and its implications for Earth and planetary science, *Earth Planet Sci Lett* 189, 85-91
- [87] Rickard D, Luther GW III (2007) The chemistry of iron sulfides. *Chem Revs* 107, 514-562
- [88] Rickard D, Morse JW (2005) Acid volatile sulfide (AVS) *Marine Chem* 97, 141-197
- [89] Righter K, Drake MJ, Yaxley G (1997) Prediction of siderophile element metal-silicate partition coefficients to 20GPa and 2,800 ° C: the effects of pressure, temperature, oxygen fugacity, and silicate and metallic melt compositions. *Phys Earth Planet Interiors* 100, 115-134
- [90] Roche S, Fujiwara T (1998) Fermi surfaces and anomalous transport in quasicrystals. *Phys Rev B* 58(17), 11338-11344
- [91] Rosensweig RE (1997) *Ferrohydrodynamics*. Dover, New York
- [92] Roth J, Ghler F (1998) Atomic self-diffusion in dodecagonal quasicrystals. *Eur Phys J B* 6, 425-445
- [93] Russell MJ (1986) Reflections on the genesis of the Irish Carboniferous lead+zinc+baryte deposits. In: Nesbitt RW, Nichol I (eds) *Geology in the Real World - the Kingsley Dunham Volume*, Inst Min Metall, London, pp 375-386
- [94] Russell MJ, Arndt NT (2005) Geodynamic and metabolic cycles in the Hadean. *Biogeosciences* 2, 97-111

- [95] Russell MJ, Daniel RM, Hall AJ, Sherringham JA (1994) A hydrothermally precipitated catalytic iron sulphide membrane as a first step toward life. *J Mol Evol* 39, 231-243
- [96] Russell MJ, Hall AJ (1997) The emergence of life from iron monosulphide bubbles at a submarine hydrothermal redox and pH front. *J Geol Soc London* 154(pt 3) 377-402
- [97] Russell MJ, Hall AJ (2008) A hydrothermal source of energy and materials at the origin of life. In press
- [98] Russell MJ, Hall AJ, Turner D (1989) In vitro growth of iron sulphide chimneys: possible culture chambers for origin-of-life experiments. *Terra Nova* 1(3), 238-241
- [99] Russell MJ, Hall AJ, Boyce AJ et al (2005) On hydrothermal convection systems and the emergence of life. *Economic Geology* 100 (3), 419-438
- [100] Russell MJ, Hall AJ, Mellersh AR (2003) On the dissipation of thermal and chemical energies on the early Earth: The onsets of hydrothermal convection, chemiosmosis, genetically regulated metabolism and oxygenic photosynthesis. In: Ikan R (ed) *Natural and laboratory-simulated thermal geochemical processes*. Dordrecht, Kluwer Academic Publishers, pp325-388
- [101] Ryskin A, Muller HW, Pleiner H (2003) Thermal convection in binary fluid mixtures with a weak concentration diffusivity, but strong solutal buoyancy forces. *Phys Rev E* 67, 046302, p 1-8
- [102] Sawlowicz Z (1993) Pyrite framboids and their development: a new conceptual mechanism. *Int J Earth Sci* 82, 148-156
- [103] Sawlowicz Z (2000) Framboids: from their origin to application. *Prace Mineralog Pan* 88, 1-80
- [104] Schroedinger E (1944) *What is Life?* Cambridge University Press, Cambridge
- [105] Selvam AM (1998) Quasicrystalline pattern formation in fluid substrates and phyllotaxis. In: Barabe D, Jean RV (eds) *Symmetry in Plants*. World Scientific Series in Mathematical Biology and Medicine (4), Singapore, pp795-809
- [106] Sharma HR, Fourne V, Shimoda M et al (2004) Structure of the fivefold surface of the icosahedral Al-Cu-Fe quasicrystal: Experimental evidence of bulk truncations at larger interlayer spacings. *Phys Rev Lett* 93, 165502
- [107] Shechtman D, Blech I, Gratias D et al (1984) Metallic phase with long-range orientational order and no translational symmetry. *Phys Rev Lett* 53, 1951-1953
- [108] Skomski R (2008) *Simple models of magnetism*. Oxford University Press, Oxford
- [109] Stein DL (1996) Spin glasses, In: Rigden JS (ed) *Macmillan Encyclopedia of Physics*. Simon and Schuster Macmillan, New York, pp1514 - 1516

- [110] Stone DA, Goldstein RE (2004) Tubular precipitation and redox gradients on a bubbling template. *Proc Natn Acad Sci USA* 101, 11537-1154
- [111] Suck J-B, Schreiber M, Hussler P (eds) (2004) *Quasicrystals : An introduction to structure, physical properties and applications*. Springer-Verlag, New York
- [112] Sun S, Murray CB, Weller D et al (2000) Monodisperse FePt Nanoparticles and Ferromagnetic FePt Nanocrystal Superlattices. *Science* 287, 1989-1992
- [113] Taketomi S, Takahashi H, Inaba N et al (1991) Experimental and theoretical investigations on agglomeration of magnetic colloidal particles in magnetic fluids. *J Phys Soc Jpn* 60 (5), 1689-1707
- [114] Tarduno JA, Cottrell RD, Watkeys MK et al (2007) Geomagnetic field strength 3.2 billion years ago recorded by single silicate crystals. *Nature*, 446, 657-660
- [115] Tenailleau C, Pring A, Estchmann B et al (2006) Transformation of pentlandite to violarite under mild hydrothermal conditions. *Amer Mineral* 91 (4), 706-709
- [116] Thiel PA (2004) An introduction to the surface science of quasicrystals. *Prog Surf Sci* 75, 69-86
- [117] Torbet J, Ronziere M-C (1984) Magnetic alignment of collagen during self-assembly. *Biochem J* 219, 1057-1059
- [118] Traverso S (2005) Cytoskeleton as a fractal percolation cluster: Some biological remarks. In : Losa G, Merlini D, Nonnenmacher E, Weibel E (eds) *Fractals in biology and medicine*. (Part 4) Birkhauser, Basel, pp 269-275
- [119] Trebin H-R (ed) (2003) *Quasicrystals: Structure and physical properties*. Wiley-VCH, Weinham
- [120] Trevors JT, Pollack GH (2005) Hypothesis: the origin of life in a hydrogel environment. *Prog Biophys Mol Biol* 89(1), 1-8
- [121] Tnyi I, Guba P, Roth LE et al (2003) Electric discharges in the protoplanetary nebula as a source of impulse magnetic fields to promote dust aggregation. *Earth, Moon, and Planets* 93 (1), 65-74
- [122] Tuszynski JA, Hameroff S, Sataric MV et al (1995) Ferroelectric behaviour in microtubule dipole lattices: Implications for information processing, signalling and assembly/dissassembly. *J theor Biol* 174, 371-380
- [123] Uechi I, Katsuki A, Dunin-Barkovskiy L et al (2004) 3D-Morphological Chirality Induction in Zinc Silicate Membrane Tube Using a High Magnetic Field. *J Phys Chem B* 108(8), 2527 -2530
- [124] Vaughan DJ, Craig JR (1985) the crystal chemistry of iron-nickel thiospinels. *Amer Mineral* 70, 1036-1043
- [125] Vaughan DJ, Tossell JA (1981) Electronic structure of thiospinels minerals: results from MO calculations. *Amer Mineral* 66, 1250-3

- [126] Wachtershauser G (1988) Before enzymes and templates: theory of surface metabolism. *Microbiol Rev* 52, 452-484
- [127] Wang J, Zhu Y, Li W et al (2005) Necklace-shaped assembly of single-crystal $NiFe_2O_4$ nanospheres under magnetic field. *Mater Lett* 59, 2101-2103
- [128] Wang Z, Holm C (2003) Structure and magnetic properties of polydisperse ferrofluids: A molecular dynamics study. *Phys Rev E* 68, 041401, p1-11
- [129] Wasilewski P, Kletetschka G (1999) Lodestone: Nature's only permanent magnet-what it is and how it gets charged; *Geophys Res Lett*, 26(15), p. 2275-2278
- [130] Weaver JC, Vaughan TE, Astumian RD (2000) Biological sensing of small field differences by magnetically sensitive chemical reactions. *Nature* 405 (6787), 707-709
- [131] Wilkin RT, Barnes HL (1997) Formation processes of framboidal pyrite. *Geochim Cosmochim Acta* 61, 323-339
- [132] Wilkin RT, Barnes HL, Brantley SL (1996) The size distribution of framboidal pyrite in modern sediments: An indicator of redox conditions. *Geochim Cosmochim Acta* 60, 3897-3912
- [133] Winkelhofer M (2005) Biogenic magnetite and magnetic sensitivity in organisms -from magnetic bacteria to pigeons. *Magneto hydrodynamics* 41(4), 295-304
- [134] Wolthers M, Van Der Gaast SJ, Rickard D (2003) The structure of disordered mackinawite. *Amer Mineral* 88, 2007-2015
- [135] Wu M, Xiong Y, Jia Y et al (2005) Magnetic field-assisted hydrothermal growth of chain-like nanostructure of magnetite. *Chem Phys Lett* 401, 374-379
- [136] Zhang L, Luo J, Chen Q (2005) Magnetic properties of assembled ferrite nanostructures induced by magnetic fields. *J Phys Condens Matter* 17, 5095-5100
- [137] Zubarev AY, Fleischer J, Odenbach S (2005) Towards a theory of dynamical properties of polydisperse magnetic fluids: Effect of chain-like aggregates. *Physica A* 358 475491
- [138] Zubarev AY, Iskakova LY (2004) To the theory of rheological properties of ferrofluids: influence of drop-like aggregates. *Physica A* 343, 65-80