

A Non-Fermi Liquid from a Charged Black Hole; A Critical Fermi Ball

Sung-Sik Lee

Department of Physics & Astronomy, McMaster University,

Hamilton, Ontario L8S 4M1, Canada

(Dated: February 21, 2019)

Abstract

Using the AdS/CFT correspondence, we calculate the single particle spectral function in a 2+1 dimensional non-Fermi liquid state which is dual to a gravitational theory in the AdS_4 background with a charged black hole. In the non-Fermi liquid state, there is no quasiparticle but the Fermi surface is still well defined. Interestingly, all momentum points inside the Fermi surface are critical and the gapless modes are defined in a *critical Fermi ball* in the momentum space.

arXiv:0809.3402v1 [hep-th] 19 Sep 2008

Landau Fermi liquid theory is a low energy effective theory for degenerate many-body Fermi systems. In Fermi liquid states, the curvature of a Fermi surface suppresses non-forward scatterings, which makes it possible for well defined quasiparticles to exist in the low energy limit[1, 2]. Then non-interacting Fermi gases essentially capture the qualitative nature of interacting Fermi systems. Although the Fermi liquid theory provides correct descriptions for most metals, strongly correlated systems including the high temperature superconductors[3] and heavy fermion compounds[4, 5] are not described by the Fermi liquid theory. These so called non-Fermi liquid states are different states of conducting matters and an understanding of those states is very important not only for applications but also from a fundamental physics point of view. However, theoretical understandings of non-Fermi liquid states are still limited. In particular, there is no general theoretical tool which enables one to make quantitative predictions. The fundamental difficulty lies in the fact that non-Fermi liquid states are strong coupling phenomena for which the well developed standard perturbation theories usually break down.

On the other hand, there have been significant developments in understanding a class of strongly coupled quantum field theories. In the AdS/CFT correspondence, a gravitational theory in the anti-de Sitter (AdS) space is dual to a strongly coupled conformal field theory (CFT) defined on the boundary of the AdS space[6, 7, 8]. In a large N limit, the gravitational theory is reduced to a classical gravity from which one can understand non-trivial strong coupling physics of the boundary CFT. Recently the AdS/CFT correspondence has been applied to various phenomena which arise in the context of condensed matter systems[9, 10, 11, 12, 13, 14, 15, 16]. The goal of this paper is to apply the AdS/CFT correspondence to construct a non-Fermi liquid state and examine dynamical properties of the state by calculating the single particle spectral function. For this, we consider a gravitational background of a charged black hole where the charge induces a non-zero density of fermions in the boundary quantum field theory.

We consider the action,

$$\begin{aligned}
S = & \frac{1}{\kappa_4^2} \int d^4x \sqrt{-g} \left[\frac{1}{4} R - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{3}{2} \right] \\
& + \int d^4x \sqrt{-g} \left[\bar{\psi} \gamma^\mu \left(\partial_\mu + \frac{1}{2} \omega_\mu^{bc} \Sigma_{bc} - i A_\mu \right) \psi - m \bar{\psi} \psi \right] \\
& + \int d^3x \sqrt{-g_\epsilon} \bar{\psi} \psi.
\end{aligned} \tag{1}$$

Here $x^\mu = (t, x, y, z)$ is the space-time coordinate with signature $(-1, 1, 1, 1)$. R is the scalar curvature and $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the field strength tensor of a U(1) gauge field A_μ . ψ is a

four-component Dirac spinor, ω_μ^{bc} is the spin connection and $\Sigma_{bc} = \frac{1}{4}[\Gamma_b, \Gamma_c]$ is the generator of the local Lorentz transformation with Γ_a , the gamma matrices. $\gamma^\mu = e_a^\mu \Gamma^a$ where e_a^μ is the tetrad. The action describe the U(1) gauge field and the Dirac spinor coupled with gravity in the background with a negative cosmological constant which is set to be -1 in our unit. The last term in the action is a boundary term defined at $z = \epsilon$ where g_ϵ is the determinant of the induced metric on the 2+1D space. Although the boundary term does not affect the equation of motion in the bulk, it is important for obtaining a non-trivial dependence of the saddle point action on the boundary value of the spinor field[17].

The above action has an AdS_4 black hole solution given by

$$\begin{aligned} ds^2 &= \frac{1}{z^2} \left[\alpha^2 (-f(z)^2 dt^2 + dx^2 + dy^2) + \frac{dz^2}{f(z)^2} \right], \\ A_0 &= q\alpha(z-1), \end{aligned} \tag{2}$$

where $f(z) = \sqrt{1 + q^2 z^4 - (1 + q^2)z^3}$. This is a special case of a more general dyonic black hole solution considered in Ref. [9]. In this coordinate system, the horizon of the black hole is at $z = 1$. The metric describes the AdS_4 space near the boundary at $z = 0$. The Hawking temperature of the black hole is $T_H = \frac{\alpha}{4\pi}(3 - q^2)$ [9]. For nonzero q and α , the black hole carries a nonzero charge. Non-vanishing components of the spin connection are $\omega_{\hat{t}\hat{z}}^{\hat{t}\hat{z}} = z^2 \left(\frac{f}{z}\right)'$ and $\omega_{\hat{x}\hat{z}}^{\hat{x}\hat{z}} = \omega_{\hat{y}\hat{z}}^{\hat{y}\hat{z}} = -f$, where $(\hat{t}, \hat{x}, \hat{y}, \hat{z})$ represents the local Lorentz coordinate.

According to the AdS/CFT correspondence, we can view this classical gravitational theory as a strongly coupled 2+1D quantum field theory in a large N limit. The Hawking temperature corresponds to the temperature of the boundary field theory. This theory can be motivated from the M-theory defined in $AdS_4 \times S^7$ which describes the low energy physics of the 2+1D supersymmetric Yang-Mills theory with 16 supercharges. If the infinite tower of the Kaluza-Klein modes are truncated self consistently in the M-theory, the resulting theory would contain the above theory with particular values of the fermionic mass. Here, we will not restrict ourselves to the M-theory and we will regard the fermionic mass as a free parameter which characterizes the corresponding 2+1D quantum field theory. In particular, we will focus on the case with $m = 0$ where the chiral symmetry simplifies the calculation significantly. However, the qualitative features may be similar for other values of the mass.

The Dirac spinor is a source field which is linearly coupled with the fermionic field of the boundary theory and the gauge field is coupled with the conserved current of the fermion. The electrostatic potential induces a nonzero density of the boundary fermions. The chemical potential

of the boundary theory is given by $\mu = A_0(z = 0) = -q\alpha$. This is crucial in obtaining a system of non-relativistic fermions with a Fermi surface.

To study dynamical properties of the fermion in the boundary field theory, we calculate the single fermion spectral function. For $m = 0$, the chiral symmetry enables us to focus on one chiral mode. Due to the 2+1 dimensional translational symmetry, we can assume a plane wave solution for the left chiral modes ψ_- and $\bar{\psi}_-$,

$$\begin{aligned}\psi(t, x, y, z) &= e^{-i(\omega t - \mathbf{k} \cdot \mathbf{r})} \psi_-(z), \\ \bar{\psi}(t, x, y, z) &= e^{i(\omega t - \mathbf{k} \cdot \mathbf{r})} \bar{\psi}_-(z),\end{aligned}\tag{3}$$

where $\mathbf{r} = (x, y)$ and $\mathbf{k} = (k_x, k_y)$. In the chiral representation with $\Gamma^0 = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$, $\Gamma^i = \begin{pmatrix} 0 & \sigma^i \\ \sigma^i & 0 \end{pmatrix}$, where σ^i are the Pauli matrices with $i = x, y, z$ and I is the 2×2 identity matrix, the equation of motion for the two-component the chiral spinors becomes

$$\begin{aligned}\left[\frac{i\omega z}{\alpha f} + \frac{iqz(z-1)}{f} + \frac{i\mathbf{k} \cdot \boldsymbol{\sigma}}{\alpha} z + \left(\frac{zf'}{2} - \frac{3f}{2} + zf\partial_z \right) \sigma^z \right] \psi_-(z) &= 0, \\ \bar{\psi}_-(z) \left[\frac{i\omega z}{\alpha f} + \frac{iqz(z-1)}{f} - \frac{i\mathbf{k} \cdot \boldsymbol{\sigma}}{\alpha} z + \left(\frac{zf'}{2} - \frac{3f}{2} + \overleftarrow{\partial}_z z f \right) \sigma^z \right] &= 0.\end{aligned}\tag{4}$$

At $T = 0$ ($q^2 = 3$), the solution near the horizon ($z \rightarrow 1$) reads $\psi_-(z) \sim e^{-i\omega \frac{1}{6\alpha(1-z)} \sigma^z} C$, $\bar{\psi}_-(z) \sim \bar{C} e^{-i\omega \frac{1}{6\alpha(1-z)} \sigma^z}$ in the leading order of $(1-z)$ for some fixed spinors C, \bar{C} . To maintain the causality in the boundary theory, we impose the ingoing boundary condition,

$$\psi_-(z) \sim e^{i\omega \frac{1}{6\alpha(1-z)}} \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad \bar{\psi}_-^T(z) \sim e^{-i\omega \frac{1}{6\alpha(1-z)}} \begin{pmatrix} 1 \\ 0 \end{pmatrix}\tag{5}$$

as $z \rightarrow 1$. At a finite temperature with $q^2 < 3$, the boundary condition is modified to be $\psi_- \sim e^{-i\omega \frac{\ln(1-z)}{\alpha(3-q^2)}} \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ and $\bar{\psi}_-^T \sim e^{i\omega \frac{\ln(1-z)}{\alpha(3-q^2)}} \begin{pmatrix} 1 \\ 0 \end{pmatrix}$. Therefore there is one-parameter family of solutions that satisfy the ingoing boundary condition near the horizon. Near the boundary of the AdS space ($z \rightarrow 0$), the spinors behave as $\psi_-(z) \sim z^{3/2} \chi$ and $\bar{\psi}_-(z) \sim z^{3/2} \bar{\chi}$, where χ and $\bar{\chi}$ should be chosen so that the solution satisfies the ingoing boundary condition near the horizon.

We represent the solution near the boundary as

$$\begin{aligned}\psi_-(z) &\sim \left(\frac{z}{\alpha}\right)^{3/2} \begin{pmatrix} P(\omega, \mathbf{k}) \\ 1 \end{pmatrix} \eta, \\ \bar{\psi}_-^T(z) &\sim \left(\frac{z}{\alpha}\right)^{3/2} \begin{pmatrix} Q(\omega, \mathbf{k}) \\ 1 \end{pmatrix} \eta^*,\end{aligned}\tag{6}$$

where η and η^* are Grassmann numbers (not spinors) that we use to impose boundary data. Once the second components of the spinors are chosen to be η and η^* , the complex functions $P(\omega, k)$ and $Q(\omega, k)$ are uniquely determined from the boundary condition near the horizon. It is noted that we could have chosen the boundary condition near $z = 0$ in different ways. This is because the two components of the spinors decay in the same power[18]. For example, we could have chosen the first component as boundary data and let the second component to be determined by the boundary condition near the horizon. Different choices of boundary condition may correspond to different field theories on the boundary[19]. However, we note that Eq. (6) is a natural choice for the following reasons. First, the vector $(0, 1)$ along which we impose boundary data is an eigenvector of σ^z , the generator of the rotation in the $x - y$ plane. Therefore, this prescription is independent of momentum direction, which guarantees that the propagator that we will calculate below is invariant under the rotation. Second, once the second component of ψ_- is chosen as boundary data, it is natural to choose the second component of $\bar{\psi}_-$ as boundary data. This can be seen by turning on a small fermion mass which mixes ψ_- and ψ_+ , and identifying $\bar{\psi}_- \sim \psi_+^\dagger$.

From the AdS/CFT dictionary, the Green's function of the fermion in the boundary theory is given by $G(\omega, \mathbf{k}) = i \frac{\partial^2 S[\eta^*, \eta]}{\partial \eta \partial \eta^*}$, where $S[\eta^*, \eta]$ is the action evaluated for the saddle configuration of the spinor fields which satisfy the boundary conditions, Eqs. (5) and (6). The bulk spinor action in Eq. (1) vanishes at saddle points. Only the boundary term contribute to the action and we obtain the Green's function

$$G(\omega, \mathbf{k}) = i (P(\omega, \mathbf{k})Q(\omega, \mathbf{k}) + 1).\tag{7}$$

The quantity of physical importance is the spectral function, $A(\omega, \mathbf{k}) = \lim_{\delta \rightarrow 0^+} \text{Im}G(\omega + i\delta, \mathbf{k})$ which measures how much spectral weight a fermion with momentum \mathbf{k} has at energy ω .

We numerically integrate the equation of motion (4) to obtain the spectral function as a function of ω and $k = \sqrt{k_x^2 + k_y^2}$. Due to the rotational symmetry, $A(\omega, \mathbf{k})$ does not depend on momentum direction. In Fig. 1, we show the spectral function at zero temperature. For a large momentum,




FIG. 1: The contour plot of the zero temperature spectral function as a function of energy and momentum for $q = -\sqrt{3}$ and $\alpha = 10$. The darkest region represents the area with no spectral weight and the brightest region, the highest value of the spectral function.

the spectral function as a function of energy shows a broad peak which is centered at a negative energy ω_0 . The broad peak does not disperse significantly as momentum changes. However, the width of the broad peak becomes larger as momentum increases and the edges of the broad peak disperse as $\omega_{edge} \approx \pm k + \omega_0$. There is no quasiparticle peak, which implies that the fermions are in a non-Fermi liquid state.

Although there is no delta function peak, the spectral function shows sharp peaks at zero energy for momenta smaller than a critical momentum $k_c \approx 9$. To closely examine the low energy structure, we display the spectral function as a function of energy for fixed values of momentum in Fig. 2. Within the numerical accuracy, the sharp peak has an algebraic singularity at $\omega = 0$ [20]. We emphasize that this is not a quasiparticle peak. As is shown in Fig. 2, the zero energy peak is more pronounced at a smaller momentum and the size of the peak decreases as momentum increases. The critical momentum k_c above which the zero energy peak disappears coincides with the momentum at which the edge of the broad peak cross the Fermi energy $\omega = 0$. Therefore, we interpret k_c as Fermi momentum. The most striking feature is that the algebraic singularities at zero energy exist for all momenta below the Fermi momentum. Namely, all momentum points




FIG. 2: The energy distribution curves of the spectral function for $q = -\sqrt{3}$ and $\alpha = 10$ at momenta (a) $k = 3$, (b) $k = 5$, (c) $k = 7$, (d) $k = 9$, (e) $k = 11$ and (f) $k = 13$.

inside a two dimensional disk with $|\mathbf{k}| < k_c$ has the singular peak at zero energy. We call the set of these momentum points a *critical Fermi ball*. Although not shown here, the Fermi momentum increases as α increases. If we switch the sign of q , the broad peak is centered at a positive energy. However, the position of the critical Fermi ball does not change.

Unlike the Fermi liquid state or a non-Fermi liquid state with a critical Fermi surface[21] where low energy excitations exist only near a Fermi surface, in the present non-Fermi liquid state all momentum points below the Fermi surface are important at low energies. Therefore we expect

that the low energy properties of this state to be drastically different from a Fermi liquid state or a non-Fermi liquid state with a critical Fermi surface. For example, low temperature thermodynamic properties of this 2+1D non-Fermi liquid state with a critical Fermi ball will behave like a 3+1D critical Fermi surface. In a sense, this ‘dimensional lift’ is not surprising because the 2+1D non-Fermi liquid theory is described by the 3+1D gravity.

Although the occurrence of the critical Fermi ball is somewhat counter-intuitive, one may understand the critical Fermi ball as a consequence of strong interactions. Since the gravitational description is valid in the strong coupling limit, where the interaction energy scale is presumably larger than the Fermi energy, even those fermions which are deep inside the Fermi surface can participate in the low energy physics, overcoming the kinetic energy penalty.




FIG. 3: Temperature dependence of the spectral function at $k = 6$. With a fixed $\alpha = 10$, q is changed to tune temperature to $T = 0$ (solid line), $T = \frac{1}{4\pi}$ (dashed line) and $T = \frac{10}{4\pi}$ (dotted line).

Until now, we have examined the zero temperature spectral function. In Fig. 3, we compare the spectral function at zero temperature and finite temperatures. As expected, the singular zero energy peak is rounded at finite temperature due to thermal fluctuations. If temperature is high enough, the sharp peak completely disappears.

In summary, we obtain a non-Fermi liquid state from the AdS/CFT correspondence. The single fermion spectral function reveals a critical Fermi ball in the momentum space where all momentum points inside the Fermi ball are critical.

In the future, it would be interesting to study physical properties of the critical Fermi ball in more details, such as possible instabilities and thermodynamic/transport properties. Due to the presence of extensive gapless modes, it is expected that there are infinitely many singular channels of particle-hole and particle-particle excitations with different momenta which compete with each

other.

I thank Pavel Kovtun and Subir Sachdev for helpful comments. This work has been supported by NSERC.

- [1] R. Shankar, *Rev. Mod. Phys.* **66**, 129 (1994).
- [2] J. Polchinski, *Recent directions in particle theory*, Proc. 1992 TASI, eds. J. Harvey and J. Polchinski (World Scientific, Singapore, 1993).
- [3] P. A. Lee, N. Nagaosa and X.-G. Wen, *Rev. Mod. Phys.* **78**, 17 (2006); references there-in.
- [4] H. v. Lohneysen, A. Rosch, M. Vojta and P. Wolfle, *Rev. Mod. Phys.* **79**, 1015 (2007)
- [5] P. Gegenwart, Q. Si and F. Steglich, *Nature Physics* **4**, 186 (2008).
- [6] J. M. Maldacena, *Adv. Theor. Math. Phys.* **2**, 231 (1998).
- [7] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, *Phys. Lett. B* **428**, 105 (1998).
- [8] E. Witten, *Adv. Theor. Math. Phys.* **2**, 253 (1998).
- [9] S. A. Hartnoll and P. K. Kovtun, *Phys. Rev. D* **76**, 066001 (2007).
- [10] S. A. Hartnoll, P. K. Kovtun, M. Muller and S. Sachdev, *Phys. Rev. B* **76**, 144502 (2007).
- [11] C. P. Herzog, P. K. Kovtun, S. Sachdev and D. T. Son, *Phys. Rev. D* **75**, 085020 (2007).
- [12] A. Karch and A. O'Bannon, *J. High Energy Phys.* **09**, 024 (2007).
- [13] S. S. Gubser, arXiv:0801.2977.
- [14] S. A. Hartnoll, C. P. Herzog and G. T. Horowitz, *Phys. Rev. Lett.* **101**, 031601 (2008).
- [15] D. T. Son, *Phys. Rev. D* **78**, 046003 (2008).
- [16] A. Karch, D. T. Son and A. O. Starinets, arXiv:0806.3796.
- [17] M. Henningson and K. Sfetsos, *Phys. Lett. B* **431**, 63 (1998).
- [18] W. Muck and K. S. Viswanathan, *Phys. Rev. D* **58**, 106006 (1998).
- [19] I.R. Klebanov and A.M. Polyakov, *Phys. Lett. B* **550**, 213 (2002).
- [20] In the numerical calculation, the peak is not divergent because we use a finite $\delta = 10^{-3}$.
- [21] T. Senthil, *Phys. Rev. B* **78**, 035103 (2008).