

# A KUROSU-TYPE THEOREM FOR TYPE III FACTORS

JASON ASHER

## 1. INTRODUCTION

In [Oz], Ozawa obtained analogues of the Kurosh subgroup theorem (and its consequences) in the setting of free products of semiexact  $\text{II}_1$  factors with respect to the canonical tracial states. In particular he was able to prove a certain unique-factorization theorem that distinguishes, for example, the  $n$ -various  $L(F_\infty) * (L(F_\infty) \otimes R)^{*n}$ . The paper was a continuation of the joint work [OP] with Popa that proved various unique-factorization results for tensor products of  $\text{II}_1$  factors.

These papers shared a particular combination of ideas from [Oz2] and [Po3]. The first was a  $C^*$ -algebraic method for detecting injectivity of a von Neumann algebra, adopted from its original context of proving solidity of finite von Neumann algebras with the Akemann-Ostrand property. This method was used in concert with Popa's intertwining-by-bimodules technique that (roughly speaking) allows one to conclude unitary conjugacy results from a spatial condition. More precisely, if  $A$  and  $B$  are diffuse subalgebras of a finite von Neumann algebra  $M$ , the presence of an  $A$ - $B$  sub-bimodule of  $L^2(M)$  with finite right  $B$ -dimension implies that a 'corner' of  $A$  can be conjugated into  $B$  by a partial isometry in  $M$ . The technique was actually shown by Popa to apply to the case of  $(M, \varphi)$  not necessarily finite (but with discrete decomposition), though one needs the additional assumption that  $A$  and  $B$  are subalgebras of the centralizer  $M^\varphi$ .

In this paper we will demonstrate how the above results can be extended to the case of free products of semiexact  $\text{II}_1$  factors with respect to arbitrary (non-tracial) states. The primary difficulty this setting presents is that the resulting factors will be necessarily of type III. Following the same basic outline of the proof from [Oz], we will therefore utilize a generalization of Popa's intertwining technique that allows the ambient algebra  $M$  to be completely arbitrary, and makes no assumption about the relative position of the subalgebras  $A$  and  $B$  in  $M$ . One of the fruits of our labors will be the result that for  $M_i$  semiexact non-prime non-injective  $\text{II}_1$  factors with faithful normal states  $\varphi_i$ , the reduced free product  $\star_{i=1}^m (M_i, \varphi_i)$  is uniquely written in such a way as the free product of such factors, up to permutation of indices and stable isomorphism.

Acknowledgement: The author would like to thank Dimitri Shlyakhtenko for initially suggesting this investigation, and for his invaluable subsequent advice and direction.

## 2. REDUCED FREE PRODUCT

We will first review the reduced free product construction [Vo] in our particular case, retaining the notation from [Oz]. For  $i \in \{1, \dots, m\}$ , let  $M_i$  be semiexact  $\text{II}_1$  factors with faithful normal states  $\varphi_i$ , and let  $H_i = L^2(M_i, \varphi_i)$ . Denote by  $\xi_i$  the  $M_i$ -cyclic-and-separating state vector  $\hat{1} \in H_i$ , and let  $J_i$  be the modular conjugation on  $H_i$ . Finally, let  $H_i^0 = H_i \ominus \mathbb{C}\xi_i$ ,

and construct a new Hilbert space  $H$  as follows:

$$H = \mathbb{C}\xi \oplus \bigoplus_{n \geq 1} \bigoplus_{\substack{i_1, \dots, i_n \in \{1, \dots, m\} \\ i_1 \neq i_2 \neq \dots \neq i_n}} H_{i_1}^0 \otimes \dots \otimes H_{i_n}^0.$$

Next, let  $H(i)$  be the subspace of  $H$  given by the direct sum of  $\mathbb{C}\xi$  and those terms in the direct sum above with  $i_1 \neq i$ . For each  $i$  there is a unitary isomorphism

$$U_i : H \rightarrow H_i \otimes H(i)$$

which identifies  $H(i)$  with  $\mathbb{C}\xi \otimes H(i)$  and  $H(i)^\perp$  with  $H_i^0 \otimes H(i)$ . Define a representation  $\lambda_i : \mathbb{B}(H_i) \rightarrow \mathbb{B}(H)$  by

$$\lambda_i(a) = U_i^*(a \otimes 1_{H(i)})U_i.$$

The reduced free-product  $(M, \varphi) = \ast_{i=1}^m (M_i, \varphi_i)$  is then the von Neumann algebra in  $\mathbb{B}(H)$  generated by  $\lambda_i(M_i)$  for  $i \in \{1, \dots, m\}$ , with faithful normal state  $\varphi$  given by

$$\varphi(x) = \langle x\xi, \xi \rangle_H.$$

Note that  $\xi$  is an  $M$ -cyclic-and-separating state vector, and so  $H = L^2(M, \varphi)$  with the associated modular conjugation  $J$  given by

$$J(\eta_{i_1} \otimes \dots \otimes \eta_{i_n}) = J_{i_n} \eta_{i_n} \otimes \dots \otimes J_{i_1} \eta_{i_1}$$

where  $\eta_{i_1} \otimes \dots \otimes \eta_{i_n} \in H_{i_1}^0 \otimes \dots \otimes H_{i_n}^0$  as in the direct sum decomposition above. Furthermore, one can thus see that  $JH(i)$  is the direct sum of  $\mathbb{C}\xi$  and those terms in the direct sum above with  $i_n \neq i$ .

For each  $i$  there is thus another unitary isomorphism

$$V_i : H \rightarrow JH(i) \otimes H_i$$

which identifies  $JH(i)$  with  $JH(i) \otimes \mathbb{C}\xi$  and  $JH(i)^\perp$  with  $JH(i) \otimes H_i^0$ . Then, precisely as in [Oz] Lemma 3.1, we have the following

**Lemma 2.1.** *For  $a \in \mathbb{B}(H_i)$ ,  $j \neq i$ , and with  $P_\xi$  the projection of  $H$  onto  $\mathbb{C}\xi$  we have*

$$\begin{aligned} \lambda_i(a) &= J V_i^* (1_{JH(i)} \otimes J_i a J_i) V_i J \\ &= V_i^* (P_\xi \otimes a + \lambda_i(a)|_{JH(i) \ominus \mathbb{C}\xi} \otimes 1_{H_i}) V_i \\ &= V_j^* (\lambda_i(a)|_{JH(i)} \otimes 1_{H_i}) V_j \end{aligned}$$

This shows that  $V_i$  is a right  $M_i$ -module isomorphism of  $L^2(M, \varphi)$  with  $K_i \otimes H_i$ , where  $K_i = JH(i)$ . Furthermore, if we let  $D_i = V_i^*(\mathbb{K}(K_i) \otimes_{\min} \mathbb{B}(H_i))V_i$  then this lemma also shows that  $\lambda_j(a)D_i \subseteq D_i$  for  $i, j \in \{1, \dots, m\}$ .

Let  $B_i \subseteq M_i$  be a  $\sigma$ -weakly dense exact  $C^*$  algebra, and let  $B$  be the  $C^*$  algebra in  $\mathbb{B}(H)$  generated by  $\lambda_i(B_i)$  for  $i \in \{1, \dots, m\}$ . Then for  $C = JBJ$  we have the following lemma, as in [Oz], Prop. 3.2.

**Lemma 2.2.** *Suppose that  $\Psi : \mathbb{B}(H) \rightarrow \mathbb{B}(H)$  is a  $C$ -bimodule unital completely positive map. If  $D_i \subset \ker \Psi$  for  $i \in \{1, \dots, m\}$ , then the unital completely positive map*

$$\begin{aligned} \tilde{\Psi} : B \otimes C &\rightarrow \mathbb{B}(H) \\ \sum_{k=1}^n a_k \otimes x_k &\mapsto \Psi \left( \sum_{k=1}^n a_k x_k \right) \end{aligned}$$

*is continuous with respect to the minimal tensor norm.*

Finally, we will need the following control of partial isometries that normalize diffuse subalgebras in free products, see [Po2] and [Oz] Lemma 2.3. For completeness, we include the proof.

**Lemma 2.3.** *Let  $(M, \varphi) = \ast_{i=1}^l (M_i, \varphi_i)$  be the reduced free product of  $II_1$  factors  $M_i$  with faithful normal states  $\varphi_i$ , and let  $Q \subseteq M_j$  be a diffuse von Neumann subalgebra. If  $v \in M$  is a nonzero partial isometry with  $vv^* \in Q' \cap M$  and such that  $v^*Qv \subseteq M_i$ , then  $i = j$  and  $v \in M_j$ .*

*Proof:* Let  $\zeta \in L^2(M_i, \varphi_i) \subseteq L^2(M, \varphi)$  be such that  $\langle \cdot, \zeta \rangle$  gives the canonical trace on  $M_i$ . Note that then  $x\zeta = J_\varphi x^* J_\varphi \zeta = \zeta x$  for all  $x \in M_i$ . If we let  $\eta = v\zeta$ , then  $\eta \neq 0$  and

$$q\eta = vv^*qv\zeta = v\zeta(v^*qv) = \eta(v^*qv)$$

for all  $q \in Q$ .

Suppose, by contradiction, that  $i \neq j$ . Then  $L^2(M, \varphi)$  as an  $M_j - M_i$  bimodule is an infinite multiple of the ‘coarse’ bimodule  $L^2(M_j, \varphi_j) \otimes L^2(M_i, \varphi_i)$ . Considering the projection of  $\eta$  onto any coarse component, we get a Hilbert-Schmidt operator  $T$  on  $L^2(M, \varphi)$  such that

$$qT = T(v^*qv)$$

for all  $q \in Q$ . As  $Q$  is diffuse, we must have that  $Tv^* = 0$ , so that  $T = Tv^*v = 0$ . Thus,  $\eta = 0$ , a contradiction.

So, we must have that  $i = j$ . Now, note that  $L^2(M, \varphi)$  as an  $M_j - M_j$  bimodule is isomorphic to the direct sum of  $L^2(M_j, \varphi_j)$  and an infinite multiple of the coarse bimodule  $L^2(M_j, \varphi_j) \otimes L^2(M_j, \varphi_j)$ . By the above argument, we must have that  $\eta \in L^2(M_j, \varphi_j)$ . Thus, considering these  $L^2$ -vectors as unbounded operators affiliated with  $M_j$ , we get that  $v = \eta\zeta^{-1} \in M_j$ .  $\square$

### 3. INTERTWINING-BY-BIMODULES

We present here a generalization of the intertwining-by-bimodules technique of Popa, see Theorem 2.1 of [Po3]. The technical advantage is that we allow the ambient algebra to be arbitrary, and we make no assumption about the location of the subalgebras.

The following lemma is standard - for a reference on this and other aspects of the modular theory of Tomita and Takesaki see [Ta].

**Lemma 3.1.** *Let  $M$  be a von Neumann algebra in a standard representation  $(M, H, J, P)$ , where  $J$  is the modular conjugation, and  $P$  is the self-dual cone. Then any vector  $\eta \in H$  has a unique polar decomposition as  $\eta = u|\eta|$  for  $|\eta| \in P$  and a partial isometry  $u \in M$  such that the source of  $u$  is  $[M'|\eta|]$ , and the target of  $u$  is  $[M'\eta]$ . Furthermore if  $\eta$  satisfies an*

equation of the form  $x\eta = Jx^*J\eta$  for all  $x$  in a (possibly nonunital) subalgebra  $A$  of  $M$ , then  $|\eta|$  satisfies the same equation, and  $u$  satisfies  $xu = ux$  for all  $x \in A$ .

Proof: Note that  $(M', H, J, P)$  is a standard representation for  $M'$ , and consider the positive normal functional  $\varphi_\eta$  on  $M'$  given by

$$\varphi_\eta(z) = \langle z\eta, \eta \rangle.$$

Then there is a unique vector  $|\eta| \in B$  such that  $\varphi_\eta(z) = \langle z|\eta|, |\eta| \rangle$ . Defining  $u$  on  $M'|\eta|$  by  $uz|\eta| = z\eta$  is well-defined (and when extended by continuity gives a partial isometry) because  $\|z\eta\|^2 = \varphi_\eta(z^*z) = \|z|\eta|\|^2$ . Then  $u$  is clearly in  $M$  and furthermore, uniqueness is now obvious.

Suppose further that  $\eta$  satisfies  $x\eta = Jx^*J\eta$  for all  $x \in A$ . By considering  $A + \mathbb{C}(1 - p)$  for  $p$  the unit of  $A$ , we may assume that  $A$  is unital. Then for any unitary  $w \in A$  one has that

$$(wuw^*)(wJwJ|\eta|) = wJwJ\eta = \eta.$$

Also,  $\zeta = wJwJ|\eta| \in B$ , and  $wuw^*$  is a partial isometry with source  $[M'\zeta]$  and target  $[M'\eta]$ . Thus, by uniqueness, we obtain the result.

□

**Proposition 3.2.** *Let  $M$  be a von Neumann algebra with faithful normal state  $\varphi$ , and let  $A, B \subseteq M$  be two factors of type  $II_1$ . Suppose that there is a nonzero  $A$ - $B$  bimodule  $H \subset L^2(M, \varphi)$  such that the von Neumann dimension of  $H$  as a right  $B$  module satisfies  $\dim_B H_B < \infty$ . Then there are nonzero projections  $q \in A$  and  $p \in B$ , an injective \*-homomorphism  $\theta : qAq \rightarrow pBp$ , and a nonzero partial isometry  $v \in M$  such that  $v^*v \leq p$ ,  $vv^* \leq q$ , and  $xv = v\theta(x)$  for all  $x \in qAq$ .*

Proof:

We can pick nonzero projections  $q \in A$  and  $p \in B$  such that  $K = qJ_\varphi pJ_\varphi H$  is a nonzero  $qAq$ - $pBp$  bimodule with  $\dim_{pBp} K_{pBp} = 1$ . Then, we can find an right Hilbert  $pBp$ -module isomorphism  $\alpha : L^2(pBp, \tau_{pBp}) \rightarrow H$ , and define  $\theta : qAq \rightarrow pBp$  by

$$\theta(x)\zeta = \alpha^{-1}(x\alpha(\zeta)) \text{ for all } \zeta \in L^2(pBp, \tau_{pBp}).$$

Note that  $\theta$  is a nonzero, and hence injective, \*-homomorphism.

Furthermore, if we let  $\xi = \alpha(\hat{p})$ , then for  $x \in qAq$ ,

$$x\xi = \alpha(\theta(x)\hat{p}) = \alpha(\hat{p}\theta(x)) = J_\varphi\theta(x)^*J_\varphi\xi.$$

Consider now  $M_2 \otimes M$  with the faithful normal state  $\tilde{\varphi} = \text{Tr} \otimes \varphi$ , and define for  $x \in qAq$ ,

$$\tilde{x} = \begin{bmatrix} x & 0 \\ 0 & \theta(x) \end{bmatrix}.$$

Also, let  $\tilde{\xi} \in L^2(M_2 \otimes M, \tilde{\varphi})$  be defined as

$$\tilde{\xi} = \begin{bmatrix} 0 & \xi \\ 0 & 0 \end{bmatrix}$$

and note that we then have the equation

$$\tilde{x}\tilde{\xi} = J_{\tilde{\varphi}}\tilde{x}^*J_{\tilde{\varphi}}\tilde{\xi}$$

for all  $x \in qAq$ .

By the lemma above, if we let  $\tilde{\xi} = u|\tilde{\xi}|$  be the polar decomposition of  $\tilde{\xi}$  then  $u$  in  $M_2 \otimes M$  is a partial isometry satisfying

$$\tilde{x}u = u\tilde{x}$$

for all  $x \in qAq$ . It is easy to see that  $u$  has the form

$$u = \begin{bmatrix} 0 & w \\ 0 & 0 \end{bmatrix}$$

for a partial isometry  $w \in M$  that satisfies  $xw = w\theta(x)$  for all  $x \in qAq$ . In fact  $w$  is the polar part of  $\xi$ , and so letting  $v = qw = qw\theta(q) = w\theta(q)$ , we obtain the desired result.

□

#### 4. KUROSH-TYPE THEOREM

We will now present the main technical result, a generalization of Theorem 3.3 in [Oz]. Note that much of the argument remains the same, but we include it for narrative coherence. Also, in everything that follows, all von Neumann algebras are assumed to be separable.

**Theorem 4.1.** *Let  $M_i$ , for  $i \in \{1, \dots, l\}$ , be semiexact  $II_1$  factors with faithful normal states  $\varphi_i$ , and let  $(M, \varphi) = \ast_{i=1}^l (M_i, \varphi_i)$  be their reduced free product. Suppose that  $Q \subseteq M$  is an injective  $II_1$  factor with a normal conditional expectation  $E_Q : M \rightarrow Q$ . If  $Q' \cap M$  is non-injective, then there exists an index  $i$  and a maximal partial isometry  $u \in M$  with  $uu^* \in Q' \cap M$  and such that  $u^*Qu \subseteq M_i$ .*

Proof: Recall all of the notation from the reduced free product construction above. Let

$$\Psi_Q : \mathbb{B}(L^2(M, \varphi)) \rightarrow Q' \cap \mathbb{B}(L^2(M, \varphi))$$

be a proper conditional expectation, i.e. such that for all  $x \in \mathbb{B}(L^2(M, \varphi))$  we have

$$\Psi_Q(x) \in \overline{\text{co}}^\omega \{uxu^* : u \in U(Q)\}.$$

Note that  $\Psi_Q$  is a  $Q' \cap \mathbb{B}(L^2(M, \varphi))$ -bimodule map.

As  $Q$  is in the range of a normal conditional expectation from  $M$ , we can find a faithful normal state  $\nu$  on  $M$  which centralizes  $Q$ . Then  $\Psi_{Q|_M}$  is  $\nu$ -preserving, and so  $\Psi_{Q|_M}$  is the unique  $\nu$ -preserving normal conditional expectation  $E_{Q' \cap M} : M \rightarrow Q' \cap M$ . Lemma 5 from [Oz2] thus implies that, as  $Q' \cap M$  is non-injective, the map

$$\begin{aligned} \tilde{\Psi}_Q : B \otimes C &\rightarrow \mathbb{B}(L^2(M, \varphi)) \\ \sum_{k=1}^n a_k \otimes x_k &\mapsto \Psi_Q \left( \sum_{k=1}^n a_k x_k \right) \end{aligned}$$

is not continuous with respect to the minimal tensor norm. In turn, Lemma 2.2 above implies that there must be an index  $i$  such that  $D_i \not\subseteq \ker(\Psi_Q)$ .

By continuity of  $\Psi_Q$ , there must be a finite rank projection  $f \in \mathbb{B}(K_i)$  such that  $x = \Psi_Q(f \otimes 1) \neq 0$ , where we have identified  $L^2(M, \varphi)$  with  $K_i \otimes H_i$  via  $V_i$ . Note that  $x$  commutes with the right  $M_i$ -action, and so  $x \in (\mathbb{B}(K_i) \bar{\otimes} M_i) \cap Q'$ . Furthermore,  $\text{Tr} \otimes \tau_{M_i}(x) \leq \text{Tr}(f) < \infty$ , and so by taking a suitable spectral projection  $e$  of  $x$  we get that  $eL^2(M, \varphi)$  is a nonzero  $Q$ - $M_i$  bimodule with  $\dim_{M_i} eL^2(M, \varphi) < \infty$ .

Thus, by Proposition 3.2, there are nonzero projections  $q \in Q$  and  $p \in M_i$ , a  $*$ -homomorphism  $\theta : qQq \rightarrow pM_i p$ , and a nonzero partial isometry  $v \in M$  such that  $v^*v \leq p$ ,  $vv^* \leq q$  and for all  $a \in qQq$  we have

$$av = v\theta(a).$$

Note that this implies that  $vv^* \in (qQq)' \cap qMq = q(Q' \cap M)q$  and so we can write  $vv^* = qq'$  for  $q' \in Q' \cap M$ . Furthermore, this also implies that  $v^*v \in \theta(qQq)' \cap pM_i p$ , and so by Lemma 2.3 we have that  $v^*v \in pM_i p$ . Thus for all  $a \in Q$ ,

$$v^*av = v^*v\theta(qaq) \in M_i.$$

Also, note that all of this remains true after trimming  $q$  and  $q'$  (and  $v$ ,  $\theta$ , in turn). So, by restricting  $q$  (and hence  $v$  and  $\theta$ ) we may assume that  $\tau_Q(q) = \frac{1}{n}$  for some integer  $n$ . Then, by restricting  $q'$  (and hence  $v$ ), we may assume that  $\tau_{M_i}(v^*v) = \frac{1}{mn}$  for some integer  $m$ . We can then find  $n$  partial isometries  $u_i \in Q$  and  $w_i \in M_i$ , respectively, such that  $\sum_i u_i u_i^* = 1$ ,  $u_i^* u_i = q$ ,  $\sum_i w_i^* w_i \leq 1$ , and  $w_i w_i^* = v^*v$ . Then, letting  $w = \sum_i u_i v w_i$  we get that for  $a \in Q$ ,

$$w^*aw = \sum_i \sum_j w_i^* v^* u_i^* a u_j v w_j \in M_i$$

while

$$ww^* = \sum_i u_i q' u_i^* = q' \in Q' \cap M.$$

Note that  $\tau_{M_i}(w^*w) = \frac{1}{m}$ . If  $Q' \cap M$  is infinite, then by repeating an argument like the one above, we can enlarge  $w$  to a partial isometry  $u$  with  $u^*u = 1$ ,  $uu^* \in Q' \cap M$  and such that  $u^*Qu \subseteq M_i$ . Furthermore, if  $Q' \cap M$  is type III (or if  $q'$  is an infinite projection and  $Q' \cap M$  is type  $\text{II}_\infty$ ) we can easily dilate  $u$  to a unitary.

Otherwise, if  $Q' \cap M$  is type  $\text{II}_1$  we have two cases. First, if  $\tau_{Q' \cap M}(q') \leq \frac{1}{m}$ , then we can proceed precisely as in the infinite situation and obtain  $u$  with the same properties. If on the other hand  $\tau_{Q' \cap M}(q') > \frac{1}{m}$ , we can restrict  $w$  and assume that  $\tau_{Q' \cap M}(ww^*) = \frac{1}{m} \geq \tau_{M_i}(w^*w)$ . Then, proceeding as above we can enlarge  $w$  to a partial isometry  $u$  with  $uu^* = 1$  and such that  $u^*Qu \subseteq M_i$ .  $\square$

The following lemma follows easily from some well-known results of Popa, see A.1.1 and A.1.2 of [Po]. The proof in the case that the ambient algebra  $M$  is finite can be found as Proposition 13 of [OP], and the general case follows by the exact same argument where one replaces trace-preserving conditional expectations with those that preserve a state  $\varphi$  on  $M$  that centralizes  $N$ .

**Lemma 4.2.** *Let  $N \subseteq M$  be an inclusion of factors with  $N$  type  $\text{II}_1$  and in the range of a normal conditional expectation  $E_N : M \rightarrow N$ . Then there is an injective  $\text{II}_1$  factor  $Q \subseteq N$  such that  $Q' \cap M = N' \cap M$ .*

**Corollary 4.3.** *Let  $M_i$ , for  $i \in \{1, \dots, l\}$ , be semireact  $\text{II}_1$  factors with faithful normal states  $\varphi_i$ , and let  $(M, \varphi) = \ast_{i=1}^l (M_i, \varphi_i)$  be their reduced free product. Suppose that  $N \subseteq M$  is a non-prime non-injective  $\text{II}_1$  factor with a normal conditional expectation  $E_N : M \rightarrow N$ . Then there is an index  $i$  and a maximal partial isometry  $u$  such that  $u^*Nu \subseteq M_i$ .*

Proof: Write  $N = N_1 \otimes N_2$  for  $\text{II}_1$  factors  $N_i$  with  $N_2$  non-injective. By Lemma 4.2, there is an injective  $\text{II}_1$  factor  $Q \subseteq N_1$  such that  $Q' \cap M = N_1' \cap M$ . Note that  $N_2 \subseteq Q' \cap M$ , and thus  $Q' \cap M$  is non-injective. Furthermore,  $Q$  is in the range of a normal conditional expectation

from  $M$ . Thus, by Theorem 4.1, there is an index  $i$  and a maximal partial isometry  $u$  such that  $uu^* \in Q' \cap M = N'_1 \cap M$ , and  $u^*Qu \subseteq M_i$ .

Now, note that if  $x \in N_2$  and  $q \in Q$ , then

$$(u^*xu)(u^*qu) = u^*xqu = u^*qxu = (u^*qu)(u^*xu)$$

and so by Lemma 2.3,  $u^*xu \in M_i$ .

Furthermore, if  $y \in N_1$  and  $x \in N_2$ , then as  $uu^* \in N'_1 \cap M$ ,

$$(u^*yu)(u^*xu) = u^*yxu = u^*xyu = (u^*xu)(u^*yu)$$

and so by Lemma 2.3 again,  $u^*yu \in M_i$ .

Thus,  $u^*N_1N_2u = u^*N_1uu^*N_2u \subseteq M_i$ , and so  $u^*Nu \subseteq M_i$ .  $\square$

**Corollary 4.4.** *Let  $\{M_i\}_{i=1}^m$ ,  $\{N_j\}_{j=1}^n$  be semiexact non-prime non-injective  $\text{II}_1$  factors with faithful normal states  $\varphi_i$ ,  $\psi_j$ , and let  $M, \varphi$ ,  $N, \psi$  be their respective reduced free products. If  $M \simeq N$ , then  $m = n$  and after reordering indices  $M_i$  is isomorphic to an amplification of  $N_i$ .*

Proof: Identify  $M$  and  $N$  via the implied isomorphism. By the proof of Corollary 4.3, for each index  $i$  there is an injective  $\text{II}_1$  factor  $Q_i \subseteq M_i$ , an index  $\alpha(i)$ , and a maximal partial isometry  $u_i \in M$  such that  $u_iu_i^* \in Q'_i \cap M$  and  $u_i^*M_iu_i \subseteq N_{\alpha(i)}$ . Similarly for each index  $j$  there is an injective  $\text{II}_1$  factor  $P_j \subseteq N_j$ , an index  $\beta(j)$ , and a maximal partial isometry  $v_j \in M$  such that  $v_jv_j^* \in P'_j \cap M$  and  $v_j^*N_jv_j \subseteq M_{\beta(j)}$ .

Consider the projections  $p_i = u_i^*u_i$  and  $q_i = v_{\alpha(i)}v_{\alpha(i)}^*$  in  $N_{\alpha(i)}$  (Lemma 2.3 implies that  $q_i \in N_{\alpha(i)}$ ). First, by restricting the target projection of  $u_i$  in  $Q'_i \cap M$ , we may assume that  $\tau_{N_{\alpha(i)}}(p_i) \leq \tau_{N_{\alpha(i)}}(q_i)$ . Furthermore, by factoriality of  $N_{\alpha(i)}$  we can rotate  $v_i$  by a partial isometry to assure that  $p_i \leq q_i$ . Then, if we let  $w_i = u_iv_{\alpha(i)}$ ,  $w_i$  is a partial isometry such that  $w_i^*Q_iw_i \subseteq M_{\beta(\alpha(i))}$ , and  $w_iw_i^* = u_iu_i^* \in Q'_i \cap M$ . Thus, by Lemma 2.3,  $\beta(\alpha(i)) = i$  and  $w_i \in M_i$ . So,  $u_i$  implements an isomorphism of  $(u_iu_i^*)A(u_iu_i^*)$  with  $pN_{\alpha(i)}p$ .

A completely symmetric argument shows that  $\alpha(\beta(j)) = j$ .  $\square$

Note that this last corollary implies that, for example, if  $d_i \in M = L(\mathbb{F}_n) \otimes L(\mathbb{F}_m)$  are positive and  $\varphi_i(\cdot) \sim \tau_M(d_i \cdot)$ , then  $\star_{i=1}^l(M, \varphi_i)$  are non-isomorphic for different  $l$ .

## REFERENCES

- [Oz] N. Ozawa, *A Kurosh type theorem for type  $\text{II}_1$  factors*. Int. Math. Res. Not. (2006), doi:10.1155/IMRN/2006/97560
- [Oz2] N. Ozawa, *Solid von Neumann algebras*. Acta Math. **192** (2004), 111–117.
- [OP] N. Ozawa and S. Popa, *Some prime factorization results for type  $\text{II}_1$  factors*. Invent. Math. **156** (2004), 223–234.
- [Po] S. Popa, *Classification of amenable subfactors of type II*. Acta Math. **172** (1994) 163–255.
- [Po2] S. Popa, *Orthogonal pairs of  $*$ -subalgebras in finite von Neumann algebras*. J. Operator Theory **9** (1983), 253–268.
- [Po3] S. Popa, *Strong rigidity of  $\text{II}_1$  factors coming from malleable actions of weakly rigid groups, I*. Invent. Math. **165** (2006), 369–408.
- [Ta] M. Takesaki, *Theory of Operator Algebras II*. Encyclopedia of Mathematical Sciences **125**, Springer–Verlag, 2000.
- [Vo] D. Voiculescu, *Symmetries of some reduced free product  $C^*$ -algebras*. Operator Algebras and Their Connections with Topology and Ergodic Theory, Lecture Notes in Mathematics, vol. 1132, Springer–Verlag, 1985, 556–588.