

Generating Static, Spherically Symmetric Black-holes in Third Order Lovelock Gravity.

S. Habib Mazharimousavi,^{*} O. Gurtug,[†] and M. Halilsoy[‡]

Department of Physics, Eastern Mediterranean University,

G. Magusa, North Cyprus, Mersin 10 - Turkey.

Abstract

Generalization of the Salgado's theorem (*Class. Quant. Grav.* **20** (2003) 4551) to generate static, spherically symmetric black-hole solutions in higher dimensional ($N \geq 7$) third order Lovelock gravity is presented. Particular limits, such as Gauss-Bonnet (GB) and/or Einstein-Hilbert (EH) in any dimension N yield all the solutions known to date with an energy-momentum. The occurrence of naked singularities are studied and it is found that, it remains nonsingular in quantum mechanical sense when it is probed with quantum test particles. Finally we extend the theorem to cover Bertotti-Robinson (BR) type solutions which arise from the extremal black holes.

^{*}Electronic address: habib.mazhari@emu.edu.tr

[†]Electronic address: ozay.gurtug@emu.edu.tr

[‡]Electronic address: mustafa.halilsoy@emu.edu.tr

I. INTRODUCTION

One of the most interesting features of the string theory is to provide an arena for higher dimensional space-times. String theory together with higher dimensions supports also the description of objects known as branes. There is no doubt that, the most intriguing solution in higher dimensional space-times is the one that is associated with black holes. The pioneering work in this regard belong to Boulware and Deser [1]. They obtained the most general static black hole solutions in Einstein - Gauss - Bonnet (EGB) theory. Recent studies show that there is a growing interest to find black-hole solutions in higher dimensional gravity. This task is accomplished by the use of the most general action that describes black-hole solutions in Einstein-Lovelock theory[2]. This is the most general theorem that hosts higher order invariants in particular combinations so that field equations remain second order and therefore do not contain ghosts. Physical properties of the Einstein - Lovelock theory that admits black hole solutions is analysed in detail in the Ref.[3]. Black hole solutions in Lovelock theory is important because the higher order curvature terms contributes to the inner structure of black holes. For example, in 4- dimensional general relativity, inner (Cauchy) and outer (event) horizons arise only in the Reissner-Nordstrom black hole in which the corresponding space-time admits two Killing vectors orthogonal to each other. However, it is demonstrated in [12] that, two horizons may arise in the Einstein - Lovelock theory without the matter fields. Another remarkable aspects of the Lovelock theory is to provide the existence of topological black holes in which the curvature constant is not positive[4, 5, 6].

In general theory of relativity, static, spherically symmetric space-times constitute one of the simplest and tractable model among the others as far as the analytic, exact solutions are concerned. Recently, Salgado [7], has developed a theorem, to generate exact black-hole solutions by imposing some conditions on the energy-momentum tensor. This theorem has been extended to the higher dimensional spacetimes [8] in which Schwarzschild, Reissner-Nordstrom and global monopole solutions in higher dimensions are particular cases. Later on, another extension of the theorem has been introduced to find exact radiative (dynamic) black-hole solutions [9]. This can be interpreted as the dynamic extension of the Salgado's theorem. In this study, the generalized, spherically symmetric Type II fluid solution is obtained. Many known solutions are shown to be the particular limits of this generalized solution. The extension of this non-static radiative solution in the EGB and Born-Infeld

nonlinear electrodynamic theory is also considered in [10]. More recently, it has been shown that the N -dimensional extension of radiative, dynamic black-hole solutions are possible [11].

In this paper, we wish to extend Salgado's theorem to the arbitrary dimensional, third order Lovelock gravity coupled with matter fields. In our case, matter fields couple to the system through an arbitrary parameter k . It is shown that, our general solution includes the well-known solutions in particular limits, namely, in the GB and Einstein limits in any lower dimensional solutions. Besides all of these, we present a black hole solution with interesting properties that, depending on the constant parameters α_2 , α_3 and C , one or two horizons may develop. We discussed also the behaviour of the naked singularity when it is probed with quantum particles. It is found that the classical timelike curvature singularity at the origin remains nonsingular quantum mechanically.

The paper is organised as follows: In Sec. *II* we give a brief summary of the third order Lovelock gravity. We present the static spherically symmetric solution in third order Lovelock theory in general form and explicit solutions in 7- dimension is discussed in Sec. *III*. In Sec. *IV* we introduce the general form of the Bertotti-Robinson (BR) type solutions in any dimension. The paper ends with a conclusion in Sec. *V*.

II. N -DIMENSIONAL THIRD ORDER EINSTEIN - LOVELOCK GRAVITY.

The action describing N -dimensional third order Einstein - Lovelock gravity coupled with matter fields is given by;

$$S = \int dx^N \sqrt{-g} \{L_{EH} + \alpha_2 L_{GB} + \alpha_3 L_L\} + S_{matter}, \quad (1)$$

where L_{EH} , L_{GB} and L_L represents the Einstein-Hilbert, Gauss-Bonnet and third order Lovelock Lagrangians, respectively, and S_{matter} is the action of the matter fields coupled to this system. We recall that for $S_{matter} = 0$, L_{GB} and L_L terms become meaningful only for $N \geq 5$ and $N \geq 7$, respectively. Our spherically symmetric metric ansatz is given in Eq. (4) below. Variation of the action with respect to the metric tensor $g_{\mu\nu}$ yields the equations of motion in the form

$$G_{\mu\nu}^{EH} + \alpha_2 G_{\mu\nu}^{GB} + \alpha_3 G_{\mu\nu}^L = T_{\mu\nu}, \quad (2)$$

where $T_{\mu\nu}$ is the energy-momentum tensor representing the matter fields. Among the others, rr -component of the Eq. (2) is the simplest one and can be written as,

$$T_r^r = \frac{n}{2r^6} \{ [r^5 - 2\tilde{\alpha}_2 r^3 g(r) + 3\tilde{\alpha}_3 r g(r)^2] g'(r) + (n-1) r^4 g(r) - (n-3) \tilde{\alpha}_2 r^2 g(r)^2 + (n-5) \tilde{\alpha}_3 g(r)^3 \} \quad (3)$$

in which a prime denotes derivative with respect to r , $g(r) = f(r) - 1$, $\tilde{\alpha}_2 = (n-1)(n-2)\alpha_2$, $\tilde{\alpha}_3 = (n-1)(n-2)(n-3)(n-4)\alpha_3$ and $n = N - 2$.

III. THE STATIC SOLUTION.

Generalization of the Salgado's theorem to the third order Lovelock theory together with arbitrary matter fields is as follows:

Theorem 1 *Let (M, g_{ab}) be a N -dimensional space-time with $\text{sign}(g_{ab}) = N-2$, $N \geq 3$, such that : (1) it is static and spherically symmetric, (2) it satisfies the Einstein field equations, (3) the energy momentum tensor T^{ab} satisfies the conditions $T_r^r = T_t^t$ and $T_{\theta_i}^{\theta_i} = kT_r^r$ ($1 \leq i \leq n-2$, $k = \text{const} \in \mathbb{R}$), (4) it possess a regular Killing horizon or a regular origin. Then, the metric of the space-time is given by*

$$ds^2 = -f(r)dt^2 + f(r)^{-1}dr^2 + r^2 d\Omega_n^2, \quad (4)$$

where

$$d\Omega_n^2 = d\theta_1^2 + \sum_{i=2}^n \prod_{j=1}^{i-1} \sin^2 \theta_j d\theta_i^2, \quad 0 \leq \theta_n \leq 2\pi, \quad 0 \leq \theta_i \leq \pi, \quad 1 \leq i \leq n-1, \quad (5)$$

and the energy momentum tensor in general is in the following form

$$T_{b(Diag.)}^a = \frac{C}{r^{n(1-k)}} [1, 1, k, \dots, k], \quad (6)$$

in which C is an integration constant.

The general solution for the Eq.(3) in any dimension is obtained for two different classes as follows

$$f(r) = 1 + \frac{\tilde{\alpha}_2}{3\tilde{\alpha}_3} r^2 - 2G \left(1 - \frac{\tilde{\alpha}_2^2}{3\tilde{\alpha}_3}\right) r^{n-1} \xi_2^{-1/3} + \frac{1}{3\tilde{\alpha}_3 G r^{n-5}} \xi_2^{1/3}, \quad \text{for } k \neq -\frac{1}{n}, \quad (7)$$

in which,

$$\begin{aligned} \xi_2 &= r^{n-9} \left[216C\tilde{\alpha}_3^2 r^{n(1+k)} + 8G\tilde{\alpha}_2 \left(\tilde{\alpha}_2^2 - \frac{9}{2}\tilde{\alpha}_3 \right) r^{2n} + \right. \\ &\quad \left. 12\tilde{\alpha}_3 r^4 \left(\sqrt{3\xi_1} - 9\frac{mG\tilde{\alpha}_3}{n} r^{n-5} \right) \right] G^2, \\ \xi_1 &= 108C^2\tilde{\alpha}_3^2 r^{2n(1+k)-8} + \frac{G}{n} \left\{ r^{n(3+k)-8} \left(8n\tilde{\alpha}_2^2 C \left(\tilde{\alpha}_2^2 - \frac{9}{2}\tilde{\alpha}_3 \right) + 108mC\tilde{\alpha}_3^2 r^{-n-1} \right) - \right. \\ &\quad \left. \frac{G}{n} r^{2(n-2)} \left[4mn\tilde{\alpha}_2 r^{n-2} \left(\tilde{\alpha}_2^2 - \frac{9}{2}\tilde{\alpha}_3 \right) + n^2 (\tilde{\alpha}_2^2 - 4\tilde{\alpha}_3) r^{2(n-2)} - 27\tilde{\alpha}_3^2 m^2 r^{-6} \right] \right\}, \\ G &= n(nk + 1). \end{aligned} \quad (8)$$

The other possible solution is

$$f(r) = 1 + \frac{\tilde{\alpha}_2}{3\tilde{\alpha}_3} r^2 - 2 \left(1 - \frac{\tilde{\alpha}_2^2}{3\tilde{\alpha}_3}\right) r^{n+5} \zeta_2^{-1/3} + \frac{1}{6\tilde{\alpha}_3 n r^{n+1}} \zeta_2^{1/3}, \quad \text{for } k = -\frac{1}{n}, \quad (9)$$

where

$$\begin{aligned} \zeta_2 &= n^2 r^{2(n+3)} \left[216\tilde{\alpha}_3^2 r^2 \left(C \ln r - \frac{m}{2} \right) + 8\tilde{\alpha}_2 \left(\tilde{\alpha}_2^2 - \frac{9}{2}\tilde{\alpha}_3 \right) r^{n+3} + 12\tilde{\alpha}_3 \sqrt{3\xi_1} \right], \\ \zeta_1 &= -n^2 (\tilde{\alpha}_2^2 - 4\tilde{\alpha}_3) r^{2(n+3)} + 27(m - 2C \ln r) \times \\ &\quad \left[-\frac{4}{27} n\tilde{\alpha}_2 r^{n+5} \left(\tilde{\alpha}_2^2 - \frac{9}{2}\tilde{\alpha}_3 \right) + r^4 \tilde{\alpha}_3^2 (m - 2C \ln r) \right]. \end{aligned} \quad (10)$$

We note that in $(n+2)$ dimensions for $k = -1$ and $C = -q^2$ this solution overlaps with the Dehghani's[12] solution. Also by setting $k = \frac{n-4}{n}$ and $C = -\frac{1}{2}n(n-1)Q^2$ one recovers the Einstein - Yang - Mills (EYM) black hole solutions [13]. Another case is to take $k = 1$ which leads us to a perfect-fluid type energy momentum tensor with constant energy density (ρ), tension (τ) and pressure (p), such that $C = \rho = \tau = p$.

We observe from Eqs.(9-10) that, in any dimension it is possible to have a logarithmic term. This type of solutions are encountered in 5-dimensional EYM [13] theories. As we demonstrate in Eqs.(9-10) this type of solutions are also possible in Einstein- Third Order Lovelock gravity. However, its physical interpretation higher than 5- dimensional case needs further investigation.

A. Properties of the General Solution.

Since the obtained solution is too complicated for physical interpretation, we prefer to constraint the constants $\tilde{\alpha}_2$ and $\tilde{\alpha}_3$ in such a way that $3\tilde{\alpha}_3 = \tilde{\alpha}_2^2$. This choice simplifies the above results to,

$$\begin{aligned}
f(r) &= 1 + \frac{1}{2\tilde{\alpha}_2^2 G r^{n-5}} \left(\xi_2^{1/3} + 2G\tilde{\alpha}_2 r^{n-3} \right) \quad \text{for } k \neq -\frac{1}{n} \\
\xi_2 &= 4\tilde{\alpha}_2^2 \left[6r^{n(2+k)-9} C \tilde{\alpha}_2^2 - 3\tilde{\alpha}_2^2 \frac{G}{n} m r^{2(n-5)} - G\tilde{\alpha}_2 r^{3(n-3)} + \sqrt{3\xi_1} r^{n-5} \right] G^2 \\
\xi_1 &= 3 \left\{ 4r^{2(n-4+nk)} C^2 \tilde{\alpha}_2^2 + [4\tilde{\alpha}_2 C r^{n(k+2)} \left(\frac{n r^{n-8}}{3} + r^{-9} \tilde{\alpha}_2 m \right) + \right. \\
&\quad \left. \left(\frac{2m}{3} n \tilde{\alpha}_2 r^{3n-9} + \tilde{\alpha}_2^2 r^{2n-10} m^2 + \frac{n^2 r^{4(n-2)}}{9} \right) (nk+1) \right\} (nk+1).
\end{aligned} \tag{11}$$

and

$$\begin{aligned}
f(r) &= 1 + \frac{r^2}{\tilde{\alpha}_2} + \frac{1}{2n\tilde{\alpha}_2^2 r^{n+1}} \zeta_2^{1/3}, \quad \text{for } k = -\frac{1}{n}, \\
\zeta_2 &= 4n^2 \tilde{\alpha}_2^2 r^{2(n+3)} \left[6\tilde{\alpha}_2^2 r^2 \left(C \ln r - \frac{m}{2} \right) - n\tilde{\alpha}_2 r^{n+3} + \sqrt{3\zeta_1} \right], \\
\zeta_1 &= \frac{\tilde{\alpha}_2^2}{3} \left\{ n^2 r^{2(n+3)} + 9(m - 2C \ln r) \tilde{\alpha}_2 \left[\frac{2}{3} n r^{n+5} + r^4 \tilde{\alpha}_2 (m - 2C \ln r) \right] \right\},
\end{aligned} \tag{12}$$

respectively. The matter fields are coupled to the system through the constant parameters C and k . The inclusion of the matter fields must satisfy certain energy conditions as far as the physically acceptable solutions are concerned. These energy conditions are given in Appendix. According to these conditions our general solution satisfies the Dominant Energy Condition (DEC) and restricts the constant parameters as $C \leq 0$ and $-1 \leq k \leq 0$, while the causality condition bounds the parameter k further to $-1 \leq k \leq -\frac{1}{n}$.

We wish to underline the interesting special case when $k = \frac{1}{n}$. The general solution for this particular case is,

$$f(r) = 1 + \frac{r^2}{\tilde{\alpha}_2} \left\{ 1 - \sqrt[3]{1 - \frac{3C\tilde{\alpha}_2}{nr^{n-1}} + \frac{3m\tilde{\alpha}_2}{nr^{n+1}}} \right\}. \tag{13}$$

This is a black hole solution with radius of horizon r_h which can be found from $f(r_h) = 0$. This implies,

$$nr_h^{n-5}\tilde{\alpha}_2^2 + 3nr_h^{n-3}\tilde{\alpha}_2 + 3nr_h^{n-1} + 3Cr_h^2 - 3m = 0. \quad (14)$$

It is important to note that this solution (13) does satisfy the Weak Energy Condition (WEC), and therefore the limitations on the constant parameters are given by $C \leq 0$ and $k \leq 1$. This particular case will be investigated in detail in the following subsection

1. Seven - Dimensional Case.

As it was clarified in Ref.[12], the physically acceptable description of the solution obtained in the third order Lovelock gravity is to reduce the general solution to a seven - dimensional spacetime. The seven - dimensional spacetimes represent the most general Lagrangian producing second order field equations, as in the four - dimensional gravity the Einstein - Hilbert action constitutes the most general such Lagrangian. Hence the solution transforms to;

$$\begin{aligned} f(r) &= 1 + \frac{r^2}{\tilde{\alpha}_2} + \frac{\xi_2^{1/3}}{10(5k+1)\tilde{\alpha}_2^2} \quad \text{for} \quad k \neq -\frac{1}{5} \quad (15) \\ \xi_2 &= 100\tilde{\alpha}_2^2(5k+1)^2 \left[\sqrt{3\xi_1} + 6C\tilde{\alpha}_2^2 r^{5k+1} - (5k+1)(5r^6 + 3m\tilde{\alpha}_2) \right], \\ \xi_1 &= -12\tilde{\alpha}_2^2 \left\{ (5k+1) \left(m\tilde{\alpha}_2 + \frac{5}{3}r^6 \right) \left[C\tilde{\alpha}_2 r^{5k+1} - \frac{1}{4} \left(m\tilde{\alpha}_2 + \frac{5}{3}r^6 \right) (5k+1) \right] \right. \\ &\quad \left. - \tilde{\alpha}_2^2 C^2 r^{2(5k+1)} \right\}, \end{aligned}$$

This result generalizes the formerly obtained solutions for any choice of matter fields upon choosing specific values for k and C . For example, if we choose $k = -1$ and $C = -q^2$, we recover the solution obtained by Dehghani [12].

The other solution is

$$\begin{aligned} f(r) &= 1 + \frac{r^2}{\tilde{\alpha}_2} + \frac{\zeta_2^{1/3}}{10\tilde{\alpha}_2^2 r^6}, \quad \text{for} \quad k = -\frac{1}{5}, \quad (16) \\ \zeta_2 &= 100\tilde{\alpha}_2^2 r^{16} \left[6\tilde{\alpha}_2^2 r^2 \left(C \ln r - \frac{m}{2} \right) - 5\tilde{\alpha}_2 r^8 + \sqrt{3\zeta_1} \right], \\ \zeta_1 &= 3\tilde{\alpha}_2^2 r^4 \left[\frac{5}{3}r^6 + \tilde{\alpha}_2 (m - 2C \ln r) \right]^2. \end{aligned}$$

2. *The case for $k = \frac{1}{5}$.*

Another interesting case occurs for $k = \frac{1}{5}$: In order to study its physical properties we first look for the location of horizons. For this particular case, the metric function given in Eq.(13) becomes,

$$f(r) = 1 + \frac{r^2}{\tilde{\alpha}_2} \left\{ 1 - \sqrt[3]{1 - \frac{3C\tilde{\alpha}_2}{5r^4} + \frac{3m\tilde{\alpha}_2}{5r^6}} \right\}, \quad (17)$$

and the radius of horizon is obtained from Eq.(14) as

$$r_h = \sqrt{\frac{-(5\tilde{\alpha}_2 + C) \pm \sqrt{(5\tilde{\alpha}_2 + C)^2 + 60m - 100\tilde{\alpha}_2^2}}{10}}. \quad (18)$$

We wish to remind that, with this particular choice of k , the resulting solution satisfies only the WEC. This condition further implies that $C \leq 0$. These limitations induce three possible cases on the constant parameters C and $\tilde{\alpha}_2$ appeared in the Eq.(18).

Case 1: One possible case is when $5\tilde{\alpha}_2 + C = 0$. The location of the horizon from Eq.(18) is found as,

$$r_h = \sqrt{\frac{\sqrt{60m - 100\tilde{\alpha}_2^2}}{10}}. \quad (19)$$

Case 2: Another possibility is when $5\tilde{\alpha}_2 + C > 0$. As in the previous case there is only one horizon and its location is given by

$$r_h = \sqrt{\frac{-(5\tilde{\alpha}_2 + C) + \sqrt{(5\tilde{\alpha}_2 + C)^2 + 60m - 100\tilde{\alpha}_2^2}}{10}}. \quad (20)$$

For a physically acceptable solution, the mass of the black hole is constrained in each of the cases above as

$$60m \geq 100\tilde{\alpha}_2^2, \quad (21)$$

which imposes the critical mass value,

$$m_{critical} = \frac{5}{3}\tilde{\alpha}_2^2. \quad (22)$$

Case 3: For the case $5\tilde{\alpha}_2 + C < 0$; we have two horizons whose locations are given by,

$$r_+ = \sqrt{\frac{-(5\tilde{\alpha}_2 + C) + \sqrt{(5\tilde{\alpha}_2 + C)^2 + 60m - 100\tilde{\alpha}_2^2}}{10}}, \quad (23)$$

$$r_- = \sqrt{\frac{-(5\tilde{\alpha}_2 + C) - \sqrt{(5\tilde{\alpha}_2 + C)^2 + 60m - 100\tilde{\alpha}_2^2}}{10}}. \quad (24)$$

In this particular case the mass of the black hole is constrained by

$$\frac{5}{3}\tilde{\alpha}_2^2 - \frac{1}{60} |5\tilde{\alpha}_2 + C|^2 \leq m \leq \frac{5}{3}\tilde{\alpha}_2^2. \quad (25)$$

Therefore the critical mass for this particular case is $m_{critical} = \frac{5}{3}\tilde{\alpha}_2^2 - \frac{1}{60} |5\tilde{\alpha}_2 + C|^2$. This constraint condition implies that the naked singularity develops at $r = 0$, if

$$m < m_{critical}. \quad (26)$$

The type of the singularity at $r = 0$ can be found, if one introduce a new radial coordinate $r_* = \int \frac{dr}{f}$, and take its limit as $r \rightarrow 0$. If r_* takes a finite value as $r \rightarrow 0$, the singularity is timelike. If $r_* = -\infty$ as $r \rightarrow 0$, then it is null. In our case, r_* is finite, therefore it has a timelike character. Since the spacetime is static and the type of the singularity is timelike, in the following subsection we shall study the behaviour of this singularity in quantum mechanical sense.

The Hawking temperature at the radius of horizon is given by,

$$T_H = \frac{\kappa}{2\pi} = \frac{1}{4\pi} |f'(r_h)| = \frac{\sqrt{A}}{30\tilde{\alpha}_2 B^{2/3}} \left\{ B^{2/3} + 5C\tilde{\alpha}_2 - \frac{A^2}{32400} \right\}, \quad (27)$$

where κ denotes the surface gravity and

$$\begin{aligned} B &= -\frac{A^3}{5832000} + \frac{AC\tilde{\alpha}_2}{12} - 75m\tilde{\alpha}_2, \\ A &= -450\tilde{\alpha}_2 - 90C + 30\sqrt{S}, \\ S &= -75\tilde{\alpha}_2^2 + 90C\tilde{\alpha}_2 + 9C^2 + 180m. \end{aligned} \quad (28)$$

3. Naked Singularities

In this subsection, we wish to emphasize another important property of the Lovelock theory. We stated previously that if the mass of the black hole $m < m_{critical}$, no horizon forms so that the singularity at $r = 0$ becomes naked with a timelike character. In classical 4- dimensional general relativity, this is a curvature singularity, indicating timelike geodesic incompleteness. However, our main concern here is to analyse this naked singularity when probed with quantum test particles. In other words, we are aiming to see whether this singular spacetime (in classical sense) is nonsingular or not quantum mechanically. To achieve this we adopt the method which was initiated by Wald [15] and developed by Horowitz and Marolf [16], for static spacetimes having timelike curvature singularities. This method states that a spacetime is quantum mechanically nonsingular if the time evolution of any wave packet is uniquely determined by the initial wave function. The method is briefly as follows:

A scalar quantum particle with mass M is described by the Klein-Gordon equation $(\nabla^\mu \nabla_\mu - M^2) \psi = 0$. This equation can be written by splitting the temporal and spatial portion as $\frac{\partial^2 \psi}{\partial t^2} = -A\psi$, such that the spatial operator A is defined by $A = -\sqrt{f} D^i (\sqrt{f} D_i) + f M^2$, where $f = -\xi^\mu \xi_\mu$ with ξ^μ the timelike Killing field, while D_a is the spatial covariant derivative defined on the static slice Σ . The method requires that the spatial operator A is essentially self-adjoint. That is, the extension of the operator A_E is always possible with the condition that this extension must be unique. Then the Klein-Gordon equation for a free relativistic particle satisfies $i \frac{\partial \psi}{\partial t} = \sqrt{A_E} \psi$, with the solution $\psi(t) = \exp(it\sqrt{A_E}) \psi(0)$. The ambiguity occurs in these equations if A is not essentially self-adjoint. Consequently, a sufficient condition for the operator A to be essentially self-adjoint is to satisfy

$$A\psi \pm i\psi = 0. \quad (29)$$

The separable solution is assumed for Eq.(29) in the form of $\psi = \phi(r)Y(\text{angles})$. The radial part of the Eq(29) becomes,

$$\frac{\partial^2 \phi}{\partial r^2} + \frac{1}{fr^5} \frac{\partial(fr^5)}{\partial r} \frac{\partial \phi}{\partial r} - \frac{c}{fr^2} \phi - \frac{M^2}{f} \phi \pm i \frac{\phi}{f^2} = 0, \quad (30)$$

in which $c \geq 0$ is the eigenvalue of the Laplacian on the 5-sphere. Equation (30) can be solved with the help of Fuchsian equation [17] by assuming the massless case (i.e $M = 0$) and

ignoring the term $\pm i \frac{\phi}{f^2}$ (since it is negligible near the origin, $r = 0$). The Fuchsian equation is $\frac{\partial^2 \phi}{\partial r^2} + r^{-1} p(r) \frac{\partial \phi}{\partial r} + r^{-2} q(r) \phi = 0$, such that $p(r)$ and $q(r)$ are analytic at the origin. This equation admits solution in the form of $\phi(r) = r^\beta F(r)$, where $F(r)$ is analytic function and β is a complex number that solves the indicial equation $\beta(\beta - 1) + \beta p(0) + q(0) = 0$. Substituting Eq.(17) in Eq(30), we find that $p(0) = 5$ and $q(0) = -\frac{c}{1 - \left(\frac{3m}{5\alpha_2}\right)^{1/3}}$. For $c = 0$, one of the two solutions to indicial equation, solves the Eq.(30) and the resulting solution diverges as fast as $|\phi(r)|^2 = r^{-8}$. This solution always has infinite norm near $r = 0$ since

$$\langle \phi | \phi \rangle = \int \frac{|\phi(r)|^2 r^5}{f} dr. \quad (31)$$

Consequently, $\phi(r)$ fails to be square integrable near the origin. According to this result we conclude that the timelike curvature singularity at the origin turns out to be quantum mechanically nonsingular when probed with quantum test particles. Similar analysis is also shown in Ref. [3] for the 5- dimensional Boulware-Deser metric which also remains regular when tested by quantum probes. A similar proof of quantum regularity applies for $N > 5$ as well.

IV. GENERATING BR TYPE SOLUTIONS IN EGB THEORY

Closely related with the black hole solutions is the class of BR type solutions in GB gravity. This class arises as a limiting case of extremal black holes so that an analogous theorem can be stated to cover this class as well. In $N = 4$, the BR solution is the unique, conformally flat EM solution. In higher dimensions ($N > 4$) we found that both, conformal and asymptotical flatness fail[14]. Being almost as important as black holes, specifically in supergravity, we wish to present conditions on the energy momentum involved in order to generate solutions of BR form in the EYMGB theory.

To this end we adopt the metric in the form

$$ds^2 = -f dt^2 + f^{-1} dr^2 + h^2 d\Omega_{N-2}^2, \quad (32)$$

where h is a constant to be specified and $f = f(r)$ is a function to be found. The energy momentum tensor is assumed in the form

$$T_b^a = C [-1, -1, k, k, \dots] \quad (33)$$

where C and k are constants that characterize the matter fields. The Einstein's tensor in N -dimension is given by,

$$G_b^a = \left[-\frac{(N-3)(N-2)}{2h^2}, -\frac{(N-3)(N-2)}{2h^2}, -\frac{(N-3)(N-4) + f''h^2}{2h^2}, \dots \right]. \quad (34)$$

in which the higher terms repeat the third one. From the Einstein's equation $G_{ab} = T_{ab}$, we have,

$$C = \frac{(N-3)(N-2)}{2h^2}, \quad (35)$$

and

$$k = \frac{f''h^2 - (N-3)(N-4)}{2Ch^2}. \quad (36)$$

Solving the Eq.(36) with the help of Eq.(35) we get,

$$f(r) = \frac{1}{2} [(2Ck + (N-3)(N-4))] r^2 + C_1 r + C_2, \quad (37)$$

where C_1 and C_2 are integration constants. This general result includes some of the well-known solutions for particular choices of C and k . It can be anticipated that for $C_1 \neq 0 \neq C_2$, $f(r) = 0$, admits roots resulting in non-asymptotically flat black hole solutions. Beside this, for $C = q^2$ (q is the electric charge) and $k = 1$ it corresponds to N - dimensional BR like solution in the Einstein-Maxwell theory. Note that we choose the integration constants, $C_1 = C_2 = 0$ for this particular case. Another interesting solution is obtained for the EYM theory if one takes $C = \frac{(N-3)(N-2)}{2Q^2}$ and $k = -\frac{N-6}{N-2}$.

V. CONCLUSION

In this paper, we have extended the Salgado's theorem to generate static, spherically symmetric black hole solutions in higher dimensional ($N \geq 7$) third order Lovelock gravity with matter fields. We have shown explicitly that our general solution recovers formerly obtained solutions in particular limits. We have also obtained a new class of black hole

solutions in 7 - dimensions with arbitrary matter fields. The arbitrary matter fields couple to the system through the constant parameters C and k . It is shown that these parameters are restricted as a result of energy conditions.

Firstly we generate N - dimensional black hole solutions in general form. However, because of the complexity of the resulting expressions, we constrain, the parameters $\tilde{\alpha}_2$ and $\tilde{\alpha}_3$ as in Ref. [12], so that we obtain solutions that overlap with the known solutions in 7- dimensions.

A new black hole solution in 7-dimensions with $k = \frac{1}{5}$ is studied in detail. It is shown that depending upon on the values of C and $\tilde{\alpha}_2$, one or two horizons may develop. If the mass of the black hole $m < m_{critical}$, the naked singularity arises at the origin. In classical 4- dimensional general relativity, this singularity is a timelike curvature singularity. The structure of this singularity is further analysed with quantum test particles according to the method developed by Horowitz and Marolf [16]. Our analyses has revealed that, although $r = 0$ is singular in classical sense, it becomes nonsingular when probed with quantum test particles. In the light of this remarkable result, the Lovelock theory of gravity is important in the sense that, it provides an arena to investigate the contribution of higher order curvature terms at short distances; especially for the solutions that incorporate black holes.

Secondly, we extend the theorem to cover Bertotti-Robinson type solutions which arise from the extremal black holes.

-
- [1] D. Boulware and S. Deser, *Phys. Rev. Lett.* **55** 2656 (1985).
 - [2] D. Lovelock, *J. Math. Phys.* **12**, 498 (1971).; N. Deruelle and L. Farina-Busto, *Phys. Rev. D* **41**, 3696 (1990).; G. A. MenaMarugan, *ibid.* **46**, 4320 (1992); 4340 (1992).
 - [3] C. Garraffo and G. Giribet, *Mod. Phys. Lett.* **A23**, 1801 (2008).
 - [4] R. Aros, R. Troncoso and J. Zanelli, *Phys. Rev.* **D63** 084015 (2001), [arXiv:hep-th/0011097].
 - [5] Rong-Gen Cai, *Phys. Rev.* **D65** 084014 (2002), [arXiv:hep-th/0109133].
 - [6] M. H. Dehghani, N. Alinejadi and S.H. Hendi, *Phys. Rev.* **D77** 104025 (2008).
 - [7] M. Salgado, *Class. Quant. Grav.* **20** 4551 (2003).
 - [8] E. Gallo, *Gen. Rel. Grav.* **36** 1463 (2004).
 - [9] A. K. Dawood and S. G. Ghosh, *Phys. Rev.* **D70** 104010 (2004).
 - [10] A.E. Dominguez and E. Gallo, *Phys. Rev.* **D73** 064018 (2006).

- [11] S. G. Ghosh and A. K. Dawood, *Gen. Rel. Grav.* **40** 9 (2008).
- [12] M. H. Dehghani and M. Shamirzaie, *Phys. Rev.* **D72** 124015 (2005).
- [13] S. H. Mazharimousavi and M. Halilsoy, *Phys. Rev.* **D76** 087501 (2007).; S. H. Mazharimousavi and M. Halilsoy, *Phys. Lett.* **B659** 471 (2008).
- [14] S. H. Mazharimousavi and M. Halilsoy, arXiv:0801.2110.
- [15] R. M. Wald, *J. Math. Phys.*, 21, 2802 (1980).
- [16] G. Horowitz and D. Marolf, *Phys. Rev.* **D52** 5670 (1995).
- [17] S. Hassani, *Foundations of Mathematical Physics*, (Allyn and Bacon, 1991).

APPENDIX: Energy Conditions

When a matter field couples to any system, energy conditions must be satisfied for a physically acceptable solutions. We follow the steps as given in [7].

A. Weak Energy Condition (WEC)

The WEC states that,

$$\rho \geq 0 \quad \text{and} \quad \rho + p_i \geq 0 \quad (i = 1, 2, \dots, n+1) \quad (\text{A1})$$

in which ρ is the energy density and p_i is the principal pressures are given by

$$\rho = -T_t^t = -T_r^r = -\frac{C}{r^{n(1-k)}}, \quad p_i = T_i^i \quad (\text{no sum convention}) \quad (\text{A2})$$

The WEC imposes the following conditions on the constant parameters C and k ;

$$C \leq 0 \quad \text{and} \quad k \leq 1, \quad (\text{A3})$$

B. Strong Energy Condition (SEC)

This condition states that;

$$\rho + \sum_{i=1}^{n+1} p_i \geq 0 \quad \text{and} \quad \rho + p_i \geq 0. \quad (\text{A4})$$

This condition together with the WEC constrains the parameters as,

$$C \leq 0 \quad \text{and} \quad k \leq 0. \quad (\text{A5})$$

C. Dominant Energy Condition (DEC)

In accordance with DEC, the effective pressure p_{eff} which is defined as in Eq. (A6) should not be negative i.e $p_{eff} \geq 0$.

$$p_{eff} = \frac{1}{n+1} \sum_{i=1}^{n+1} T_i = -\frac{(1+nk)}{1+n} \rho \quad (\text{A6})$$

One can show that DEC, together with SEC and WEC conditions impose the following conditions on the parameters

$$C \leq 0 \quad \text{and} \quad -1 \leq k \leq 0. \quad (\text{A7})$$

D. Causality Condition

In addition to the energy conditions one can impose the causality condition

$$0 \leq \frac{p_{eff}}{\rho} < 1, \quad (\text{A8})$$

which implies

$$C \leq 0 \quad \text{and} \quad -1 \leq k \leq -\frac{1}{n} \quad (\text{A9})$$