

# Monitoring the radio spectra of selected blazars in the Fermi-GST era

The Effelsberg 100 m telescope covering the cm band

E. Angelakis, L. Fuhrmann, N. Marchili, T.P. Krichbaum and J.A. Zensus

Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany email: angelaki@mpifr-bonn.mpg.de

**Abstract.** The analysis of the spectral energy distribution variability at frequencies from radio to TeV is a powerful tool in the investigation of the dynamics, the physics and the structure evolution occurring in the most exotic flavour of active galaxies, the blazars. In particular, the presence of *Fermi-GST* is providing a unique opportunity for such studies delivering  $\gamma$ -ray data of unprecedented quality. Here we introduce a monitoring program that has been running at the Effelsberg 100 m telescope since January 2007, underpinning a broad multi-frequency collaboration of facilities that cover the band from radio to infrared. Sixty one selected blazars are observed monthly between 2.64 GHz and 43 GHz. The calibration accuracy is better than a few percent as it is demonstrated with some preliminary examples.

Key words. Galaxies: active – Galaxies: nuclei – Radio continuum: galaxies – BL Lacertae objects: general

# 1. Introduction

Being among the most dramatic manifestations of the activity induced in the nuclei of active galaxies, blazars comprise a unique probe of the exotic physics at play in such systems. Their phenomenology is dominated by extreme characteristics such as high degree of linear polarisation, intense variability – both in total power and polarisation – at all wavebands, apparent motions of features in their radio structure, which are often highly superluminal and brightness temperatures exceeding the Compton limit (see e.g. Urry 1999). This violent behaviour is attributed to relativistic jets oriented very close ( $\leq 20^{\circ}$  to  $30^{\circ}$ ) to the line-ofsight (see e.g. Urry & Padovani 1995). Their spectral energy distribution (SED) is characterised by the presence of two peaks. The lower energy one, spreading from radio to far ultraviolet and soft X-rays, is believed to be due to relativistic electrons emitting via the synchrotron mechanism. The high energy part, reaching up to TeV energies is believed to be produced by synchrotron self-Compton or external Compton scattering.

Despite the overall understanding, many details remain unclear and are subject to models attempting their clarification. For instance, several ideas have been put forth to explain the origin of their variability. Two of them are like

Send offprint requests to: E. Angelakis

 Table 1. The monitored sources.

Target sources							
PKS 0003-066	TXS 0814+425	OE 355					
PKS 0215+015	1E0317.0+18	OJ 248					
PKS 0235+164	H 1426+428	OJ 287					
PKS 0238-084	1ES 1959+650	OP 313					
PKS 0336-019	1ES 2344+514	3C 84					
PKS 0420-014	1ES 1544+820	3C 111					
PKS 0528+134	1ES 0502+675	3C 120					
PKS 0735+178	0059+581	3C 273					
PKS 0748+126	0219+428	3C 279					
PKS 1038+064	1128+592	3C 345					
PKS 1127-145	Mkn 180	3C 371					
PKS 1406-076	Mkn 421	3C 446					
PKS 1510-089	Mkn 501	3C 454.3					
PKS 1730-130	4C 21.35	CTA 102					
PKS 2155-152	4C 28.07	Cyg A					
PKS 2155-304	4C 31.63	BLLAC					
PKS 2345-16	4C 38.41	WCom					
S50836+71	4C 56.27	M87					
S51803+78	4C 47.08	OS 319					
S50716+71	TON 0599						
S40954+65	NRAO 150						

the shock-in-jet model (e.g. Marscher & Gear 1985; Aller et al. 1985; Marscher 1996) and the relativistic plasma shells (e.g. Spada et al. 2001; Guetta et al. 2004). Alternatively, it has been suggested that light-house effect could be causing variability in cases of rotating helical jets or the helical trajectories of plasma elements (e.g. Begelman et al. 1980; Camenzind & Krockenberger 1992).

A powerful tool in the investigation of these details and for understanding the dynamics, the physics and the structure of the radiating regions is the analysis of the variability of their SEDs at frequency ranges as broad as possible (from radio to TeV). Such an approach can shed light on linear scales inaccessible even to interferometric techniques and most importantly discriminate between different variability scenarios. Particularly, the presence of *Fermi-GST* is providing a unique opportunity for these studies providing the  $\gamma$ -ray data of unprecedented quality.

To benefit fully from the newly flying *Fermi-GST* telescope, a broad collaboration of facilities that aim at understanding the blazar phenomenon via the broad-band SED variability monitoring has been initiated by Fuhrmann et al. (2007). A sample of 61 selected sources are monitored monthly and in tight coordination with other facilities covering the cm, mm, optical and infrared bands (Fuhrmann et al. in prep.). Here we present a very brief introduction of the activities in the centimetric band with the 100 m Effelsberg telescope and show some preliminary examples.

# 2. The 100 m telescope monitoring program

Observationally, the goal of the currently discussed project is the monthly radio spectrum monitoring for a sample of the 61 sources shown in table 1 with the Effelsberg 100 m telescope. The broad collaboration involves also mainly the 30 m IRAM telescope and the 40 m Owens Valley telescope (Fuhrmann et al. 2007). The former is covering the band between roughly 86 GHz and 270 GHz whereas the latter is focusing on 15 GHz alone to monitor a few hundred sources per day and acquire information about their variability index. The time difference between the Effelsberg and IRAM observations is of order of one to maximum a few days. The monitoring program at Effelsberg has been running continuously since January 2007 providing a remarkable flow of data.

#### 3. The sample

The sources in table 1 have been selected from the "high-priority blazars" list of the *Fermi-GST* AGN team. There is a remarkable overlap with other studies such as 2 cm-VLBA survey/MOJAVE program (Kellermann et al. 1998; Zensus et al. 2002; Kellermann et al. 2004), the Boston 43 GHz VLBI survey (Jorstad et al. 2001), the IRAM 30-m telescope polarisation monitoring program, the OVRO monitoring program at 15 GHz and other multi-frequency campaigns.

#### 4. Observations and data reduction

For our monitoring program the receivers at 2.64 GHz, 4.85 GHz, 8.35 GHz, 10.45 GHz, 14.60 GHz, 23.05 GHz, 32.00 GHz and 42.00 GHz have been employed. Almost all of them (except for 32.00 GHz) deliver polarisation information, which provides another useful tool in the exploration of blazar physics.

The measurements are conducted with the newly installed adaptive secondary reflector characterised by low surface RMS that induces higher sensitivity (up to 50 % increase at 42.00 GHz). The so-called "cross-scan" observing technique has been applied. The advantage of this method is mainly the fact that it allows the direct detection of cases of confusion as well as the correction for pointing offset errors. The individual spectra are measured quasi-simultaneously within  $\leq$  40 min to guarantee that they are free of source variability of time-scales longer than that.

Considerable effort is put in applying some necessary post-observational corrections to the raw data. Namely, (a) pointing offset correction, (b) gain correction, (c) opacity correction and (d) sensitivity correction (see Angelakis et al. 2008). The sensitivity correction is done with reference to standard calibrators (e.g. table 2). There, the average flux densities are given along with the modulation index, m, over a number of measurements (m = $100 \cdot \frac{rms}{\langle S \rangle}$ ). The modulation index serves as a measure of the source variability. Hence, this table is a demonstration of the system repeatability and the achievable precision on the assumption that the calibrators are intrinsically non-variable.

## 5. Spectra

In figure 1 are collected two preliminary examples of spectra measured with Effelsberg. The comparison of these spectra with that of 3C 286 in figure 1(a) illustrates the significance of the observed variability. Remarkably, intense variability appears already in the cm band. Its correlation with that in other bands or possible changes in the structure of interferometric imaging is under examination. It is

**Table 2.** Average flux densities *S* in Jy for two of our main calibrators after preliminary analysis along with their modulation index *m* in percentage, as a function of *v* in GHz. *m* serves as a pragmatic measure of the system repeatability. For  $v \ge 23.05$ , NGC 7027 appears slightly extended for the Effelsberg beam. In column 4 we give its observed flux density *S'* convolved with the observing beam. In column 6 we show the beam correction factor  $\alpha$  in percentage, for a  $10 \times 12''$  disk-like brightness distribution. In column 7 we list the total flux density  $S = (1 + \alpha/100) \cdot S'$ .

ν	3C 2	286	NGC 7027			
	S	т	S'	т	α	S
2.64	10.7	0.7	3.7	1.1		
4.85	7.5	0.6	5.5	0.2		
8.35	5.2	0.7	5.9	0.6		
10.45	4.5	1.8	5.9	2.3		
14.60	3.5	1.9	5.8	1.5		
23.05	2.5	2.8	5.3	2.6	6.6	5.7
32.00	1.9	2.8	5.2	2.3	5.6	5.5
42.00	1.4	2.1	5.0	5.0	7.0	5.4

important to mention that the admittedly larger uncertainties at the higher frequencies is solely due to atmospheric variations.

# 6. Discussion

Provided that the calibrators are presumably not intrinsically variable, figure 1(a) serves as an excellent means of estimating the system repeatability. In table 2 the calibration precision is of the order of a few percent. Factors that induce such variations are primarily tropospheric variations, opacity effects, confusion and system instabilities. At higher frequencies (23.05 GHz and above), the atmospheric opacity becomes particularly important.

The plots in figure 1 show how intensive and fast the variability of the spectrum may be. This very fact urges for dense frequency coverage especially of the millimetre regime from where the evolution of a flaring event in the



**Fig. 1.** Examples of sources that exhibit intense spectrum variability in the cm band. The top panel shows the spectrum variability of one of the main calibrators namely 3C 286 for comparison. 3C 286 is calibrated with calibration factor averaged over all used calibrators.

radio band is expected to start. The evolution of the spectrum over time is among the firstpriority studies. That requires tight coordination of the different participating stations as has been managed so far. The activity seen in the centimetre band is noteworthy. Whether this activity is correlated with that at other bands and especially that at high energies, or with changes in the structure is one of the first questions to be addressed. The behaviour of the polarisation when such an event occurs is also among the most important issues to be investigated.

Acknowledgements. Based on observations with the 100 m telescope of the MPIfR (Max-Planck-Institut für Radioastronomie).

# References

- Aller, H. D., Aller, M. F., & Hughes, P. A. 1985, ApJ, 298, 296
- Angelakis, E., Kadler, M., Lewis, K., Sambruna, R. M., Eracleous, M., & Zensus, J. A. 2008, these proceedings
- Begelman, M. C., Blandford, R. D., & Rees, M. J. 1980, Nature, 287, 307
- Camenzind, M. & Krockenberger, M. 1992, A&A, 255, 59
- Fuhrmann, L., Zensus, J. A., Krichbaum, T. P., Angelakis, E., & Readhead, A. C. S. 2007, in American Institute of Physics Conference Series, Vol. 921, ed. S. Ritz, P. Michelson, & C. A. Meegan, 249–251
- Guetta, D., Ghisellini, G., Lazzati, D., & Celotti, A. 2004, A&A, 421, 877
- Jorstad, S. G., Marscher, A. P., Mattox, J. R., et al. 2001, ApJS, 134, 181
- Kellermann, K. I., Lister, M. L., Homan, D. C., et al. 2004, ApJ, 609, 539
- Kellermann, K. I., Vermeulen, R. C., Zensus, J. A., & Cohen, M. H. 1998, AJ, 115, 1295
- Marscher, A. P. 1996, in ASP Conference Series, Vol. 100, ed. P. E. Hardee, A. H. Bridle, & J. A. Zensus, 45-+
- Marscher, A. P. & Gear, W. K. 1985, ApJ, 298, 114
- Spada, M., Ghisellini, G., Lazzati, D., & Celotti, A. 2001, MNRAS, 325, 1559
- Urry, C. M. 1999, Astroparticle Physics, 11, 159
- Urry, C. M. & Padovani, P. 1995, PASP, 107, 803
- Zensus, J. A., Ros, E., Kellermann, K. I., et al. 2002, AJ, 124, 662