Cohomological Dimension of Laumon 1-motives up to Isogenies

Nicola Mazzari

October 26, 2018

Dipartimento di Matematica "Federigo Enriques" Università degli Studi di Milano Via Saldini, 50 20133 - Milano E-Mail: mazzari@mat.unimi.it

Abstract

We prove that the category of Laumon 1-motives up isogenies over a field of characteristic zero is of cohomological dimension ≤ 1 . As a consequence this implies the same result for the category of formal Hodge structures of level ≤ 1 (over \mathbb{Q}). MSC: 14C99, 14L15.

Contents

1	Lau	imon 1-motives	2
	1.1	fppf sheaves	5
2 Extensions		ensions	6
	2.1	The group of n -extensions \ldots \ldots \ldots \ldots \ldots	6
		2.1.1 A lemma on 2-fold extensions	6
	2.2	Ext of 1-motives up to isogenies	7
References			10

Introduction

In [6] Deligne defined 1-motive over a field k as $\operatorname{Gal}(k^{\operatorname{sep}}|k)$ -equivariant morphism $[u: \mathbf{X} \to \mathbf{G}(k^{\operatorname{sep}})]$ where \mathbf{X} is a free $\operatorname{Gal}(k^{\operatorname{sep}}|k)$ -module and \mathbf{G} is

a semi-abelian algebraic group over k. They form a category that we shall denote by $\mathcal{M}_{1,k}$ or \mathcal{M}_1 .

Deligne's definition was motivated by Hodge theory. In fact the category of 1-motives over the complex numbers is equivalent, via the so called *Hodge realization* functor, to the category MHS_1 of mixed Hodge structures of level ≤ 1 . It is known the the category MHS_1 is of cohomological dimension 1 (see [5]) and the same holds for $\mathcal{M}_{1,\mathbb{C}}$.

F. Orgogozo proved more generally that for any field k, the category $\mathcal{M}_{1,k} \otimes \mathbb{Q}$ is of cohomological dimension ≤ 1 (see [14, Prop. 3.2.4]).

Over a field of characteristic 0 it is possible to define the category $\mathcal{M}_{1,k}^{a}$ of Laumon 1-motives generalizing that of Deligne 1-motives (See [11]). In [3] L. Barbieri-Viale generalized the Hodge realization functor to Laumon 1-motives. He defined the category FHS_1 of formal Hodge structures of level ≤ 1 containing MHS_1 and proved that FHS_1 is equivalent to the category of Laumon 1-motives over \mathbb{C} (compatibly with the Hodge realization).

In this paper we prove that the category of Laumon 1-motives up to isogenies is of cohomological dimension 1.

Acknowledgments

The author would like to thank L. Barbieri-Viale for pointing his attention to this subject and for helpful discussions. The author also thanks A. Bertapelle for many useful comments and suggestions.

1 Laumon 1-motives

In this paper k is a field of characteristic 0 and \bar{k} is its algebraic closure. As explained in the §1.1 we can assume that the categories of formal and algebraic groups are full sub-category of Ab_k , i.e. the category of abelian sheaves on the category of affine k-schemes w.r.t. the fppf topology.

Definition 1.1. A Laumon 1-motive over k (or an effective free 1-motive over k, cf. [2, 1.4.1]) is the data of

i) A (commutative) formal group \boldsymbol{F} over k, such that Lie \boldsymbol{F} is a finitely generated and $\boldsymbol{F}(\bar{k}) = \lim_{[k':k] < \infty} \boldsymbol{F}(k')$ is finitely generated and torsion-free $\operatorname{Gal}(\bar{k}/k)$ -module.

ii) A connected commutative algebraic group scheme G over k.

iii) A morphism $u: \mathbf{F} \to \mathbf{G}$ in the category Ab_k .

Note that we can consider a Laumon 1-motive (over k) $M = [u : \mathbf{F} \to \mathbf{G}]$ as a complex of sheaves in Ab_k concentrated in degree 0, 1. It is known that any formal k group \mathbf{F} splits appendix $\mathbf{F}^{o} \times \mathbf{F}$.

It is known that any formal k-group \boldsymbol{F} splits canonically as product $\boldsymbol{F}^{o} \times \boldsymbol{F}_{et}$ where \boldsymbol{F}^{o} is the identity component of \boldsymbol{F} and is a connected formal k-group, and $\boldsymbol{F}_{et} = \boldsymbol{F}/\boldsymbol{F}^{o}$ is étale. Moreover, \boldsymbol{F}_{et} admits a maximal sub-group scheme \mathbf{F}_{tor} , étale and finite, such that the quotient $\mathbf{F}_{et}/\mathbf{F}_{tor} = \mathbf{F}_{fr}$ is constant of the type \mathbb{Z}^r over \bar{k} . One says that \mathbf{F} is torsion-free if $\mathbf{F}_{tor} = 0$.

By a theorem of Chevalley any connected algebraic group scheme G is extension of an abelian variety A by a linear k-group scheme L that is product of its maximal sub-torus T with a vector k-group scheme V. (See [7] for more details on algebraic and formal groups)

Definition 1.2. A morphism of Laumon 1-motives is a commutative square in the category Ab_k . We denote by $\mathcal{M}_1^a = \mathcal{M}_{1,k}^a$ the category of Laumon k-1-motives, i.e. the full sub-category of $C^b(Ab_k)$ whose objects are Laumon 1-motives.

Remark 1.3. The category of Deligne 1-motives (over k) is the full subcategory \mathcal{M}_1 of \mathcal{M}_1^a whose objects are $M = [u : \mathbf{F} \to \mathbf{G}]$ such that $\mathbf{F}^o = 0$ and \mathbf{G} is semi-abelian (cf. [6, §10.1.2]).

Proposition 1.4. The category \mathcal{M}_1^{a} of Laumon 1-motives (over k) is an additive category with kernels and co-kernels.

Proof. See [11, Prop. 5.1.3].

Remark 1.5. i) Let



be a morphism from $M = [u : \mathbf{F} \to \mathbf{G}]$ to $M' = [u' : \mathbf{F}' \to \mathbf{G}']$. Then from previous proof we get

$$\operatorname{Ker}(f,g) = [u^* \operatorname{Ker}(g)^o \to \operatorname{Ker}(g)^o]$$
(1)

where $u : \operatorname{Ker}(f) \to \operatorname{Ker}(g)$, and

$$\operatorname{Coker}(f,g) = [\operatorname{Coker}(f)_{\mathrm{fr}} \to \operatorname{Coker}(g)] \tag{2}$$

ii) The category of Laumon 1-motives is not abelian. In fact consider a surjective morphism of connected algebraic groups $g: \mathbf{G} \to \mathbf{G}'$. Then $\operatorname{Ker}(g)$ is not necessarily connected. Hence in the category of connected algebraic groups the canonical map

$$\operatorname{Coim}(g) = \mathbf{G} / \operatorname{Ker}(g)^o \to \operatorname{Im}(g) = \mathbf{G}'$$

is not an isomorphism in general.

Note that the category of connected algebraic groups is fully embedded in \mathcal{M}_1^a .

According to [14] we define the category $\mathcal{M}_1^a \otimes \mathbb{Q}$ of Laumon 1-motives up to isogenies: the objects are the same of \mathcal{M}_1^a ; the Hom groups are $\operatorname{Hom}_{\mathcal{M}_1^a}(M, M') \otimes_{\mathbb{Z}} \mathbb{Q}$.

Remark 1.6. Note that a morphism $(f,g): M \to M'$ is an isogeny (i.e. an isomorphism in $\mathcal{M}_1^{\mathbf{a}} \otimes \mathbb{Q}$) if and only if f is injective with finite co-kernel and g is surjective with finite kernel.

Proposition 1.7. The category of Laumon 1-motives up to isogenies is abelian.

Proof. By construction $\mathcal{M}_1^{\mathbf{a}} \otimes \mathbb{Q}$ is an additive category. Let $(f,g) : M \to M'$ be a morphism of Laumon 1-motives. We know that the group $\pi_0(\operatorname{Ker}(g)) = \operatorname{Ker}(g)/\operatorname{Ker}(g)^o$ is a finite group scheme, hence there exists an integer n such that the following diagram commutes in Ab_k



Hence $n \cdot u$ factors through $\operatorname{Ker}(g)^o$. Then it is easy to check that $\operatorname{Ker}((f,g)) = [(u^* \operatorname{Ker}(g)^o) \to \operatorname{Ker}(g)^o]$ is isogenous to $[\operatorname{Ker}(f) \to \operatorname{Ker}(g)^o]$.

It follows that $\operatorname{Coim}(f,g)$ is isogenous to $[(F/\operatorname{Ker}(f))_{\mathrm{fr}} \to G/\operatorname{Ker}(g)]$. As $G/\operatorname{Ker}(g)^o \to G/\operatorname{Ker}(g)$ is an isogeny we get that the canonical map $\operatorname{Coim}(f,g) \to \operatorname{Im}(f,g)$ an isogeny too.

This is enough to prove that the category $\mathcal{M}_1^a \otimes \mathbb{Q}$ is abelian.

Remark 1.8. We can define the category ${}^{t}\mathcal{M}_{1}^{a}$ of 1-motives with torsion (over k) as the full sub-category of $D^{b}(\mathsf{Ab}_{k})$ with objects complexes $[u: \mathbf{F} \to \mathbf{G}]$ concentrated in degree 0, 1 such that

i) A (commutative) formal group F over k, such that Lie F is a finitely generated and $F(\bar{k}) = \lim_{[k':k] < \infty} F(k')$ is finitely generated (non necessarily torsion-free!) Gal $(\bar{k}|k)$ -module.

ii) A connected commutative algebraic group scheme G over k.

iii) A morphism $u: \mathbf{F} \to \mathbf{G}$ in the category Ab_k .

Also we denote by ${}^{t}\mathcal{M}_{1} \subset {}^{t}\mathcal{M}_{1}^{a}$ the full sub-category whose object are of the form $[u: \mathbf{F} \to \mathbf{G}]$ with $\mathbf{F}^{o} = 0$ and \mathbf{G} semi-abelian.

In [4, C.7.3] is proven that the canonical functor $\mathcal{M}_1 \to {}^t\mathcal{M}_1$ induces an equivalence of the same categories up to isogeny, i.e. $\mathcal{M}_1 \otimes \mathbb{Q} \cong {}^t\mathcal{M}_1 \otimes \mathbb{Q}$. The same result holds for Laumon 1-motives, in fact all the arguments given in [4] work also in this setting. Hence there is an equivalence of categories

$$\mathcal{M}_1^{\mathrm{a}} \otimes \mathbb{Q} \xrightarrow{\sim} {}^t \mathcal{M}_1^{\mathrm{a}} \otimes \mathbb{Q}$$
.

A Deligne 1-motive is endowed with an increasing filtration (of sub-1motives) called the weight filtration $([6, \S10.1.4])$ defined as follows

$$W_{i} = W_{i}M := \begin{cases} [u: \mathbf{X} \to \mathbf{G}] & i \ge 0\\ [0 \to \mathbf{G}] & i = -1\\ [0 \to \mathbf{T}] & i = -2\\ [0 \to 0] & i \le -3 \end{cases}$$

hence we get

$$\operatorname{gr}_{i}^{W} M = \begin{cases} [\boldsymbol{X} \to 0] & i = 0\\ [0 \to \boldsymbol{A}] & i = -1\\ [0 \to \boldsymbol{T}] & i = -2\\ [0 \to 0] & \text{otherwise} \end{cases}$$

According to [4, C.11.1] we extend the weight filtration to Laumon 1motives. Let $M = [u : \mathbf{F} \to \mathbf{G}]$ be an Laumon 1-motive, then

$$W_{-3} = 0 \ \subset W_{-2} = [0 \to L] \ \subset W_{-1} = [0 \to G] \ \subset W_0 = M$$

1.1 fppf sheaves

Let Sch_k be the category of schemes over k and Aff_k be the full sub-category of affine schemes. According to [1, Exp. IV §6.3] the fppf topology on Sch_k is the one generated by: the families of jointly surjective open immersions in Sch_k ; the finite families of jointly surjective, flat, of finite presentation and quasi-finite morphisms in Aff_k .

Recall that Ab_k is the category of abelian sheaves on Aff_k w.r.t. the fppf topology.

Proposition 1.9. *i)* The category of commutative group schemes over k is a full sub-category of Ab_k *via the functor of points* $\boldsymbol{G} \mapsto h_{\boldsymbol{G}} := Hom_{sch_k}(-, \boldsymbol{G})$.

ii) Let $\operatorname{char}(k) = 0$. The category of formal group schemes is a full subcategory of Ab_k via the functor of points $\mathbf{F} = \operatorname{Spf}(A) \mapsto h_{\mathbf{F}} := \operatorname{Hom}_{\operatorname{alg}_k}^{\operatorname{cont}}(A, -)$ (where $\operatorname{Hom}_{\operatorname{alg}_k}^{\operatorname{cont}}(-, -)$ denote the set of continuous homomorphisms of kalgebras).

Proof. By a result of Grothendieck ([8, Part I, §2.3.6]) every scheme (over k) is a sheaf (on sets) w.r.t. the fppf topology on Sch_k . Hence it is also a fppf-sheaf on the sub-category $\mathsf{Aff}_k \subset \mathsf{Sch}_k$. From this fact (i) and (ii) follow for étale formal groups.

By the decomposition theorem for formal groups over a perfect field [7] it remains to prove that any connected (or local) formal group is a sheaf. It is sufficient to note that

$$\widehat{\mathbb{G}}_a \cong \operatorname{colim}_n \operatorname{Spec}(k[t]/(t^{n+1}))$$

is a direct limit of affine schemes, hence a direct limit of sheaves of sets w.r.t. the fppf topology. $\hfill \Box$

2 Extensions

2.1 The group of *n*-extensions

Let A be any abelian category (we don't suppose it has enough injective objects), then we can define its derived category D(A) and the group of n-fold extension classes

$$\operatorname{Ext}_{\mathsf{A}}^{n}(A,B) := \operatorname{Hom}_{D(\mathsf{A})}(A,B[n]) \qquad A,B \in \mathsf{A}$$

As usual we identify this group with the group of classes of *Yoneda extensions*, i.e. the set of exact sequences

 $0 \to B \to E_1 \to \dots \to E_n \to A \to 0$

modulo congruences (See [10] or [9]).

2.1.1 A lemma on 2-fold extensions

Consider a 2-fold extension $\gamma \in \operatorname{Ext}^2_{\mathsf{A}}(M, M')$. It is represented by an exact sequence

$$0 \to M' \to E_1 \to E_2 \to M \to 0 . \tag{3}$$

This can be written as the product of two 1-fold extensions as follows. Let $E := \operatorname{Ker}(E_2 \to M) = \operatorname{Coker}(M' \to E_1)$, then let $\gamma_1 \in \operatorname{Ext}^1_{\mathsf{A}}(E, M')$, $\gamma_2 \in \operatorname{Ext}^1_{\mathsf{A}}(M, E)$ be the classes represented by

$$0 \to M' \to E_1 \to E \to 0 , \quad 0 \to E \to E_2 \to M \to 0 \tag{4}$$

respectively. Then $\gamma = \gamma_1 \cdot \gamma_2$.

As a particular case, consider $W_{-2} \subset W_{-1} \subset W_0$ a sequence of objects of A. We have the following exact sequences

$$\begin{split} \gamma : & 0 \to W_{-2} \to W_{-1} \to W_0/W_{-2} \to W_0/W_{-1} \to 0 \\ & \gamma_1 : & 0 \to W_{-2} \to W_{-1} \to W_{-1}/W_{-2} \to 0 \\ & \gamma_2 : & 0 \to W_{-1}/W_{-2} \to W_0/W_{-2} \to W_0/W_{-1} \to 0 \end{split}$$

and $\gamma = \gamma_1 \cdot \gamma_2 \in \operatorname{Ext}^2_{\mathsf{A}}(W_0/W_{-1}, W_{-2})$. In this particular case we get

Lemma 2.1. $\gamma = 0$ in $\operatorname{Ext}^{2}_{A}(W_{0}/W_{-1}, W_{-2})$.

Proof. See [14, Lemma 3.2.5], or [9, p. 184].

2.2 Ext of 1-motives up to isogenies

From now on we call 1-motive a Laumon 1-motive (over k) and $\operatorname{Ext}^{i}_{\mathbb{Q}}(M, M')$ is the group of classes of i-fold extensions in $\mathcal{M}^{a}_{1} \otimes \mathbb{Q}$.

We are going to prove that $\mathcal{M}_1^a \otimes \mathbb{Q}$ is of cohomological dimension ≤ 1 . We start with the following result.

Lemma 2.2. Let $\operatorname{gr}_{i}^{W} \mathcal{M}_{1}^{a}$ be the full sub-category of \mathcal{M}_{1}^{a} with objects 1motives pure of weight i, i = 0, -1, -2. Then $\operatorname{gr}_{i}^{W} \mathcal{M}_{1}^{a} \otimes \mathbb{Q}$ is an abelian thick sub-category of $\mathcal{M}_{1}^{a} \otimes \mathbb{Q}$ and it is of cohomological dimension 0.

Proof. First we consider the case $M = \mathbf{F}[1], M' = \mathbf{F}'[1]$ pure of weight 0 (i.e. formal groups). Let $0 \to \mathbf{F}'[1] \to E \to \mathbf{F}[1] \to 0$ an exact sequence of 1-motives modulo isogenies. Then E is also of weight 0 (this follows directly from the definitions). Hence $\operatorname{Ext}_{\mathbb{Q}}^{1}(\mathbf{F}[1], \mathbf{F}'[1])$ is isomorphic to the group of classes of extensions in the category of formal groups over k modulo isogenies. We know that of k-vector spaces Mod_k is semi-simple, and so is the \mathbb{Q} -linearized category of free $\operatorname{Gal}(\bar{k}/k)$ -modules, $\operatorname{Mod}_{\operatorname{Gal}(\bar{k}/k)}^{\operatorname{free}} \otimes \mathbb{Q}$, by the lemma of Maschke (See [15, p. 47], for the representations of finite groups; the case of pro-finite is a direct consequence). Hence the category of formal groups up to isogeny, equivalent to $\operatorname{Mod}_k \times \operatorname{Mod}_{\operatorname{Gal}(\bar{k}/k)}^{\operatorname{free}} \otimes \mathbb{Q}$, is of cohomological dimension 0.

The second case is that of abelian varieties (weight -1). Again using the definitions we get that $\operatorname{Ext}_{\mathbb{Q}}^{1}(A', A)$ correspond to the group of extensions in the category of abelian varieties modulo isogenies. This group is zero (See [13, p. 173]).

The third case is that of linear groups (weight -2). This can be reduced to the first case by Cartier duality (See [11, §5]) or proved explicitly.

Lemma 2.3. Let M, M' be pure Laumon 1-motives with weights w < w'. Then $\operatorname{Ext}^2_{\mathbb{Q}}(M, M') = 0$.

Proof. Fix a 2-fold extension $\gamma \in \operatorname{Ext}^2_{\mathbb{O}}(M, M')$ represented by

$$0 \to M' \to E_1 \to E_2 \to M \to 0$$

and take $\gamma_2 \in \operatorname{Ext}^1_{\mathbb{Q}}(M, E), \ \gamma_1 \in \operatorname{Ext}^1_{\mathbb{Q}}(E, M')$ (as in (3), (4)) such that $\gamma = \gamma_1 \cdot \gamma_2$.

We have an exact sequence

$$\operatorname{Ext}^{1}_{\mathbb{O}}(M, W_{-1}E) \to \operatorname{Ext}^{1}_{\mathbb{O}}(M, E) \to \operatorname{Ext}^{1}_{\mathbb{O}}(M, \operatorname{gr}_{0}E)$$

By assumption $-2 \leq w < w' \leq 0$, then M is pure of weight -1 or -2. In this case we get easily $\operatorname{Ext}^{1}_{\mathbb{Q}}(M, \operatorname{gr}_{0} E) = 0$ and we can lift γ_{2} to $\gamma'_{2} \in \operatorname{Ext}^{1}_{\mathbb{Q}}(M, W_{-1}E)$. Then let γ'_{1} be the image of γ_{1} via $\operatorname{Ext}^{1}_{\mathbb{Q}}(E, M') \to \operatorname{Ext}^{1}_{\mathbb{Q}}(W_{-1}E, M')$. Now using

$$\operatorname{Ext}^{1}_{\mathbb{Q}}(\operatorname{gr}_{-1} E, M') \to \operatorname{Ext}^{1}_{\mathbb{Q}}(W_{-1}E, M') \to \operatorname{Ext}^{1}_{\mathbb{Q}}(W_{-2}E, M')$$

we can reduce to consider E pure of weight -1, in fact $\operatorname{Ext}^{1}_{\mathbb{Q}}(W_{-2}E, M') = 0$ because w' > -2. From this follows that $\gamma_{1} = \gamma_{2} = 0$.

Lemma 2.4. Let M, M' be pure Laumon 1-motives with weights w > w'. Then $\operatorname{Ext}^2_{\mathbb{Q}}(M, M') = 0$.

Proof. As in the previous proof fix a 2-fold extension $\gamma \in \operatorname{Ext}^2_{\mathbb{Q}}(M, M')$ represented by

$$0 \to M' \to E_1 \to E_2 \to M \to 0$$

and take $\gamma_2 \in \operatorname{Ext}^1_{\mathbb{Q}}(M, E), \gamma_1 \in \operatorname{Ext}^1_{\mathbb{Q}}(E, M')$ (as in 3, 4) such that $\gamma = \gamma_1 \cdot \gamma_2$.

We have to consider three cases: (a) $M = \mathbf{F}[1], M' = \mathbf{A}[0];$ (b) $M = \mathbf{F}[1], M' = \mathbf{L}[0];$ (c) $M = \mathbf{A}[0], M' = \mathbf{L}[0]$. Where \mathbf{F} is a formal group, \mathbf{A} an abelian variety, \mathbf{L} a linear group.

Case (a): now $\gamma_1 \in \text{Ext}^1_{\mathbb{Q}}(E, \mathbf{A}) \ \gamma_2 \in \text{Ext}^1_{\mathbb{Q}}(\mathbf{F}[1], E)$. Then $E = [\mathbf{F}' \to \mathbf{A}']$ is such that $W_{-2}E = 0$. Consider the exact sequence

$$0 \to \operatorname{gr}_{-1} E \to E \to \operatorname{gr}_0 E \to 0$$

applying $\operatorname{Hom}_{\mathbb{Q}}(\boldsymbol{F}[1], -)$ to it we get

$$\operatorname{Ext}^{1}_{\mathbb{Q}}(\boldsymbol{F}[1],\operatorname{gr}_{-1}E) \to \operatorname{Ext}^{1}_{\mathbb{Q}}(\boldsymbol{F}[1],E) \to \operatorname{Ext}^{1}_{\mathbb{Q}}(\boldsymbol{F}[1],\operatorname{gr}_{0}E)$$

We proved that $\operatorname{Ext}_{\mathbb{Q}}^{1}(\boldsymbol{F}[1], \operatorname{gr}_{0} E) = 0$ so we can lift γ_{2} to a class $\gamma_{2}' \in \operatorname{Ext}_{\mathbb{Q}}^{1}(\boldsymbol{F}[1], \operatorname{gr}_{-1} E)$ (This lifting is not canonical). Similarly using $\operatorname{Hom}_{\mathbb{Q}}(-, \boldsymbol{A})$ to it we get an exact sequence

$$\operatorname{Ext}^{1}_{\mathbb{Q}}(\operatorname{gr}_{0} E, A) \to \operatorname{Ext}^{1}_{\mathbb{Q}}(E, A) \to \operatorname{Ext}^{1}_{\mathbb{Q}}(\operatorname{gr}_{-1} E, A)$$

and we can map $\gamma_1 \mapsto \gamma'_1 \in \operatorname{Ext}^1_{\mathbb{Q}}(\operatorname{gr}_{-1} E, \mathbf{A})$. By standard arguments it holds $\gamma'_1 \cdot \gamma'_2 = \gamma_1 \cdot \gamma_2 = \gamma$. Recalling that $\operatorname{Ext}^1_{\mathbb{Q}}(\operatorname{gr}_{-1} E, \mathbf{A}) = 0$ we get the result.

Case (c): Is similar to case (a).

Case (b): now $\gamma \in \operatorname{Ext}_{\mathbb{Q}}^{2}(\boldsymbol{F}[1], \boldsymbol{L})$. We want to reduce to the hypothesis of lemma 2.1. Thus we have to show: we can take E pure of weight 1 (i.e. an abelian variety); there exists a 1-motive N such that $\gamma_{1} \in \operatorname{Ext}_{\mathbb{Q}}^{1}(\boldsymbol{E}, \boldsymbol{L})$ is represented by $0 \to W_{-2}N \to W_{-1}N \to \operatorname{gr}_{-1}N \to 0$; $\gamma_{2} \in \operatorname{Ext}_{\mathbb{Q}}^{1}(\boldsymbol{F}[1], E)$ is represented by $0 \to \operatorname{gr}_{-1}N \to W_{0}N/W_{-2} \to \operatorname{gr}_{0}N \to 0$.

We know that $\operatorname{Ext}_{\mathbb{Q}}^{1}(\boldsymbol{F}[1], \operatorname{gr}_{0} E) = 0$, so like in case (a) we can lift γ_{2} to a class $\gamma_{2}' \in \operatorname{Ext}_{\mathbb{Q}}^{1}(\boldsymbol{F}[1], W_{-1}E)$. Let γ_{1}' the image of γ_{1} via $\operatorname{Ext}_{\mathbb{Q}}^{1}(E, \boldsymbol{L}) \to \operatorname{Ext}_{\mathbb{Q}}^{1}(W_{-1}E, \boldsymbol{L})$. Hence $\gamma_{1}' \cdot \gamma_{2}' = \gamma_{1} \cdot \gamma_{2}$.

Now we can suppose E of weight ≤ -1 . Using the same argument we can lift γ'_1 to $\gamma''_1 \in \operatorname{Ext}^1_{\mathbb{Q}}(\operatorname{gr}_{-1} E, \mathbf{L})$ (because $\operatorname{Ext}^1_{\mathbb{Q}}(\operatorname{gr}_{-2} E, \mathbf{L}) = 0$) and send $\gamma'_2 \mapsto \gamma''_2 \in \operatorname{Ext}^1_{\mathbb{Q}}(\mathbf{F}[1], \operatorname{gr}_{-1} E)$. We proved that there exists an abelian variety $\mathbf{A}, \gamma_1 \in \operatorname{Ext}^1_{\mathbb{Q}}(\mathbf{A}, \mathbf{A}), \gamma_2 \in \operatorname{Ext}^1_{\mathbb{Q}}(\mathbf{F}[1], \mathbf{A})$, such that $\gamma_1 \cdot \gamma_2 = \gamma$. We claim that γ_1, γ_2 can be represented by extensions in the category Laumon-1-motives. In fact let

$$\gamma_1: \quad 0 \to \boldsymbol{L} \stackrel{f \otimes n^{-1}}{\longrightarrow} \boldsymbol{G} \stackrel{g \otimes m^{-1}}{\longrightarrow} \boldsymbol{A} \to 0$$

be an extension in the category of 1-motives modulo isogenies: f, g are morphism of algebraic groups, $n, m \in \mathbb{Z}$. Then consider the push-forward by n^{-1} and the pull-back by m^{-1} , we get the following commutative diagram with exact rows in $\mathcal{M}_1^{\mathrm{a,fr}} \otimes \mathbb{Q}$

$$0 \longrightarrow L \xrightarrow{f/n} G \xrightarrow{g/m} A \longrightarrow 0$$

$$\downarrow n^{-1} \qquad \downarrow \text{id} \qquad \downarrow \text{id}$$

$$0 \longrightarrow L \xrightarrow{f} G \xrightarrow{g/m} A \longrightarrow 0$$

$$\uparrow \text{id} \qquad \uparrow \text{id} \qquad \uparrow m^{-1}$$

$$0 \longrightarrow L \xrightarrow{f} G \xrightarrow{g} A \longrightarrow 0$$

The exactness of the last row is equivalent to the following: Ker f is finite; let $(\text{Ker }g)^0$ be the connected component of Ker g, then $\text{Im}f \to (\text{Ker }g)^0$ is surjective with finite kernel K; g is surjective. So after replacing L, A with isogenous groups we have an exact sequence in $\mathcal{M}_1^{a, \text{fr}}$

$$0 \rightarrow \boldsymbol{L} \rightarrow \boldsymbol{G} \rightarrow \boldsymbol{A} \rightarrow 0$$

Explicitly



With similar arguments we can prove that γ_2 is represented by an extension in the category $\mathcal{M}_1^{a,\mathrm{fr}}$

$$0 \to \mathbf{A} \to N \to \mathbf{F}[1] \to 0$$

with $N = [u : \boldsymbol{F} \to \boldsymbol{A}].$

To apply lemma 2.1 we need to prove that there is lifting $u': \mathbf{F} \to \mathbf{G}$. First suppose $\mathbf{F} = \mathbf{F}_{et}$: consider the long exact sequence

$$\operatorname{Hom}_{\operatorname{Ab}_k}(F,G) \to \operatorname{Hom}_{\operatorname{Ab}_k}(F,A) \xrightarrow{O} \operatorname{Ext}^1_{\operatorname{Ab}_k}(F,L)$$

We know ([12]) that $\operatorname{Ext}^{1}_{\mathsf{Ab}_{k}}(\boldsymbol{F}, \boldsymbol{L})$ is a torsion group. So modulo replacing \boldsymbol{F} with an isogenous lattice we get $\partial u = 0$ and the lift exists.

In case $F = F^o$ is a connected formal group we have a commutative diagram in Ab_k



where $\widehat{?}$ is the formal completion at the origin of ? = G, A. The formal completion is an exact functor so $\widehat{\pi}$ is an epimorphism. The category of formal groups is of cohomological dimension 0, then we can choose a section of $\widehat{\pi}$ and lift u.

Theorem 2.5. The category $\mathcal{M}_1^a \otimes \mathbb{Q}$ is of cohomological dimension 1.

Proof. First note that we can restrict to consider pure motives M, M' (a 1-motive is pure if it is isomorphic to one of its graded pieces w.r.t. the weight filtration). In fact given M, M' 1-motives, not necessarily pure, we have the canonical exact sequences given by the weight filtration

$$0 \to W_{-1}M' \to M' \to \operatorname{gr}_0^W M' \to 0$$
$$0 \to W_{-2}M' \to W_{-1}M' \to \operatorname{gr}_{-1}^W M' \to 0$$

Hence applying $\operatorname{Hom}_{\mathbb{Q}}(M, -)$ we get two long exact sequences

$$\cdots \operatorname{Ext}^{2}_{\mathbb{Q}}(M, W_{-1}M') \to \operatorname{Ext}^{2}_{\mathbb{Q}}(M, M') \to \operatorname{Ext}^{2}_{\mathbb{Q}}(M, \operatorname{gr}^{W}_{0}M') \cdots$$

$$\cdots \operatorname{Ext}^{2}_{\mathbb{Q}}(M, W_{-2}M') \to \operatorname{Ext}^{2}_{\mathbb{Q}}(M, W_{-1}M') \to \operatorname{Ext}^{2}_{\mathbb{Q}}(M, \operatorname{gr}^{W}_{-1}M') \cdots$$

from this follows that we can reduce to prove $\operatorname{Ext}^2_{\mathbb{Q}}(M, M') = 0$ for M' pure. In the same way we reduce to consider M pure.

Then using the previous three lemmas we conclude.

References

- Schémas en groupes. I: Propriétés générales des schémas en groupes. Séminaire de Géométrie Algébrique du Bois Marie 1962/64 (SGA 3). Dirigé par M. Demazure et A. Grothendieck. Lecture Notes in Mathematics, Vol. 151. Springer-Verlag, Berlin, 1970.
- [2] Luca Barbieri-Viale and Alessandra Bertapelle. Sharp de Rham realization. arXiv:math/0607115v1, 2006.

- [3] Luca Barbieri-Viale. Formal Hodge theory. Math. Res. Lett., 14(3):385– 394, 2007.
- [4] Luca Barbieri-Viale and Bruno Kahn. On the derived category of 1motives, I. arXiv:0706.1498v1, 2007.
- [5] A. A. Beïlinson. Notes on absolute Hodge cohomology. In Applications of algebraic K-theory to algebraic geometry and number theory, Part I, II (Boulder, Colo., 1983), volume 55 of Contemp. Math., pages 35-68. Amer. Math. Soc., Providence, RI, 1986.
- [6] Pierre Deligne. Théorie de Hodge. III. Inst. Hautes Études Sci. Publ. Math., (44):5-77, 1974.
- [7] Michel Demazure. Lectures on p-divisible groups. Lecture Notes in Mathematics, Vol. 302. Springer-Verlag, Berlin, 1972.
- [8] Barbara Fantechi, Lothar Göttsche, Luc Illusie, Steven L. Kleiman, Nitin Nitsure, and Angelo Vistoli. Fundamental algebraic geometry, volume 123 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2005. Grothendieck's FGA explained.
- [9] Sergei I. Gelfand and Yuri I. Manin. Methods of homological algebra. Springer Monographs in Mathematics. Springer-Verlag, Berlin, second edition, 2003.
- [10] Birger Iversen. Cohomology of sheaves. Universitext. Springer-Verlag, Berlin, 1986.
- [11] Gerard Laumon. Transformation de Fourier generalisee. arXiv:alg-geom/9603004v1, 1996.
- [12] James S. Milne. Etale Cohomology, volume 33 of Princeton Mathematical Series. Princeton University Press, Princeton, N.J., 1980.
- [13] David Mumford. Abelian varieties. Tata Institute of Fundamental Research Studies in Mathematics, No. 5. Published for the Tata Institute of Fundamental Research, Bombay, 1970.
- [14] Fabrice Orgogozo. Isomotifs de dimension inférieure ou égale à un. Manuscripta Math., 115(3):339-360, 2004.
- [15] Jean-Pierre Serre. Linear representations of finite groups. Springer-Verlag, New York, 1977. Translated from the second French edition by Leonard L. Scott, Graduate Texts in Mathematics, Vol. 42.