

RCF 3

Map-Code Interpretation via Closure

$\in \mathcal{U}_{\tilde{\varepsilon}}$

Michael Pfender*

July 2008[†]

Abstract

For a (minimal) Arithmetical theory with higher Order Objects, i. e. a (minimal) *Cartesian closed* arithmetical theory – coming as such with the corresponding *closed evaluation* – we *interpret* here *map codes*, out of $[A, B]$ say, into these maps “themselves”, coming as elements (“names”) of hom-Objects B^A . The interpretation (family) uses a Chain of *Universal Objects* $\mathcal{U}_{\underline{n}}$, one for each Order stratum with respect to “higher” Order of the Objects. Combined with *closed*, axiomatic *evaluation*, these interpretation family gives *code-self-evaluation*. Via the usual diagonal argument, Antinomie RICHARD then can be *formalised* within our minimal higher Order (Cartesian closed) arithmetical theory, and yields this way inconsistency, for all of its extensions, in particular of **set** theories as **ZF**, of the Elementary Theory of (higher Order) Topoi with Natural Numbers Object as considered by FREYD as well as already of the Theory of Cartesian Closed Categories with NNO considered by LAMBEK.

1 Introduction

Starting point is a discussion of CANTOR’s (indirect) argument for uncountability of the real numbers (in the unit interval), i. e. of the *set* $2^{\mathbb{N}} = \mathcal{P} \mathbb{N}$ of (“actual” infinit) sequences $a = a(j) : \mathbb{N} \rightarrow 2$.

This indirect argument assumes all these $a : \mathbb{N} \rightarrow 2$ to be enumerated in form $a_i = a_i(j) : \mathbb{N} \rightarrow 2$, $i \in \mathbb{N}$. CANTOR then takes as sequence outside this enumeration of the a_i the sequence $\tilde{a} = \tilde{a}(j) =_{\text{def}} \neg a_i(i) : \mathbb{N} \rightarrow 2$.

⁰ this is part 3 of a cycle on *Recursive Categorical Foundations*

⁰ Legend of LOGO: *closed evaluation* \in (part of Cartesian Closure) to give, with the help of “stratified” code *interpretation* into *Universal Chain* \mathcal{U} , code “self”-evaluation $\tilde{\varepsilon}$.

*TU Berlin, Mathematik, pfender@math.tu-berlin.de

[†]last revised August 23, 2021

But what is this $a_i(i) \in 2$? Let us try to apply CANTOR's argument to any type of *constructive* real numbers, where in fact there *is* an enumeration, a_i of all (finite) *texts*, (Computer) *programs*, standing for – “describing” – these constructive real numbers, e.g. the primitive recursive power-series descriptions for e and π . But if you want to *change* the diagonal *values* in this CANTOR's infinit table $a_i(j)$ of the *constructive* reals, you must be able to *evaluate* the i th of these – say primitive recursive – programs at $i \in \mathbb{N}$. Now ACKERMANN has shown, that for the case of PR function *codes* (“programs”, texts) this diagonal evaluation (and then its a posteriori modification) cannot be PR any more: The related (equi-complex) “Ackermann function”, namely *diagonal evaluation*

$\varepsilon(f_n, n) : \mathbb{N} \xrightarrow{\Delta} \mathbb{N} \times \mathbb{N} \xrightarrow{\# \times \text{id}_{\mathbb{N}}} [\mathbb{N}, \mathbb{N}] \times \mathbb{N} \xrightarrow{\varepsilon} \mathbb{N}$, grows faster than any PR function; here $\#(n) : \mathbb{N} \xrightarrow{\cong} [\mathbb{N}, \mathbb{N}]$ is the PR enumeration of all PR map PR codes f_n “from” \mathbb{N} “to” \mathbb{N} . The diagonal then says: “apply” n -th PR map to – *evaluate* n th PR map code *at* – argument n .

[Presumably this *non-closedness* under code-evaluation applies to any constructive class of real numbers and power sets, such real numbers obtained e.g by (iterated) “application” of Intermediate-Value Theorem taken as axiom.]

So the possibility of *closed evaluation*, here of

$$\in_{\mathbb{N},2} = \in_{\mathbb{N},2}(\chi, n) = \chi(n) = [n \in \chi] : 2^{\mathbb{N}} \times \mathbb{N} = \mathcal{P} \times \mathbb{N} \rightarrow 2$$

is at the basis of classical set theory, with its *closure* under (iterated) *formation* of power set (and internal hom sets). This gave rise to investigation of “all” the uncountable cardinalities in set theory, a central branch of this theory proper.

The claim of present investigation is that these uncountabilities, at least a (potentially) infinit ascending chain of uncountabilities, leads to a contradiction. The idea is to interpret the *map codes*, $\ulcorner f \urcorner \in [A, B]$ say, of a (minimally presented) theory **PR** of PR Arithmetic with (“higher Order”) Cartesian Closure added, into these maps “themselves”, $f \in B^A$, out of internal hom Object B^A , in **set** theory the *map set*

$$B^A = \{f \in \mathcal{P}(A \times B) \mid \forall a \in A \exists! b \in B (a, b) \in f\}.$$

Combined with *closed*, axiomatic evaluation $\in_{A,B} : B^A \times A \rightarrow B$, $\in_{A,B}(f, a) = f(a)$, available in **set** theory and there *needed* for (generalisation of) CANTOR's argument above to establish the strictly ascending hierarchy of *cardinals*, will give a *code-self-evaluation*, $\tilde{\varepsilon}_{\mathbb{N},2} : [\mathbb{N}, 2] \times \mathbb{N} \rightarrow 2$, and from this – because of the “*self*” – an (anti-)diagonal predicate $d = d(n) : \mathbb{N} \rightarrow 2 \xrightarrow{\neg} 2$, whence a *liar* map $\text{liar} = \neg \text{liar} : \mathbb{1} \rightarrow 2$ establishing the asserted contradiction for (minimal) Cartesian Closed PR Theory **PR** and its extensions.

We now outline the sections to come and foreshadow at this occasion some of the notations to be introduced:

2 Theory Closure by Internal hom and Evaluation:

Here we extend *basic* (categorical) Theory $\mathbf{PR}_A = \mathbf{PR} + (\text{abstr})$ of Primitive Recursion with (virtual) extensions $\{A|\chi\}$ of PR predicates (see part RCF1) by Cartesian Closure, this in form of adding just new internal hom *Objects*, B^A , new *map constants* $\in_{A,B} : \text{closed evaluation}$, and $\lhd_{A,B} : \text{for Cartesian Closure front adjunctions}$, as well as suitable *equations* for then already available *conjugate* and *coconjugate* maps, but **no** new (meta) operations for maps. Resulting Theory is called $\mathbf{PR}\in$, since its decisive ingredient over Theory \mathbf{PR} is closed evaluation $\in_{A,B} : B^A \times A \rightarrow B$ with its characteristic equations.

3 Order Stratification for Closed Arithmetic $\mathbf{PR}\in$

In this section we divide higher Order Theory $\mathbf{PR}\in$ into *strata* $\mathbf{PR}\in \underline{n} \prec \mathbf{PR}\in$, Cartesian PR theories with Order of Objects up to \underline{n} . **Note:** $\text{Ord}(C^B)^A \stackrel{\text{def}}{=} \text{Ord } C^{B \times A} < \text{Ord } C + \text{Ord } B + \text{Ord } A = \text{Ord } C^{B^A} \stackrel{\text{by def}}{=} \text{Ord } C^{(B^A)}$, “since” $(C^B)^A \cong C^{B \times A}$.

4 An Ascending, *Universal Object Chain*

Based on *Universal Object* $\mathbb{X} \subset \mathbb{N}$ for Theory \mathbf{PR}_A , \mathbb{X} made out of all (codes of) singletons $\langle n \rangle$ and (possibly nested) pairs $\langle a; b \rangle$ of natural numbers – it contains all Objects A of \mathbf{PR}_A coretractively embedded – we obtain an ascending *Chain* $\mathcal{U} : \mathcal{U}_1 = \mathbb{X} \xrightarrow{\sqsubseteq} \mathcal{U}_2 = \mathbb{X}^{\mathbb{X}} \dots$ of Objects and coretractions, each $\mathcal{U}_{\underline{n}}$ *universal* for its stratum $\mathbf{PR}\in \underline{n} : A \sqsubset \mathcal{U}_{\underline{n}}$ coretractive for each (pointed) Object A of Order up to \underline{n} .

5 Map-Code Interpretation

This section develops the central idea of present investigation: An *interpretation map family*

$$\text{int}^{\underline{n}} = [\text{int}_{A,B}^{\underline{n}} : [A, B]_{\mathbf{PR}\in \underline{n}} \rightarrow B^A]_{A,B}, \underline{n} \in \mathbb{N} \text{ external, “naive”}$$

is constructed, stratum by stratum, the $\text{int}^{\underline{n}}$ leading into Universal Object (at most) $\mathcal{U}_{2\underline{n}}$.

Technically, these Object-pairs indexed families (must and) can be “derived” from a stratum specific “global” Interpretation $\text{Int}^{\underline{n}} = \text{Int}^{\underline{n}}(u) : V_{\underline{n}} \rightarrow \mathcal{U}_{2\underline{n}}$, $V_{\underline{n}}$ the map code set of (whole) stratum $\mathbf{PR}\in \underline{n}$: reason for considering *Universal Objects*, here: $\mathcal{U}_{2\underline{n}}$.

What we have to do is to “*interpret*” code constants and *code operations*, namely (formal) composition, Cartesian product and iteration of map codes into the *objective correspondants*, e.g.– *plugged* into \mathbf{ZF} –

$$\ulcorner g \urcorner \odot \ulcorner f \urcorner = \ulcorner g \urcorner \ulcorner \circ \urcorner \ulcorner f \urcorner \stackrel{\text{by def}}{=} \ulcorner g \circ f \urcorner \stackrel{\text{int}}{\mapsto} \text{int}(g) \circ \text{int} f = g \circ f.$$

In our “formally minimal” context, this interpretation is based on the *name* $\lfloor f \rfloor : \mathbb{1} \rightarrow B^A$ of a map $f : A \rightarrow B$, $\lfloor f \rfloor$ easily defined via *conjugation*, in **set** theoretical terms: $\lfloor f \rfloor = \{(\emptyset, f)\} : \mathbb{1} \rightarrow B^A$.

Interpretation int works by the correspondence of operations $\odot = \ulcorner \circ \urcorner$, $\ulcorner \times \urcorner$, and $\ulcorner \S \urcorner$ on map *codes* for composition, Cartesian product and *iteration* one hand, and associated *internal closed* operations, called $\sqcup \circ \sqcup$, $\sqcup \times \sqcup$, as well as $\sqcup \S \sqcup$ on the other. These latter are all defined out of set theoretically motivated “coconjugated” ones, by *conjugation*.

Straightforward but technically complicated calculations then give the central **Interpretation Theorem**, saying essentially that (stratum specific) *interpretation*

$$\begin{aligned} \text{int}_{A,B}^{\underline{n}} : [A, B]_{\mathbf{PR} \in \underline{n}} &\rightarrow B^A \xrightarrow{\sqsubseteq} \mathcal{U}_{2\underline{n}} \text{ is } \textit{objective}, \text{ i. e.:} \\ \text{int}_{A,B}^{\underline{n}}(\ulcorner f \urcorner) &= \lfloor f \rfloor : \mathbb{1} \rightarrow B^A, \text{ for } f : A \rightarrow B \text{ in } \mathbf{PR} \in \underline{n}. \end{aligned}$$

6 Self-Evaluation

With interpretation properties above it is now easy to give a *sound, objective code-self-evaluation* for “minimal” Cartesian Closed PR Theory $\mathbf{PR} \in$, namely

$$\begin{aligned} \tilde{\varepsilon}_{A,B}(u, a) &=_{\text{def}} \in_{A,B}(\text{int}_{A,B}(u), a) : \\ [A, B]_{\mathbf{PR} \in} \times A &\xrightarrow{\text{int} \times A} B^A \times A \xrightarrow{\varepsilon} B, \text{ with} \\ \tilde{\varepsilon}_{A,B}(\ulcorner f \urcorner, a) &= f(a) : A \rightarrow B. \end{aligned} \quad (\text{Objectivity}).$$

This then gives immediately formalisation of Antinomie RICHARD for $\mathbf{PR} \in$ by the usual diagonal argument.

Notions and results for *basic* Theory $\mathbf{PR}_{\mathbf{A}} = \mathbf{PR} + (\text{abstr})$ of Free-Variables (categorical) Theory of Primitive Recursion with schema of predicate abstraction – and its Universal Object – are given in PFENDER/KRÖPLIN/PAPE 1994 and in PFENDER 2008 RCF1, RCFX.

2 Theory Closure by Internal hom and Evaluation

We **extend** here categorical Theory $\mathbf{PR}_{\mathbf{A}} = \mathbf{PR} + (\text{abstr})$ of Primitive Recursion – with *predicate abstraction* $\chi \mapsto \{A \mid \chi\}$ – into a Theory $\mathbf{PR} \in =_{\text{def}} \mathbf{PR}_{\mathbf{A}} + (\text{hom})$, with – in addition – *internal hom* $\langle A, B \rangle \mapsto B^A$ given by **axiom**, as well as theory internal – axiomatic, *closed* – evaluation

$$\in = [\in_{A,B} : B^A \times A \rightarrow B]_{A,B \in \mathbf{PR} \in}.$$

This in – logical – contrast to *constructive*, ACKERMANN type, formally *partial* – but still “*constructive*” – evaluation family

$$\varepsilon = [\varepsilon_{A,B} : [A, B] \times A \rightarrow B]_{A,B \in \mathbf{PR}_{\mathbf{A}}}$$

for theories $\pi_O \mathbf{R}$ (strengthening $\mathbf{PR}_{\mathbf{A}}$) above – *family* obtained out of one single (formally *partial* PR) map

$$\varepsilon = \varepsilon(u, x) : \mathbf{PR}_{\mathbf{A}} \times \mathbb{X} = [\mathbb{X}, \mathbb{X}]_{\mathbf{PR}_{\mathbf{A}}} \times \mathbb{X} \rightarrow \mathbb{X}.$$

Comment on Notation: *Closed evaluation* reads e.g.

$$\in_{A,2} = \in_{A,2}(\chi, a) = \chi(a) = [a \in \{A \mid \chi\}] : 2^A \times A = \mathcal{P}A \times A \rightarrow 2.$$

This motivates notation for closed evaluation. The “other” use of symbol “ \in ” is – in Cartesian Theories – “ $a \in A$ free”: a is a (*free*) *variable* on

A , categorical meaning: a is (identity of A) or a *projection* onto A . This legitimates free-variables diagram chase below categorically.

Theories \mathbf{PR}_A and $\mathbf{PR} \in$ fixed, we explain now some (known) basic concepts and results, in the language of Primitive Recursion and Higher Order Arithmetic sketched above.

Basic for our *Universal Chain* of Objects – *upwards open (!)* – is the *First Order Universal Object* $\mathbb{X} \subset \mathbb{N}$ of all (codes of) *singletons*, $\langle n \rangle$, and (possibly nested) *pairs*, $\langle a; b \rangle$, of natural numbers.

Each *fundamental* \mathbf{PR} -Object $\mathbb{1}, \mathbb{N}, (\mathbb{N} \times \mathbb{N})$ etc. is *coretractively embedded* into \mathbb{X} , for example

$$(\mathbb{N} \times \mathbb{N}) \ni (m, n) \xrightarrow{\sqsubseteq} \langle m; n \rangle \in (\mathbb{N} \times \mathbb{N}) \subset \mathbb{X}.$$

Extension of Theory \mathbf{PR}_A into *Cartesian Closed* Theory $\mathbf{PR} \in$ **presented** equationally – by HORN inferences – via additional (formal) *exponential* Objects (Object terms) of form (B^A) for A, B “already there”, examples: $\mathbb{N}^{\mathbb{N}}, \mathcal{U}2 = \mathbb{X}^{\mathbb{X}}, \mathcal{U}3 = \mathbb{X}^{(\mathbb{X}^{\mathbb{X}})}$ etc., as well as (additional) families of *map constants*

$$\in_{A,B} : B^A \times A \rightarrow B \text{ (axiomatic, closed evaluation),}$$

$$[\text{within set theory: } \in(f, a) =_{\text{by def}} f(a),] \text{ as well as}$$

$$\angle_{A,B} : A \rightarrow (A \times B)^B, \text{ closed front adjunction, “} A \ni a \xrightarrow{\angle} [b \mapsto (a, b)] \text{”}.$$

These two families are to satisfy the *adjointness* equations for (covariant) Functors, $A \times B \dashv B^A : \mathbf{PR} \in \longrightarrow \mathbf{PR} \in$, (A “fixed”), namely defining *conjugation* and *coconjugation* below as mutually inverse (meta) bijections.

These HORN schemata are merged with those of \mathbf{PR}_A , here: with forming *Cartesian products* of Objects, with *iteration* schema (and Freyd’s uniqueness of *initialised iterated*), as well as schema (abstr) of forming (*virtual*) *extensions*, cf part RCF 1.

Taken together the above internal hom structure with endo map iteration – and FREYD’s uniqueness of the *initialised iterated* – as well as with (virtual) *predicate abstraction* – we arrive at Theory $\mathbf{PR} \in = \mathbf{PR}_A + (\text{hom}) = \mathbf{PR} + (\text{abstr}) + (\text{hom})$, of *Primitive Recursion* with *Object exponentiation* and *closed evaluation*: Evaluation within the Theory itself.

[The latter in contrast to availability of “only” – ACKERMANN type, *not* \mathbf{PR} , (still) *constructive* – evaluation of Theory $\mathbf{PR}_A = \mathbf{PR} + (\text{abstr})$ within “only” Theory \mathbf{PR}_A – of formally *partial* \mathbf{PR} maps, theory equivalent to Theory $\mu\mathbf{R}$ of (partial) *mu*-recursive maps, see RCF1.]

Remark: Theory **Fin** of *finite* (number) sets *has* internal hom – exponentiation – coming with closed evaluation family $\in_{A,B} : B^A \times A \rightarrow B$. But if you want to **define** this – infinitely indexed family – made out of (finite) maps, you need Primitive Recursive case distinction on $\mathbb{N} \supset B^A$, and this “global”, *mother* evaluation

$$\hat{\in} : \mathbb{N} \times \mathbb{N} \supset \bigoplus_{A,B} (B^A \times A) \rightarrow B \subset \mathbb{N}$$

is necessarily *genuine* PR, *not finite*.

Internal hom – and “closed” evaluation \in – give, within Theory **PR** of *Higher order Arithmetic*, cf. EILENBERG & KELLY 1966 for internal hom structure, as well as FREYD 1972 and LAMBEK & SCOTT 1986 for the combined structure, the following **defined** map families *conjugation* and *coconjugation*:

Conjugation is **given** by **schema**

$$\begin{array}{l} \text{(conj)} \quad \frac{f : A \times B \rightarrow C \text{ in } \mathbf{PR} \in}{\overline{f} = \text{conj}[f] \quad =_{\text{def}} \quad f^B \circ \angle_{A,B} : A \rightarrow (A \times B)^B \rightarrow C^B,} \\ \text{in } \mathbf{set} \text{ theory conjugate } \overline{f} \text{ realised as} \\ a \xrightarrow{\overline{f}} [b \mapsto (a, b) \mapsto f(a, b) \in C], \end{array}$$

and *coconjugation* is introduced by **schema**

$$\begin{array}{l} \text{(coconj)} \quad \frac{g : A \rightarrow C^B \text{ in } \mathbf{PR} \in}{\overline{g} = \text{coconj}[g] \quad =_{\text{def}} \quad \in_{B,C} \circ (g \times \text{id}_B) : A \times B \rightarrow C^B \times B \rightarrow C,} \\ \text{in } \mathbf{set} \text{ theory coconjugate } \overline{g} \text{ realised as} \\ [(a, b) \mapsto (g(a), b) \mapsto \overline{g}(a, b) \quad =_{\text{by def}} \quad (g(a))(b) \in C]. \end{array}$$

These two families are to satisfy – by **axiom**, and do so (already within *finite*) **set** theory and the *Elementary Theory of Topoi* **ETT** in place of Theory **PR** around to be “constructed” – the following *higher order meta-bijection* equations:

$$\begin{array}{l} \text{(co/conj)} \quad \frac{f : A \times B \rightarrow C \text{ in } \mathbf{PR} \in}{\mathbf{PR} \in \vdash \quad \overline{\overline{f}} = \text{coconj}[\text{conj}[f]] = f : A \times B \rightarrow C} \end{array}$$

as well as

$$\begin{array}{l} \text{(conj/co)} \quad \frac{g : A \rightarrow C^B \text{ in } \mathbf{PR} \in}{\mathbf{PR} \in \vdash \quad \overline{\overline{g}} = \text{conj}[\text{coconj}[g]] = g : A \rightarrow C^B.} \end{array}$$

The above data, in particular (axiomatically given) *families* \angle and \in , **define** the following meta-map, and make it into a *covariant functor* hom – the *covariant internal hom functor* – via the following schema:

$$\begin{array}{l} \text{(hom-co)} \quad \frac{A, g : B \rightarrow C \text{ in } \mathbf{PR} \in}{\overline{g \circ \in} : B^A \times A \xrightarrow{\in} B \xrightarrow{g} C} \\ g^A \quad =_{\text{def}} \quad \overline{g \circ \in_{A,B}} : B^A \rightarrow C^A \end{array}$$

Analogous schema **defining** the *contravariant* (closed) internal hom *functor*:

$$\begin{array}{l} \text{(hom-contr)} \quad \frac{A, g : B \rightarrow C \text{ in } \mathbf{PR} \in}{\overline{\in \circ (A^C \times g)} : A^C \times B \xrightarrow{A^C \times g} A^C \times C \xrightarrow{\in} A} \\ A^g \quad =_{\text{def}} \quad \overline{\in \circ (A^C \times g)} : A^C \rightarrow A^B. \end{array}$$

All four: *Universal* property, the two Functor properties, and *right adjointness*, of covariant closed internal hom $g \mapsto g^A$ – namely right adjointness to *Cylindrification*

$$\langle g : B \rightarrow C \rangle \mapsto \langle A \times g : A \times B \rightarrow A \times C \rangle, \quad \text{Object } A \text{ fixed,}$$

are consequences of the pair $\text{conj}/\text{coconj}$ above to be a pair of meta-bijections, inverse to each other.

Remark; $\angle_{A,B}$ and $\in_{A,B}$ are natural transformations, but we will not rely on these properties here.

3 Order Stratification for Closed Arithmetic $\mathbf{PR} \in$

Definition: The – formal – Order Ord A of a *higher order Object* – of Theory $\mathbf{PR} \in$ – is **defined** externally \mathbf{PR} as follows:

$$\text{Ord } \mathbb{1}, \text{Ord } \mathbb{N} =_{\text{def}} 1,$$

$$\text{Ord } (A \times B) =_{\text{def}} \max\{\text{Ord } A, \text{Ord } B\},$$

$$\text{Ord } \{A \mid \chi : A \rightarrow 2\} =_{\text{def}} \text{Ord } A,$$

$$\text{in particular } \text{Ord } 2 = \text{Ord } \{n \in \mathbb{N} \mid n < 2\} = 1,$$

$$\text{Ord } \mathbb{X} = \text{Ord } \{\mathbb{N} \mid \mathbb{X} : \mathbb{N} \rightarrow 2\} = 1 \text{ (}\mathbb{X} \text{ is a predicative subset of } \mathbb{N}\text{.)}$$

For B in \mathbf{PR}_A and A in $\mathbf{PR} \in$ ($\text{Ord } A$ “already known”) :

$$\text{Ord } B^A = 1 + \text{Ord } A;$$

finally: for $C \in \mathbf{PR}_A$, B, C in $\mathbf{PR} \in$:

$$\begin{aligned} \text{Ord } (C^B)^A &=_{\text{def}} \text{Ord } C^{B \times A} =_{\text{by def}} 1 + \text{Ord } (B \times A) \\ &=_{\text{by def}} 1 + \max(\text{Ord } B, \text{Ord } A). \end{aligned}$$

The latter clause takes in account the (canonical) $\mathbf{PR} \in$ *reduction* isomorphism $(C^B)^A \cong C^{B \times A}$.

With this **definition**, we have in particular $\text{Ord } B^A \leq \text{Ord } B + \text{Ord } A$ for all $\mathbf{PR} \in$ Objects A, B , as well as $\text{Ord } \underline{U}\underline{n} = \underline{n}$, e.g. $\text{Ord } \underline{U}3 =_{\text{by def}} \text{Ord } \mathbb{X}^{\mathbb{X}^{\mathbb{X}}} = \text{Ord } \mathbb{X}^{(\mathbb{X}^{\mathbb{X}})} = 3$.

So subSystem \mathbf{PR}_A of Theory $\mathbf{PR} \in$ has all its (presenting) Objects of Order 1, it is our basic, “1st” Order, subSystem of Theory $\mathbf{PR} \in$ – not a priori an (“embedded”) subCategory, since the higher-order axioms of $\mathbf{PR} \in$ may entail – within $\mathbf{PR} \in$ – *new* equations between map terms of \mathbf{PR}_A viewed as map terms of $\mathbf{PR} \in$, in *logical* terms: The *Extension* $\mathbf{PR} \in$ of \mathbf{PR}_A may be not *conservative*.

Broadening to Theories Extension Chain: We **define** an exhaustive Chain of subSystems $\mathbf{PR} \in \underline{n} \preceq \mathbf{PR} \in$, $\underline{n} \in \mathbb{N}$, \mathbf{PR} as follows:

- $\mathbf{PR} \in 1 =_{\text{def}} \mathbf{PR}_A$;
- Assume $\mathbf{PR} \in \underline{n} \preceq \mathbf{PR} \in$ to be known via its (canonical) *presentation*:
Object terms, *map* terms, schemata for *map* (term) *equations*.

Then subSystem $\mathbf{PR} \in \underline{n}+ = \mathbf{PR} \in [\underline{n}+1]$ is **defined** to be the *Cartesian-PR-Category Closure* of subSystem $\mathbf{PR} \in \underline{n}$ merged with Closure under *formal adjunction* of

- all Objects of Order $\underline{n} + 1$
- the canonical isomorphisms $(C^B)^A \xrightarrow{\cong} C^{B \times A}$ given in $\mathbf{PR} \in$ for C in \mathbf{PR}_A , A, B in $\mathbf{PR} \in \underline{n}$, and their inverses $C^{B \times A} \xrightarrow{\cong} (C^B)^A$
- $\mathbf{PR} \in$ families $\angle_{A,B} : A \rightarrow (A \times B)^B$ as well as $\in_{A,B} : B^A \times A \rightarrow B$, this for $\text{Ord } A + \text{Ord } B$, $2 \text{Ord } B \leq \underline{n} + 1$, and $\text{Ord } A + \text{Ord } B \leq \underline{n} + 1$ respectively.

Additional (merged) **equations** come in, for the maps of $\mathbf{PR} \in \underline{n}+$, via schemata (co/conj) as well as (conj/co) of $\mathbf{PR} \in$ (above), which are to establish the *conjugation/coconjugation bijection* for all those of their instances, for which all formal ingredients – Object terms and map terms – are enumerated so far within $\mathbf{PR} \in \underline{n}+$.

Corollary to this **Definition**:

(i) *Conjugation upgrade*:

$$\text{(upgrade)} \quad \frac{f : A \times B \rightarrow C \text{ in } \mathbf{PR} \in \underline{n} \prec \mathbf{PR} \in,}{\overline{f} = \text{conj}[f] = f^B \circ \angle_{A,B} : A \rightarrow (A \times B)^B \rightarrow C^B \text{ lives in } \mathbf{PR} \in 2\underline{n} \prec \mathbf{PR} \in.}$$

(ii) *Coconjugation upgrade*:

$$\text{(co-upgrade)} \quad \frac{g : A \rightarrow C^B \text{ in } \mathbf{PR} \in \underline{n} \prec \mathbf{PR} \in}{\overline{g} = \text{coconj}(g) = \in_{B,C} \circ (g \times \text{id}_B) : A \times B \rightarrow C^B \times B \rightarrow C \text{ lives already in } \mathbf{PR} \in \underline{n} :}$$

Critical exponential Object C^B is presupposed to belong already to Theory $\mathbf{PR} \in \underline{n}$.

- (iii) Theory $\mathbf{PR} \in \underline{n}$ contains Objects up to Order \underline{n} , and in fact some of its Objects *have* this Order.
- (iv) External ascending “union” of all subSystems $\mathbf{PR} \in \underline{n}$, $\underline{n} \in \mathbb{N}$, *exhausts* Theory $\mathbf{PR} \in$, i.e. gives a – *stratified* – **presentation** of Theory $\mathbf{PR} \in$: Objects, maps, and equations.

4 An Ascending, *Universal* Object Chain

Basic – 1st Order – Arithmetical Theory $\mathbf{PR} \in 1 = \mathbf{PR}_A$ has a *Universal Object* in itself, a *first-Order Universal Object*, namely the Object $\mathbb{X} \subset \mathbb{N}$ – of (codes of) all singleton (lists) and of pairs, possibly nested: binary bracketed NNO tuples.

\mathbb{X} is a Universal Object – of Theory \mathbf{PR}_A and therefore also of its *stengthenings*, as for example for the full first order subcategory $\mathbf{PR} \in 1$ of $\mathbf{PR} \in$. Object \mathbb{X} is *universal* in the following sense:

\mathbb{X} admits – for each \mathbf{PR}_A -Object A , an *embedding* (here an *injective map*), even a *coretractive* map (see below), $\sqsubset_A: A \xrightarrow{\sqsubset} \mathbb{X}$, **defined** externally PR in the obvious way.

All these embeddings $\sqsubset_A: A \xrightarrow{\sqsubset} \mathbb{X}$ – *disjoint* as far as *fundamental* Objects A are concerned, namely binary bracketed powers of \mathbb{N} , no genuine *abstracted* sets – come with canonical *retractions* $\sqsupset_A: \mathbb{X} \xrightarrow{\sqsupset} A$, the latter equally for abstracted Objects $\{A | \chi\}$ having a *point*, $a_0: \mathbb{1} \rightarrow \{A | \chi\}$, as in particular $\mathbb{X} \subset \mathbb{N}$, coming with “its” *zero* $\langle 0 \rangle: \mathbb{1} \rightarrow \mathbb{X}$.

Graded-Universal-Object Chain: Each of our Theories $\mathbf{PR} \in \underline{n}$ in the hierarchy – except (!) “roof” Theory $\mathbf{PR} \in$ itself – comes with a *canonical* Universal Object, $\mathcal{U}_{\underline{n}} = \mathbb{X}^{\uparrow \underline{n}}$, externally PR **defined** as follows, as an internal version of a GROTHENDIECK-*Universe* (?):

$$\begin{aligned} \mathcal{U}_1 &= \mathbb{X}^{\uparrow 1} =_{\text{def}} \mathbb{X}^1 = \mathbb{X}, \\ \mathcal{U}_{\underline{n}+1} &=_{\text{def}} \mathbb{X}^{\mathcal{U}_{\underline{n}}} = \mathbb{X}^{\mathbb{X}^{\uparrow \underline{n}}} =_{\text{by def}} \mathbb{X}^{\uparrow \underline{n}+1} \end{aligned}$$

For opening the possibility that a *higher, later* Universal Object in the chain is good also as Universal Object for a *lower, earlier* Theory in the hierarchy, we establish first the *Universal Chain* \mathcal{U} as a chain of *embeddings* $\sqsubset = \sqsubset_{\underline{n}}: \mathcal{U}_{\underline{n}} \rightarrow \mathcal{U}_{\underline{n}+1}$ coming each with a retraction $\sqsupset = \sqsupset_{\underline{n}}: \mathcal{U}_{\underline{n}+1} \rightarrow \mathcal{U}_{\underline{n}}$, as follows:

Universal Chain \mathcal{U} begins with (commutative) DIAGRAM

$$\begin{array}{ccc} & \mathbb{X}^1 & \\ \swarrow \scriptstyle \sqsupset_{\mathbb{X}} & & \nwarrow \scriptstyle \mathbb{X}^1 \xrightarrow{\langle 0 \rangle} \mathbb{X} \\ \mathcal{U}_1 = \mathbb{X} & \xleftrightarrow[\sqsubset]{\sqsubset} & \mathbb{X}^{\mathbb{X}} = \mathcal{U}_2 \\ \searrow \scriptstyle \bar{\ell}_{\mathbb{X},1} & & \swarrow \scriptstyle \mathbb{X}^{\mathbb{X}} \xrightarrow{!} \mathbb{1} \\ & \mathbb{X}^1 & \end{array}$$

Diagram chase in case of set theory:

$$\begin{array}{ccc} & [0 \mapsto \langle 0 \rangle \mapsto x] & \\ \swarrow \scriptstyle \sqsupset_{\mathbb{X}} & & \nwarrow \scriptstyle \mathbb{X}^1 \xrightarrow{\langle 0 \rangle} \mathbb{X} \\ x & \xleftrightarrow[\sqsubset]{\sqsubset} & [y \mapsto 0 \mapsto x] \\ \searrow \scriptstyle \bar{\ell}_{\mathbb{X},1} & & \swarrow \scriptstyle \mathbb{X}^{\mathbb{X}} \xrightarrow{!} \mathbb{1} \\ & [0 \mapsto x] & \end{array}$$

The general Universal Chain member then is recursively **defined** by commutativity of DIAGRAM

$$\begin{array}{ccc}
& \mathbb{X}^{\mathbb{U}_{\underline{n}-1} \sqsubset \mathcal{U}_{\underline{n}}} & \\
\mathcal{U}_{\underline{n}} = \mathbb{X}^{\mathbb{U}_{\underline{n}-1}} & \begin{array}{c} \xleftarrow{\sqsupset} \\ \xrightarrow{\sqsubset} \end{array} & \mathbb{X}^{\mathcal{U}_{\underline{n}}} = \mathcal{U}_{\underline{n}+1} \\
& \mathbb{X}^{\mathcal{U}_{\underline{n}} \sqsupset \mathbb{U}_{\underline{n}-1}} &
\end{array}$$

Easy Diagram chase for verifying section/retraction property e.g. in **set** theory.

Generalising the above to the case of B^A instead of $\mathbb{X}^{\mathbb{U}_{\underline{n}-1}}$ we now **define** recursively the (coretractive) embeddings

$$\sqsubset = \sqsubset_{B^A} : B^A \rightarrow \mathcal{U}_{\underline{n}+1}, \quad B^A \text{ in } \mathbf{PR} \in \underline{n}+ = \mathbf{PR} \in [\underline{n} + 1],$$

based on the (coretractive) embeddings $\sqsubset_B : B \rightarrow \mathbb{X} = \mathbb{U}_1$ above, as follows, “but” first only for Object B in $\mathbf{PR} \in 1 = \mathbf{PR}_A$:

– Anchor: for A in \mathbf{PR}_A , (natural) embedding $\sqsubset_A : A \rightarrow \mathcal{U}_1 =_{\text{by def}} \mathbb{X}^1 = \mathbb{X}$ has been **defined** above by converting natural numbers n in singleton codes $\langle n \rangle$, and – recursively – pairs in code pairs, out of \mathbf{PR}_A *Universal Object* $\mathbb{X} \subset \mathbb{N}$. Furthermore, a canonical *retraction* $\sqsupset_A : \mathbb{X} \rightarrow A$ for the embedding has been mentioned above, for Object A coming with a *point*, $a_0 : 1 \rightarrow A$ say.

– Step: Assume embedding $\sqsubset_A : A \rightarrow \mathcal{U}_{\underline{n}}$ to be given, together with retraction $\sqsupset_A : \mathcal{U}_{\underline{n}} \rightarrow A$, for “each” Object A of Order \underline{n} – in $\mathbf{PR} \in \underline{n}$.

Consider then a (genuine) Object in $\mathbf{PR} \in \underline{n}+$, of form B^A , A in $\mathbf{PR} \in \underline{n}$, B in \mathbf{PR}_A (!). Then the DIAGRAM below – simplified one of the former one above – **defines** “universal” embedding and retraction for Object B^A into/from $\mathcal{U}_{\underline{n}+1} =_{\text{by def}} \mathbb{X}^{\mathcal{U}_{\underline{n}}}$:

$$\begin{array}{ccc}
& B^{\mathcal{U}_{\underline{n}}} & \\
B^{\sqsupset A} \swarrow & & \searrow \sqsupset_B^{\mathcal{U}_{\underline{n}}} \\
B^A & \begin{array}{c} \xleftarrow{\sqsupset_{B^A}} \\ \xrightarrow{\sqsubset_{B^A}} \end{array} & \mathbb{X}^{\mathcal{U}_{\underline{n}}} = \mathcal{U}_{\underline{n}+1} \\
\searrow \sqsubset_B^A & & \swarrow \mathbb{X}^{\mathcal{U}_{\underline{n}} \sqsupset A} \\
& \mathbb{X}^A &
\end{array}$$

Again easy Diagram chase for verifying section/retraction property in case of **set** theory.

The general, not *normal form* case, of a $\mathbf{PR} \in \underline{n}+$ Object of form B^A , $B = D^C$ not basic, not in \mathbf{PR}_A , is reduced to the above one via (natural) isomorphism $(D^C)^A \cong D^{(C \times A)}$ – such isomorphism possibly applied several times –, to a *normal form* case Object to be embedded, by a map *within* $\mathbf{PR} \in \underline{n}+$ (or lower) *into* $\mathcal{U}_{\underline{n}+1}$ or lower, by the method above for the case of Object B in \mathbf{PR}_A . Embedding into $\mathcal{U}_{\underline{n}+1}$ in the latter case then is by composition with embedding $\mathcal{U}_{\underline{m}} \xrightarrow{\sqsubset} \mathcal{U}_{\underline{n}+1}$, $\underline{m} < \underline{n} + 1$.

Taken together the above – including the modification for the non-normal-form case – we have $\mathbf{PR} \in \underline{n} +$ embedded all Objects of $\mathbf{PR} \in \underline{n} +$ into $\mathcal{U}_{\underline{n}+1}$, namely all $\mathbf{PR} \in$ Objects of – up to – Order $\underline{n} + 1$. This **proves**

Embedding Theorem for Chain \mathcal{U} :

- (i) Each single of our Theories $\mathbf{PR} \in \underline{n}$ admits *coretractive* embeddings $\sqsubset_A: A \xrightarrow{\sqsubset} \mathcal{U}_{\underline{n}}$ for each of its (pointed) Objects A , into “its” *Universal Object* within the *section/retraction Chain*

$$\mathcal{U} : \quad \mathcal{U}_1 = \mathbb{X} \xrightleftharpoons{\quad} \xrightleftharpoons{\quad} \mathcal{U}_{\underline{n}} \xrightleftharpoons{\quad} \mathcal{U}_{\underline{n}+1} \xrightleftharpoons{\quad}$$

of these “Universal” Objects, the Chain \mathcal{U} hosted as an ascending chain in *global, higher Order* Theory $\mathbf{PR} \in$.

- (ii) By the above discussion of – canonical – natural retractions $\sqsupset_{\underline{n}}: \mathcal{U}_{\underline{n}+1} \rightarrow \mathcal{U}_{\underline{n}}$, retractions to embeddings $\sqsubset_{\underline{n}}: \mathcal{U}_{\underline{n}} \rightarrow \mathcal{U}_{\underline{n}+1}$, the above coretractive embedding for all Objects of $\mathbf{PR} \in \underline{n}$, into $\mathcal{U}_{\underline{n}}$, gives also (canonical) embeddings into *later* Objects of chain \mathcal{U} , i. e. if $\mathcal{U}_{\underline{n}}$ is replaced by $\mathcal{U}_{\underline{m}}$, $\underline{m} > \underline{n}$, and (coretractive) *embedding* $A \rightarrow \mathcal{U}_{\underline{m}}$ is taken as $\sqsubset_A: A \xrightarrow{\sqsubset} \mathcal{U}_{\underline{n}} \xrightarrow{\sqsubset} \dots \xrightarrow{\sqsubset} \mathcal{U}_{\underline{m}}$.

5 Map-Code Interpretation

Using Order Stratification above – of *higher order Cartesian Closed Theory* $\mathbf{PR} \in = \mathbf{PR}_{\mathbf{A}} + (\text{hom})$ – we now **define** – via \mathbf{PR} – a Theory-internal *interpretation map family*

$$\text{int}^{\underline{n}} = [\text{int}_{A,B}^{\underline{n}} : [A, B]_{\underline{n}} =_{\text{def}} [A, B]_{\mathbf{PR} \in \underline{n}} \rightarrow B^A]_{A,B}, \quad \underline{n} \in \mathbb{N},$$

A, B Objects of *stratum* $\mathbf{PR} \in \underline{n}$; interpretation $\text{int}_{A,B}^{\underline{n}}$ will be **defined** inside stratum $\mathbf{PR} \in 2\underline{n}$.

Example: $\text{int}_{\mathbb{N},2}^1 : [\mathbb{N}, 2]_{\mathbf{PR}_{\mathbf{A}}} = [\mathbb{N}, 2]_{\mathbf{PR} \in 1} \rightarrow 2^{\mathbb{N}}$ will live inside stratum $\mathbf{PR} \in 2$ – and higher –, see discussion in foregoing section.

[Such a *stratum* is a \mathbf{PR} Cartesian Theory, but it is *truncated* what concerns (exponential) Order of Objects and (axiomatic) evaluation. We will see below – in particular for our *interpretation of constructive*, \mathbf{PR} defined “internal” hom sets $[A, B]$ into *closed* ones B^A , that it is sufficient to climb up to stratum $2\underline{n}$ for interpretation of stratum \underline{n} .]

In our present – categorical – context, family $\text{int}_{A,B} = \text{int}_{A,B}^{\underline{n}}$, $\underline{n} \in \mathbb{N}$ fixed, *can* and *must* (?) be **defined** formally as (a family) derived from *one* single $\mathbf{PR} \in 2\underline{n}$ map. So, as one Interpretation for all – on stratum $\mathbf{PR} \in \underline{n}$ fixed – we are lead to **define** – \mathbf{PR} over $\mathbf{PR} \in 2\underline{n}$ – this *global* Interpretation as a $\mathbf{PR} \in 2\underline{n}$ map, with suitable, *universal*, *Domain* and *CoDomain*.


We start by *type-description* of this family – to be defined, later, as a family of *Domain/Codomain* restrictions of the *one single* map $\text{Int}^{\underline{n}}$ of Theory $\mathbf{PR} \in \underline{n}$ to be (objectively) \mathbf{PR} **defined** – of following type:

$\text{Int}^{\underline{n}} = \text{Int}^{\underline{n}}(u) : V_{\underline{n}} =_{\text{def}} \bigoplus_{A,B} |A, B|_{\underline{n}} \dashrightarrow \mathcal{U}_{2\underline{n}}$, where $|A, B|_{\underline{n}}$ is an abbreviation for internal, *syntactical* $\mathbf{PR} \in \underline{n}$ -map code (!) set $[A, B]_{\mathbf{PR} \in \underline{n}} \subset V \subset \mathbb{N}$, from A to B , A, B both Objects of $\mathbf{PR} \in \underline{n}$.

We turn now our “typifying” proposal (!) above, into a DIAGRAM which displays a special – central – countable sum (“disjoint union”), and its (litteral) component-inclusions. This “special” *sum*-DIAGRAM is available within $\mathbf{PR} \in \underline{n}$ – as *litteral*, disjoint union of *predicates*, disjoint by definition.

Global $\mathbf{PR} \in \underline{n}$ Interpretation $\text{Int}^{\underline{n}}$, (\underline{n} fixed), to be **defined** following actual type-discussion, then will be **characterised** a posteriori (!) as $\mathbf{PR} \in 2\underline{n}$ map, *induced* map out of the (countable) sum, induced by its *components* $\text{int}_{A,B}^{\underline{n}} : |A, B|_{\underline{n}} \rightarrow B^A \sqsubset \mathcal{U}_{2\underline{n}}$, A, B in stratum $\mathbf{PR} \in \underline{n}$.


In other words: $\text{Int}^{\underline{n}}$ will be *PR* “constructed” – “over” $\mathbf{PR} \in \underline{n}$, “but only” within $\mathbf{PR} \in 2\underline{n}$ – in such a way that it becomes the (unique) $\mathbf{PR} \in 2\underline{n}$ map out of **sum** $V_{\underline{n}} \subset \mathbb{N}$, which makes *commute* the following (externally) countable DIAGRAM, this diagram *available* within $\mathbf{PR} \in 2\underline{n}$:



Interpretation map DIAGRAM (A, B in $\mathbf{PR} \in \underline{n}$)

\mathbf{PR} Construction of $\mathbf{PR} \in \underline{n}$ map $\text{Int}^{\underline{n}} : \mathbb{N} \supset V_{\underline{n}} = \bigoplus_{A,B} [A, B] \rightarrow \mathcal{U}_{2\underline{n}}$

is recursively *merged* with that of maps $\text{int}_{A,B}^{\underline{n}} : |A, B|_{\underline{n}} \rightarrow B^A$, the latter being (recursively) **defined** as Domain/Codomain restrictions of *universal* \mathbf{PR} defined Interpretation map $\text{Int}^{\underline{n}}$ within $\mathbf{PR} \in 2\underline{n}$, in fact by the followig **defining** commutative DIAGRAM (B pointed):



This type of restriction becomes *possible* – at least easier – by the fact that “all” maps considered come as *section/retraction* pairs. This is in particular the case for all *injections-into-sums* embeddings here to be treated.

Constructive Internalisation of meta operations for our Theories $\mathbf{PR} \in \underline{n}$ and subSystems $\mathbf{PR} \in \underline{n}$:

Composition $\circ : \mathbf{T} \times \mathbf{T} \longrightarrow \mathbf{T}(A, C)$ of \mathbf{T} – Theory \mathbf{T} any (categorical) theory – *constructively* internalises to

$\odot = \ulcorner \circ \urcorner : [B, C]_{\mathbf{T}} \times [A, B]_{\mathbf{T}} \rightarrow [A, C]_{\mathbf{T}}$, $(v, u) \mapsto \langle v \odot u \rangle \in [A, C]_{\mathbf{T}}$. As Objects A, B here all Objects of \mathbf{T} are allowed, for $\mathbf{T} := \mathbf{PR} \in \underline{n}$ in particular Object $\mathcal{U}_{\underline{n}}$ and its (embedded) subobjects.

Analogously Cartesian product “ \times ” has as *coded version* family $[A, B]_{\mathbf{T}} \times [C, D]_{\mathbf{T}} \ni (u, v) \xrightarrow{\ulcorner \times \urcorner} \langle u \ulcorner \times \urcorner v \rangle \in [A \times C, B \times D]_{\mathbf{T}}$, for (arbitrary) \mathbf{T} -Objects A, B, C, D , including in particular Objects $\mathcal{U}_{\underline{n}}$ in case of theory $\mathbf{T} := \mathbf{PR} \in \underline{n}$.

Analogously for *iteration* “ \S ” within (Cartesian) PR theories in particular “again” for extension $\mathbf{PR} \in \underline{n}$ of PR Theory $\mathbf{PR}_{\mathbf{A}} = \mathbf{PR} + (\text{abstr}) : [A, A] \ni v \xrightarrow{\ulcorner \S \urcorner} v \ulcorner \S \urcorner \in [A \times \mathbb{N}, A]$, here e.g. for iteration of $\mathbf{PR} \in \underline{n}$ endo maps with Domain $A := \mathcal{U}_{\underline{n}}$ and their internalisations.

Definition: The *constructive* $\mathbf{PR} \in \underline{n}$ -codes in $V_{\underline{n}}$ are – **first** – the *constructive internal map-constants*

$$\ulcorner \wedge_{A,B} \urcorner : \mathbb{1} \rightarrow |A, (A \times B)^B|_{\underline{n}}, \text{ and } \ulcorner \in_{A,B} \urcorner : \mathbb{1} \rightarrow |B^A \times A, B|_{\underline{n}}$$

for $\wedge_{A,B}, \in_{A,B}$ in $\mathbf{PR} \in \underline{n}$.

Second: the “derived” *Cartesian map constants* for the new Objects and their Cartesian products – with the “old” ones and with the new ones –: identities, terminal maps, (left and right) projections, and

Third: “Closure” under *composition* and *cylindrification* (Cartesian product with an identity) as well as under *iteration* of endo maps.

Next we **define**, for $f : A \rightarrow B$ in $\mathbf{PR} \in \underline{n}$ – and hence in particular Objects A, B in $\mathbf{PR} \in \underline{n}$, the notion *name of* $f : A \rightarrow B$, symbolised as $[f] = [f : A \rightarrow B] : \mathbb{1} \rightarrow B^A$, available in stratum $2\underline{n}$.

This up-to- $2\underline{n}$ th Order construct $[f]$ is defined simply by *conjugation*, as $[f] = \overline{f \circ r_{\mathbb{1}, A}} = \text{conj}[f \circ r : \mathbb{1} \times A \xrightarrow{\cong} A \rightarrow B] : \mathbb{1} \rightarrow B^A$.

Name $[f]$ of f represents, meta-bijectively, map $f : A \rightarrow B$ *within* – as *defined element* of – *closed* internal hom set B^A . In **set** theory: $[f] =_{\text{def}} \{(\emptyset, f)\} : \mathbb{1} \rightarrow B^A \subset \mathcal{P}(A \times B)$.

By its **definition** via *conjugation*, $[f]$ has characteristic property $\in_{A,B}([f], a) = f(a) = f : A \rightarrow B$, $a \in A$ free. Verification of this *closed Objectivity* from definition is trivial for set theoretic environment, and straight forward for the general higher Order case.

Definition of *global Interpretation* $\text{Int}^{\underline{n}} = \text{Int}^{\underline{n}}(u) : \mathbb{N} \supset V_{\underline{n}} \rightarrow \mathcal{U}_{2\underline{n}}$, of (\underline{n} -truncated), internal *map-code-set* $V_{\underline{n}}$ of Theory $\mathbf{PR} \in \underline{n}$ into $\mathbf{PR} \in 2\underline{n}$ ’s *Universal Object* $\mathcal{U}_{2\underline{n}}$ – within (the language of) Theory $\mathbf{PR} \in 2\underline{n}$, is by *recursive case distinction* on the *structure* of the *map code* $u \in V_{\underline{n}}$ to be *interpreted*. (At beginning we do not typify into types A, B for $|A, B|_{\underline{n}} \subset V_{\underline{n}}$.) This PR **case distinction** for Definition of *Interpretation* $\text{Int}^{\underline{n}}(u) : V_{\underline{n}} \rightarrow \mathcal{U}_{2\underline{n}}$ runs as follows:

– **Case of $\mathbf{PR} \in \underline{n}$ map constants** “bas”, namely $0 : \mathbb{1} \rightarrow \mathbb{N}$ and $s : \mathbb{N} \rightarrow \mathbb{N}$ as well as *all* Cartesian map constants of $\mathbf{PR} \in \underline{n}$: identities, terminal maps, diagonals, (binary) projections, as well as case of the additional – *closed* – map constants $\angle_{A,B}, \in_{A,B}$ of $\mathbf{PR} \in \underline{n}$:

For all of these *anchor cases*, we **define Interpretation** $\text{Int}^{\underline{n}} = \text{Int}^{\underline{n}}(u) : V_{\underline{n}} \rightarrow \mathcal{U}_{2\underline{n}}$ in the below – PR – by “codes to names:”

$$\text{Int}(\ulcorner \text{bas} \urcorner) =_{\text{def}} \lfloor \text{bas} \rfloor : \mathbb{1} \rightarrow \mathcal{U}_{2\underline{n}}, \text{ e.g.}$$

$$\text{Int}(\ulcorner \ell : A \times B \rightarrow A \urcorner) =_{\text{def}} \lfloor \ell : A \times B \rightarrow A \rfloor : \mathbb{1} \rightarrow A^{A \times B} \xrightarrow{\sqsubset} \mathcal{U}_{2\underline{n}},$$

Objects A, B in stratum $\mathbf{PR} \in \underline{n}$.

This gives in particular for the “extra” $\mathbf{PR} \in \underline{n}$ basic codes, with appropriate $\mathbf{PR} \in \underline{n}$ Objects as *types*:

$$\text{Int}(\ulcorner \angle_{A,B} : A \rightarrow (A \times B)^B \urcorner) =_{\text{def}} \lfloor \angle_{A,B} \rfloor :$$

$$\mathbb{1} \rightarrow ((A \times B)^B)^A \xrightarrow{\sqsubset} \mathcal{U}_{2\underline{n}}, \text{ as well as}$$

$$\text{Int}(\ulcorner \in_{A,B} : B^A \times A \rightarrow B \urcorner) =_{\text{def}} \lfloor \in_{A,B} \rfloor : \mathbb{1} \rightarrow B^{B^A \times B} \xrightarrow{\sqsubset} \mathcal{U}_{2\underline{n}},$$

The latter two “inclusions” $\sqsubset \mathcal{U}_{2\underline{n}}$ are available by the fact that $\angle_{A,B}$ and $\in_{A,B}$ were supposed to live “already” within $\mathbf{PR} \in \underline{n}$, and that conjunction – at the base of *name* $\lfloor f \rfloor$ – at most doubles Order of (minimal) “receiving” stratum, here Order \underline{n} .

What we still have to worry about is *self-referential (!)* Interpretation of family members $\text{int}_{A,B}^{\underline{n}} : [A, B] \rightarrow B^A$, obtained from $\text{Int}^{\underline{n}} : V_{\underline{n}} = \mathbf{PR} \in \underline{n} \rightarrow \mathcal{U}_{2\underline{n}}$ by *Domain/CoDomain restriction*.

For these *injections* into *sum* $V_{\underline{n}}$, we *will* obtain (!), out of our PR **case-definition** of *global* Interpretation $\text{Int}^{\underline{n}} : V_{\underline{n}} \rightarrow \mathcal{U}_{2\underline{n}}$, by **definition** – below – of families

$\text{int}_{A,B}^{\underline{n}} : [A, B]_{\underline{n}} \rightarrow B^A$ as Domain/CoDomain *restrictions* of $\text{Int}^{\underline{n}} :$

$$\text{Int}^{\underline{n}}(\ulcorner \text{int}_{A,B}^{\underline{n}} \urcorner) = \lfloor \text{int}_{A,B}^{\underline{n}} \rfloor : \mathbb{1} \rightarrow (B^A)^{[A,B]} \cong B^{A \times [A,B]} \text{ within } \mathbf{PR} \in 2\underline{n}.$$

The latter map will lead in fact – Order verification – into $\mathbf{PR} \in 2\underline{n}$ by our **definition** of

$$\begin{aligned} \text{Ord}(B^A)^{[A,B]} &= \text{Ord } B^{A \times [A,B]} \\ &\leq \text{Ord } B + \max(\text{Ord } A, \text{Ord } [A, B]) \\ &= \text{Ord } B + \max(\text{Ord } A, \text{Ord } \mathbb{N}) \\ &\leq \text{Ord } B + \text{Ord } A \leq 2\underline{n}, \end{aligned}$$

and since the isomorphism pair $(B^A)^{[A,B]} \cong B^{A \times [A,B]}$ is included in $\mathbf{PR} \in 2\underline{n}$ by **definition** of stratum $\mathbf{PR} \in \underline{n}+ = \mathbf{PR} \in [\underline{n} + 1]$.

Based on the anchor cases above, we define by *genuine primitive recursion* stratum Interpretation $\text{Int}^{\underline{n}}$ of (constructively) *composed* codes, *Cartesian* “parallelised” as well as of *iterated* ones, as follows by PR

case distinction on *Iteration Domain* for **PR definition** of $\text{Int}^{\underline{n}} : V_{\underline{n}} \rightarrow \mathcal{U}_{2\underline{n}}$, PR case distinction on the *disjoint components* $|A, B|_{\underline{n}}$ of “syn-tactic (code) universe” $V_{\underline{n}} \subset \mathbb{N}$, which in turn is a PR defined predicative subObject of \mathbb{N} within Theory **PR_A** – in the rôle of (internal) Metamathematics – **PR_A subSystem** of $\text{PR} \in \underline{n} \prec \text{PR} \in [= “\text{PR} \in \infty”]$.

With – always below – **abbreviation**

$|A, B|_{\underline{n}} =_{\text{by def}} [A, B]_{\text{PR} \in \underline{n}} \subset V_{\underline{n}} = \text{PR} \in \underline{n} \subset \mathbb{N}$, we introduce **PR_{in} \underline{n}** map (map-family, indexed on $\underline{n} \in \mathbb{N}$)

$$\text{Int}^{\underline{n}} = \text{Int}^{\underline{n}}(u) : V_{\underline{n}} = \bigoplus_{A, B} |A, B|_{\underline{n}} \rightarrow \mathcal{U}_{2\underline{n}},$$

merged with its *Domain/Codomain* restrictions, recursively as follows:

Interpretation of *constructive internal composition*: For A, B, C in stratum **PR_{in} \underline{n}** :

$$\begin{aligned} & \text{for } u \in |A, B|_{\underline{n}} \subset V_{\underline{n}}, \ v \in |B, C|_{\underline{n}} \subset V_{\underline{n}} \\ & \quad [\implies \langle v \odot u \rangle \in |A, C|_{\underline{n}} \subset V_{\underline{n}}] : \\ & \text{Int}^{\underline{n}} \langle v \odot u \rangle \quad [= \text{int}_{A, C}^{\underline{n}}(\langle v \odot u \rangle) \subset V_{\underline{n}}] \\ & =_{\text{def}} \text{Int}^{\underline{n}}(v) \circ_{\perp} \text{Int}^{\underline{n}}(u) \\ & =_{\text{by def}} \circ_{\perp} (\text{Int}^{\underline{n}}(v), \text{Int}^{\underline{n}}(u)) : \\ & V_{\underline{n}} \times V_{\underline{n}} \xrightarrow{\supset} |B, C|_{\underline{n}} \times |A, B|_{\underline{n}} \xrightarrow{\text{int}_{B, C}^{\underline{n}} \times \text{int}_{A, B}^{\underline{n}}} C^B \times B^A \\ & \xrightarrow{\circ_{\perp}} C^A \sqsubseteq \mathcal{U}_{2\underline{n}}. \end{aligned}$$

This is a formally defined **PR_{in} $2\underline{n}$** map, in particular since $V_{\underline{n}} \times V_{\underline{n}} \xrightarrow{\supset} |B, C|_{\underline{n}} \times |A, B|_{\underline{n}}$ is – obviously – a *retraction*. We recall further that “embedding” $C^A \xrightarrow{\sqsubseteq_{C^A}} \mathcal{U}_{2\underline{n}}$ also comes with a retraction, $\mathcal{U}_{2\underline{n}} \xrightarrow{\supset_{C^A}} C^A$.

Axiomatic internal composition – *competing* with *constructive internal composition* $\odot = \lceil \circ \rceil$, gets a similar symbol, \circ_{\perp} , which may be read *Closed internal composition*, similarly: \times_{\perp} : *Closed internal Cartesian product*, as well as \S_{\perp} for *Closed internal iteration*: For the general background on *Closed Categories* see EILENBERG & KELLY 1966.

Closed internal composition $\circ_{\perp} : \mathcal{U}_{2\underline{n}} \xrightarrow{\supset} C^B \times B^A \xrightarrow{\circ_{\perp} A, B, C} C^A$, (retraction $\sqsupset = \sqsupset_{C^B \times B^A}$ just cares on – feasible – case distinction)

is **defined** via *conjugate* $\circ_{\perp} = \overline{\overline{\circ_{\perp}}}$ of

$$\begin{aligned} & \overline{\overline{\circ_{\perp}}}_{A, B, C} : (C^B \times B^A) \times A \rightarrow C \\ & =_{\text{def}} [\in \circ (C^B \times \in) : C^B \times B^A \times A \xrightarrow{C^B \times \in} C^B \times B \xrightarrow{\in} C], \end{aligned}$$

with Cartesian *associativity* (natural) isomorphisms of form

$$A \times B \times C =_{\text{def}} (A \times B) \times C \xrightarrow[\cong]{\text{ass}} A \times (B \times C) \text{ omitted.}$$

This **case** of **PR**∈-map $\text{Int}^{\underline{n}} : V_{\underline{n}} \xrightarrow{\supseteq} |A, C|_{\underline{n}} \rightarrow C^A \xrightarrow{\sqsubseteq} \mathcal{U}_{2\underline{n}}$ describes in fact a **PR**∈ $2\underline{n}$ map:

In its chain of Objects – and in its Order minimal presentation of maps – it is at most of Order $2\underline{n}$ – for Objects A, B, C all of Order at most \underline{n} .

– *Interpretation* of constructive internal *product of maps*: This is analogous to the above, even easier, since the two components of a Cartesian product are completely independent of each other, “exercise”.

– **Case** of an internally *iterated* $v^{\ulcorner \S \urcorner} \in |A \times \mathbb{N}, \mathbb{N}|_{\underline{n}}$, $v \in |A, A|_{\underline{n}}$ free, Object A in **PR**∈ \underline{n} . **Define** in this case

$$\begin{aligned} V_{\underline{n}} \supset |A, A|_{\underline{n}} \ni v &\xrightarrow{\text{Int}^{\underline{n}}} \text{Int}^{\underline{n}}(v^{\ulcorner \S \urcorner}) \in A^{A \times \mathbb{N}} \xrightarrow{\sqsubseteq} \mathcal{U}_{2\underline{n}} \text{ by} \\ \text{Int}^{\underline{n}}(v^{\ulcorner \S \urcorner}) &=_{\text{def}} \ulcorner \S \urcorner (\text{Int}^{\underline{n}}(v)) : V_{\underline{n}} \xrightarrow{\supseteq} |A, A|_{\underline{n}} \xrightarrow{\ulcorner \S \urcorner} A^{A \times \mathbb{N}} \xrightarrow{\sqsubseteq} \mathcal{U}_{2\underline{n}}. \end{aligned}$$

Here **PR**∈ $\underline{n}2$ map $\ulcorner \S \urcorner : A^A \rightarrow A^{A \times \mathbb{N}}$ is defined as *conjugate* to

$$\begin{aligned} \overline{\ulcorner \S \urcorner} &= \overline{\ulcorner \S \urcorner}_A : A^A \times (A \times \mathbb{N}) \rightarrow A, \\ \text{this in turn } \mathbf{defined} &- \text{PR} - \text{by} \\ \overline{\ulcorner \S \urcorner}_A(v, (a, 0)) &=_{\text{def}} a : A^A \times (A \times \mathbb{1}) \rightarrow A, \\ \overline{\ulcorner \S \urcorner}_A(v, (a, s n)) &=_{\text{def}} \in_{A,A}(v, \overline{\ulcorner \S \urcorner}(v, (a, n))) : A^A \times (A \times \mathbb{N}) \rightarrow A. \end{aligned}$$

With the above, in particular with definition of *Interpretation* map $\text{Int}^{\underline{n}}$ on *map constants* – among them (the codes of) \ltimes and \in , $\text{Int}^{\underline{n}}$ is (PR) defined on all of its arguments, in particular on *conjugated* and hom-functor values, since these are definable in terms of Composition, Cartesian Product and Iteration out of the *basics*. Furthermore, the above *type insertions* show that **PR**∈ \underline{n} map

$$\text{Int}^{\underline{n}} = \text{Int}^{\underline{n}}(u) : \mathbb{N} \supset V_{\underline{n}} = \bigoplus |A, B|_{\underline{n}} \rightarrow \mathcal{U}_{2\underline{n}}, \text{ Ord } A \leq \underline{n},$$

is – as expected – induced by *Object-pair typified family*

$$\text{int}_{A,B}^{\underline{n}} : |A, B|_{\underline{n}} \rightarrow B^A \xrightarrow{\sqsubseteq_A^{\underline{n}}} \mathcal{U}_{2\underline{n}},$$

(\underline{n} still *fixed*), more precisely: it is the *induced* out of countable sum:

$$\begin{aligned} \text{Int}^{\underline{n}} &= (\sqsubseteq_{B^A} \circ \text{int}_{A,B}^{\underline{n}} : |A, B|_{\underline{n}} \rightarrow B^A \sqsubseteq \mathcal{U}_{2\underline{n}})_{A,B} : \\ &\bigoplus |A, B|_{\underline{n}} - - \succ \mathcal{U}_{2\underline{n}}. \end{aligned}$$

By **Definition** of *constructive coding* – namely by definition of code *composition* $v \odot u = v^{\ulcorner \circ \urcorner} u$, of code *product* $u^{\ulcorner \times \urcorner} v$, and of code *iteration* $u^{\ulcorner \S \urcorner}$, all simply given by concatenation of ASCII strings – we have the following

Structure Preservation by *Constructive Coding*:

Composition: for $f : A \rightarrow B$ and $g : B \rightarrow C$ in $\mathbf{PR} \in \underline{n}$:

$$\lceil g \circ f \rceil = \lceil g \rceil \odot \lceil f \rceil =_{\text{by def}} \lceil g \rceil \lceil \circ \rceil \lceil f \rceil :$$

$$\mathbb{1} \rightarrow |A, C|_{\underline{n}} =_{\text{by def}} [A, C]_{\mathbf{PR} \in \underline{n}};$$

Cartesian product: for $f : A \rightarrow C$ and $g : B \rightarrow D$ in $\mathbf{PR} \in \underline{n}$:

$$\lceil (f \times g) \rceil = \langle \lceil f \rceil \lceil \times \rceil \lceil g \rceil \rangle : \mathbb{1} \rightarrow |A \times B, C \times D|_{\underline{n}},$$

as well as *Iteration*: for $f : A \rightarrow A$ in $\mathbf{PR} \in \underline{n}$:

$$\lceil f^{\S} \rceil = \lceil f \rceil \lceil \S \rceil : \mathbb{1} \rightarrow |A \times \mathbb{N}, A|_{\underline{n}}.$$

For *closed* internalisation we have an analogous result, namely

Structure Preservation by *Closed Internalisation*: *Naming*

$$\langle f : A \rightarrow B \rangle \mapsto \langle \lfloor f \rfloor : \mathbb{1} \rightarrow B^A \rangle$$

preserves Composition, map-Product and iteration *into* the corresponding closed families $\lceil \circ \rceil_{A,B,C}$, $\lceil \times \rceil_{A,B,C,D}$, as well as $\lceil \S \rceil_A$, in detail:

– *Composition*: For $A \xrightarrow{f} B \xrightarrow{g} C$ in $\mathbf{PR} \in \underline{n}$ we have:

$$\begin{aligned} \lfloor g \circ f \rfloor &= \lfloor g \rfloor \lfloor \circ \rfloor \lfloor f \rfloor =_{\text{by def}} \lfloor \circ \rfloor (\lfloor g \rfloor, \lfloor f \rfloor) : \\ &\mathbb{1} \xrightarrow{(\lfloor g \rfloor, \lfloor f \rfloor)} C^B \times B^A \xrightarrow{\lfloor \circ \rfloor} C^A, \end{aligned}$$

it lives within stratum $\mathbf{PR} \in 2\underline{n}$.

– *Cartesian product*: For $f : A \rightarrow C$, and $g : B \rightarrow D$ in $\mathbf{PR} \in \underline{n}$:

$$\begin{aligned} \lfloor (f \times g) \rfloor &= \langle \lfloor f \rfloor \lfloor \times \rfloor \lfloor g \rfloor \rangle =_{\text{def}} \lfloor \times \rfloor (\lfloor f \rfloor, \lfloor g \rfloor) : \\ &\mathbb{1} \rightarrow C^A \times D^B \xrightarrow{\lfloor \times \rfloor} (C \times D)^{A \times B}, \end{aligned}$$

this again lives in stratum $\mathbf{PR} \in 2\underline{n}$.

– *Iteration*: For $f : A \rightarrow A$ in $\mathbf{PR} \in \underline{n}$,

$$\lfloor f^{\S} \rfloor = \lfloor \S \rfloor (\lfloor f \rfloor) = \lfloor \S \rfloor \circ \lfloor f \rfloor : \mathbb{1} \rightarrow A^A \rightarrow A^{A \times \mathbb{N}},$$

it is likewise a $\mathbf{PR} \in 2\underline{n}$ map.

Proof:

– (Central), *Composition case*: We consider first *coconjugated* composition, namely

$$\begin{aligned} \overline{\lfloor \circ \rfloor} \circ ((\lfloor g \rfloor, \lfloor f \rfloor) \times A) &: \mathbb{1} \times A \rightarrow C^B \times B^A \times A \xrightarrow{\overline{\lfloor \circ \rfloor}} C \\ &=_{\text{by def}} \in \circ (C^B \times \in) \circ ((\lfloor g \rfloor, \lfloor f \rfloor) \times A) : \\ &\mathbb{1} \times A \rightarrow C^B \times B^A \times A \xrightarrow{C^B \times \in} C^B \times B \xrightarrow{\in} C \\ &= g \circ f \circ r : \mathbb{1} \times A \rightarrow B \rightarrow C. \end{aligned} \tag{*}$$

The latter equation follows from the evaluation properties of *closed evaluation* instances $\in : B^A \times A \rightarrow B$, and $\in : C^B \times B \rightarrow C$, by *Free Variable chasing* – namely free variable $a := r_{\mathbb{1}, A} : \mathbb{1} \times A \rightarrow A$.

By *conjugation* of (both sides of) the above equation we get the assertion in the present composition case:

$$\begin{aligned}
 & \overline{\lfloor \circ \rfloor} \circ ((\lfloor g \rfloor, \lfloor f \rfloor) \times A) : \mathbb{1} \times A \rightarrow C^B \times B^A \times A \rightarrow C \\
 (conj) \quad & \frac{= (g \circ f) \circ r : \mathbb{1} \times A \rightarrow A \rightarrow C}{\lfloor \circ \rfloor \circ (\lfloor g \rfloor, \lfloor f \rfloor) : \mathbb{1} \rightarrow C^B \times B^A \rightarrow C^A} \\
 & = \lfloor g \circ f \rfloor : \mathbb{1} \rightarrow C^A.
 \end{aligned}$$

– Case of *Cartesian product*: analogous, “exercise”.

– *Iteration case*: We start again with the *conjugate* side: For a $\mathbf{PR} \in \underline{n}$ endo $f : A \rightarrow A$, we want to show

$$\begin{aligned}
 & \overline{\lfloor f^\S \rfloor} : \mathbb{1} \times (A \times \mathbb{N}) \rightarrow A \\
 & =_{\text{by def}} f^\S \circ \cong : \mathbb{1} \times (A \times \mathbb{N}) \xrightarrow{\cong} A \times \mathbb{N} \xrightarrow{f^\S} A \\
 & = \overline{\lfloor \S \rfloor} \circ (\lfloor f \rfloor \times (A \times \mathbb{N})) : \quad \quad \quad (***) \\
 & \mathbb{1} \times (A \times \mathbb{N}) \rightarrow A^A \times (A \times \mathbb{N}) \xrightarrow{\overline{\lfloor \S \rfloor}} A.
 \end{aligned}$$

For **Proof** of $(***)$ we use the **definition** above, of $\overline{\lfloor \S \rfloor}$, *coconjugate* of $\lfloor \S \rfloor : A^A \rightarrow A^{A \times \mathbb{N}}$, making commute the lower two rectangles of the following diagram:

$$\begin{array}{ccccc}
 & & \xrightarrow{\cong} & A \times \mathbb{N} & \xrightarrow{f^\S} \\
 & & & \text{(***)} & \\
 \mathbb{1} \times (A \times \mathbb{N}) & \xrightarrow{\lfloor f \rfloor \times (A \times \mathbb{N})} & A^A \times (A \times \mathbb{N}) & \xrightarrow{\overline{\lfloor \S \rfloor}} & A \\
 \downarrow \cong & & \downarrow \cong & & \uparrow r \\
 (\mathbb{1} \times A) \times \mathbb{N} & \xrightarrow{(\lfloor f \rfloor \times A) \times \mathbb{N}} & (A^A \times A) \times \mathbb{N} & \xrightarrow{(\ell, \epsilon)^\S} & A^A \times A
 \end{array}$$

=by def

For showing $(***)$, we show commutativity of the frame diagram, by free variables diagram chasing, with free variables $a := \ell_{A, \mathbb{N}} n := r_{A, \mathbb{N}}$:

$$\begin{array}{ccccc}
 & & \xrightarrow{\cong} & (a, n) & \xrightarrow{f^\S} \\
 & & & & \\
 (0, (a, n)) & & & & f^\S(a, n) \\
 \downarrow \cong & & & & \uparrow r \\
 ((0, a), n) & \xrightarrow{(\lfloor f \rfloor \times A) \times \mathbb{N}} & ((\lfloor f \rfloor, a), n) & \xrightarrow{(\ell, \epsilon)^\S} & (\lfloor f \rfloor, f^\S(a, n)) \\
 & & & (\mapsto) &
 \end{array}$$

Remains to show (\mapsto) , i. e. to show:

$$\begin{aligned}
 & (\ell, \epsilon)^\S((\lfloor f \rfloor, a), n) = (\lfloor f \rfloor, f^\S(a, n)) : \quad \quad \quad (\bullet) \\
 & (\mathbb{1} \times A) \times \mathbb{N} \rightarrow A^A \times A.
 \end{aligned}$$

We show this by external Peano Induction, i.e. by *uniqueness of the iterated*, as follows:

$$(\ell, \in)^\S((\lfloor f \rfloor, a), 0) = (\lfloor f \rfloor, a) = (\lfloor f \rfloor, f^\S(a, 0)) \quad (\text{anchor})$$

as well as

$$\begin{aligned} (\ell, \in)^\S((\lfloor f \rfloor, a), n+1) &= (\ell, \in)^\S((\lfloor f \rfloor, \in_{A,A}(\lfloor f \rfloor, a)), n) \\ &= (\ell, \in)^\S((\lfloor f \rfloor, f(a)), n) \end{aligned}$$

by *evaluation property* of $\in_{A,A} : A^A \times A \rightarrow A$

$$= (\lfloor f \rfloor, f^\S(f(a), n)) \text{ by induction hypothesis on } n$$

$$= (\lfloor f \rfloor, f^\S(a, n+1)) : A \times \mathbb{N} \rightarrow A. \quad (\text{step})$$

This shows (\bullet) , i.e. (\mapsto) in the DIAGRAM: Map $(\ell, e)^\S$ – diagram – throws in fact $((\lfloor f \rfloor, a), n)$ into $(\lfloor f \rfloor, f^\S(a, n))$. So assertion $(***)$ above has been **shown**. Whence, by *conjugation*:

$$\begin{aligned} \lfloor f^\S \rfloor &= \text{coconj}[f^\S \circ \cong : \mathbb{1} \times (A \times \mathbb{N}) \rightarrow A \times \mathbb{N} \xrightarrow{f^\S} A] \\ &= \lfloor \S \rfloor \circ \lfloor f \rfloor : \mathbb{1} \xrightarrow{\lfloor f \rfloor} A^A \xrightarrow{\lfloor \S \rfloor} A^{A \times \mathbb{N}}, \end{aligned}$$

and that **proves** the remaining case of *Structure Preservation* via Closed Internalisation **q.e.d.**

We now come to our central result, the

Interpretation Theorem:

- (i) *CoDomain Suitability* of interpretation family: PR defined $\mathbf{PR} \in 2\underline{n}$ *interpretation family* $\text{int}_{A,B}^{\underline{n}} : |A, B|_{\underline{n}} \rightarrow \mathcal{U}_{2\underline{n}}$ – indexed by Object-pairs, *stratum* (strata) $\mathbf{PR} \in \underline{n}$ (and $\mathbf{PR} \in 2\underline{n}$) – restricts in its (single) *CoDomains* to

$$\text{int}_{A,B}^{\underline{n}} : |A, B|_{\underline{n}} \rightarrow B^A \quad [\sqsubset_{\rightarrow} \mathcal{U}_{2\underline{n}}] \quad (*)$$

within $\mathbf{PR} \in 2\underline{n}$, in form of a commuting DIAGRAM, for B having a point:

$$\begin{array}{ccccc} & & |A, B|_{\underline{n}} & \xrightarrow{\text{int}_{A,B}^{\underline{n}}} & B^A \\ & \swarrow \subset & \updownarrow \subset & & \updownarrow \sqsubset_{B^A} \\ \bigoplus_{A,B} |A, B|_{\underline{n}} & \xrightarrow{\quad \quad} & V_{\underline{n}} & \xrightarrow{\text{Int}^{\underline{n}}} & \mathcal{U}_{2\underline{n}} \end{array}$$

Interpretation DIAGRAM: stratum by stratum,
global/individual with respect to map-code sets

- (ii) *Objectivity* within one *stratum*: For $f : A \rightarrow B$ in $\mathbf{PR} \in \underline{n} \subseteq \mathbf{PR} \in$, we have

$$\mathbf{PR} \in \underline{n} \vdash \text{int}_{A,B}^{\underline{n}}(\ulcorner f \urcorner) =_{\text{by def}} \text{int}_{A,B}^{\underline{n}} \circ \ulcorner f \urcorner = [f] : \mathbb{1} \rightarrow B^A \quad (**)$$


Codes “originating from” Objective level are interpreted into names.

- (iii) *Stratum-Globalisation* of Interpretation: Stratum-indexed family $[\text{int}_{A,B}^{\underline{n}} : |A, B|_{\underline{n}} \rightarrow B^A]_{\underline{n} \in \mathbb{N}}$ admits, within Theory

$\mathbf{PR} \in = \bigcup_{\underline{n}} \uparrow \mathbf{PR} \in \underline{n}$, Object $[A, B]$ of $\mathbf{PR} \in$ again as an *ascending Union*, written $[A, B] =_{\text{by def}} [A, B]_{\mathbf{PR} \in} = \bigcup_{\underline{n}} \uparrow |A, B|_{\underline{n}}$, predica-

tively, and has the universal property of an *inductive limit* by \mathbf{PR} “construction”.

In particular, family $\text{int}_{A,B}^{\underline{n}} : |A, B|_{\underline{n}} \rightarrow B^A$ above induces a – *unique* – *strata-global* map $\text{int}_{A,B} : [A, B] \rightarrow B^A$ (***) making commute the following DIAGRAM:



Strata-global interpretation DIAGRAM

- (iv) Strata-global *Objectivity* of Interpretation, “Codes to names”:

For an arbitrary $\mathbf{PR} \in \underline{n}$ map $f : A \rightarrow B$ we have:

$$\begin{aligned} \mathbf{PR} \in \vdash \text{int}_{A,B}(\ulcorner f \urcorner) &=_{\text{by def}} \text{int}_{A,B} \circ \ulcorner f \urcorner \\ &= [f] : \mathbb{1} \rightarrow B^A \xrightarrow{\sqsubseteq} \mathcal{U}_{2\underline{n}}. \end{aligned} \quad (\bullet)$$

Proof:

- (i) Type control $V_{\underline{n}} \supset |A, B|_{\underline{n}} \ni u \mapsto \text{int}_{A,B}^{\underline{n}}(u) \in B^A \sqsubset \mathcal{U}_{2\underline{n}}$:

This is **proved** by structural induction on u , i.e. on $\text{depth}(u) : [A, B] \supset |A, B|_{\underline{n}}, \underline{n}$ “suitable” such that all the finitely many *building blocks* v, w, \dots are in finitely many components of *sum*

$V_{\underline{n}} = \bigoplus_{A,B} |A, B|_{\underline{n}}$. This type assertion has been (pre-) discussed already above.

- (ii) **Proof** of second assertion (**) on *Objectivity* of each member of the $\underline{n} \in \underline{\mathbb{N}}$ and Object-pair A, B indexed family is now as expected, namely by external structural induction on (external) depth $[f]$ of map $f : A \rightarrow B$ in $\mathbf{PR} \in$ in question, with $\ulcorner f \urcorner \in |A, B|_{\underline{n}}$, suitable $\underline{n} \in \underline{\mathbb{N}}$: Each such f comes with such a “suitable” \underline{n} , since obviously the $\mathbf{PR} \in \underline{n}, \underline{n} \in \underline{\mathbb{N}}$, exhaust all of Theory $\mathbf{PR} \in$ here considered.

Now here is the **Proof** of *Interpretation-Objectivity*, by structural induction on depth $[f : A \rightarrow B]$ “to be interpreted”:

For f one of the *map constants* of $\mathbf{PR} \in = \mathbf{PR}_A + (\text{hom})$ with depth $[f] = 1$ say, in particular for the members of adjunction map families $\in_{A,B}$ and $\angle_{A,B}$, the assertion is trivial, by definition of interpretation Int, and corresponding $\text{int}_{A,B}^{\underline{n}}$ in these cases. We now consider $\mathbf{PR} \in$ maps with greater depth:

For $f : A \rightarrow B, g : B \rightarrow C$ in $\mathbf{PR} \in$

$$\begin{aligned} \mathbf{PR} \in \underline{n} \vdash \text{int}_{A,C}^{\underline{n}} (\ulcorner g \circ f \urcorner) &= \text{int}_{A,C}^{\underline{n}} \langle \ulcorner g \urcorner \odot \ulcorner f \urcorner \rangle \\ &[\odot = \ulcorner \circ \urcorner \text{ constructively internalises } \circ] \\ &=_{\text{by def}} \text{int}_{B,C}^{\underline{n}} (\ulcorner g \urcorner) \lrcorner \odot \lrcorner \text{int}_{A,B}^{\underline{n}} (\ulcorner f \urcorner) \\ &= [g] \lrcorner \odot \lrcorner [f] \text{ by } \underline{\text{hypothesis}} \text{ on } f \text{ and } g \\ &= [g \circ f] : \mathbb{1} \rightarrow C^A, \end{aligned}$$

the latter by the *composition case* of *Structure preservation by Axiomatic Internalisation* above.

Similar (recursive) **Proof** for the assertion in case of the other binary meta-operation, the Cartesian product of maps.

– Remains the case of an *iterated* $f^{\S} : A \times \mathbb{N} \rightarrow A$, given by the unary meta-operation \S : In this case we have

$$\begin{aligned} \mathbf{PR} \in \underline{n} \vdash \text{int}_{A \times \mathbb{N}, A}^{\underline{n}} (\ulcorner f^{\S} \urcorner) &= \text{int}_{A \times \mathbb{N}, A}^{\underline{n}} (\ulcorner f \urcorner^{\ulcorner \S \urcorner}) \\ &\text{by definition of constructive code of an iterated} \\ &=_{\text{by def}} \lrcorner \S \lrcorner (\text{int}_{A,A}^{\underline{n}} (\ulcorner f \urcorner)) \\ &(\text{“homomorphic” PR definition of interpretation Int}) \\ &= \lrcorner \S \lrcorner \circ [f] \text{ by hypothesis on } \underline{\text{depth}}[f] \\ &= [f^{\S}] : \mathbb{1} \rightarrow A^{A \times \mathbb{N}}, \end{aligned}$$

the latter, eventually, by the *iteration case* of *Structure Preservation of Closed Internalisation*.

The last two **assertions** of the **Theorem** – (**) and (•) – follow straightforward from the former two, by the inductive-limit property of our Universal Chain \mathcal{U} .

Comment: The members of family $\text{Int}^{\underline{n}} : V_{\underline{n}} \rightarrow \mathcal{U}_{2\underline{n}}$ are special maps – Objective map terms – of Theory $\mathbf{PR} \in 2\underline{n} \prec \mathbf{PR} \in$, and are therefore covered “themselves” by the – in this regard *self-related Interpretation*

Theorem above. This is the reason why I have choosen as a *Universal Class* not a single Object or “super-Object” for Theory $\mathbf{PR} \in$, but an *ascending chain* of “Universal Objects” $\mathcal{U}_{\underline{n}}$, such that Object $\mathcal{U}_{2\underline{n}}$, of $\mathbf{PR} \in 2\underline{n}$, hosts in particular *interpretation* of all map codes of stratum $\mathbf{PR} \in \underline{n}$: Chain \mathcal{U} is “upwards open”, think at HILBERT’s hotel.

6 Self-Evaluation

Here is the key **Consequence** of the two last assertions $(***)$ and (\bullet) of the **Interpretation Theorem**, namely possibility for a *constructive self-evaluation* of Theory $\mathbf{PR} \in$:

Define code-*self-evaluation* family for Theory $\mathbf{PR} \in$, called $\tilde{\varepsilon} = \tilde{\varepsilon}_{A,B}$: $[A, B]_{\mathbf{PR} \in} \times A \rightarrow B$, within Theory $\mathbf{PR} \in$ itself as

$$\begin{aligned} \tilde{\varepsilon}_{A,B} &= \tilde{\varepsilon}_{A,B}(u, a) \stackrel{\text{def}}{=} \varepsilon_{A,B}(\text{int}_{A,B}(u), a) : \\ V \times A \supset [A, B]_{\mathbf{PR} \in} \times A &\xrightarrow{\text{int}_{A,B} \times A} B^A \times A \xrightarrow{\varepsilon} B. \end{aligned}$$

Comment: Here we used assertion $(***)$ for availability of suitable Order-global interpretation family

$$\text{int}_{A,B} : [A, B]_{\mathbf{PR} \in} = \bigcup_{\underline{n}} \uparrow [A, B]_{\underline{n}} \rightarrow B^A.$$

We get further, by last assertion (\bullet) – of the Theorem, *objectivity* of self-evaluation $\tilde{\varepsilon}$, namely: for (any) $f : A \rightarrow B$ in $\mathbf{PR} \in$

$$\begin{aligned} \mathbf{PR} \in \vdash \tilde{\varepsilon}_{A,B}(\ulcorner f \urcorner, a) &= \varepsilon_{A,B}(\text{int}_{A,B}(\ulcorner f \urcorner), a) \\ &= \varepsilon_{A,B}(\lfloor f \rfloor, a) = f(a) : A \rightarrow B. \end{aligned} \quad (*)$$

For this latter equation see introduction – and discussion – of *name* of f above, $\lfloor f \rfloor = \lfloor f : A \rightarrow B \rfloor : \mathbb{1} \rightarrow B^A$ – in **set** theory: $\lfloor f \rfloor = \{(\emptyset, f)\} : \mathbb{1} \rightarrow B^A \subset \mathcal{P}(A \times B)$.

Based on this *self-evaluation* family of Theory $\mathbf{PR} \in$, we now find within $\mathbf{PR} \in$ the following (anti) diagonal $d = d(n) : \mathbb{N} \rightarrow 2 : \mathbf{PR} \in$ -map $d = d(n) : \mathbb{N} \rightarrow 2$ is **defined** as

$$d \stackrel{\text{def}}{=} \neg \circ \tilde{\varepsilon}_{\mathbb{N},2} \circ (\#, \text{id}_{\mathbb{N}}) : \mathbb{N} \longrightarrow [\mathbb{N}, 2]_{\mathbf{PR} \in} \times \mathbb{N} \xrightarrow{\tilde{\varepsilon}} 2 \xrightarrow{\neg} 2,$$

with $\# = \#(n) : \mathbb{N} \xrightarrow{\cong} [\mathbb{N}, 2]_{\mathbf{PR} \in}$ the – isomorphic – \mathbf{PR} *count* of all (internal) predicate codes (“Klassenzeichen” in Gödel’s sense), of Theory $\mathbf{PR} \in$. As expected in such *diagonal argument* – Antinomie Richard quoted by GÖDEL – we *substitute*, within Theory $\mathbf{PR} \in$, the counting index $q \stackrel{\text{def}}{=} \#^{-1}(\ulcorner d \urcorner) = \#^{-1} \circ \ulcorner d \urcorner : \mathbb{1} \rightarrow [\mathbb{N}, 2]_{\mathbf{PR} \in} \xrightarrow{\cong} \mathbb{N}$, of d ’s *code* into $\mathbf{PR} \in$ -map $d : \mathbb{N} \rightarrow 2$ itself, and get a “liar” map $liar : \mathbb{1} \rightarrow 2$, called

liar because it turns out that this map is its own negation, as follows:

$$\begin{aligned}
\mathbf{PR} \in \vdash \text{liar} &=_{\text{def}} d \circ q : \mathbb{1} \rightarrow \mathbb{N} \rightarrow 2 \\
&=_{\text{by def}} d \circ \#^{-1} \circ \ulcorner d \urcorner \\
&=_{\text{by def}} \neg \circ \tilde{\varepsilon}_{\mathbb{N},2} \circ (\#, \text{id}_{\mathbb{N}}) \circ \#^{-1} \circ \ulcorner d \urcorner \\
&= \neg \circ \tilde{\varepsilon}_{\mathbb{N},2} \circ (\ulcorner d \urcorner, \#^{-1} \circ \ulcorner d \urcorner) \\
&=_{\text{by def}} \neg \circ \tilde{\varepsilon}_{\mathbb{N},2} (\ulcorner d \urcorner, q) \\
&= \neg \circ d(q) = \neg \circ d \circ q & (**) \\
&=_{\text{by def}} \neg \text{liar} : \mathbb{1} \rightarrow 2 \rightarrow 2,
\end{aligned}$$

a contradiction: The argument is equation marked (**), which is a special instance of *objectivity* equation (*) above, objectivity of *self-evaluation* $\tilde{\varepsilon}$, which has been **defined** within theory $\mathbf{PR} \in$ out of *closed evaluation* \in composed with interpretation family *int*, of map *codes* into *names*.

Conclusion: The argument shows incompatibility of (even just *potential*) *infinity* with (formally, axiomatically given) *Cartesian Closed* “Higher Order” structure of Theory $\mathbf{PR} \in$.

We obtain this way inconsistency of all extensions of Theory $\mathbf{PR} \in$, in particular of – higher order – **set** theories, and also of any type of higher Order Arithmetic, even when given in a categorical setting, as in particular in LAWVERE 1963, and then in FREYD’s 1972 setting of (higher Order) Topos Theory with NNO, and in that of LAMBEK & SCOTT 1986.

The present argument does not depend on *quantification* nor on availability of a subobject classifier: the (equality) *predicates* we rely on here are given by the Cartesian PR Arithmetic of theories considered.

Disclaimer: The argument does *not* apply to *Closed Categories* in the sense of EILENBERG & KELLY, since there is no NNO required for the theory. In the applications, e.g. Categories of Modules, there is an NNO only *downstairs*, in a suitably conceived category of sets. But that NNO does not bear (naturally) the structure of an abelian group.

Even if you consider the category of abelian *semi*-groups which includes semigroup $\mathbb{N} = \langle \mathbb{N}, 0, + \rangle$: an *iterated* $f^{\S} : A \times \mathbb{N} \rightarrow A$ will not become linear, even not bilinear, and hence even not linear when converted into a map $f^{\S} : A \otimes \mathbb{N} \rightarrow A$ from the tensor product into A . So this category cannot have \mathbb{N} as an NNO in any suitable way.

Analogously, the original Elementary Theory **ETT** of Topoi seems me to be not concerned, **ETT** in the sense explained by WRAITH 1973 on the base of mainly (?) LAWVERE 1970, 1972, and TIERNEY 1971, as well as more recently explained in LAWVERE & SHANUEL 1991:

The data and axioms for this *genuine* Theory of Topoi do not include an NNO. The motivating examples for Topoi are Categories of *sheaves* over a topological space. Question: Do these – Cartesian Closed – Categories come with an NNO on sheaf level? By the above, they cannot come so, except they are based on an – inconsistent – Cartesian Closed set Theory with NNO.

Problem: Diagonal map above is a map within subSystem $\mathbf{PR} \in \underline{n}$, subSystem of Theory $\mathbf{PR} \in$, for \underline{n} from some \underline{n}_0 upwards. Presumably an upper bound for such contradictory Order \underline{n}_0 can be calculated. It would be certainly interesting to know a lower bound \underline{n}_0 making $\mathcal{P}^{\underline{n}}$ contradictory, incompatible with (potential) *infinity*, in the sense of availability of a Natural Numbers Object \mathbf{N} .

References

- S. EILENBERG, C. C. ELGOT 1970: *Recursiveness*. Academic Press.
- S. EILENBERG, G. M. KELLY 1966: Closed Categories. *Proc. Conf. on Categorical Algebra*, La Jolla 1965, pp. 421-562. Springer.
- S. EILENBERG, S. MAC LANE 1945: General Theory of Natural Equivalences. *Trans. AMS* 58, 231-294.
- P. J. FREYD 1972: Aspects of Topoi. *Bull. Australian Math. Soc.* **7**, 1-76.
- K. GÖDEL 1931: Über formal unentscheidbare Sätze der Principia Mathematica und verwandter Systeme I. *Monatsh. der Mathematik und Physik* 38, 173-198.
- R. L. GOODSTEIN 1971: *Development of Mathematical Logic*, ch. 7: Free-Variable Arithmetics. Logos Press.
- D. HILBERT: Mathematische Probleme. Vortrag Paris 1900. *Gesammelte Abhandlungen*. Springer 1970.
- P. T. JOHNSTONE 1977: *Topos Theory*. Academic Press
- A. JOYAL 1973: Arithmetical Universes. Talk at Oberwolfach.
- J. LAMBEK, P. J. SCOTT 1986: *Introduction to higher order categorical logic*. Cambridge University Press.
- F. W. LAWVERE 1964: An Elementary Theory of the Category of Sets. *Proc. Nat. Acad. Sc. USA* **51**, 1506-1510.
- F. W. LAWVERE 1970: Quantifiers and Sheaves. *Actes du Congrès International des Mathématiciens*. Nice, pp. 329-334.
- F. W. LAWVERE, S. H. SHANUEL 1997 (1991): *Conceptual Mathematics, A first introduction to categories*. Cambridge University Press.
- S. MAC LANE 1972: *Categories for the working mathematician*. Springer.
- B. PAREIGIS 1969: *Kategorien und Funktoren*. Teubner.
- R. PÉTER 1967: *Recursive Functions*. Academic Press.
- M. PFENDER 2008 RCF1: Theories of PR Maps and Partial PR Maps. pdf file. TU Berlin.
- M. PFENDER 2008 RCFX: Universal Objects and Theory Embedding. pdf file. TU Berlin.
- M. PFENDER, M. KRÖPLIN, D. PAPE 1994: Primitive Recursion, Equality, and a Universal Set. *Math. Struct. in Comp. Sc.* **4**, 295-313.
- W. RAUTENBERG 1995/2006: *A Concise Introduction to Mathematical Logic*. Universitext Springer 2006.
- R. REITER 1980: Mengentheoretische Konstruktionen in arithmetischen Universen. Diploma Thesis. TU Berlin.

L. ROMÀN 1989: Cartesian categories with natural numbers object. *J. Pure and Appl. Alg.* **58**, 267-278.

W. W. TAIT 1996: Frege versus Cantor and Dedekind: on the concept of number. Frege, Russell, Wittgenstein: *Essays in Early Analytic Philosophy (in honor of Leonhard Linsky)* (ed. W. W. Tait). Lasalle: Open Court Press (1996): 213-248. Reprinted in *Frege: Importance and Legacy* (ed. M. Schirn). Berlin: Walter de Gruyter (1996): 70-113.

A. TARSKI, S. GIVANT 1987: *A formalization of set theory without variables*. AMS Coll. Publ. vol. 41.

M. TIERNEY 1973: Axiomatic Sheaf Theory. *C. I. M. E. Conf. on Categories and Commutative Algebra, Varenna*, pp. 249-326. Edizione Cremonese. Roma. (Quoted in WRAITH 1973.)

G. WRAITH 1973: Lectures on Elementary Topoi. In *Model Theory and Topoi*, LN in Math. **445**, 114-206.

Address of the author:

M. PFENDER

D-10623 Berlin

Institut für Mathematik

Technische Universität Berlin

pfender@math.TU-Berlin.DE