# CLASSIFICATION OF LOCAL SINGULARITIES ON TORUS CURVES OF TYPE (2,5)

M. KAWASHIMA

ABSTRACT. In this paper, we consider curves of degree 10 of torus type (2,5), C:  $f_5(x,y)^2 + f_2(x,y)^5 = 0$ . Assume that  $f_2(0,0) = f_5(0,0) = 0$ . Then O = (0,0) is a singular point of C which is called an inner singularity. In this paper, we give a topological classification of singularities of (C, O).

# 1. INTRODUCTION

A plane curve  $C \subset \mathbb{P}^2$  is called a *curve of torus type* (p,q) if there is a defining polynomial F of C which can be written as  $F = F_p^q + F_q^p$ , where  $F_p$ ,  $F_q$  are homogeneous polynomials of X, Y, Z of degree p and q respectively. In [7], D.T. Pho classified the local and global configurations of the singularities of sextics of torus type (2,3). In this paper, we will classify local inner singularities of torus type (2,5). We assume C is a reduced curve of torus type (2,5). Using affine coordinates x = X/Z, y = Y/Z, C is defined as

$$C: f_2(x,y)^5 + f_5(x,y)^2 = 0,$$

where  $f_2(x,y) = F_2(x,y,1)$ ,  $f_5(x,y) = F_5(x,y,1)$ . Put  $C_2 := \{f_2(x,y) = 0\}$  and  $C_5 := \{f_5(x,y) = 0\}$ . We assume that the origin O = (0,0) is an intersection point of  $C_2$  and  $C_5$  and O is an isolated singularity of C. We classify the local singularity type of (C,O) following the method of [7]. If  $f_2(x,y) = \ell(x,y)^2$  for some linear form  $\ell$ , the curve C is called a *linear torus curve* and it consists of two quintics, as  $f_2^5 + f_5^2 = (f_5 + \ell^5 \sqrt{-1})(f_5 - \ell^5 \sqrt{-1})$ .

First we recall the following notations.

$$A_n : x^{n+1} + y^2 = 0 \quad (n \ge 1)$$
$$D_n : x^{n-1} + xy^2 = 0 \quad (n \ge 4)$$
$$E_6 : x^3 + y^4 = 0, \quad E_7 : x^3 + xy^3 = 0, \quad E_8 : x^3 + y^5 = 0$$
$$B_{n,m} : x^n + y^m = 0 \quad (\text{Brieskorn-Pham type})$$

In the case gcd(m,n) > 1, the equation of  $B_{n,m}$  can contain other monomials on the Newton boundary. For example,  $B_{2,4}$  has the form  $C_t : x^2 + txy^2 + y^4 = 0$ , for  $t \neq \pm 2$ .  $C_t$  is topologically equivalent to  $B_{2,4}$  (Oka, [3]). In these notations, we note that  $A_n = B_{n+1,2}$  and  $E_6 = B_{3,4}$ .

We remark that every non-degenerate singularity in the sense of Newton boundary is a union of Brieskorn-Pham type singularities. For example,  $C_{p,q}: x^p + x^2y^2 + y^q$ ,  $p,q \ge 4$ ,  $p+q \ge 9$  is the union of two singularities:  $x^{p-2} + y^2 = 0$  and  $x^2 + y^{q-2} = 0$ . So we introduce the notation:  $B_{p-2,2} \circ B_{2,q-2}$  to express  $C_{p,q}$ .

<sup>1991</sup> Mathematics Subject Classification. 14H20,14H45.

Key words and phrases. Torus curve, Newton boundary, non-degenerate.

For the classification, we use the local intersection multiplicity  $\iota := I(C_2, C_5; O)$  effectively. The complete classifications is given by Theorem 1 in §5.

This paper consists of the following sections:

- §2. Preliminaries.
- §3. Some lemmas for tours curves of type (p,q).
- §4. Calculation of the local singularities.
- §5. List of classification.
- §6. Linear torus curve of type (2, 5).
- §7. Appendix.

### 2. Preliminaries

2.1. Toric modification. Throughout this paper, we follow the notations of Oka [5]. First we recall a toric modification. Let

$$\sigma = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$$

be a unimodular integral  $2 \times 2$  matrix. We associate to  $\sigma$  a birational morphism  $\pi_{\sigma}$ :  $\mathbb{C}^{*2} \to \mathbb{C}^{*2}$  by  $\pi_{\sigma}(x, y) = (x^{\alpha}y^{\beta}, x^{\gamma}y^{\delta})$ . If  $\alpha, \gamma \geq 0$  (respectively.  $\beta, \delta \geq 0$ ), this map can be extend to x = 0 (resp. y = 0). Note that  $\{\pi_{\sigma}\}$ 's satisfy the properties  $\pi_{\sigma} \circ \pi_{\tau} = \pi_{\sigma\tau}$ and  $(\pi_{\sigma})^{-1} = \pi_{\sigma^{-1}}$ .

Let N be a free Z-module of rank two with a fixed basis  $\{E_1, E_2\}$ . Through this basis, we identify  $N_{\mathbb{R}} := N \otimes \mathbb{R}$  with  $\mathbb{R}^2$ . Thus N can be understood as the set of integral points in  $\mathbb{R}^2$ . We denote a vector in N by a column vector. Hereafter we fix two special vectors  $E_1 = {}^t(1,0)$  and  $E_2 = {}^t(0,1)$ . Let  $N^+$  be the space of positive vectors of N. Let  $\{P_1, \dots, P_m\}$  be given positive primitive integral vectors in  $N^+$ . Let  $P_i = {}^t(a_i, b_i)$ and assume that det $(P_i, P_{i+1}) > 0$  for each  $i = 0, \dots, m$ . Here  $P_0 = E_1, P_{m+1} = E_2$ . We associated to  $\{P_0, P_1, \dots, P_{m+1}\}$  a simplicial cone subdivision  $\Sigma^*$  of  $N_{\mathbb{R}}$  which has m + 1cones Cone $(P_i, P_{i+1})$  of dimension two where

$$Cone(P_i, P_{i+1}) := \{tP_i + sP_{i+1} \mid t, s \ge 0\}.$$

We call  $\{P_0, \dots, P_{m+1}\}$  the vertices of  $\Sigma^*$ . We say that  $\Sigma^*$  is a regular simplicial cone subdivision of  $N^+$  if det $(P_i, P_{i+1}) = 1$  for each  $i = 0, \dots, m$ .

Assume that  $\Sigma^*$  is a given regular simplicial cone subdivision with vertices  $\{P_0, P_1, \dots, P_{m+1}\}$ ,  $(P_0 = E_1, P_{m+1} = E_2)$  and let  $P_i = {}^t(a_i, b_i)$ . For each cone,  $\operatorname{Cone}(P_i, P_{i+1})$ , we associate the unimodular matrix

$$\sigma_i := \begin{pmatrix} a_i & a_{i+1} \\ b_i & b_{i+1} \end{pmatrix}$$

We identify  $\operatorname{Cone}(P_i, P_{i+1})$  with the unimodular matrix  $\sigma_i$ . Let (x, y) be a fixed system of coordinates of  $\mathbb{C}^2$ . Then we consider, for each  $\sigma_i$ , an affine space  $\mathbb{C}^2_{\sigma_i}$  of dimension two with coordinates  $(x_{\sigma_i}, y_{\sigma_i})$  and the birational map  $\pi_{\sigma_i} : \mathbb{C}^2_{\sigma_i} \to \mathbb{C}^2$ . First we consider the disjoint union of  $\mathbb{C}^2_{\sigma_i}$  for  $i = 0, \ldots, m$  and we define the variety X as the quotient of this union by the following identification: Two points  $(x_{\sigma_i}, y_{\sigma_i}) \in \mathbb{C}^2_{\sigma_i}$  and  $(x_{\sigma_j}, y_{\sigma_j}) \in \mathbb{C}^2_{\sigma_j}$  are identified if and only if the birational map  $\pi_{\sigma_j^{-1}\sigma_i}$  is well defined at the point  $(x_{\sigma_i}, y_{\sigma_i})$ and  $\pi_{\sigma_j^{-1}\sigma_i}(x_{\sigma_i}, y_{\sigma_i}) = (x_{\sigma_j}, y_{\sigma_j})$ . As can be easily checked, X is non-singular and the maps  $\{\pi_{\sigma_i} : \mathbb{C}^2_{\sigma_i} \to \mathbb{C}^2 \mid 0 \le i \le m\}$  glue into a proper analytic map  $\pi : X \to \mathbb{C}^2$ .

 $\mathbf{2}$ 

**Definition 1.** The map  $\pi : X \to \mathbb{C}^2$  is called the toric modification associated with  $\{\Sigma^*, (x, y), O\}$  where is a regular simplicial cone subdivision of  $N^+$  and (x, y) is a coordinate system of  $\mathbb{C}^2$  centered at the origin O.

Recall that this modification has the following properties.

1.  $\{\mathbb{C}^2_{\sigma_i}, (x_{\sigma_i}, y_{\sigma_i})\}, (0 \leq i \leq m)$  give coordinate charts of X and we call them the toric coordinate charts of X.

2. Two affine divisors  $\{y_{\sigma_{i-1}} = 0\} \subset \mathbb{C}^2_{\sigma_{i-1}}$  and  $\{x_{\sigma_i} = 0\} \subset \mathbb{C}^2_{\sigma_i}$  glue together to make a compact divisor isomorphic to  $\mathbb{P}^1$  for  $0 \leq i \leq m$ . We denote this divisor by  $\hat{E}(P_i)$ .

3.  $\pi^{-1}(O) = \bigcup_{i=1}^{m} \hat{E}(P_i)$  and  $\pi : X - \pi^{-1}(O) \to \mathbb{C}^2 - \{O\}$  is isomorphism. The noncompact divisor  $x_{\sigma_0} = 0$  (respectively.  $y_{\sigma_m} = 0$ ) is mapped isomorphically onto the divisor x = 0 (resp. y = 0).

4.  $\hat{E}(P_i) \cap \hat{E}(P_j) \neq \emptyset$  if and only if  $i - j = \pm 1$ . If  $i - j = \pm 1$ , they intersect transversely at a point.

2.2. Toric modification with respect to an analytic function. Let  $\mathcal{O}$  be the ring of germs of analytic functions at the origin. Recall that  $\mathcal{O}$  is isomorphic to the ring of convergent power series in x, y. Let  $f \in \mathcal{O}$  be a germ of complex analytic function and suppose that f(O) = 0. Let  $f(x, y) = \sum c_{i,j} x^i y^j$  be the Taylor expansion of f at the origin. We assume that f(x, y) is reduced as a germ. The Newton polygon  $\Gamma_+(f; x, y)$  of f, with respect to the coordinate system (x, y), is the convex hull of the union  $\bigcup_{i,j} \{(i, j) + \mathbb{R}^2_+\}$ where the union is taken for (i, j) such that  $c_{i,j} \neq 0$  and the Newton boundary  $\Gamma(f; x, y)$ is the union of compact faces of the Newton polygon  $\Gamma_+(f; x, y)$ . For each compact face  $\Delta$  of  $\Gamma(f; x, y)$ , the face function  $f_{\Delta}(x, y)$  is defined by  $f_{\Delta}(x, y) := \sum_{(i,j)\in\Delta} c_{i,j} x^i y^j$ .

In the space M where the Newton polygon  $\Gamma_+(f; x, y)$  is contained, we use  $(\nu_1, \nu_2)$  as the coordinates. For any positive weight vector  $P = {}^t(a, b) \in N$ , we consider P as a linear function on M by  $P(\nu_1, \nu_2) = a\nu_1 + b\nu_2$ . We define d(P; f) to be the smallest value of the restriction of P to the Newton polygon  $\Gamma_+(f; x, y)$  and  $\Delta(P; f)$  be the face where this smallest value is taken. For simplicity we shall write  $f_P$  instead of  $f_{\Delta(P;f)}$ . By the definition,  $f_P$  is weighted homogeneous polynomial of degree d(P; f) with weight  $P = {}^t(a, b)$ . For each face  $\Delta \in \Gamma(f; x, y)$  there is a unique primitive integral vector  $P = {}^t(a, b)$  such that  $\Delta = \Delta(P; f)$ . The Newton boundary  $\Gamma(f; x, y)$  has a finite number of faces.



Let  $\Delta_1, \ldots, \Delta_m$  these faces and let  $P_i = {}^t(a_i, b_i)$  be the corresponding positive primitive integral vector, i.e.,  $\Delta_i = \Delta(P_i; f)$ . We call  $P_i$  the weight vector of the face  $\Delta_i$ . Then we can write

$$f_{P_i}(x,y) = cx^{r_i}y^{s_i}\prod_{j=1}^{k_i}(y^{a_i} + \gamma_{i,j}x^{b_i})^{\nu_{i,j}}, \quad c \neq 0$$

with distinct non-zero complex numbers  $\gamma_{i,1}, \cdots, \gamma_{i,k_i}$ . We define

$$\tilde{f}_{P_i}(x,y) = f_{P_i}(x,y)/x^{r_i}y^{s_i} = c \prod_{j=1}^{k_i} (y^{a_i} + \gamma_{i,j}x^{b_i})^{\nu_{i,j}}.$$

The polynomial  $\sum_{(i,j)\in\Gamma(f;x,y)} c_{i,j}x^iy^j$  is called the Newton principal part of f(x,y) and we denote by  $\mathcal{N}(f;x,y)$ . We say that f(x,y) is convenient if the intersection  $\Gamma(f;x,y)$ with each axe is non-empty. Note that  $\tilde{f}_{\Delta_i}$  is always convenient. We say that f is nondegenerate on a face  $\Delta_i$  if the function  $f_{\Delta_i} : \mathbb{C}^{*2} \to \mathbb{C}$  has no critical points. This is equivalent to  $\nu_{i,j} = 1$  for all  $j = 1, \dots, k_i$ . We say that f is non-degenerate if f is non-degenerate on any face  $\Delta_i$   $(i = 1, \dots, m)$ .

We introduce an equivalence relation  $\sim$  in  $N^+$  which is defined by  $P \sim Q$  if and only if  $\Delta(P; f) = \Delta(Q; f)$ . The equivalence classes define a conical subdivision of  $N^+$ . This gives a simplicial cone subdivision of  $N^+$  with m+2 vertices  $\{P_0, \cdots, P_{m+1}\}$  with  $P_0 = E_1$ ,  $P_{m+1} = E_2$ . We denote this subdivision by  $\Gamma^*(f; x, y)$  and we call it the dual Newton diagram of f with respect to the system of coordinates (x, y).  $\Gamma^*(f; x, y)$  has m + 1 twodimensional cones  $\text{Cone}(P_i, P_{i+1}), i = 0, \cdots, m$ . Note that these cones are not regular in general.

**Definition 2.** A regular simplicial cone subdivision  $\Sigma^*$  is admissible for f(x, y) if  $\Sigma^*$  is a subdivision of  $\Gamma^*(f; x, y)$ . The corresponding toric modification  $\pi : X \to \mathbb{C}^2$  is called an admissible toric modification for f(x, y) with respect to the system of coordinate (x, y).

There exists a unique canonical regular simplicial cone subdivision (Lemma 3,3 of [4]). We call the corresponding toric modification the canonical toric modification with respect to f(x, y). Let C be a germ of a reduced curve defined by f(x, y) = 0 and let  $\pi : X \to \mathbb{C}^2$ be a good resolution. Recall that the dual graph of the resolution is defined as follows. Let  $E_1, \ldots, E_r$  be the exceptional divisors and put  $\pi^* f^{-1}(0) = \sum_{i=1}^r m_i E_i + \sum_j^s \widetilde{C_j}$ . To each  $E_i$ , we associate a vertex  $v_i$  of  $\mathcal{G}(\pi)$  denoted it by a black circle. We give an edge joining  $v_i$  and  $v_j$  if  $E_i \cap E_j \neq \emptyset$ . For the extended dual graph  $\widetilde{\mathcal{G}}(\pi)$ , we add vertices  $w_i$ to each irreducible components  $C_i$ ,  $i = 1, \cdots, s$  and we join  $w_j$  and  $v_i$  if  $E_i \cap C_j \neq \emptyset$  by a dotted arrow line. It is also important to remember the multiplicities of  $\pi^* f$  along  $E_i$ , which we denote by  $m_i$ . So we put weights  $m_i$  to each vertex and call  $\mathcal{G}(\pi)$  with weight the weighted dual graph of the resolution  $\pi : X \to \mathbb{C}^2$ .

**Example 1.** Consider the curve  $C: y^2 - x^3 = 0$ . The Newton boundary consists of one face  $\Delta$ . Then  $P = {}^t(2,3)$  is the weight vector correspond to the face  $\Delta$  and  $\Gamma^*(f; x, y)$  has three vertices  $\{E_1, P, E_2\}$ . We take a regular simplicial cone subdivision  $\Sigma^*$  of  $\Gamma^*(f; x, y)$  and we consider an admissible toric modification  $\pi: X \to \mathbb{C}^2$  associated to  $\{\Sigma^*, (x, y), O\}$ . Vertices of  $\Sigma^*$  are  $\{E_1, T_1, P, T_2, E_2\}$  where  $T_1 = {}^t(1, 1)$  and  $T_2 = {}^t(1, 2)$  are new vertices which are added to  $\Gamma^*(f; x, y)$  in order to make  $\Sigma^*$  regular.

The proper transform  $\tilde{C}$  intersects transversely with the exceptional divisor  $\hat{E}(P)$  at 1-point. Take a toric coordinate (u, v) so that u = 0 defines  $\hat{E}(P)$  and let  $\xi$  be the intersection point. Then  $\tilde{C}$  is defined by v - 1 = 0 and  $\tilde{C}$  is smooth at  $\xi$ .  $(\pi^* f) = 2\hat{E}(T_1) + 6\hat{E}(P) + 3\hat{E}(T_2) + \tilde{C}$ . LOCAL SINGULARITIES ON TORUS CURVES OF TYPE (2,5)



# 3. Some Lemmas for tours curves of type (P,Q).

3.1. Notations. Through out this paper, we use the same notations in [3], unless otherwise stated. We also use the fact that the topological equivalence class of a non-degenerate germ depends only on its Newton boundary (Theorem 2.1 of [3]). In the process of classification of topological type of an inner singularity of torus curve of type (2.5), we have the following possibilities:

(1) (C, O) is non-degenerate and  $\Gamma(f)$  has one face,

(2) (C, O) is non-degenerate (in some local coordinate system) but the Newton boundary has several faces, or

(3) (C, O) has some degenerate faces for any choice of coordinate systems (x, y).

To make the expression of these singularity classes simpler, we introduce some notations. The class of singularities (1) can be expressed by the class of  $B_{n,m}$ :

$$B_{n,m}$$
 :  $x^n + y^m + (\text{higher terms}) = 0$ 

The class of singularities (2) can be understand a union of singularities of type (1). For example,  $x^n + x^2y^2 + y^m = 0$  is topologically equivalent to  $(x^{n-2} + y^2)(x^2 + y^{m-2}) = 0$  and thus we denote this class by  $B_{n-2,2} \circ B_{2,m-2}$ , as is already introduced in §1.

The last class (3) is most complicated. For example, we consider the singularity germ (C, O) is defined by  $f(x, y) = (\lambda x^3 + y^2)^2 + x^3 y^3$  then  $\mathcal{N}(f; x, y) = (\lambda x^3 + y^2)^2$  and  $\Gamma(f; x, y)$  consists of one face  $\Delta$  with weight vector is  $P = {}^t(2, 3)$  and f is degenerate on  $\Delta$ . We take a regular simplicial subdivision  $\Sigma^*$  as in example 1. Then we take a toric modification  $\pi_1 : X_1 \to \mathbb{C}^2$  we can see that  $\widetilde{C_1}$  intersects transversely with  $\hat{E}(P)$  at  $\xi_1 = (0, -\lambda)$  in toric coordinate (u, v) correspond to  $\operatorname{Cone}(P, T_2)$  and  $\hat{E}(P) = \{u = 0\}$  with multiplicity 12. (Figure 2).



FIGURE 2.

To express the strict transform  $\tilde{C}$  at  $\xi_1$ , we choose the coordinate  $(u, v_1)$  where  $v_1 = v + \lambda$ , we call this coordinates  $(u, v_1)$  the translated toric coordinates for  $(\tilde{C}, \xi_1)$ . Now we found that  $(\tilde{C}, \xi_1)$  is defined by  $B_{3,2} : v_1^2 - \lambda u^3 + (\text{higher terms}) = 0$  where u = 0 defines the exceptional divisor which contains  $\xi_1$ . Observe that  $\hat{E}(P)$  is defined by u = 0 and the tangent cone of  $\tilde{C}$  is  $v_1^2 = 0$ . Thus the tangent cone is transverse to  $\hat{E}(P)$  at  $\xi_1$ . Again

we take a toric blow-up  $\pi_2 : X_2 \to (X_1, \xi_1)$ . This is essentially the same as the one for the cusp singularity  $v_1^2 - \lambda u^3 = 0$ . As  $\pi_1^* f(0)$  is non-degenerate in  $(u, v_1)$ ,  $\pi_2$  gives a good resolution of  $(\tilde{C}, \xi_1)$  and therefore the composition  $\pi_1 \circ \pi_2 : X_2 \to \mathbb{C}^2$  gives a good resolution of (C, O). The resolution graph is simply obtained by adding a bamboo for this blowing up (See (1), Figure 3). We denote this class of singularity by  $(B_{3,2}^2)^{B_{3,2}}$ .

Sometime, we may need to take a coordinate change of the type  $(u, v_2)$  where  $v_2 = v_1 + h(u)$  for some polynomial h(u). The important point here is that we do not change the first coordinate u, as it defines the exceptional divisor  $\hat{E}(P)$ . We call such a coordinate system  $(u, v_2)$  admissible translated toric coordinates at  $\xi_1$ .

**Example 2.** Consider  $f(x,y) = (y^5 - xy^2 + x^2y + x^5)^2 + (x - 2y)^5(x - 3y)^5$ . Then  $\mathcal{N}(f;x,y) = -31y^{10} - 2xy^7 + x^2y^2(x - y) + 2x^7y + 2x^{10}$  and  $\Gamma(f;x,y)$  consists three faces  $\Delta_i$  (i = 1, 2, 3) with weight vectors  $P_1 = {}^t(3, 1), P_2 = {}^t(1, 1)$  and  $P_3 = {}^t(1, 3)$ . Note that f(x,y) is degenerate on  $\Delta(P_2; f)$ . By adding vertices  $T_1 = {}^t(2, 1)$  and  $T_2 = {}^t(1, 2)$ , we get the canonical regular subdivision. We take the associated toric modification and we can see easily that the strict transform  $\widetilde{C}$  splits into three germs  $\widetilde{C}_1, \widetilde{C}_2, \widetilde{C}_3$  so that for  $i = 1, 3, \widetilde{C}_i$ , intersects transversely with exceptional divisor  $\hat{E}(P_i)$  at two points and  $\widetilde{C}_i$  are smooth at these points. The germ  $\widetilde{C}_2$  which intersects with  $\hat{E}(P_2)$  is still singular and intersects with  $\hat{E}(P_2)$  transversely at  $\xi_1 = (0, 1)$  in the toric coordinate (u, v) corresponding to  $\operatorname{Cone}(P_2, T_2)$ . Thus taking the translated toric coordinate  $(u, v_1), v_1 = v - 1$  at  $\xi_1$ , we can write the defining equation of  $\widetilde{C}_2$  as  $v_1^2 - 4u^2v_1 - 239u^4 + (\text{higher terms}) = 0$ , while the exceptional divisor  $\hat{E}(P_2)$  is defined by u = 0. Note that  $(\widetilde{C}_2, \xi_1) = B_{4,2}$ . Thus  $\pi_1^*f$  is non-degenerate and we need one more toric modification centered at  $\xi_1$ . Then the resolution is given by (2), Figure 3. We denote this class of singularity as  $(C, O) \sim B_{6,2} \circ (B_{1,1}^2)^{B_{4,2}} \circ B_{2,6}$ .



3.2. Some lemmas for general torus curves. First we prepare some lemmas for the general torus curves of type (p,q). Let  $C = \{f = 0\}$  be a curve of torus type (p,q) which can be written as  $f = f_q^p + f_p^q$  where  $f_p$  and  $f_q$  are polynomial of degree p and q respectively. We assume that  $p \ge q \ge 2$ . We put  $C_q := \{f_q = 0\}$  and  $C_p := \{f_p = 0\}$ . Suppose that  $O \in C_q \cap C_p$  and let  $\iota$  be the local intersection multiplicity  $I(C_p, C_q; O)$ . We recall a following key lemmas.

**Lemma 1** (Lemma 1 of [1]). Suppose that  $C_p$  is non-singular at O and then the singularity (C, O) is topologically equivalent to the Brieskorn-Pham singularity  $B_{pl,q}$ .

**Lemma 2** (Lemma 4.3 of [6]). Suppose that  $(C_p, O)$  is singular with multiplicity m and  $(C_q, O)$  is smooth. If  $C_q$  intersects transversely with  $C_p$  and p < qm, the singularity (C, O)is topologically equivalent to  $B_{mq,p}$ .

**Lemma 3.** Suppose that  $(C_p, O)$  is singular with multiplicity m and the tangent cone of  $C_p$  consist of m distinct lines. Then  $C_p$  consists of m smooth components at O. Consider the local factorization  $f_p = \prod_{i=1}^m g_i$ .

- (1) Suppose that  $C_q$  is smooth at O and p < qm.
  - (a) If  $\iota < \frac{p}{p-q}(m-1)$ , then  $(C, O) \sim B_{q\iota, p}$ . (b) If  $\iota > \frac{p}{p-q}(m-1)$ ,  $(C, O) \sim B_{\beta, q} \circ B_{q(m-1), p-q}$  where  $\beta = p(\iota m + 1) 1$ (c) If  $\iota = \frac{p}{p-q}(m-1)$  and the coefficients are generic, then  $(C,O) \sim B_{q\iota,p}$ .
- (2) Suppose that m = 2,  $C_q$  is smooth and p > 2q. Then  $(C, O) \sim B_{p(\iota-1)-q,q} \circ B_{q,p-q}$ .
- (3) Suppose that  $(C_q, O)$  is singular with multiplicity 2 and the tangent cone  $T_O C_q$ consists of distinct two lines and let  $f_q = h_1 h_2$  be the local factorization and we put  $\ell_i = \{h_i = 0\}, i = 1, 2$ . Let  $\nu_i = I(\ell_i, C_p; O), i = 1, 2$ . We assume that  $g_1(x_2, y_2) = x_2, g_2(x_2, y_2) = y_2$  for a local coordinate system  $(x_2, y_2)$  and

$$I(\ell_1, g_i; O) = 1, \quad (i \neq 1) \quad I(\ell_2, g_j; O) = 1, \quad (j \neq 2)$$

and put  $\nu_1 = I(\ell_1, q_1; O)$  and  $\nu_2 = I(\ell_2, q_2; O)$ . Assume that 2p > qm. Then

$$(C,O) \sim B_{\beta_{1},2} \circ (B^{q}_{m-2,\ m-2})^{(m-2)B_{2p-qm,q}} \circ B_{2,\beta_{2}}, \quad \beta_{i} = p(\nu_{i}+1) - q(m-1)$$

*Proof.* By the assumption, the tangent cone of  $C_p$  consists of m distinct lines. This implies  $(C_p, O)$  has m smooth components which are transverse each other.

First we consider the case (1). First we take a local coordinate system  $(x_1, y_1)$  so that  $C_q$  is defined by  $y_1 = 0$ . Let  $f_p(x_1, y_1) = \prod_{i=1}^m g_i(x_1, y_1)$  be the factorization in  $\mathcal{O}$  such that  $g_1(x_1, y_1) = y_1 + \alpha_1 x_1^{\nu} + (\text{higher terms}), g_i(x_1, y_1) = y_1 - \alpha_i x_1 + (\text{higher terms})$  with  $\alpha_i \neq 0$   $(2 \leq i \leq m)$ . In this expression, we have  $\iota = \nu + m - 1$ . In the case of (a):  $\iota < \frac{p}{p-q}(m-a)$ , the Newton boundary of  $f_p(x_1, y_1)^q$ ,  $f_q(x_1, y_1)^p$  are as the left hand side of Figure 4. Thus it is easy to see that  $f(x_1, y_1)$  is non-degenerate in this coordinate and we have  $(C, O) \sim B_{q\iota,p}$ .

In the case of (b):  $\iota > \frac{p}{p-q}(m-a)$ ,  $f(x_1, y_1)$  are degenerate in this coordinate. We take another coordinate:  $(x_2, y_2)$  with  $x_2 = x_1, y_2 = g_1(x_1, y_1)$  so that  $y_2|f_p$ . Then in this coordinate  $(x_2, y_2)$ , the Newton boundaries of  $f_p(x_2, y_2)$  and  $f_q(x_2, y_2)$  are given as the right hand side of Figure 4 so that  $f(x_2, y_2)$  are now non-degenerate and the assertion follows.



For the proof of (2), we take a local coordinate system so that  $f_p(x, y) = c x_1 y_1, c \neq 0$ . Then we may assume that  $f_q(x_1, y_1) = y_1 + c' x^{\iota-1} + (\text{higher terms})$ . The the assertion is immediate from Newton boundary argument.

Next we consider the case (3). We have chosen a local coordinate system  $(x_2, y_2)$  so that  $g_1(x_2, y_2) = c x_2 \ (c \neq 0)$  and  $g_2(x_2, y_2) = y_2$ . Put  $g_i(x_2, y_2) = y_2 - \alpha_i x_2 + (\text{higher terms})$ ( $3 \leq i \leq m$ ). By the assumption, we can write

 $h_1(x_2, y_2) = x_2 + d_1 y_2^{\nu_1} + (\text{higher terms}), \quad h_2(x_2, y_2) = y_2 + d_2 x_2^{\nu_2} + (\text{higher terms})$ 

with  $d_1, d_2 \neq 0$ . Then the Newton principal part of f can be written as:

$$\mathcal{N}(f, x_2, y_2) = d_1^p y_2^{p(\nu_1 + 1)} + c^q x_2^q y_2^q \prod_{i=3}^m (y_2 - \alpha_i x_2)^q + d_2^p x_2^{p(\nu_2 + 1)}$$

Newton boundary of f consists of three faces  $\Delta_1$ ,  $\Delta_2$  and  $\Delta_3$  and f is non-degenerate on  $\Delta_1$ ,  $\Delta_3$  but degenerate on  $\Delta_2$ . We take an admissible toric blowing-up  $\pi : X \to \mathbb{C}^2$  with respect to some regular simplicial cone subdivision  $\Sigma^* = \{P_0, \ldots, P_k\}$ . Assume that  $P_{\gamma} = {}^t(1,1)$ , the weight vector of the homogeneous face  $\Delta_2$  and put  $P_{\gamma+1} = {}^t(n, n+1)$ . Then the pull-back of  $f_p$  and  $f_q$  to the coordinate chart  $(\mathbb{C}^2_{\sigma}, (u_{\gamma}, v_{\gamma}))$  with  $\sigma = \text{Cone}(P_{\gamma}, P_{\gamma+1})$  are given by

$$\begin{aligned} \pi_{\sigma}^* f_p(u_{\gamma}, v_{\gamma}) &= u_{\gamma}^m \, v_{\gamma} \prod_{i=3}^m \left\{ g_i(u_{\gamma}v_{\gamma}^n, u_{\gamma}v_{\gamma}^{n+1})/u_{\gamma} \right\} \\ \pi_{\sigma}^* f_q(u_{\gamma}, v_{\gamma}) &= u_{\gamma}^2(h_1(u_{\gamma}v_{\gamma}^n, u_{\gamma}v_{\gamma}^{n+1})/u_{\gamma}) \times (h_2(u_{\gamma}v_{\gamma}^n, u_{\gamma}v_{\gamma}^{n+1})/u_{\gamma}) \end{aligned}$$

where

$$g_i(u_{\gamma}v_{\gamma}^n, u_{\gamma}v_{\gamma}^{n+1})/u_{\gamma} \equiv v_{\gamma} - \alpha_i \mod u_{\gamma}$$
$$h_1(u_{\gamma}v_{\gamma}^n, u_{\gamma}v_{\gamma}^{n+1})/u_{\gamma} \equiv 1 \mod u_{\gamma}$$
$$h_2(u_{\gamma}v_{\gamma}^n, u_{\gamma}v_{\gamma}^{n+1})/u_{\gamma} \equiv v_{\gamma} \mod u_{\gamma}.$$

Thus we get

$$\pi^*_{\sigma}(f(u_{\gamma}, v_{\gamma})) = u^{qm}_{\gamma}(\bar{f}^q_p(u_{\gamma}, v_{\gamma}) + u^{2p-qm}_{\gamma}\bar{f}^p_q(u_{\gamma}, v_{\gamma}))$$
$$= u^{qm}_{\gamma}\left(\prod_{i=3}^m (v_{\gamma} - \alpha_i)^q + u^{2p-qm}\bar{f}_q(u_{\gamma}, v_{\gamma}) + (\text{higher terms})\right)$$

and  $\overline{f}_q(\alpha_i, 0) \neq 0$ . We put  $\xi_i = (0, \alpha_i)$  and we take a local coordinates  $(u_\gamma, v_{\gamma,i})$  at  $\xi_i$  with  $v_{\gamma,i} = v_\gamma - \alpha_i$ . 6The above expression implies  $(\widetilde{C}, \xi_i) \sim B_{2p-qm,q}$ .

The assertion (1) of Lemma 3 can be generalized for the case where the tangent cone of  $C_p$  may have some factors with multiplicity as follows.

**Lemma 4.** Suppose that  $C_p$  is singular at O and  $C_q$  is smooth at O. Let m be the multiplicity of  $(C_p, O)$ . We take a local coordinate system  $(x_1, y_1)$  so that  $C_q$  is defined by  $y_1 = 0$ . We assume that p < qm and

-either  $y_1 \not\downarrow (f_p)_m$  (this implies that  $C_q$  intersects transversely with  $T_O C_p$ ), or  $-y_1 = 0$  is a simple tangent line of  $T_O C_p$  (this is equivalent to  $y_1 | (f_p)_m, y_1^2 \not\downarrow (f_p)_m$ ). where  $(f_p)_m$  is the homogeneous part of degree m of  $f_p$ . Then we have:

(1) If 
$$\iota < \frac{p}{p-q}(m-1)$$
, then  $(C, O) \sim B_{q\iota, p}$ 

(2) If 
$$\iota > \frac{p}{p-q}(m-1)$$
,  $(C,O) \sim B_{\beta, q} \circ B_{q(m-1), p-q}$  where  $\beta = p(\iota - m + 1) - q(m-1)$ .  
(3) If  $\iota = \frac{p}{p-q}(m-1)$  and the coefficients are generic, then  $(C,O) \sim B_{q\iota,p}$ .

*Proof.* The proof is completely parallel to that of Lemma 3.

We come back to our original situation of a torus curve of type (2, 5):

$$C: f_5(x,y)^2 + f_2(x,y)^5 = 0, \quad C_5: f_5(x,y) = 0, \ C_2: f_2(x,y) = 0.$$

First we consider the case that  $C_2$  is reduced in section 4 and 5. Next consider the case of C being a linear torus curve in section 6. For the classification, we start from the following generic equations:

$$f_2(x,y) = \sum_{i+j \le 2} a_{ij} x^i y^j, \quad f_5(x,y) = \sum_{i+j \le 5} b_{ij} x^i y^j.$$

Hereafter x, y are the affine coordinates x = X/Z, y = Y/Z on  $\mathbb{C}^2 := \mathbb{P}^2 \setminus \{Z = 0\}$ . As we assume that  $C_2, C_5$  pass through the origin, we have  $a_{00} = b_{00} = 0$ . We study the inner singularity  $O \in C_2 \cap C_5$ . We denote hereafter the multiplicities of  $C_2$  and  $C_5$  at the origin O by  $m_2$  and  $m_5$  respectively and the intersection multiplicity  $I(C_2, C_5; O)$  of  $C_2$  and  $C_5$  at O by  $\iota$ . By Bézout theorem, we have the inequalities:

$$1 \le \iota \le 10, \quad m_2 m_5 \le \iota$$

For the classification of possible topological types of the singularity (C, O), we divide the situations into the 5 cases, corresponding to the values of  $m_5$ . Then for a fixed  $m_5$ , we consider the subcases, corresponding to  $\iota$ ,  $m_5 \leq \iota \leq 10$  taking the geometry of the intersection of  $C_2$  and  $C_5$  at O into account. And each case has several sub-cases by type of the singularity of  $(C_5, O)$ .

- (1) Case I.  $m_5 = 1$ . The quintic  $C_5$  is smooth.
- (2) Case II.  $m_5 = 2$ . We divide this case into two subcases (a) and (b) by type of tangent cone of  $C_5$ .
  - (a) The tangent cone of  $C_5$  is distinct two lines i.e.,  $(C_5, O) \sim A_1$ .
  - (b) The tangent cone of  $C_5$  is a single line with multiplicity 2.
- (3) Case III.  $m_5 = 3$ . We divide this case into three subcases by type of tangent cone of  $C_5$ .
  - (a) The tangent cone of  $C_5$  is distinct three lines.
  - (b) The tangent cone of  $C_5$  is a double line and another line.
  - (c) The tangent cone of  $C_5$  is a single line with multiplicity 3.
- (4) Case IV.  $m_5 = 4$ . We divide this case into five subcases by type of tangent cone of  $C_5$ .
  - (a) The tangent cone of  $C_5$  is distinct four lines.
  - (b) The tangent cone of  $C_5$  is a double line and distinct two line.
  - (c) The tangent cone of  $C_5$  is a triple line and another line.
  - (d) The tangent cone of  $C_5$  is a single line with multiplicity 4.
  - (e) The tangent cone of  $C_5$  is two double line.

(5) Case V.  $m_5 = 5$ . The quintic  $C_5$  is five lines.

4.1. Case I:  $m_5 = 1$ . Quintic  $C_5$  is smooth. This case is determined by Lemma 1 as follows.

**Proposition 1.** Suppose that  $C_5$  is smooth at O and let  $\iota = I(C_5, C_2; O)$  be the local intersection multiplicity. Then  $(C, O) \sim B_{5\iota,2}$  for  $\iota = 1, \cdots, 10$ .

4.2. Case II:  $m_5 = 2$ . We divide Case II into two subcases (a) and (b) by type of tangent cone  $T_OC_5$  of  $C_5$ .

(a)  $T_O C_5$  consists of distinct two lines i.e.,  $(C_5, O) \sim A_1$ .

(b)  $T_O C_5$  consists of a line with multiplicity 2.

4.2.1. Case II-(a). We consider the subcase (a). In this case, we assume that the tangent cone of  $C_5$  is given xy = 0,  $f_5(x, y) = xy + (\text{higher terms})$ .

**Proposition 2.** Under the situation in (a), we have the following possibilities.

- (1) We assume that the conic is smooth and the tangent cone of  $C_2$  is y = 0 ( $a_{10} = 0$ ,  $a_{01} \neq 0$ ) for simplicity. The possibility of (C, O) are  $B_{5\iota-7,2} \circ B_{2,3}$  for  $2 \le \iota \le 10$ .
- (2) For a conic which consists of two lines  $\ell_1, \ell_2$  (i.e.,  $a_{01} = a_{10} = 0$ ), the generic singularity is  $B_{8,2} \circ B_{2,8}$ . Further degeneration occurs when these lines are tangent to one or both tangent cones of  $C_5$ . Put  $\iota_i = I(\ell_i, C_5; O)$  (i = 1, 2). Then  $4 \leq \iota_1 + \iota_2 \leq 10$  and the corresponding singularity is  $B_{5\iota_2-2,2} \circ B_{2,5\iota_1-2}$ .

*Proof.* Both assertions are immediate from Lemma 2 and Lemma 3

4.2.2. Case II-(b). Now we consider the case (II-b). So we assume that  $m_5 = 2$  and the tangent cone of  $C_5$  is given by  $L : y^2 = 0$  (with multiplicity 2). We divide this subcase (II-b) into two subcases: (b-1)  $m_2 = 1$  and (b-2)  $m_2 = 2$ .

(b-1) Assume that  $m_5 = 2$ ,  $m_2 = 1$  and the tangent cone of  $C_5$  is  $y^2 = 0$ .

**Proposition 3.** Suppose that the tangent cone of  $C_5$  is a line with multiplicity 2 and  $C_2$  is smooth. Then we have the following possibilities.

- (1) If  $C_2$  and  $C_5$  are transverse at  $O(\iota = 2)$ , then  $(C, O) \sim B_{5,4}$ .
- (2) If  $(C_5, O) \sim B_{3,2}$  and  $\iota = 3$ , we have  $(C, O) \sim (B_{3,2}^2)^{B_{3,2}}$ .
- (3) If  $(C_5, O) \sim B_{4,2}$ , we have  $(C, O) \sim (B_{4,2}^2)^{(B_{5\iota-18,2}+B_{2,2})}$  for  $\iota = 4, \cdots, 10$ . (Here the upper  $(B_{5\iota-18,2}+B_{2,2})$  implies we have two non-degenerate singularities  $B_{5\iota-18,2}, B_{2,2}$ sitting on two different points on the exceptional divisor  $\widehat{E}(P)$ ,  $P = {}^t(1,2)$ . after one toric modification.)
- (4) If  $(C_5, O) \sim B_{5,2}$ , we have (a)  $(C, O) \sim B_{10,4}$  generically and  $B_{k,2} \circ B_{5,2}$ ,  $6 \le k \le 15$  for  $\iota = 4$ , (b)  $(B_{5,2}^2)^{B_{5,2}}$  for  $\iota = 5$ .
- (5) If  $(C_5, O) \sim B_{6,2}$ , we have (a)  $(C, O) \sim B_{10,4}$  for  $\iota = 4$  and (b)  $(C, O) \sim (B_{6,2}^2)^{(B_{5\iota-27,2}+B_{3,2})}$  for  $\iota = 6, \ldots, 10$ .
- (6) If  $(C_5, O) \sim B_{7,2}$ , we have
  - (a)  $(C, O) \sim B_{10,4}$  for  $\iota = 4$  and
  - (b)  $(C,O) \sim (B_{7,2}^2)^{B_{5\iota-28,2}}$  for  $\iota = 6, 7$ .
- (7) If  $(C_5, O) \sim B_{8,2}$ , we have (a)  $(C, O) \sim B_{10,4}$  for  $\iota = 4$ ,

(b) 
$$(C, O) \sim B_{15,4}$$
 for  $\iota = 6$ ,  
(c)  $(B_{8,2}^2)^{(B_{5\iota-36,2}+B_{4,2})}$  for  $\iota = 8, 9, 10$ .  
(8) If  $(C_5, O) \sim B_{9,2}$ , we have  
(a)  $(C, O) \sim B_{10,4}$  for  $\iota = 4$ ,  
(b)  $(C, O) \sim B_{15,4}$  for  $\iota = 6$ ,  
(c)  $(C, O) \sim (B_{9,2}^2)^{B_{5\iota-35,2}}$  for  $\iota = 8, 9$ .  
(9) If  $(C_5, O) \sim B_{10,2}$ , we have  
(a)  $(C, O) \sim B_{10,4}$  for  $\iota = 4$ ,  
(b)  $(C, O) \sim B_{10,4}$  for  $\iota = 6$ ,  
(c)  $(C, O) \sim B_{20,4}$  and  $B_{k,2} \circ B_{10,2}$   $(k = 11, 12)$  for  $\iota = 8$ ,  
(d)  $(B_{10,2}^2)^{2B_{5,2}}$  for  $\iota = 10$ .  
(10) If  $(C_5, O) \sim B_{11,2}$ , we have  
(a)  $(C, O) \sim B_{10,4}$  for  $\iota = 4$ ,  
(b)  $(C, O) \sim B_{15,4}$  for  $\iota = 6$ ,  
(c)  $(C, O) \sim B_{20,4}$  for  $\iota = 8$ ,  
(d)  $(C, O) \sim B_{12,2}$ , we have  
(a)  $(C, O) \sim B_{12,2}$ , we have  
(a)  $(C, O) \sim B_{12,2}$ , we have  
(a)  $(C, O) \sim B_{12,4}$  for  $\iota = 4$ ,  
(b)  $(C, O) \sim B_{15,4}$  for  $\iota = 8$ ,  
(c)  $(C, O) \sim B_{15,4}$  for  $\iota = 8$ ,  
(d)  $(C, O) \sim B_{13,2}$ , we have  
(a)  $(C, O) \sim B_{13,2}$ , we have  
(a)  $(C, O) \sim B_{13,2}$ , we have  
(b)  $(C, O) \sim B_{13,2}$ , we have  
(c)  $(C, O) \sim B_{13,2}$ , we have  
(c)  $(C, O) \sim B_{13,2}$ , we have  
(c)  $(C, O) \sim B_{13,4}$  for  $\iota = 4$ ,  
(c)  $(C, O) \sim B_{13,4}$  for  $\iota = 4$ ,  
(c)  $(C, O) \sim B_{13,4}$  for  $\iota = 4$ ,  
(c)  $(C, O) \sim B_{13,4}$  for  $\iota = 4$ ,  
(c)  $(C, O) \sim B_{13,4}$  for  $\iota = 4$ ,  
(c)  $(C, O) \sim B_{13,4}$  for  $\iota = 4$ ,  
(c)  $(C, O) \sim B_{13,4}$  for  $\iota = 4$ ,  
(c)  $(C, O) \sim B_{13,4}$  for  $\iota = 4$ ,  
(c)  $(C, O) \sim B_{13,4}$  for  $\iota = 4$ ,  
(c)  $(C, O) \sim B_{13,4}$  for  $\iota = 4$ ,  
(c)  $(C, O) \sim B_{20,4}$  for  $\iota = 8$ ,  
(c)  $(C, O) \sim B_{20,4}$  for  $\iota = 8$ ,  
(d)  $(C, O) \sim B_{23,4}$  for  $\iota = 10$ .

*Proof.* We can proceed the classification mainly using the local intersection multiplicity  $\iota = I(C_2, C_5; O)$  and geometry of  $C_2$  and  $C_5$ . By the assumption  $m_5 = 2$ , we have  $\iota \geq 2$ . When  $\iota = 2, C_2$  intersects transversely with  $C_5$  at the origin and then  $(C, O) \sim B_{5,4}$ . Thus hereafter we assume that  $\iota \geq 3$ .

Assume that  $(C_5, O) \sim A_{\ell-1}$ . Then by a triangular change of local coordinates  $(x, y_1)$  where  $y_1 = y + c_2 x^2 + \cdots + c_{k-1} x^{k-1}$ , we can write  $f_5$  as

$$f_5(x, y_1) = \alpha y_1^2 + \beta x^{\ell} + (\text{higher terms}), \quad \ell \ge 2(k-1), \, \alpha, \beta \ne 0.$$

A simple computation shows that  $\ell \leq 13$ . Now we can write  $f_2$  in this coordinates as

$$f_2(x, y_1) = \gamma y_1 + \delta x^{\nu} + (\text{higher terms}), \ \nu \ge 2, \ \gamma, \delta \ne 0.$$

(Here  $\delta = 0$  if  $\nu = \infty$ , i.e.,  $y_1|f_2$ .) First we notice that

(\*):  $\iota \geq \min(2\nu, \ell)$  and the equality holds except the case  $\ell = 2\nu$  and  $\alpha\delta^2 + \beta\gamma^2 = 0$ .

We observe that

1. If  $5\nu < 2\ell$ , f is non-degenerate in this coordinates and  $(C, O) \sim B_{5\nu,4}$ .

2. If  $5\nu = 2\ell$  (in this situation, the possible pairs which satisfy this condition are  $(\nu, \ell) = (2, 5), (4, 10)$ ), we have  $\mathcal{N}(f, x, y) = \alpha^2 y_1^4 + 2\alpha\beta x^\ell y_1^2 + (\beta^2 + \delta^5) x^{2\ell}$  and if  $(\beta^2 + \delta^5) \neq 0$ , we have  $(C, O) \sim B_{2\ell,4}$ .

If  $(\beta^2 + \delta^5) = 0$ , we can see that the Newton boundary has two faces with  $R = (\ell, 2)$  as the common vertex of the faces and f can be non-degenerate on these faces, after taking a suitable triangular change of coordinates  $(x, y_2)$ .

3. If  $5\nu > 2\ell$ , it is easy to see that the Newton principal part of  $f(x, y_1)$  is given by  $(\alpha y_1^2 + \beta x^\ell)^2$  which implies that  $f(x, y_1)$  is degenerate in this coordinate. So we need to take first a toric modification  $\pi : X \to \mathbb{C}^2$  with respect to the canonical regular simplicial cone subdivision  $\{P_0, \ldots, P_m\}$  and we have to see the total transform equation in X.

The weight vector of  $A_{\ell-1}$  is given as  $P = {}^{t}(2,\ell), {}^{t}(1,\ell/2)$  for  $\ell$  is odd or even respectively. Note also the germ  $A_{\ell-1}$  has two smooth components if  $\ell$  is even. Thus the description for the toric modification has to be divided in two cases.

**Case 1.**  $\ell$  is odd. The weight vector of  $\Gamma(f_5; x, y_1)$  is given by  $P = {}^t(2, \ell)$ . We may assume that  $P = P_s$  and we consider the cone  $\sigma := \text{Cone}(P_s, P_{s+1})$  and we consider the corresponding unimodular matrix

$$\sigma := \begin{pmatrix} 2 & a \\ \ell & b \end{pmatrix}, \quad 2b - a\ell = 1$$

Let (u, v) be the toric coordinates of this chart. Then in this coordinates, we have  $x = u^2 v^a$ ,  $y_1 = u^\ell v^b$ . Then we can write

$$\pi^* f_5(u,v) = u^{2\ell} v^{a\ell} \tilde{f}_5(u,v), \quad \tilde{f}_5(u,v) = \alpha v + \beta + h_5(u,v)$$
  
$$\pi^* f_2(u,v) = u^{\mu} v^{\mu'} \tilde{f}_2(u,v), \quad \mu = \min(\ell, 2\nu), \quad \mu' = \min(b, a\nu)$$

Putting  $\xi = (0, -\beta/\alpha), \eta = 5\mu - 4\ell$ , we can write as

$$\pi^* f(u,v) = u^{4\ell} v^{2a\ell} f(u,v)$$
$$\tilde{f}(u,v) = \tilde{f}_5(u,v)^2 + u^\eta v^{5\mu'-2a\ell} \tilde{f}_2(u,v)^5$$

Thus using admissible translated toric coordinates  $(u, v_2)$ ,  $v_2 = v_1 + h(u)$ ,  $v_1 = v + \beta/\alpha$  for some polynomial h, the strict transform is defined as

$$\alpha^2 v_2^2 + \epsilon u^{\eta'} + (\text{higher terms}) = 0, \ \epsilon \neq 0$$

which implies  $(\tilde{C},\xi) \sim B_{\eta',2}$  and the tangent cone is transverse to the exceptional divisor u = 0 where  $\eta' \geq \eta$ .

**Case 2.**  $\ell$  is even. The weight vector of  $\Gamma(f_5; x, y_1)$  is given by  $P = {}^t(1, k)$  where  $\ell = 2k$ . We may consider that there is a cone corresponding to a unimodular matrix

$$\sigma := \begin{pmatrix} 1 & 0 \\ k & 1 \end{pmatrix}$$

Let (u, v) be the toric coordinates of this chart. Then in this coordinates, we have  $x = u, y_1 = u^k v$ . Then we can write

$$\pi^* f_5(u,v) = u^{\ell} \tilde{f}_5(u,v), \quad \tilde{f}_5(u,v) = \alpha(v+\alpha_1)(v+\alpha_2) + uvh_5(u,v)$$
  
$$\pi^* f_2(u,v) = u^{\mu} \tilde{f}_2(u,v), \ \mu = \min(k,\nu)$$

Putting  $\xi_i = (0, \alpha_i), (1 = 1, 2), \eta = 5\mu - 4\ell$ , we can write as

$$\pi^* f(u, v) = u^{2\ell} \left( \tilde{f}_5(u, v)^2 + u^\eta \tilde{f}_2(u, v)^5 \right)$$

Then the strict transform  $\widetilde{C}$  has two components. Thus using admissible translated toric coordinates  $(u, v'_i)$ ,  $v'_i = v_i + h(u)$ ,  $v_i = v + \alpha_i$  (i = 1, 2) for some polynomial h, the total

transform  $\pi^* f$  is described as

$$\pi^* f(u, v_i') = u^{2\ell} \Big( \alpha^2 (\alpha_1 - \alpha_2)^2 v_i'^2 + \epsilon u^{\eta'} + (\text{higher terms}) \Big) = 0, \ \epsilon \neq 0$$

which implies  $(\tilde{C}, \xi_i) \sim B_{\eta', 2}$  where  $\eta' \geq \eta$ .

Putting the above strategy into consideration, we will explain some more detail for several cases.

First we consider Case (2):  $(C_5, O) \sim B_{3,2}, \iota = 3 \ (\ell = 3, \nu \ge 2).$ 

We have  $f_5(x, y) = \alpha y^2 + \beta x^3 + (\text{higher terms})$ . We have to consider the toric modification in the toric coordinate chart:

$$\sigma = \begin{pmatrix} 2 & 1 \\ 3 & 2 \end{pmatrix}, \ \pi(u, v) = (u^2 v, u^3 v^2)$$

in above observation. Then taking the translated coordinates  $(u, v_1)$ ,  $v_1 = v + \beta/\alpha$ , we have

$$\pi^* f_5(u, v_1) = u^6 (v_1 - \beta/\alpha)^3 (\alpha v_1 + cu + h_5(u, v_1))$$
  

$$\pi^* f_2(u, v_1) = u^3 (v_1 - \beta/\alpha)^2 (\delta u + \gamma + h_2(u, v_1))$$
  

$$\pi^* f(u, v_1) = u^{12} (v_1 - \beta/\alpha)^6 \tilde{f}(u, v_1)$$
  

$$\tilde{f}(u, v_1) = (\alpha v_1 + cu + h_5(u, v_1))^2 + u^3 (v_1 - \beta/\alpha)^4 (\delta u + \gamma + h_2(u, v_1))^5$$

Now we can see that

$$\tilde{f}(u, v_2) = v_2^2 + c'u^3 + (\text{higher terms}), \quad c' \neq 0 \quad v_2 = \alpha v_1 + cu$$

which implies the corresponding singularity is  $(B_{3,2}^2)^{B_{3,2}}$ .

Next we consider the case (3):  $(C_5, O) \sim B_{4,2}$  and  $\iota \geq 4$ . Thus

$$f_5(x, y_1) = \alpha y_1^2 + \beta x^4 + \text{(higher terms)}$$
  
$$f_2(x, y_1) = \gamma y_1 + \delta x^{\nu} + \text{(higher terms)}, \ \nu \ge 2.$$

Note that  $\nu \ge 2$  and the case  $\iota > 4$  only if  $\nu = 2$  and  $\alpha \, \delta^2 + \beta \, \gamma^2 = 0$ . Thus for simplicity, we assume that  $\nu = 2$ . For the simplicity of the calculation, we put:

$$f_5(x, y_1) = \alpha y_1^2 + \beta x^4 + (\text{higher terms}) = \alpha (y_1 + \alpha_1 x)(y_1 + \alpha_2 x) + (\text{higher terms}),$$
  
$$f_2(x, y_1) = \gamma y_1 + \delta x^2 + (\text{higher terms})).$$

The corresponding toric chart is associated with:

$$\sigma = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}, \ \pi(u, v) = (u, u^2 v)$$

by the above consideration. Note that  $\iota = 4$  if and only if  $\alpha \delta^2 + \beta \gamma^2 \neq 0$ . Then taking the translated coordinates  $(u, v_1), v_1 = v + \alpha_1$  (respectively  $(u, v_2), v_2 = v + \alpha_1$ ), we have

$$\begin{aligned} \pi^* f_5(u, v_1) &= u^4 (\alpha (\alpha_1 - \alpha_2) v_1 + c_1 u + h_5(u, v_1)), \\ \pi^* f_2(u, v_1) &= u^2 ((\gamma v_1 - \gamma \alpha_1 + \delta) + u h_2(u, v_1)), \\ \pi^* f(u, v_1) &= u^8 \tilde{f}(u, v_1), \quad (resp. \ \pi^* f(u, v_2) = u^8 \tilde{f}(u, v_2)) \\ \tilde{f}(u, v_1) &= ((\alpha (\alpha_1 - \alpha_2) v_1 + c_1 u)^2 + (\delta - \gamma \alpha_1)^5 u^2 + (\text{higher terms})) \\ (resp. \ \tilde{f}(u, v_2) &= (\alpha (\alpha_2 - \alpha_1) v_1 + c_2 u)^2 + (\delta - \gamma \alpha_2)^5 u^2 + (\text{higher terms})) \end{aligned}$$

where  $c_i$  (i = 1, 2) are constant. Then if  $\iota = 4$ , we have  $\alpha \delta^2 + \beta \gamma^2 \neq 0$  and we see that  $(\tilde{C}, \xi_i) = A_1$  for i = 1, 2. Thus  $(C, O) \sim (B_{4,2}^2)^{2B_{2,2}}$  and resolution graph is given by Figure 5.



The case  $\nu > 2$  gives the same conclusion as above.

If  $\nu = 2$ ,  $\iota > 4$  and assuming  $\alpha \, \delta^2 + \beta \, \gamma^2 \neq 0$ , we have  $(\tilde{C}, \xi_2) = A_1$  but  $(\tilde{C}, \xi_1)$  is bigger that  $A_1$ . Thus we have to take a triangular change of coordinates  $(u, v'_1)$  so that  $\tilde{C}$  is defined at  $\xi_1$  as  $(v'_1)^2 + c' \, u^k + (\text{higher terms}) = 0$ . The explicit computation shows that the possible k are  $5\iota - 18$ ,  $4 \leq \iota \leq 10$ .

Next we consider the assertion (4):  $(C_5, O) \sim B_{5,2} \ \iota \geq 4$ . We put as above

$$f_5(x, y_1) = \alpha y_1^2 + \beta x^5 + (\text{higher terms}), \quad f_2(x, y_1) = \gamma y_1 + \delta x^{\nu} + (\text{higher terms}).$$

Note that  $\iota = 4$  if and only if  $\nu = 2$ . If  $\nu > 2$ ,  $\iota = 5$  and  $\mathcal{N}(f, x, y_1) = (\alpha y_1^2 + \beta x^5)^2$ and we have to take a toric modification. If  $\nu = 2$ , then  $\iota = 4$  and as  $\mathcal{N}(f, x, y_1) = y_1^4 + 2\beta x^5 y_1^2 + (\beta^2 + \delta^5) x^{10}$ , we see that  $(C, O) \sim B_{10,4}$  if  $\iota = 4$  and  $\beta^2 + \delta^5 \neq 0$ . If  $\beta^2 + \delta^5 = 0$ , the Newton boundary has two faces.

-First we consider the case  $\iota = 4$  (so  $\delta \neq 0$ ) and  $\beta^2 + \delta^5 = 0$ . Then  $\mathcal{N}(f, x, y_1) = y_1^4 + 2\beta x^5 y_1^2 + \gamma_8 x^8 y_1 + \gamma_{11} x^{11}$  and  $\Gamma(f; x, y_1)$  consists of two face  $\Delta_1$  and  $\Delta_2$ . Clearly f is non-degenerate on  $\Delta_1$ . If f is degenerate on  $\Delta_2$ , we take a suitable triangulated change of coordinates  $(x, y_2)$  so that  $\mathcal{N}(f; x, y_2) = \alpha^2 y_2^4 + 2\alpha \beta x y_2^2 + \gamma' x^{11+k}, k = 0, \ldots, 9$ . This implies  $(C, O) \sim B_{k+6,2} \circ B_{5,2}$ .

–Secondly, we consider the case  $\iota \geq 5$  (i.e.,  $\nu > 2$ ). This case, by the previous consideration, we see that  $\iota = 5$ . Then  $\mathcal{N}(f, x, y_1) = (\alpha y_1^2 + \beta x^5)^2$  and  $\Gamma(f; x, y_1)$  consists of one face  $\Delta$  with the weight vector  $P = {}^t(2, 5)$  and f is degenerate on  $\Delta$ . We consider the toric modification with respect to the canonical regular subdivision  $\Sigma^*$  of  $\Gamma^*(f; x, y_1)$ . The toric coordinate chart which intersects the strict transform  $\widetilde{C}$  is described by a unimodular matrix

$$\sigma = \begin{pmatrix} 2 & 1 \\ 5 & 3 \end{pmatrix}, \ \pi(u, v) = (u^2 v, u^5 v^3).$$

Then we taking admissible translated toric coordinates  $(u, v_2)$ ,  $v_2 = v_1 + h(u)$ ,  $v_1 = \alpha v + \beta$  for a suitable polynomial h, we have

$$\pi^* f(u, v_2) = u^{20} (v_2 - \beta/\alpha + h(u)) \Big( \alpha^2 v_2^2 + \beta'' u^5 + (\text{higher terms}) \Big), \quad \beta'' \neq 0.$$

Thus we can get  $(C, O) \sim (B_{5,2}^2)^{B_{5,2}}$ . Hence we have the assertion (4) of Proposition 3. The assertions (5),  $\cdots$ , (12) can be shown in a similar manner.

**Remark 1.** Note that the singularity  $B_{25,4}$  in case (12) has Milnor number 72 and 72 is the maximum Milnor number of irreducible curve of degree 10. Thus in this case, C is a rational curve.

The classification Proposition 3 can be rewritten as follows from the viewpoint of  $\iota$ .

- (1) If  $\iota = 2$ , we have  $(C, O) \sim B_{5,4}$ .
- (2) If  $\iota = 3$ , we have  $(C, O) \sim (B_{3,2}^2)^{B_{3,2}}$ .

- (3) If  $\iota = 4$ , we have  $(C, O) \sim (B_{4,2}^2)^{2B_{2,2}}$ ,  $B_{10,4}$  and  $B_{k,2} \circ B_{5,2}$  ( $6 \le k \le 15$ ).
- (4) If  $\iota = 5$ , we have  $(C, O) \sim (B_{4,2}^2)^{(B_{7,2}+B_{2,2})}$  and  $(B_{5,2}^2)^{B_{5,2}}$ .
- (5) If  $\iota = 6$ , we have  $(C, O) \sim (B_{4,2}^2)^{(B_{12,2}+B_{2,2})}$ ,  $(B_{6,2}^2)^{2B_{3,2}}$ ,  $(B_{7,2}^2)^{B_{2,2}}$  and  $B_{15,4}$ .
- (6) If  $\iota = 7$ , we have  $(C, O) \sim (B_{4,2}^2)^{(B_{17,2}+B_{2,2})}, (B_{6,2}^2)^{(B_{8,2}+B_{3,2})}$  and  $(B_{7,2}^2)^{B_{7,2}}$ .
- (7) If  $\iota = 8$ , we have  $(C, O) \sim (B_{4,2}^2)^{(B_{22,2}+B_{2,2})}$ ,  $(B_{6,2}^2)^{(B_{13,2}+B_{3,2})}$ ,  $(B_{8,2}^2)^{2B_{4,2}}$ ,  $(B_{9,2}^2)^{B_{5,2}}$ ,  $B_{20,4}$ ,  $B_{11,2} \circ B_{10,2}$  and  $B_{12,2} \circ B_{10,2}$ .
- (8) If  $\iota = 9$ , we have  $(C, O) \sim (B_{4,2}^2)^{(B_{27,2}+B_{2,2})}$ ,  $(B_{6,2}^2)^{(B_{18,2}+B_{3,2})}$ ,  $(B_{8,2}^2)^{(B_{9,2}+B_{4,2})}$  and  $(B_{9,2}^2)^{B_{10,2}}$ .
- (9) If i = 10, we have  $(C, O) \sim (B_{4,2}^2)^{(B_{32,2}+B_{2,2})}$ ,  $(B_{6,2}^2)^{(B_{23,2}+B_{3,2})}$ ,  $(B_{8,2}^2)^{(B_{14,2}+B_{4,2})}$ ,  $(B_{10,2}^2)^{2B_{5,2}}$ ,  $(B_{11,2}^2)^{B_{6,2}}$ ,  $(B_{12,2}^2)^{2B_{1,2}}$  and  $B_{25,4}$ .

(b-2) Assume that  $C_2$  is a union of two lines meeting at the origin  $(m_2 = 2)$  and the tangent cone of  $C_5$  is  $y^2 = 0$ .

**Proposition 4.** Suppose that the tangent cone of  $C_5$  is a line with multiplicity 2 and  $C_2$  is a union of two lines meeting at the origin. Then the germ (C, O) can take following singularities.

(1) If  $(C_5, O) \sim B_{3,2}$ , we have (a)  $(C,O) \sim (B_{3,2}^2)^{B_{8,2}}$  for  $\iota = 4$  and (b)  $(C,O) \sim (B_{3,2}^2)^{B_{13,2}}$  for  $\iota = 5$ . (2) If  $(C_5, O) \sim B_{4,2}$ , we have (a)  $(C,O) \sim (B_{4,2}^2)^{2B_{2,2}}$  for  $\iota = 4$ , (b)  $(C,O) \sim (B_{4,2}^2)^{2B_{7,2}}$  for  $\iota = 6$ , and (c)  $(C, O) \sim B_{16,2} \circ (B_{2,1}^2)^{B_{7,2}}$  for  $\iota = 7$ . (3) If  $(C_5, O) \sim B_{5,2}$ , we have (a)  $(C, O) \sim B_{10,4}$  or  $B_{k,2} \circ B_{5,2} (6 \le k \le 15)$  for  $\iota = 4$ , (b)  $(C, O) \sim (B_{5,2}^2)^{B_{10,2}}$  for  $\iota = 6$  and (c)  $(C, O) \sim (B_{5,2}^2)^{B_{15,2}}$  for  $(\iota = 7)$ . (4) If  $(C_5, O) \sim B_{6,2}$ , we have (a)  $(C, O) \sim B_{10,4}$  for  $\iota = 4$  and (b)  $(C,O) \sim (B_{6,2}^2)^{2B_{3,2}}$  for  $\iota = 6$ . (5) If  $(C_5, O) \sim B_{7,2}$ , we have (a)  $(C, O) \sim B_{10,4}$  for  $\iota = 4$  and (b)  $(C, O) \sim (B_{7,2}^2)^{B_{2,2}}$  for  $\iota = 6$ . (6) If  $(C_5, O) \sim B_{k,2}$ , we have (a)  $(C, O) \sim B_{10,4}$  for  $\iota = 4$  and (b)  $(C, O) \sim B_{15,4}$   $(8 \le k \le 13)$  for  $\iota = 6$ 

*Proof.* By taking a local coordinates  $(x, y_1)$ , we can assume

 $f_5(x,y_1) = \alpha y_1^2 + \beta x^\ell + (\text{higher terms}), \quad \alpha, \, \beta \neq 0.$ 

Now we assume that  $f_2(x, y_1) = \ell_1(x, y_1)\ell_2(x, y_1)$  where

 $\ell_1 = y_1 + c_\nu x^\nu + (\text{higher terms}), \ \ell_2 = c_2(y_1 + \gamma x) + (\text{higher terms}), \ c_\nu, c_2 \neq 0$ 

We put  $\iota_1 = I(\ell_1, C_5; O) \le 5$ ,  $\iota_2 = I(\ell_2, C_5; O)$ . As  $\gamma \ne 0$ , we have  $\iota_2 = 2$ . Hence  $\iota = \iota_1 + 2$ and we have  $4 \le \iota \le 7$ .

Comparing Newton boundaries of  $f_5^2$  and  $f_2^5$ , and applying an similar argument as in case (b-1), we get assertions of Proposition 4.

4.3. Case III :  $m_5 = 3$ . We divide Case III into three cases by type of the tangent cone of  $C_5$ .

(a)  $T_O C_5$  consists of distinct three lines.

(b)  $T_O C_5$  consists of a double line and a simple line.

(c)  $T_O C_5$  consists of is a line with multiplicity 3.

First we remark that if  $C_2$  is smooth and  $C_2$  intersects transversely with  $C_5$  at the origin  $(\iota = 3)$ , we have  $(C, O) \sim B_{6,5}$  by Lemma 2 in §3. So hereafter, we consider the case  $C_2$  and the tangent cone of  $C_5$  does not intersect transversely.

4.3.1. Case III-(a). : We first consider Case III-(a). We assume that the tangent cone of  $C_5$  consists of distinct three lines.

**Proposition 5.** Under the situation of Case III-(a),  $(C, O) \sim B_{6,5}$  if  $\iota = 3$ . For  $\iota \ge 4$  we have the following possibilities of (C, O).

- (1) Assume that  $C_2$  is smooth and tangent to y = 0. Then (C, O) can be  $B_{5\iota-14,2} \circ B_{4,3}$  for  $\iota = 4, 5, \cdots 10$ .
- (2) Assume that  $C_2$  consists of two distinct lines  $\ell_1, \ell_2$ . Put  $\iota_i = I(\ell_i, C_5; O) \ge 3$ (i = 1, 2). Then  $(C, O) \sim B_{5\iota_2 - 9, 2} \circ (B_{1,1}^2)^{B_{4,2}} \circ B_{2,5\iota_1 - 9}$  with  $\iota_1 + \iota_2 = 6, 7, \cdots, 10$ .

*Proof.* The assertion is immediate from Lemma 3.

4.3.2. Case III-(b). In this case, we may assume that the tangent cone of 
$$C_5$$
 consists of a double line  $\{y = 0\}$  and a single line  $\{x = 0\}$ .

(b-1) Assume first that  $C_2$  is smooth  $(m_2 = 1)$ . If  $\iota = 3$ ,  $(C, O) \sim B_{6,5}$  by Lemma 2. Therefore we consider the case  $\iota \geq 4$ . The common tangent line of  $C_2$  is either  $\{x = 0\}$  or  $\{y = 0\}$ . When the common tangent line is  $\{x = 0\}$ , (C, O) is described by Lemma 4.

So we assume that common tangent cone is  $\{y = 0\}$ . If  $\iota = 4$ , we have  $f_5(x, y) = b_{12}xy^2 + b_{40}x^4 + (\text{higher terms})$  and  $\iota = 4$  if and only if  $b_{40} \neq 0$ . Hence we have  $(C, O) \sim B_{8,5}$ .

Next we consider the case  $\iota \ge 5$  and we take a local coordinates system  $(x, y_1)$  so that  $C_2$  is defined by  $y_1 = 0$  and we have  $f_5(x, y_1) = \beta_{12}xy_1^2 + \beta_{31}x^3y_1 + \beta_{50}x^5 + (\text{higher terms})$  with  $\beta_{12} \ne 0$ . First we assume that  $\iota = 5$ . Then  $\beta_{50} \ne 0$  and we factor  $\mathcal{N}(f; x, y_1)$  as

$$\mathcal{N}(f;x,y_1) = y_1^5 + x^2(\beta_{12}y_1^2 + \beta_{31}x^2y_1 + \beta_{50}x^4)^2 = \prod_{i=1}^5 (y_1 + \alpha_i x^2)$$

and we see that  $\Gamma(f; x, y)$  consists of one face with weight vector  $P = {}^{t}(1, 2)$ . Then we have several cases:

- (1)  $\alpha_1, \ldots, \alpha_5$  are all distinct.
- (2)  $\alpha_1 = \alpha_2$  and  $\alpha_3, \alpha_4, \alpha_5$  are mutually distinct and different from  $\alpha_1$ .
- (3)  $\alpha_1 = \alpha_2 = \alpha_3$  and  $\alpha_4, \alpha_5$  are mutually distinct and different from  $\alpha_1$ .
- (4)  $\alpha_1 = \alpha_2, \alpha_3 = \alpha_4$  and  $\alpha_1 \neq \alpha_3$  and  $\alpha_5$  is different from  $\alpha_1, \alpha_3$ .

We can see by an easy computation that the other cases are not possible. (By a direct computation, we see that if  $\mathcal{N}(f; x, y_1) = 0$  has a root with multiplicity 4,  $\beta_{50} = 0$  and the intersection number jumps up to 6.)

**Lemma 5.** Under the above situation, we further assume that  $\iota = 5$ .

- (1) If  $\alpha_1, ..., \alpha_5$  are all distinct,  $(C, O) \sim B_{10,5}$ .
- (2) If  $\alpha_1 = \alpha_2$  and  $\alpha_3, \alpha_4, \alpha_5$  are mutually distinct and different from  $\alpha_1$ ,  $(C, O) \sim B_{k,2} \circ B_{6,3}$ ,  $(5 \le k \le 12)$ .
- (3) If  $\alpha_1 = \alpha_2 = \alpha_3$  and  $\alpha_4, \alpha_5$  are mutually distinct and different from  $\alpha_1$ ,  $(C, O) \sim B_{k,3} \circ B_{4,2}$   $(k = 7, \cdots, 11)$  or  $B_{3,1} \circ B_{5,2} \circ B_{4,2}$  or  $B_{3,1} \circ B_{7,2} \circ B_{4,2}$  or  $B_{k,2} \circ B_{3,1} \circ B_{4,2}$  (k = 7, 8, 9).
- (4) If  $\alpha_1 = \alpha_2$ ,  $\alpha_3 = \alpha_4$  and  $\alpha_1 \neq \alpha_3$  and  $\alpha_5$  is different from  $\alpha_1$ ,  $\alpha_3$ ,  $(C,O) \sim B_{k_2+4,2} \circ B_{2,1} \circ (B_{2,1}^2)^{B_{k_1,2}}$  where  $(k_1, k_2)$  moves in the set  $\{(k_1, k_2); 13 k_2 \geq k_1 \geq k_2 4, k_2 = 5, \dots, 7\} \cup \{(4, 8), (5, 9)\}.$

Proof. The case (1) is clear. We consider the case (2) and we may assume  $\alpha_1 = \alpha_2$ . Then we see that the Newton boundary  $\Gamma(f; x, y_1)$  has two faces  $\Delta_1$  and  $\Delta_2$  and f is non-degenerate on  $\Delta_1$ . Taking a suitable triangular change of coordinates, we can make f non-degenerate on  $\Delta_2$ . Hence this gives series  $(C, O) \sim B_{k,2} \circ B_{6,3}, k = 5, \cdots, 12$ . We can consider the cases (3) and (4) similarly.

**Remark 2.** In (4) of Lemma 5, we have the following symmetry. Let

$$f(x, y_1) = (y_1 + \alpha_1 x^2)^2 (y_1 + \alpha_3 x^2)^2 (y_1 + \alpha_5 x^2) + \text{(higher terms)}$$

be a defining polynomial of (C, O). First we take a change of coordinates  $(x, y_2) = (x, y_1 + \alpha_1 x^2)$  and we take further changes of coordinates of type  $y_2 \rightarrow y_2 + c x^j$ ,  $2 \leq j \leq [k_2/2]$  if necessary and we can assume

$$f(x, y_2) = y_2^2 (y_2 + (\alpha_3 - \alpha_1)x^2)^2 (y_2 + (\alpha_5 - \alpha_1)x^2) + \beta x^{k_2 + 6} + \text{(higher terms)}$$

and its Newton boundary consists of two faces  $\Delta_1$  and  $\Delta_2$  and f is non-degenerate on  $\Delta_2$ but degenerate on  $\Delta_1$ . (Here "higher terms" are linear combinations of monomials above the Newton boundary.) Let  $P_1 = {}^t(1,2)$  and  $P_2$  be the weight vectors corresponding to  $\Delta_1, \Delta_2$  respectively. The Cone $(E_1, P_1)$  needs one vertex  $T_1 = {}^t(1,1)$  to be regular. For the cones Cone $(P_1, P_2)$  and Cone $(P_2, E_2)$ , the subdivision changes for  $k_2$  being even or odd.

- For  $k_2 = 2m + 1$ ,  $P_2 = {}^t(2, 2m + 1)$  and  $\operatorname{Cone}(P_1, P_2)$  and  $\operatorname{Cone}(P_2, E_2)$  are subdivided into regular fans by adding vertices  $\{T_i = {}^t(1, i), 3 \leq i \leq m\}$  and  $S = {}^t(1, m + 1)$ .

- For  $k_2 = 2m$ ,  $P_2 = t(1,m)$  and  $\operatorname{Cone}(P_1, P_2)$  is subdivided into a regular fan by adding vertices  $\{T_i = t(1,i), 3 \leq i \leq m-1\}$ . The  $\operatorname{Cone}(P_2, E_2)$  is already regular. Note that in any case, the corresponding resolution is minimal. In the second case,  $\hat{E}(P_2)^2 = -1$  but it intersects with two components of C.



After taking a toric modification with respect to the canonical subdivision, we have  $(\tilde{C}, \xi_1) \sim B_{k_1,2}$ . Hence  $(C, O) \sim B_{k_2,2} \circ B_{2,1} \circ (B_{2,1}^2)^{B_{k_1,2}}$ .

Using canonical subdivision for the second toric modification, we see that resolution graph has three branches with center  $\hat{E}(P_1)$ : one branch with a single vertex which corresponds to  $\hat{E}(T_1)$ . The second branch corresponds to the vertices in Cone $(P_1, P_2)$  and

 $\operatorname{Cone}(P_2, E_2)$  (respectively  $\operatorname{Cone}(P_1, P_2)$ ) for  $k_2$  is odd (resp. even). The third branch corresponds to the vertices for the second toric modification.

To see the relation between the second and third branches, we take change of coordinates  $(x, y_3) = (x, y_2 + (\alpha_3 - \alpha_1)x^2)$  from the beginning. After a finite number of triangular changes of coordinates, we arrive to a expression  $(C, O) \sim B_{k'_2, 2} \circ B_{2,1} \circ (B^2_{2,1})^{B_{k'_1, 2}}$ . By an easy calculation and the minimality of the resolution graph so obtained, we see that  $k'_1 = k_2 - 4, k'_2 = k_1 + 4$ . Thus  $B_{k_2, 2} \circ B_{2, 1} \circ (B^2_{2, 1})^{B_{k_1, 2}} \sim B_{k_1 + 4, 2} \circ B_{2, 1} \circ (B^2_{2, 1})^{B_{k_2 - 4, 2}}$ . Therefore in the classification, we can assume that  $k_1 + 4 \ge k_2$ .

Next we consider the case  $\iota \geq 6$ .

**Lemma 6.** Under case III-b, we assume that  $\iota = 6$ . Then the topological type of (C, O) is generically  $B_{9,2} \circ B_{6,3}$  and it can degenerate into  $(B_{5,2}^2)^{B_{1,2}} \circ B_{2,1}$  or  $B_{9,2} \circ B_{2,1} \circ (B_{2,1}^2)^{B_{k,2}}$ ,  $(k = 1, \dots, 8)$ .

*Proof.* We take a local coordinates system  $(x, y_1)$  so that  $C_2$  is defined by  $y_1 = 0$  and  $f_5(x, y_1) = \beta_{12}xy_1^2 + \beta_{31}x^3y_1 + \beta_{60}x^6 + (\text{higher terms})$ . Then  $f(x, y_1)$  is written as

$$f(x, y_1) = y_1^5 + (\beta_{12}xy_1^2 + \beta_{31}x^3y_1 + \beta_{60}x^6)^2 + \text{(higher terms)}$$
  
=  $y_1^5 + (\beta_{12}xy_1^2 + \beta_{31}xy_1^3)^2 + (\beta_{31}x^3y_1 + \beta_{60}x^6)^2 - \beta_{31}^2x^6y_1^2 + \text{(higher terms)}.$ 

If  $\beta_{31} \neq 0$ ,  $\Gamma(f; x, y_1)$  consists of two faces  $\Delta_1$ ,  $\Delta_2$  and  $f_{\Delta_1}(x, y_1) = y_1^5 + x^2 y_1^2 (\beta_{12} y_1 + \beta_{31} x^2)^2$  and  $f_{\Delta_2}(x, y) = (\beta_{31} x^3 y_1 + \beta_{60} x^6)^2$ . In this case, we first take a triangular change of coordinates of type  $(x, y_2) = (x, y_1 + c_3 x^3 + c_4 x^4)$  so that the face  $\Delta$  changes into a non-degenerate face  $\Delta'_2$  (new face after change of the coordinate) and

$$f_{\Delta_2'}(x, y_2) = \beta_{31} x^6 (y_2^2 + c_3' x^9), \ c_3 \neq 0.$$

If f is non-degenerate on  $\Delta_1$ , we have  $(C, O) \sim B_{9,2} \circ B_{6,3}$ . If f is degenerate on  $\Delta_1$   $(\beta_{31} = 4\beta_{12}^3/27, \beta_{31} \neq 0)$ , then  $f_{\Delta_1}(x, y_1) = \alpha^2 y_1^2 (9y_1 + \beta_{12}^2 x^2)(9y_1 + 4\beta_{12}^2 x^2)^2$  where  $\alpha$  is a non-zero constant. To analyze the singularity on the face  $\Delta_1$ , we take a toric modification: let  $P = {}^t(1, 2)$  be the weight vector corresponding to  $\Delta_1$  and we take a toric modification with respect to an admissible regular simplicial cone subdivision  $\Sigma^*, \pi : X \to \mathbb{C}^2$ . We may assume that  $\sigma = \operatorname{Cone}(P_1, T_1)$  is a cone in  $\Sigma^*$  where  $T_1 = {}^t(1, 3)$ . We take the toric coordinates (u, v) of the chart  $\mathbb{C}^2_{\sigma}$ . Then we have  $\pi_{\sigma}(u, v) = (uv, u^2 v^3)$  and

$$\pi_{\sigma}^* f(u,v) = \alpha^2 u^{10} v^{12} \tilde{f}(u,v) = \alpha^2 u^{10} v^{12} \Big( (9v + \beta_{12}^2)(9v + 4\beta_{12}^2)^2 + (\text{higher terms}) \Big)$$

and the strict transform  $\tilde{C}$  splits into two components. We see that one of the components of  $\tilde{C}$  which correspond to the non-degenerate component of  $f_{\Delta_1}$  is smooth and transversely with  $\hat{E}(P) = \{u = 0\}$ . To see the other component of  $\tilde{C}$ , we take the translated toric coordinates  $(u, v_1), v_1 = 9v + 4\beta_{12}^2$ . Then  $\tilde{f}(u, v_1) = cv_1^2 + \gamma_1 u + (\text{higher terms})$  where c is a non-zero constant. Hence if  $\gamma_1 \neq 0$ , we get  $(C, O) \sim B_{9,2} \circ B_{2,1} \circ (B_{2,1}^2)^{B_{1,2}}$ . If  $\gamma_1 = 0$ , taking a triangular change of coordinates of the type  $(u, v_2) = (u, v_1 + d_1 u + \dots + c_k u^j), j = [k/2],$ we can easily see that  $(C, O) \sim B_{9,2} \circ B_{2,1} \circ (B_{2,1}^2)^{B_{k,2}}, (k = 2, \dots, 8).$ 

Next we consider the case  $\beta_{31} = 0$ . Then

$$\mathcal{N}(f; x, y_1) = y_1^5 + x^2 (\beta_{12} y_1^2 + \beta_{60} x^5)^2$$

where  $\beta_{60} \neq 0$  since  $\iota = 6$  and  $\Gamma(f; x, y_1)$  has two faces  $\Delta_1$  and  $\Delta_2$  and the corresponding face functions are given by  $f_{\Delta_1}(x, y_1) = y_1^5 + \beta_{12}^2 x^2 y_1^4$  and  $f_{\Delta_2}(x, y_1) = x^2 (\beta_{12} y_1^2 + \beta_{60} x^5)^2$ .

Thus f is non-degenerate on  $\Delta_1$  and degenerate on  $\Delta_2$ . Then taking a toric modification which is the same as (4) of Proposition 3, we get  $(C, O) \sim (B_{5,2}^2)^{B_{1,2}} \circ B_{2,1}$ .

For the remaining cases  $\iota \ge 7$ , we can carry out the classification in the exact same way. So we can summarize the result as follows.

**Proposition 6.** Suppose that  $C_2$  is smooth,  $m_5 = 3$  and the tangent cone of the quintic  $C_5$  consists of a double line  $L_1$  and a single line  $L_2$ .

If  $\iota = 3$ , we have  $(C, O) \sim B_{6,5}$ .

If  $\iota \geq 4$ ,  $C_2$  is tangent to either  $L_1$  or  $L_2$  and we have the following possibility.

(I) Assume that the common tangent cone is  $L_2$ .

Then the germ (C, O) can be of type  $B_{3,4} \circ B_{2,5\iota-14}$  for  $4 \le \iota \le 10$ . (See Lemma 4). (II) Assume that the common tangent cone is  $L_1$ .

Then the germ (C, O) can be of type  $B_{2\iota,5}$  for  $\iota = 4, 5$  and  $B_{5\iota-21,2} \circ B_{6,3}$  for  $\iota = 6, \cdots, 10$ . Further degenerations are given for fixed  $\iota$  by the following list.

(1) If  $\iota = 5$ , we have (a)  $(C, O) \sim B_{k,2} \circ B_{6,3}$   $(5 \le k \le 12)$ (b)  $(C, O) \sim B_{k,3} \circ B_{4,2}$   $(k = 7, \dots, 11), B_{3,1} \circ B_{5,2} \circ B_{4,2}, B_{3,1} \circ B_{7,2} \circ B_{4,2}$  and  $B_{k,2} \circ B_{3,1} \circ B_{4,2} \ (k = 7, 8, 9).$ (c)  $(C, O) \sim B_{k_2+4,2} \circ B_{2,1} \circ (B_{2,1}^2)^{B_{k_1,2}}$ where  $(k_1, k_2)$  is in  $\{(k_1, k_2); 13 - k_2 \ge k_1 \ge k_2 - 4, k_2 = 5, \dots, 7\} \cup \{(4, 8), (5, 9)\}.$ (2) If  $\iota = 6$ , we have (a)  $(C, O) \sim (B_{5,2}^2)^{B_{1,2}} \circ B_{2,1}$ (b)  $(C, O) \sim B_{9,2} \circ B_{2,1} \circ (B_{2,1}^2)^{B_{k,2}}$   $(k = 1, \dots, 8)$ (3) If  $\iota = 7$ , we have (a)  $(C, O) \sim (B_{6,2}^2)^{2B_{1,2}} \circ B_{2,1}$  and  $B_{13,4} \circ B_{2,1}$ . (b)  $(C,O) \sim B_{14,2} \circ B_{2,1} \circ (B_{2,1}^2)^{B_{k,2}}$   $(k = 1, \dots, 7)$ (4) If  $\iota = 8$ , we have (a)  $(C,O) \sim (B_{6,2}^2)^{B_{6,2}+B_{1,2}} \circ B_{2,1}$  and  $(B_{7,2}^2)^{B_{3,2}} \circ B_{2,1}$ (b)  $(C,O) \sim B_{19,2} \circ B_{2,1} \circ (B_{2,1}^2)^{B_{k,2}}$   $(k = 1, \dots, 6)$ (5) If  $\iota = 9$ , we have (a)  $(C,O) \sim (B_{6,2}^2)^{B_{11,2}+B_{1,2}} \circ B_{2,1}, (B_{8,2}^2)^{2B_{2,2}} \circ B_{2,1}, B_{18,4} \circ B_{2,1} and B_{11,2} \circ B_{2,1}$  $B_{9,2} \circ B_{2,1}$ (b)  $(C, O) \sim B_{24,2} \circ B_{2,1} \circ (B_{2,1}^2)^{B_{k,2}}, \ (k = 1, \dots, 4)$ (6) If  $\iota = 10$ , we have (a)  $(C, O) \sim (B_{6,2}^2)^{B_{16,2}+B_{1,2}} \circ B_{2,1}, (B_{8,2}^2)^{B_{7,2}+B_{2,2}} \circ B_{2,1} and (B_{9,2}^2)^{B_{5,2}} \circ B_{2,1}$ (b)  $(C, O) \sim B_{29,2} \circ B_{2,1} \circ (B_{2,1}^2)^{B_{k,2}}, \ (k = 1, \dots, 5, \ k \neq 4)$ (b-2) Assume that  $C_2$  is a union of two lines passing through the origin  $(m_2 = 2)$  and the

(b-2) Assume that  $C_2$  is a union of two lines passing through the origin  $(m_2 = 2)$  and the tangent cone of  $C_5$  consists of a double line  $\{y = 0\}$  and a single line is  $\{x = 0\}$ . So we assume that  $f_5(x, y) = b_{12}xy^2 + b_{40}x^4 + b_{04}y^4 + (\text{higher terms})$ . We assume two lines of  $C_2$  are defined by  $\ell_1 : y + \alpha_1 x = 0$ ,  $\ell_2 : \alpha_2 y + x = 0$  and we put  $\iota_i = I(\ell_i, C_5; O) \ge 3$ (i = 1, 2). Then we have  $\iota = \iota_1 + \iota_2 \ge 6$ . If  $\iota = 6$ , we have  $(\iota_1, \iota_2) = (3, 3)$   $(\alpha_1, \alpha_2 \ne 0)$ . If  $\iota \ge 7$ , we have several possibility of  $(\iota_1, \iota_2)$ .

$$(3,3) \xrightarrow{(4,3)} (5,3) \xrightarrow{(5,4)} (5,4) \xrightarrow{(3,4)} (3,4) \xrightarrow{(4,4)} (4,4) \xrightarrow{(5,4)} (5,5)$$

Above diagram depend only the numbers  $(\alpha_1, \alpha_2, b_{40}, b_{04})$ .

**Proposition 7.** Suppose that  $C_2$  is a union of two lines and the tangent cone of the quintic  $C_5$  consists of a double line and a single line. Then we have the following possibilities.

(1) If 
$$\iota = 6$$
, then we have  
(a)  $(C, O) \sim (B_{3,2}^2)^{B_{4,2}} \circ B_{2,6}$ ,  
(b)  $(C, O) \sim B_{8,4} \circ B_{2,6}$  and  $B_{k,2} \circ B_{4,2} \circ B_{2,6}$  ( $5 \le k \le 12$ ).  
(2) If  $\iota = 7$ , then we have two cases:  $(\iota_1, \iota_2) = (4, 3)$  or  $(3, 4)$ .  
(a) If  $(\iota_1, \iota_2) = (4, 3)$ , we have  
(i)  $(C, O) \sim (B_{3,2}^2)^{B_{4,2}} \circ B_{2,11}$ ,  
(ii)  $(C, O) \sim (B_{3,4}^2)^{B_{4,2}} \circ B_{2,11}$ ,  
(ii)  $(C, O) \sim B_{8,4} \circ B_{2,11}$  and  $B_{k,2} \circ B_{4,2} \circ B_{2,11}$  ( $5 \le k \le 10$ ).  
(b) If  $(\iota_1, \iota_2) = (3, 4)$ , we have  $(C, O) \sim (B_{3,2}^2)^{B_{9,2}} \circ B_{2,6}$ .  
(3) If  $\iota = 8$ , then we have two cases:  $(\iota_1, \iota_2) = (5, 3)$  or  $(4, 4)$  or  $(3, 5)$ .  
(a) If  $(\iota_1, \iota_2) = (5, 3)$ , we have  $(C, O) \sim (B_{3,2}^2)^{B_{9,2}} \circ B_{2,16}$ .  
(b) If  $(\iota_1, \iota_2) = (4, 4)$ , we have  $(C, O) \sim (B_{3,2}^2)^{B_{9,2}} \circ B_{2,11}$ .  
(c) If  $(\iota_1, \iota_2) = (3, 5)$ , we have  
(i)  $(C, O) \sim (B_{2,2}^2)^{B_{6,2}} \circ B_{2,6}$ ,  
(ii)  $(C, O) \sim (B_{3,2}^2)^{B_{6,2}} \circ B_{2,6}$ ,  
(ii)  $(C, O) \sim (B_{3,2}^2)^{B_{6,2}} \circ B_{2,6}$ ,  
(b) If  $(\iota_1, \iota_2) = (5, 4)$ , we have  $(C, O) \sim (B_{3,2}^2)^{B_{9,2}} \circ B_{2,16}$ .  
(c) If  $(\iota_1, \iota_2) = (5, 4)$ , we have  $(C, O) \sim (B_{3,2}^2)^{B_{9,2}} \circ B_{2,16}$ .  
(b) If  $(\iota_1, \iota_2) = (4, 5)$ , we have  
(c)  $(O, O) \sim (B_{3,2}^2)^{2B_{5,2}} \circ B_{2,11}$ ,  
(ii)  $(C, O) \sim (B_{3,2}^2)^{2B_{5,2}} \circ B_{2,11}$ ,  
(b)  $(C, O) \sim (B_{3,2}^2)^{2B_{5,2}} \circ B_{2,16}$ ,  
(c)  $(C, O) \sim (B_{3,2}^2)^{2B_{5,2}} \circ B_{2,16}$ ,  
(c)  $(C, O) \sim (B_{3,2}^2)^{2B_{5,2}} \circ B_{2,16}$ ,  
(d)  $(C, O) \sim (B_{3,2}^2)^{2B_{5,2}} \circ B_{2,16}$ ,  
(e)  $(C, O) \sim (B_{3,2}^2)^{2B_{5,2}} \circ B_{2,16}$ ,  
(f)  $(C, O) \sim (B_{3,2}^2)^{2B_{5,2}} \circ B_{2,16}$ ,  
(g)  $(C, O) \sim (B_{3,2}^2)^{2B_{5,2}} \circ B_{2,16}$ ,  
(h)  $(C, O) \sim (B_{3,2}^2)^{B_{6,2}} \circ B_{2,16}$ ,  $(B_{6,2}^2)^{2B_{1,2}} \circ B_{2,16}$  and  $B_{13,4} \circ B_{2,16}$ .

We omit the proof as it is parallel to that of Proposition 3.

4.3.3. Case III-(c). In this case, we may assume that the tangent cone of  $(C_5, O)$  is given by  $y^3 = 0$ .

(c-1) Assume that  $m_2 = 1$ . If  $\iota = 3$ , we have  $(C, O) \sim B_{6,5}$  by Lemma 2. Therefore we consider the case  $\iota \ge 4$ . If  $\iota = 4$ , we have  $(C, O) \sim B_{8,5}$  as in case (b-1). If  $\iota \ge 5$ , we get the following possibilities.

**Proposition 8.** Suppose that  $C_2$  is smooth,  $m_5 = 3$  and the tangent cone of  $C_5$  is a line with multiplicity 3. Then the germ (C, O) can be of type  $B_{2\iota,5}$  for  $\iota = 3, 4$  and if  $\iota \geq 5$ , we have the following possibilities.

- (1) If  $\iota = 5$ , we have  $(C, O) \sim B_{5,10}$  and  $B_{k,2} \circ B_{6,3}$   $(k = 5, \dots, 12)$ .
- (2) If  $\iota = 6$ , we have  $(C, O) \sim B_{9,2} \circ B_{6,3}$  and  $B_{12,5}$ .
- (3) If  $\iota = 7$ , we have  $(C, O) \sim B_{14,2} \circ B_{6,3}$ ,  $B_{7,2} \circ B_{8,3}$  and  $B_{14,5}$ .
- (4) If  $\iota = 8$ , we have  $(C, O) \sim B_{19,2} \circ B_{6,3}$ ,  $B_{12,2} \circ B_{8,3}$  and  $B_{16,5}$ .

- (5) If  $\iota = 9$ , we have  $(C, O) \sim B_{24,2} \circ B_{6,3}$ ,  $B_{17,2} \circ B_{8,3}$ ,  $B_{10,2} \circ B_{10,3}$  and  $B_{18,5}$ .
- (6) If  $\iota = 10$ , we have  $(C, O) \sim B_{29,2} \circ B_{6,3}$ ,  $B_{22,2} \circ B_{8,3}$ ,  $B_{15,2} \circ B_{10,3}$  and  $B_{20,5}$ .

We omit the proof as it is parallel to Lemma 5 and Lemma 6.

(c-2) Assume that  $C_2$  is two lines passing through the origin and the tangent cone of quintic  $C_5$  is defined by  $\{y^3 = 0\}$ . Then we have  $\iota \ge 6$  and we can list the possibilities as in the following proposition.

**Proposition 9.** Suppose that  $C_2$  is two lines passing through the origin and the tangent cone of quintic  $C_5$  is defined by  $\{y^3 = 0\}$ . Then we have:

- (1) If  $\iota = 6$ , we have  $(C, O) \sim (B_{4,3}^2)^{B_{6,2}}$ ,  $B_{4,2} \circ (B_{3,2}^2)^{B_{2,2}}$ ,  $B_{10,6}$  or  $B_{6,3} \circ B_{5,3}$ .
- (2) If  $\iota = 7$  we have  $(C, O) \sim (B_{4.3}^2)^{B_{11,2}}$ .
- (3) If  $\iota = 8$  we have  $(C, O) \sim B_{9,2} \circ (B_{3,2}^2)^{B_{7,2}}$  or  $(C, O) \sim (B_{5,3}^2)^{B_{10,2}}$ .

The proof is parallel to the previous computations.

4.4. Case IV :  $m_5 = 4$ . We divide Case IV into five subcases by type of tangent cone  $T_OC_5$  of  $C_5$ .

- (a)  $T_O C_5$  consists of distinct four lines.
- (b)  $T_O C_5$  consists of a double line and distinct two lines.
- (c)  $T_O C_5$  consists of a triple line and a simple line.
- (d)  $T_O C_5$  consists of a line with multiplicity 4.
- (e)  $T_O C_5$  consists of two double lines.

First we remark that if  $C_2$  is smooth and  $C_2$  intersects transversely with  $C_5$  at the origin  $(\iota = 4), (C, O) \sim B_{8,5}$  by Lemma 2. So hereafter, we consider the case  $C_2$  and the tangent cone of  $C_5$  does not intersect transversely. First we prepare the following Lemma.

**Lemma 7.** Suppose that the conic  $C_2$  is smooth and let  $(x, y_1)$  be a local coordinate system so that  $C_2$  is defined by  $y_1 = 0$ . We put  $f_5(x, y_1) = y_1(y_1 + c_1x)(y_1 + c_2x)(y_1 + c_3x) +$ (higher terms). Then

- (1) If  $c_i \neq 0$  for i = 1, 2, 3, then  $(C, O) \sim B_{2\iota,5}$  for  $\iota = 4, 5$  and if  $\iota \geq 5$ , we have two series  $B_{k,2} \circ B_{6,3}$ ,  $5 \leq k \leq 10$  for  $\iota = 5$  and  $B_{5\iota-21,2} \circ B_{6,3}$  for  $\iota = 6, \cdots, 10$ .
- (2) If  $c_1 = 0$  and  $c_i \neq 0$  for i = 2, 3, we have  $(C, O) \sim B_{2\iota,5}$  for  $\iota = 4, 5, 6$  and  $(C, O) \sim B_{5\iota-28,2} \circ B_{8,3}$  for  $\iota = 7, \cdots, 10$ .
- (3) If  $c_1 = c_2 = 0$  and  $c_3 \neq 0$ , we have  $(C, O) \sim B_{2\iota,5}$  for  $\iota = 4, \dots, 8$  and  $(C, O) \sim B_{5\iota-35,2} \circ B_{10,3}$  for  $\iota = 9, 10$ .
- (4) If  $c_i = 0$ , (i = 1, 2, 3), we have  $(C, O) \sim B_{2\iota, 5}$  for  $\iota = 4, \cdots, 10$ .

*Proof.* We observe to the Newton boundary  $\mathcal{N}(f, x, y_1)$ . We have

$$\mathcal{N}(f, x, y_1) = y_1^5 + c^2 x^6 y_1^2 + 2c\beta_{50} x^8 y_1 + \beta_{50}^2 x^{10}, \quad c := c_1 c_2 c_3$$

and  $\Gamma(f; x, y_1)$  consists of one face  $\Delta$  with weight vector  ${}^t(1, 2)$ . We can see that the discriminant R of the face function  $f_{\Delta}(x, y_1)$  can be written as  $R = \beta_{50} \alpha$  where  $\alpha$  is a polynomial of c and  $\beta_{50}$ . Then we have  $(C, O) \sim B_{10,5}$  if  $R \neq 0$ ,  $\iota = 5$  ( $\iota = 5$  if and only if  $\beta_{50} \neq 0$ ). We observe that  $R \neq 0$ , if c = 0 and  $\beta_{50} \neq 0$ .

We first consider the case (1):  $c_i \neq 0$ , i = 1, 2, 3. Note that  $\alpha = 0$  and  $\beta_{50} = 0$  is impossible. If  $\beta_{50}\alpha \neq 0$ , we have  $\iota = 5$  and  $(C, O) \sim B_{10,5}$  as is observed above. Thus we consider the case  $\alpha = 0$  or  $\beta_{50} = 0$ . In both cases, by taking a suitable triangular change of coordinates, we can get non-degenerate singularity. If  $\alpha = 0$ , we have  $(C, O) \sim B_{k,2} \circ B_{6,3}$ ,

 $5 \leq k \leq 10$ . If  $\beta_{50} = 0$ , we have  $B_{5\iota-21,2} \circ B_{6,3}$  for  $\iota = 6, \cdots, 10$ . Hence we have assertion (1).

To consider the cases (2) ~ (4), we may assume  $\beta_{50} = 0$ . Then we can write

$$f_5(x, y_1) = y_1^4 + (c_2 + c_3)xy_1^3 + c_2c_3x^2y_1^2 + \beta_{41}x^4y_1 + \beta_{60}x^6 + (\text{higher terms})$$

and we have  $\mathcal{N}(f; x, y_1) = y_1^5 + \beta_{60}^2 x^{12}$  and note that  $\beta_{60} \neq 0$  if and only if  $\iota = 6$ . If  $\beta_{60} \neq 0$ , we have  $(C, O) \sim B_{12,5}$  and if  $\beta_{60} = 0$ , we have  $\iota \geq 7$ . Secondly we consider that (2):  $c_1 = 0, c_i \neq 0$ , (i = 2, 3). Then

 $f_5(x,y_1) = y_1^4 + (c_2 + c_3)xy_1^3 + c_2c_3x^2y_1^2 + \beta_{41}x^4y_1 + \beta_{70}x^7 + (\text{higher terms})$ 

and we have  $\mathcal{N}(f; x, y_1) = y_1^5 + x^8 (\beta_{41}y_1 + \beta_{70}x^3)^2$ . The assertion (2) follows easily by the Newton boundary argument. For the assertions (3) and (4), we can consider similarly.  $\Box$ 

4.4.1. Case IV(a). Now we classify the singularities in case IV. In this case, we have the following.

**Proposition 10.** Suppose that the tangent cone of the quintic  $C_5$  consists of distinct four lines.

- (1) If the conic  $C_2$  is smooth,  $(C, O) \sim B_{2\iota,5}$  for  $\iota = 4$ ,  $B_{2\iota,5}$  or  $B_{k,2} \circ B_{6,3}$   $(5 \le k \le 10)$ , for  $\iota = 5$  and  $B_{5\iota-21,2} \circ B_{6,3}$  for  $\iota = 6, \cdots, 10$ .
- (2) Assume that the conic  $C_2$  is a union of two lines  $\ell_1$ ,  $\ell_2$ . Putting  $\iota_i = I(\ell_i, C_5; O)$ (i = 1, 2), we have  $(C, O) \sim B_{5\iota_1 - 16, 2} \circ (B_{2, 2}^2)^{2B_{2, 2}} \circ B_{2, 5\iota_2 - 16}$  with  $\iota_1 + \iota_2 = \iota$ .

*Proof.* The assertion (1) immediately follows from Lemma 7. The assertion (2) follows from the Lemma 3.  $\Box$ 

4.4.2. Case VI-(b). In this case, we denote components of the tangent tangent cone of  $C_5$  by  $L_1$  which is a double line  $L_2$  and  $L_3$ .

(b-1) Assume that  $C_2$  is smooth  $(m_2 = 1)$ . Then we have the following.

**Proposition 11.** Suppose that  $C_2$  is smooth and the tangent cone of the quintic  $C_5$  consists of a double line and distinct two line. The germ  $(C, O) \sim B_{8,5}$  for  $\iota = 4$ . If  $\iota \ge 5$ , we have possibility of (C, O).

- (1) We assume that  $C_2$  is tangent to  $L_2$  or  $L_3$ . Then  $(C, O) \sim B_{5,10}$  or  $(C, O) \sim B_{3,6} \circ B_{2,k}$ ,  $(5 \le k \le 10)$  for  $\iota = 5$  and  $B_{3,6} \circ B_{2,5\iota-21}$  for  $\iota = 6, \cdots, 10$ .
- (2) We assume that  $C_2$  is tangent to  $L_1$ . Then  $(C,O) \sim B_{2\iota,5}$  for  $\iota = 5,6$  and  $(C,O) \sim B_{5\iota-28,2} \circ B_{8,3}$  for  $\iota = 7, \cdots, 10$ .

*Proof.* The assertions (1) and (2) are immediate by Lemma 4 and Lemma 7.

(b-2) Assume that  $C_2$  is a union of two lines passing through the origin  $\ell_i : a_i x + b_i y$ , (i = 1, 2) and we assume that  $L_1 = \{y = 0\}, L_2 = \{x = 0\}$  and  $L_3 = \{y + cx = 0\}$ .

**Proposition 12.** Suppose that  $C_2$  is a union of two lines and the tangent cone of the quintic  $C_5$  consists of a double line and distinct two line. Then

$$\begin{split} \iota &= 8: \quad (C,O) \sim \qquad B_{6,4} \circ (B_{1,1}^{2})^{B_{2,2}} \circ B_{2,4}, \qquad B_{4,2} \circ B_{3,2} \circ (B_{1,1}^{2})^{B_{2,2}} \circ B_{2,4}, \\ \iota &= 9: \quad (C,O) \sim \qquad (B_{3,2}^{2})^{B_{5,2}} \circ (B_{1,1}^{2})^{B_{2,2}} \circ B_{2,4}, \qquad \ell_1 = L_1 \quad or \\ B_{6,4} \circ (B_{1,1}^{2})^{B_{2,2}} \circ B_{2,9}, \qquad B_{4,2} \circ B_{3,2} \circ (B_{1,1}^{2})^{B_{2,2}} \circ B_{2,9}, \quad \ell_1 = L_2 \\ \iota &= 10: \quad (C,O) \sim \qquad (B_{3,2}^{2})^{B_{5,2}} \circ (B_{1,1}^{2})^{B_{2,2}} \circ B_{2,9}, \quad \ell_1 = L_1, \quad \ell_2 = L_2 \\ B_{6,4} \circ (B_{1,1}^{2})^{B_{7,2}} \circ B_{2,9}, \qquad B_{4,2} \circ B_{3,2} \circ (B_{1,1}^{2})^{B_{7,2}} \circ B_{2,9}, \quad \ell_1 = L_2, \quad \ell_2 = L_3 \end{split}$$

We omit the proof as it follows from an easy calculation.

4.4.3. Case  $IV_{-}(c)$ . We assume that the tangent tangent cone of  $C_5$  consists of a triple line  $L_1$  and a simple  $L_2$ .

(c-1) Assume that  $C_2$  is smooth. Then we have the following.

**Proposition 13.** Suppose that  $C_2$  is smooth and  $C_5$  be as above. Then  $(C, O) \sim B_{8,5}$  for  $\iota = 4$ .

- If  $\iota \geq 5$ , the possibilities of (C, O) are:
- (1) If  $C_2$  is tangent to  $L_2$ ,  $(C, O) \sim B_{5,10}$ , or  $B_{3,6} \circ B_{2,k}$ ,  $(5 \le k \le 10)$  for  $\iota = 5$  and  $(C, O) \sim B_{3,6} \circ B_{2,5\iota-21}$  for  $\iota = 6, \cdots, 10$ .
- (2) If  $C_2$  is tangent to  $L_1$ ,  $(C, O) \sim B_{2\iota,5}$  for  $\iota = 4, \cdots, 8$  and  $(C, O) \sim B_{5\iota-35,2} \circ B_{10,3}$ for  $\iota = 9, 10$ .

The proof of (1) and (2) is immediate from Lemma 4 and Lemma 7.

(c-2) Assume that  $C_2$  is a union two lines  $\ell_i$ , i = 1, 2 passing through the origin and put  $\ell_i : a_i x + b_i y = 0$  (i = 1, 2). We assume that  $L_1 = \{y = 0\}$  and  $L_2 = \{x = 0\}$ . We have  $\iota \geq 8$ .

**Proposition 14.** Suppose that  $C_2$  is a union two lines  $\ell_1$ ,  $\ell_2$  and the tangent cone of the quintic  $C_5$  consists of a triple line  $L_1$  and a simple line  $L_2$ . Then

- (1) If  $\iota = 8$ ,  $(C, O) \sim B_{8,6} \circ B_{2,4}$  and  $B_{5,3} \circ B_{4,3} \circ B_{2,4}$ . (2) Suppose  $\iota = 9$ . (a) If  $\ell_1 = L_1$ ,  $(C, O) \sim (B_{4,3}^2)^{B_{5,2}} \circ B_{2,4}$  and  $B_{5,3} \circ B_{4,3} \circ B_{2,9}$ .
  - (b) If  $\ell_2 = L_2$ ,  $(C, O) \sim B_{8,6} \circ B_{2,9}$ .

(3) If  $\iota = 10$ , we have  $\ell_1 = L_1$ ,  $\ell_2 = L_2$  and  $(C, O) \sim (B_{4,3}^2)^{B_{5,2}} \circ B_{2,9}$ .

4.4.4. Case VI-(d).

**Proposition 15.** Suppose that the tangent cone of the quintic  $C_5$  is line with multiplicity 4.

- (1) If  $C_2$  is smooth, we have  $(C, O) \sim B_{2\iota,5}$  for  $\iota = 4, 5, \cdots, 10$ .
- (2) Suppose  $C_2$  consists of two lines.
  - (a) If  $\iota = 8$  we have  $(C, O) \sim B_{10,8}$  and  $B_{2,1} \circ B_{4,3} \circ B_{5,4}$ .
  - (b) If  $\iota = 9$ , we have  $(C, O) \sim (B_{5,4}^2)^{B_{5,2}}$ .

Note that when  $\iota = 10, C$  is linear tours curve. See §6.

4.4.5. Case VI-(e). In this case, we may assume that each double line is  $L_1$  and  $L_2$ .

(e-1) Assume that  $C_2$  is smooth case.

**Proposition 16.** Suppose that  $C_2$  is smooth and the tangent cone of the quintic  $C_5$  consists of is two double line. The germ (C, O) can be of type  $B_{8,5}$  for  $\iota = 4$ . If  $\iota \ge 5$ , we have possibility of (C, O).

(1) If  $C_2$  is tangent to  $L_1$ ,  $(C, O) \sim B_{2\iota,5}$  for  $\iota = 5, 6$   $B_{5\iota-28,2} \circ B_{8,3}$  for  $\iota = 7, \cdots, 10$ . (2) If  $C_2$  is tangent to  $L_2$ ,  $(C, O) \sim B_{5,2\iota}$  for  $\iota = 5, 6$   $B_{3,8} \circ B_{2,5\iota-28}$  for  $\iota = 7, \cdots, 10$ .

(e-2)  $C_2$  is two lines case.

**Proposition 17.** Suppose that  $C_2$  is two lines and the tangent cone of the quintic  $C_5$  consists of is two double line. Then

- (1) If  $\iota = 8$ ,  $(C, O) \sim B_{6,4} \circ B_{4,6}$ ,  $B_{4,2} \circ B_{3,2} \circ B_{4,6}$  and  $B_{4,2} \circ B_{3,2} \circ B_{2,3} \circ B_{2,4}$ .
- (2) If  $\iota = 9$ ,  $(C, O) \sim B_{6,4} \circ (B_{2,3}^2)^{B_{5,2}}$  and  $B_{4,2} \circ B_{3,2} \circ (B_{2,3}^2)^{B_{5,2}}$ .
- (3) If  $\iota = 10$ ,  $(C, O) \sim (B_{3,2}^2)^{B_{5,2}} \circ (B_{2,3}^2)^{B_{5,2}}$
- 4.5. Case V:  $m_5 = 5$ . Similarly we also divide this case into  $C_2$  is smooth or two lines. We have  $\iota \geq 5$ .

**Proposition 18.** Suppose that the multiplicity of the quintic  $C_5$  is 5, i.e.,  $C_5$  consists of five line components. Then

- (1) If  $C_2$  is an irreducible conic,  $(C, O) \sim B_{2\iota,5}$  for  $\iota = 5, \cdots 10$ .
- (2) If  $C_2$  consists of two lines, f is a homogeneous polynomial of degree 10 and  $(C, O) \sim B_{10,10}$ , i.e., C consists of 10 line components.

# 5. LIST OF CLASSIFICATION

Now we have the following list of local classification.

**Theorem 1.** Let  $C = \{f = f_2^5 + f_5^2 = 0\}$  is a (2,5)-torus curve. We assume that  $C_2 = \{f_2 = 0\}$  is a reduced conic. ( $C_2$  is non-reduced case in next section). The topological type of (C, O) can be one of the following where  $\dagger(C, O)$  has degenerate series of (C, O).

- I. If  $C_5$  is smooth,  $(C, O) \sim B_{5\iota,2}$ ,  $\iota = 1, \cdots, 10$ . II. Assume  $C_5$  is singular.
- (II-1) Assume  $C_2$  is an irreducible conic.

ι	(C, O)
2	$B_{3,2} \circ B_{2,3},  B_{5,4}$
3	$B_{8,2} \circ B_{2,3}, \ \ \left(B_{3,2}^2\right)^{B_{3,2}}, \ B_{6,5}$
4	$B_{13,2} \circ B_{2,3},  B_{6,2} \circ B_{4,3},  B_{8,5},  \dagger (B_{4,2}^2)^{2B_{2,2}}$
5	$B_{18,2} \circ B_{2,3},  B_{11,2} \circ B_{4,3},  \dagger B_{10,5},  \dagger (B_{4,2}^2)^{B_{7,2}+B_{2,2}}$
6	$B_{23,2} \circ B_{2,3}, B_{16,2} \circ B_{4,3}, \dagger B_{9,2} \circ B_{6,3}, B_{12,5}, \dagger (B_{4,2}^2)^{B_{12,2}+B_{2,2}}$
7	$B_{28,2} \circ B_{2,3}, B_{21,2} \circ B_{4,3}, \dagger B_{14,2} \circ B_{6,3}, \dagger (B_{4,2}^2)^{B_{17,2}+B_{2,2}}, B_{7,2} \circ B_{8,3}, B_{14,5}$
8	$B_{33,2} \circ B_{2,3}, \ B_{26,2} \circ B_{4,3}, \ \dagger B_{19,2} \circ B_{6,3}, \ \dagger (B_{4,2}^2)^{B_{22,2}+B_{2,2}}, \ B_{12,2} \circ B_{8,3}, \ B_{16,5}$
9	$B_{38,2} \circ B_{2,3},  B_{31,2} \circ B_{4,3},  \dagger B_{24,2} \circ B_{6,3},  \dagger (B_{4,2}^2)^{B_{27,2}+B_{2,2}}$ $B_{17,2} \circ B_{8,3},  B_{10,2} \circ B_{10,3},  B_{18,5}$
10	$ \begin{array}{cccc} B_{43,2} \circ B_{2,3}, & B_{36,2} \circ B_{4,3}, & \dagger B_{29,2} \circ B_{6,3}, & \dagger (B_{4,2}^2)^{B_{32,2}+B_{2,2}} \\ B_{22,2} \circ B_{8,3}, & B_{15,2} \circ B_{10,3}, & B_{20,5} \end{array} $

The singularities with *†* have further degenerations as is indicated below.

$$\begin{array}{l} & (B_{4,2}^2)^{2B_{2,2}} : B_{10,4}, B_{k,2} \circ B_{2,5} \quad (6 \leq k \leq 15) \\ & \dagger \quad (B_{4,2}^2)^{B_{7,2}+B_{2,2}} : (B_{5,2}^2)^{B_{5,2}} \\ & \dagger \quad (B_{4,2}^2)^{B_{12,2}+B_{2,2}} : (B_{6,2}^2)^{2B_{3,2}}, (B_{7,2}^2)^{B_{2,2}}, B_{15,4} \\ & \dagger \quad (B_{4,2}^2)^{B_{17,2}+B_{2,2}} : (B_{6,2}^2)^{B_{13,2}+B_{3,2}}, (B_{7,2}^2)^{B_{7,2}} \\ & \dagger \quad (B_{4,2}^2)^{B_{22,2}+B_{2,2}} : (B_{6,2}^2)^{B_{13,2}+B_{3,2}}, (B_{8,2}^2)^{2B_{4,2}}, (B_{9,2}^2)^{B_{5,2}} \\ & B_{20,4}, B_{k,2} \circ B_{10,2} \quad (k = 11, 12) \\ & \dagger \quad (B_{4,2}^2)^{B_{22,2}+B_{2,2}} : (B_{6,2}^2)^{B_{13,2}+B_{3,2}}, (B_{8,2}^2)^{B_{13,2}+B_{4,2}}, (B_{9,2}^2)^{2B_{5,2}}, (B_{11,2}^2)^{B_{6,2}}, \\ & (B_{12,2}^2)^{2B_{1,2}}, B_{25,4} \\ & \dagger \quad B_{10,5} : B_{k,2} \circ B_{6,3} \quad (5 \leq k \leq 12), B_{k,3} \circ B_{4,2} \quad (7 \leq k \leq 11), B_{3,1} \circ B_{5,2} \circ B_{4,2}, \\ & B_{3,1} \circ B_{7,2} \circ B_{4,2}, B_{k,2} \circ B_{3,1} \circ B_{4,2} \quad (k = 7, 8, 9), B_{k_2+4,2} \circ B_{2,1} \circ (B_{2,1}^2)^{B_{k,2,2}}, \\ & (k_1, k_2) \in \{(k_1, k_2) \mid k_2 - 4 \leq k_1 \leq 13 - k_2, 5 \leq k_2 \leq 7\} \cup \{(4, 8), (5, 9)\}. \\ & \dagger \quad B_{14,2} \circ B_{6,3} : (B_{5,2}^2)^{B_{1,2}} \circ B_{2,1}, B_{13,2} \circ B_{2,1} \circ (B_{2,1}^2)^{B_{k,2}} \quad (1 \leq k \leq 7) \\ & \dagger \quad B_{19,2} \circ B_{6,3} : B_{12,2} \circ (B_{3,1}^2)^{B_{1,2}} \circ B_{2,1}, (B_{7,2}^2)^{B_{4,2}} \circ B_{2,1} \circ (B_{2,1}^2)^{B_{k,2}} \quad (1 \leq k \leq 7) \\ & \dagger \quad B_{19,2} \circ B_{6,3} : B_{17,2} \circ (B_{3,1}^2)^{B_{1,2}} \circ B_{2,1}, (B_{2,2}^2)^{2B_{2,2}} \circ B_{1,2}, B_{18,4} \circ B_{2,1}, \\ & B_{24,2} \circ B_{6,3} : B_{17,2} \circ (B_{3,1}^2)^{B_{1,2}} \circ B_{2,1}, (B_{2,2}^2)^{2B_{2,2}} \circ B_{1,2} \circ (B_{2,1}^2)^{B_{k,2}} \quad (1 \leq k \leq 7) \\ & \dagger \quad B_{24,2} \circ B_{6,3} : B_{17,2} \circ (B_{3,1}^2)^{B_{1,2}} \circ B_{2,1}, (B_{2,2}^2)^{2B_{2,2}} \circ B_{1,2}, B_{18,4} \circ B_{2,1}, \\ & B_{24,2} \circ B_{6,3} : B_{17,2} \circ (B_{3,1}^2)^{B_{1,2}} \circ B_{2,1}, (B_{2,2}^2)^{2B_{2,2}} \circ B_{1,2}, B_{18,4} \circ B_{2,1}, \\ & B_{24,2} \circ B_{6,3} : B_{17,2} \circ (B_{3,1}^2)^{B_{1,2}} \circ B_{2,1}, (B_{2,2}^2)^{2B_{2,2}} \circ B_{1,2}, (B_{2,2}^2)^{B_{5,2}} \circ B_{2,1}, \\ & b_{24,2} \circ B_{6,3} : B_{22,2} \circ (B_{3,1}^2)^{B_{1,2}} \circ B_{2,1}, B_{22,2} \circ (B_{4,1}^2)^{B_{2,2}} \circ B_{2$$

M. KAWASHIMA

ι	(C, O)
4	$B_{8,2} \circ B_{2,8},  \dagger (B_{3,2}^2)^{B_{8,2}}$
5	$B_{13,2} \circ B_{2,8},  \left(B_{3,2}^2\right)^{B_{13,2}}$
6	$B_{18,2} \circ B_{2,8}, B_{13,2} \circ B_{2,13}, \dagger (B_{4,2}^2)^{2B_{7,2}} \\ B_{6,2} \circ (B_{1,1}^2)^{B_{4,2}} \circ B_{2,6}, \dagger (B_{3,2}^2)^{B_{4,2}} \circ B_{2,6}, \dagger (B_{4,3}^2)^{B_{6,2}}$
7	$B_{23,2} \circ B_{2,8},  B_{18,2} \circ B_{2,13},  \dagger B_{16,2} \circ \left(B_{2,1}^2\right)^{B_{7,2}} \\ B_{11,2} \circ \left(B_{1,1}^2\right)^{B_{4,2}} \circ B_{2,6},  \left(B_{3,2}^2\right)^{B_{9,2}} \circ B_{2,6},  \dagger \left(B_{3,2}^2\right)^{B_{4,2}} \circ B_{2,11},  \left(B_{4,3}^2\right)^{B_{11,2}}$
8	$ \begin{array}{c} B_{23,2} \circ B_{2,13}, \ B_{18,2} \circ B_{2,18}, \ B_{11,2} \circ \left(B_{1,1}^2\right)^{B_{4,2}} \circ B_{2,11}, \ B_{6,4} \circ B_{6,4}, B_{10,8} \\ B_{16,2} \circ \left(B_{1,1}^2\right)^{B_{4,2}} \circ B_{2,6}, \ \left(B_{3,2}^2\right)^{B_{9,2}} \circ B_{2,11}, \ \left(B_{3,2}^2\right)^{B_{4,2}} \circ B_{2,16}, \ \dagger \left(B_{4,2}^2\right)^{2B_{5,2}} \circ B_{2,6} \\ B_{4,2} \circ \left(B_{2,2}^2\right)^{2B_{2,2}} \circ B_{2,4}, \ \dagger B_{6,4} \circ \left(B_{1,1}^2\right)^{2B_{2,2}} \circ B_{2,4}, \ \dagger B_{8,6} \circ B_{2,4}, \ B_{4,2} \circ \left(B_{3,2}^2\right)^{B_{7,2}} \end{array} $
9	$ \begin{array}{ll} B_{23,2} \circ B_{2,18}, & B_{16,2} \circ \left(B_{1,1}^2\right)^{B_{4,2}} \circ B_{2,11} \\ \left(B_{3,2}^2\right)^{B_{9,2}} \circ B_{2,16}, & \dagger \left(B_{4,2}^2\right)^{2B_{5,2}} \circ B_{2,11}, & B_{9,2} \circ \left(B_{2,2}^2\right)^{2B_{2,2}} \circ B_{2,4} \\ \dagger B_{6,4} \circ \left(B_{1,1}^2\right)^{B_{2,2}} \circ B_{2,9}, & \left(B_{3,2}^2\right)^{B_{5,2}} \circ \left(B_{1,1}^2\right)^{B_{2,2}} \circ B_{2,4}, & \dagger B_{6,4} \circ \left(B_{1,1}^2\right)^{B_{7,2}} \circ B_{2,4} \\ \dagger \left(B_{4,3}^2\right)^{B_{5,2}} \circ B_{2,4}, & \left(B_{5,4}^2\right)^{B_{5,2}}, & \dagger B_{6,4} \circ \left(B_{2,3}^2\right)^{B_{5,2}}, & B_{8,6} \circ B_{2,9} \end{array} $
10	$ \begin{array}{c} B_{23,2} \circ B_{2,23},  B_{16,2} \circ (B_{1,1}^2)^{B_{4,2}} \circ B_{2,16} \\ \dagger (B_{4,2}^2)^{2B_{5,2}} \circ B_{2,16},  B_{9,2} \circ (B_{2,2}^2)^{2B_{2,2}} \circ B_{2,9},  \dagger B_{6,4} \circ (B_{1,1}^2)^{B_{7,2}} \circ B_{2,9} \\ (B_{3,2}^2)^{B_{5,2}} \circ (B_{1,1}^2)^{B_{2,2}} \circ B_{2,9},  B_{6,4} \circ (B_{1,1}^2)^{B_{7,2}} \circ B_{2,4} \\ (B_{4,3}^2)^{B_{5,2}} \circ B_{2,9},  (B_{3,2}^2)^{B_{5,2}} \circ (B_{2,3}^2)^{B_{5,2}} \end{array} $

The singularities with  $\dagger$  have further degenerations as is indicated below.

$$\begin{array}{l} \stackrel{+}{(B_{3,2}^2)}{}^{B_{8,2}} : (B_{4,2}^2)^{2B_{2,2}}, B_{10,4}, B_{k,2} \circ B_{2,5} \ (6 \leq k \leq 15) \\ \stackrel{+}{(B_{4,2}^2)}{}^{2B_{7,2}} : (B_{5,2}^2)^{B_{10,2}}, (B_{6,2}^2)^{2B_{3,2}}, (B_{7,2}^2)^{B_{2,2}}, B_{15,4} \\ \stackrel{+}{(B_{16,2} \circ (B_{2,1}^2))}{}^{B_{7,2}} : (B_{5,2}^2)^{B_{15,2}} \\ \stackrel{+}{(B_{3,2}^2)}{}^{B_{4,2}} \circ B_{2,6}, B_{8,4} \circ B_{2,6}, B_{k,2} \circ B_{4,2} \circ B_{2,6} \ (5 \leq k \leq 12) \\ \stackrel{+}{(B_{3,2}^2)}{}^{B_{4,2}} \circ B_{2,11} : B_{8,4} \circ B_{2,11}, B_{k,2} \circ B_{4,2} \circ B_{2,11} \ (5 \leq k \leq 10) \\ \stackrel{+}{(B_{4,2}^2)}{}^{2B_{5,2}} \circ B_{2,6} : (B_{5,2}^2)^{B_{6,2}} \circ B_{2,6}, (B_{6,2}^2)^{2B_{1,2}} \circ B_{2,6}, B_{13,4} \circ B_{2,6} \\ \stackrel{+}{(B_{4,2}^2)}{}^{2B_{5,2}} \circ B_{2,11} : (B_{5,2}^2)^{B_{6,2}} \circ B_{2,11}, \ (B_{6,2}^2)^{2B_{1,2}} \circ B_{2,11}, B_{13,4} \circ B_{2,11} \\ \stackrel{+}{(B_{4,2}^2)}{}^{2B_{5,2}} \circ B_{2,16} : (B_{5,2}^2)^{B_{6,2}} \circ B_{2,16}, \ (B_{6,2}^2)^{2B_{1,2}} \circ B_{2,16}, B_{13,4} \circ B_{2,16} \\ \stackrel{+}{(B_{4,3}^2)}{}^{B_{6,2}} : B_{10,6}, B_{6,3} \circ B_{5,3} \\ \stackrel{+}{(B_{6,4} \circ (B_{1,1}^2)^{B_{7,2}} \circ B_{2,9} : B_{4,3} \circ B_{3,2} \circ (B_{1,1}^2)^{B_{7,2}} \circ B_{2,4} \\ \stackrel{+}{(B_{6,4} \circ (B_{1,1}^2)^{B_{7,2}} \circ B_{2,9} : B_{4,3} \circ B_{3,2} \circ (B_{1,1}^2)^{B_{7,2}} \circ B_{2,9} \\ \stackrel{+}{(B_{6,4} \circ (B_{1,1}^2)^{B_{7,2}} \circ B_{2,9} : B_{4,3} \circ B_{3,2} \circ (B_{1,1}^2)^{B_{7,2}} \circ B_{2,9} \\ \stackrel{+}{(B_{6,4} \circ (B_{2,1}^2)^{B_{7,2}} \circ B_{2,9} : B_{4,3} \circ B_{3,2} \circ (B_{1,1}^2)^{B_{7,2}} \circ B_{2,9} \\ \stackrel{+}{(B_{6,4} \circ (B_{2,1}^2)^{B_{7,2}} \circ B_{2,9} : B_{4,3} \circ B_{3,2} \circ (B_{1,1}^2)^{B_{7,2}} \circ B_{2,9} \\ \stackrel{+}{(B_{6,4} \circ (B_{4,6}^2) : B_{5,4} \circ B_{4,3} \circ B_{2,9} \\ \stackrel{+}{(B_{6,4} \circ (B_{4,6}^2) : B_{5,4} \circ B_{4,3} \circ B_{2,9} \\ \stackrel{+}{(B_{6,4} \circ (B_{2,3}^2)^{B_{5,2}} : B_{4,2} \circ B_{3,2} \circ B_{2,3} \circ B_{2,3} \circ B_{2,3} \circ B_{2,4} \\ \stackrel{+}{(B_{6,4} \circ (B_{2,3}^2)}^{B_{5,2}} : B_{4,2} \circ B_{3,2} \circ (B_{2,3}^2)^{B_{5,2}} \\ \end{array} \right)$$

# 6. LINEAR TORUS CURVE OF TYPE (2,5)

**Definition 3.** Let C be a torus curve of type (2,5) which has a defining polynomial f which can be written as  $f(x,y) = f_2(x,y)^5 + f_5(x,y)^2$  where deg  $f_j = j$  (j = 2,5). If  $f_2(x,y) = \ell(x,y)^2$  for some linear form  $\ell$ , then the curve C is called a linear torus curve.

Let  $\ell(x, y) = (ax + by + c)^2$ . We may assume that  $f_2(x, y) = -y^2$  by a linear change of coordinates so that f(x, y) is a product of quintic forms  $f(x, y) = (f_5(x, y) + y^5))(f_5(x, y) - y^5)$ . It is easy to observe that the inner singularities of C are on  $\{y = 0\} \cap C_5$ .

6.1. Local Classification. In this section we determine local singularity of linear torus curve type (2,5). Similarly, we divide into five cases. If  $C_5$  is smooth, we already have the singularity (C, O) form Lemma 1. Hence we consider that the multiplicity of  $C_5$  is larger than 2.

6.2. Case L-II:  $m_5 = 2$ . In this case, the tangent cone of  $C_5$  has two types and we have  $\iota = 4, 6, 8, 10$ .

# **Proposition 19.** Suppose $m_5 = 2$ .

- (1) If the tangent cone of  $C_5$  consists of two distinct lines,  $(C, O) \sim B_{5\iota-12,2} \circ B_{2,8}$  for  $\iota = 4, 6, 8, 10$ .
- (2) Assume that the tangent cone of  $C_5$  is a line with multiplicity 2.

(a) If 
$$(C_5, O) \sim B_{3,2}$$
,  
(i)  $(C, O) \sim (B_{3,2}^2)^{B_{8,2}}$  for  $\iota = 4$  and  
(ii)  $(C, O) \sim (B_{3,2}^2)^{B_{1,2,2}}$  for  $\iota = 6$ .  
(b) If  $(C_5, O) \sim B_{4,2}$ ,  
(i)  $(C, O) \sim (B_{4,2}^2)^{2B_{2,2}}$  for  $\iota = 4$ ,  
(ii)  $(C, O) \sim (B_{4,2}^2)^{B_{2,2}+B_{12,2}}$  for  $\iota = 10$ .  
(c) If  $(C_5, O) \sim B_{5,2}$ ,  
(i)  $(C, O) \sim (B_{5,2}^2)^{B_{20,2}}$  for  $\iota = 10$ .  
(c) If  $(C, O) \sim (B_{5,2}^2)^{B_{20,2}}$  for  $\iota = 8$ .  
(ii)  $(C, O) \sim (B_{5,2}^2)^{B_{30,2}}$  for  $\iota = 8$ .  
(iii)  $(C, O) \sim (B_{5,2}^2)^{B_{30,2}}$  for  $\iota = 10$ .  
(d) If  $(C_5, O) \sim B_{6,2}$ ,  $(C, O) \sim (B_{6,2}^2)^{2B_{8,2}}$  for  $\iota = 8$ .  
(e) If  $(C_5, O) \sim B_{7,2}$ ,  $(C, O) \sim (B_{7,2}^2)^{B_{12,2}}$  for  $\iota = 8$ .  
(f) If  $(C_5, O) \sim B_{8,2}$ ,  $(C, O) \sim (B_{8,2}^2)^{2B_{4,2}}$  for  $\iota = 8$ .  
(g) If  $(C_5, O) \sim B_{10,2}$ ,  $(C, O) \sim B_{20,5}$  and  $B_{k,2} \circ B_{10,3}$  (11  $\le k \le 13$ ) for  $\iota = 8$ .

*Proof.* The assertion (1) is shown by the Newton boundary argument (cf. Lemma 3). The assertion (2) is mainly computational. See that Proposition 3.  $\Box$ 

6.3. Case L-III:  $m_5 = 3$ . In this case, the tangent cone of  $C_5$  has three types and we have  $\iota = 2k, (k = 3, 4, 5)$ .

### **Proposition 20.** Suppose $m_5 = 3$ .

(1) If the tangent cone of  $C_5$  consists of three distinct lines.  $(C, O) \sim (B_{3,3}^2)^{3B_{4,2}}$  and  $B_{16,2} \circ (B_{2,2}^2)^{2B_{4,2}}$  and  $B_{26,2} \circ (B_{2,2}^2)^{2B_{4,2}}$  for  $\iota = 6, 8, 10$ .

- (2) Suppose the tangent cone of  $C_5$  is a double line and a single line.
  - (a) If  $\iota = 6$ ,  $(C, O) \sim B_{6,2} \circ (B_{2,3}^2)^{B_{4,2}}$ ,  $B_{8,4} \circ B_{2,6}$  and  $B_{k,2} \circ B_{4,2} \circ B_{2,6}$ ,  $(5 \le k \le 1)^{10}$ 10).
  - (b)  $\iota = 8$ .
    - (i) If the tangent cone of  $C_5$  is  $\{xy^2 = 0\}$ , then  $(C, O) \sim (B_{3,2}^2)^{B_{14,2}} \circ B_{2,6}$ .
    - (ii) If the tangent cone of  $C_5$  is  $\{x^2y = 0\}$ , then  $(C, O) \sim B_{16,2} \circ (B_{2,3}^2)^{B_{4,2}}$  $B_{16,2} \circ B_{4,8}$  and  $B_{16,2} \circ B_{2,4} \circ B_{2,k}$ ,  $(5 \le k \le 10)$ .
  - (c)  $\iota = 10$ .
    - (i) If the tangent cone of  $C_5$  is  $\{xy^2 = 0\}$ , then  $(C, O) \sim (B_{4,2}^2)^{2B_{10,2}} \circ B_{2,6}$ ,  $(B_{5,2}^2)^{B_{16,2}} \circ B_{2,6}, \ (B_{6,2}^2)^{2B_{6,2}} \circ B_{2,6}, \ (B_{8,2}^2)^{2B_{2,2}} \circ B_{2,6}, \ B_{18,2} \circ B_{2,6} \ and$  $B_{19,2} \circ B_{2,6}$ .
    - (ii) If the tangent cone of  $C_5$  is  $\{x^2y = 0\}$ , then  $(C, O) \sim B_{26,2} \circ (B_{2,3}^2)^{B_{4,2}}$ .  $B_{26,2} \circ B_{4,8}$  and  $B_{26,2} \circ B_{2,4} \circ B_{2,k}$ ,  $(5 \le k \le 9)$ .
- (3) Suppose the tangent cone of  $C_5$  is a line with triple line.
  - (a) If  $\iota = 6$ ,  $(C, O) \sim (B_{4,3}^2)^{B_{6,2}}$ ,  $B_{4,2} \circ (B_{3,2}^2)^{B_{2,2}}$ ,  $B_{10,6}$  and  $B_{3,6} \circ B_{5,3}$ .
  - (b) If  $\iota = 8$ ,  $(C, O) \sim (B_{4,3}^2)^{B_{16,2}}$ .
  - (c) If  $\iota = 10$ ,  $(C, O) \sim B_{4,2} \circ (B_{3,2}^2)^{B_{12,2}}$  and  $(B_{5,3}^2)^{B_{20,2}}$ .
- 6.4. Case L-IV:  $m_5 = 4$ .

**Proposition 21.** Suppose that the multiplicity of the quintic  $C_5$  is 4.

- (1) Assume the tangent cone of  $C_5$  is distinct four lines then
  - (a) If  $\iota = 8$ , then  $(C, O) \sim (B_{4\,4}^2)^{4B_{2,2}}$
  - (b) If  $\iota = 10$ , then  $(C, O) \sim B_{14,2} \circ (B_{3,3}^2)^{3B_{2,2}}$
- (2) Assume the tangent cone of C<sub>5</sub> is a double line and distinct two lines.
  (a) If ι = 8, (C, O) ~ (B<sup>2</sup><sub>2,2</sub>)<sup>2B<sub>2,2</sub> ∘ B<sub>4,6</sub> and (B<sup>2</sup><sub>2,2</sub>)<sup>2B<sub>2,2</sub> ∘ B<sub>2,3</sub> ∘ B<sub>2,4</sub>
  </sup></sup>
  - (b) Suppose  $\iota = 10$ .
    - (i) If the tangent cone of  $C_5$  is  $\{x^2y(y+cx) = 0\}$ ,  $(C,O) \sim B_{14,2} \circ$  $(B_{1,1}^2)^{B_{2,2}} \circ B_{4,6}$  and  $B_{14,2} \circ (B_{1,1}^2)^{B_{2,2}} \circ B_{2,3} \circ B_{2,4}$ .
    - (ii) If the tangent cone of  $C_5$  is  $\{xy^2(y+cx)=0\}, (C,O) \sim (B_{3,2}^2)^{B_{10,2}} \circ$  $(B_{2,2}^2)^{2B_{2,2}} \circ B_{2,4}.$
- (3) Assume the tangent cone of  $C_5$  is a triple line and another line.
  - (a) If  $\iota = 8$ , then  $(C, O) \sim B_{4,2} \circ B_{6,8}$  and  $B_{4,2} \circ B_{3,4} \circ B_{2,3} \circ B_{1,2}$ .
    - (b) If  $\iota = 10$ ,
      - (i) The tangent cone of  $C_5$  is  $\{x^3y = 0\}$ , we have  $B_{14,2} \circ B_{6,8}$  and  $B_{14,2} \circ$  $B_{3,4} \circ B_{2,3} \circ B_{1,2}$ .

(ii) The tangent cone of  $C_5$  is  $\{xy^3 = 0\}$ , we have  $(B_{4,3}^2)^{B_{10,2}} \circ B_{2,4}$ .

- (4) Assume the tangent cone of  $C_5$  is a line with multiplicity 4.
  - (a) If  $\iota = 8$ , then  $(C, O) \sim B_{8,10}$  and  $B_{4,5} \circ B_{3,4} \circ B_{1,2}$ .
    - (b) If  $\iota = 10$ , we have  $(C, O) \sim (B_{4.5}^2)^{B_{10,2}}$ .
- (5) Assume the tangent cone of  $C_5$  consists of two double lines.
  - (a) If  $\iota = 8$ , then  $(C, O) \sim (B_{2,2}^4)^{2B_{10,2}}$ ,  $(B_{1,1}^4)^{B_{10,2}} \circ B_{4,6}$  and  $(B_{1,1}^4)^{B_{10,2}} \circ B_{2,3} \circ B_{2,4}$ .
  - (b) If  $\iota = 10$ , we have  $(C, O) \sim (B_{3,2}^2)^{B_{10,2}} \circ B_{4,6}$  and  $(B_{3,2}^2)^{B_{10,2}} \circ B_{2,3} \circ B_{2,4}$ .

Putting together the above classifications, we have:

**Theorem 2.** Let C be a linear torus curve type (2,5). The (C,O) is described as follows.

	ι	(C,O)
	2	B <sub>10,2</sub>
	4	$B_{20,2},  B_{8,2} \circ B_{2,8},  \dagger (B^2_{3,2})^{B_{8,2}}$
	6	$B_{30,2},  B_{18,2} \circ B_{2,8},  \dagger (B_{3,2}^2)^{B_{18,2}} \\ \dagger B_{6,2} \circ (B_{2,3}^2)^{B_{4,2}},  \dagger (B_{4,3})^{B_{6,2}},  \dagger (B_{3,3}^2)^{3B_{4,2}}$
	8	$ \begin{array}{c} B_{40,2},  B_{28,2} \circ B_{2,8},  \dagger (B_{4,2}^2)^{2B_{12,2}},  \dagger B_{8,10},  (B_{4,3}^2)^{B_{16,2}},  (B_{4,4}^2)^{4B_{2,2}} \\ (B_{3,2}^2)^{B_{14,2}} \circ B_{2,6},  \dagger B_{16,2} \circ (B_{2,3}^2)^{B_{4,2}},  \dagger (B_{2,2}^2)^{2B_{2,2}} \circ B_{4,6} \\ \dagger B_{4,2} \circ B_{6,8},  \dagger (B_{2,2}^4)^{2B_{10,2}},  B_{16,2} \circ (B_{2,2}^2)^{2B_{4,2}} \end{array} $
	10	$B_{50,2}, B_{38,2} \circ B_{2,8}, \dagger (B_{4,2}^2)^{B_{22,2}+B_{12,2}}, (B_{4,3}^2)^{B_{10,2}} \circ B_{2,4}$ $\dagger (B_{4,2}^2)^{2B_{10,2}} \circ B_{2,6}, B_{26,2} \circ (B_{2,3}^2)^{B_{4,2}}, B_{10,10}, \dagger B_{14,2} \circ (B_{1,1}^2)^{B_{2,2}} \circ B_{4,6}$ $\dagger B_{4,2} \circ (B_{3,2}^2)^{B_{12,2}}, B_{14,2} \circ (B_{3,3}^2)^{3B_{2,2}}, (B_{4,5}^2)^{B_{10,2}}, \dagger B_{14,2} \circ B_{6,8}$ $\dagger (B_{3,2}^2)^{B_{10,2}} \circ (B_{2,2}^2)^{2B_{2,2}} \circ B_{2,4}, B_{26,2} \circ (B_{2,2}^2)^{2B_{4,2}}, (B_{4,5}^2)^{B_{10,2}}$
† † †	$(B_3^2) (B_3^2) (B_3^2) (B_2^2) (B_4^2) (B_4$	
†	$B_{6,2}$	$B_{2,3}^{B_{4,2}} : B_{8,4} \circ B_{2,6},  B_{k,2} \circ B_{4,2} \circ B_{2,6} \ (5 \le k \le 10)$
†	$B_{16}$	$B_{2,3} \circ (B_{2,3}^2)^{B_{4,2}} : B_{16,2} \circ B_{4,8},  B_{16,2} \circ B_{2,4} \circ B_{2,k} \ (5 \le k \le 10)$
†	$B_{26}$	$B_{2,3}^{2} \circ (B_{2,3}^2)^{D_{4,2}} : B_{26,2} \circ B_{4,8},  B_{26,2} \circ B_{2,4} \circ B_{2,k} \ (5 \le k \le 9)$
†	$(B_4^2)$	$(B_{2,2}^{2D_{10,2}} \circ B_{2,6} : (B_{5,2}^2)^{D_{16,2}} \circ B_{2,6}, (B_{6,2}^2)^{2D_{6,2}} \circ B_{2,6}, (B_{8,2}^2)^{2D_{2,2}} \circ B_{2,6})$
+	$B_{18}$ ( $B_4^2$	$a_{2} \circ B_{2,6},  B_{19,2} \circ B_{2,6}$ $a_{2} \circ B_{2,6},  B_{4,2} \circ (B_{2,2}^{2})^{B_{2,2}},  B_{10,6},  B_{6,3} \circ B_{5,3}$
†	$B_{42}$	$(B_{2,2}^2)^{B_{12,2}} : (B_{5,2}^2)^{B_{20,2}}$
†	$(B_2^2)$	$(3,2)^{2B_{2,2}} \circ B_{4,6} : B_{2,4} \circ B_{2,3} \circ (B_{2,2}^2)^{2B_{2,2}} \circ B_{2,3} \circ B_{2,4}$
†	$B_{14}$	$(2^{2} \circ (B_{11}^{2})^{B_{2,2}} \circ B_{4,6} : B_{14,2} \circ (B_{11}^{2})^{B_{2,2}} \circ B_{2,3} \circ B_{2,4}$
†	$B_{4,2}$	$B \circ B_{6,8}$ : $B_{4,2} \circ B_{3,4} \circ B_{2,3} \circ B_{1,2}$ .
†	$B_{14}$	$B_{2} \circ B_{6,8} : B_{14,2} \circ B_{3,4} \circ B_{2,3} \circ B_{1,2}$
T	$B_{8,1}$	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
Ť	$(B_2^4)$	$ \begin{array}{c} 2 \\ 2 \\ B_{10,2} \\ B_{10,2} \\ B_{10,2} \\ B_{11,1} \\ B_{11,1} \\ B_{10,2} \\ B_{11,1} \\ B_{11,$
†	$(B_{3}^{2})$	$_{2})^{B_{10,2}} \circ B_{4,6}  :  (B_{3,2}^2)^{B_{10,2}} \circ B_{2,3} \circ B_{2,4}$

# 7. Appendix

In this section, we give some examples of singularities which were obtained in the previous section.

**Example-Case I.** We assume that the quintic  $C_5$  is smooth at O. Then we have  $(C, O) \sim B_{5\iota,2}$ ,  $1 \leq \iota \leq 10$ . The following example

$$C: f(x,y) = (y^5 + y + x^2)^2 + (y + x^2)^5$$

corresponds to  $\iota = 10$  and  $(C, O) \sim A_{49}$  and Milnor number  $\mu = 49$ . This is due to [1].

**Example-Case II-(a).** We assume that the tangent cone of the quintic  $C_5$  consists of distinct two lines.

(a-1) 
$$C: f(x,y) = (y-x^2)^5 + (y^5 + xy - x^3)^2$$
,  $(C,O) \sim B_{43,2} \circ B_{2,3}$ ,  $\iota = 10, \ \mu = 51$ .  
(a-2)  $C: f(x,y) = x^5y^5 + (y^5 + yx + x^5)^2$ ,  $(C,O) \sim B_{23,2} \circ B_{2,23}$ ,  $\iota = 10, \ \mu = 51$ .

**Example-Case II-(b).** We assume that the tangent cone of the quintic  $C_5$  consists of a line of multiplicity 2.

(b-1) 
$$C_t: f(x,y) = (y+x^2)^5 + (-y^5 + 2xy^3 + (2x^3 + 1)y^2 + (2+t)x^2y + (1+t)x^4)^2,$$
  
 $(C_t, O) \sim (B_{4,2}^2)^{B_{32,2}+B_{2,2}}, \ \iota = 10, \ \mu = 55.$  This singularity degenerate into:  
 $C_0: f(x,y) = (y+x^2)^5 + (-y^5 + 2xy^3 + (2x^3 + 1)y^2 + 2x^2y + x^4)^2,$   
 $(C_0, O) \sim B_{25,4}, \ \iota = 10, \ \mu = 72.$ 

**(b-2)**  $C: f(x,y) = (y^2 + xy)^5 + (y^2 + x^5)^2, \quad (C,O) \sim (B_{5,2}^2)^{B_{15,2}}, \quad \iota = 7, \ \mu = 42.$ 

**Example-Case III-(a).** We assume that the tangent cone of the quintic  $C_5$  consists of distinct three lines.

(a-1) 
$$C: f(x,y) = (y+x^2)^5 + (y^5 + xy^3 + (x^3 + x)y^2 + (x^3 + x^2)y + x^4)^2$$
  
 $(C,O) \sim B_{36,2} \circ B_{4,3}, \quad \iota = 10, \quad \mu = 56.$   
(a-2)  $C: f(x,y) = x^5y^5 + (y^5 + xy^2 + x^2y + x^5)^2$   
 $(C,O) \sim B_{16,2} \circ (B_{1,1}^2)^{B_{4,2}} \circ B_{2,16}, \quad \iota = 10, \quad \mu = 57.$ 

**Example-Case III-(b).** We assume that the tangent cone of the quintic  $C_5$  consists of a double lines and a single line.

(b-1) 
$$C: f(x,y) = (y+x^2)^5 + (y^5 + y^2x - x^5)^2, (C,O) \sim B_{29,2} \circ B_{6,3}, \ \iota = 10, \ \mu = 61.$$
  
(b-2)  $C: f(x,y) = x^5y^5 + (y^5 + y^2x + x^5)^2, \ (C,O) \sim (B_{4,2}^2)^{2B_{5,2}} \circ B_{2,16}, \ \iota = 10, \ \mu = 61.$ 

**Example-Case III-(c).** We assume that the tangent cone of the quintic  $C_5$  consists of a line with multiplicity 3.

(c-1) 
$$C: f(x,y) = (y+x^2)^5 + (y^5 + y^3 + y^2x^2 + yx^3 + x^5)^2,$$
  
 $(C,O) \sim B_{29,2} \circ B_{6,3}, \quad \iota = 10, \ \mu = 61.$   
(c-2)  $C: f(x,y) = x^5y^5 + (y^3 + x^5)^2, \quad (C,O) \sim (B_{5,3}^2)^{B_{10,2}}, \quad \iota = 8, \ \mu = 55.$ 

**Example-Case IV-(a).** We assume that the tangent cone of the quintic  $C_5$  consists of distinct four lines.

(a-1) 
$$C: f(x,y) = (y^5 + y^4x + xy^3 + 2y^2x^3 + x^3y + x^5)^2 + (y^2 + y + x^2)^5,$$
  
 $(C,O) \sim B_{29,2} \circ B_{6,3}, \quad \iota = 10, \quad \mu = 61.$   
(a-2)  $C: f(x,y) = x^5y^5 + (y^5 + y^3x + yx^3 + x^5)^2,$   
 $(C,O) \sim B_{14,2} \circ (B_{2,2}^2)^{2B_{2,2}} \circ B_{2,14}, \quad \iota = 10, \quad \mu = 67.$ 

**Example-Case IV-(b).** We assume that the tangent cone of the quintic  $C_5$  consists of a double line and distinct two lines.

(b-1) 
$$C: f(x,y) = (y^5 + y^4 + x^2y^3 + x^2y^2 + x^4y)^2 + (y+x^2)^5,$$
  
 $(C,O) \sim B_{22,2} \circ B_{6,3}, \quad \iota = 10, \ \mu = 66.$   
(b-2)  $C: f(x,y) = (2y^5 + (x^2 + x)y^3 - y^2x^2 + 2x^5)^2 + (yx - x^2)^5,$   
 $(C,O) \sim B_{6,4} \circ (B_{1,1}^2)^{B_{7,2}} \circ B_{2,9}, \quad \iota = 10, \ \mu = 69.$ 

**Example-Case IV-(c).** We assume that the tangent cone of the quintic  $C_5$  consists of a triple line and a single line.

(c-1) 
$$C: f(x,y) = (2y^5 + y^4x + (x^2 + x)y^3 + 3y^2x^3)^2 + (y^2 + (x+1)y + 3x^2)^5$$
  
 $(C,O) \sim B_{15,2} \circ B_{10,3}, \quad \iota = 10, \ \mu = 71.$ 

(c-2) 
$$C: f(x,y) = (y^5 + y^3 x + x^5)^2 + y^5 x^5, (C,O) \sim (B_{4,3}^2)^{B_{5,2}} \circ B_{2,9}, \iota = 10, \mu = 71.$$

**Example-Case IV-(d).** We assume that the tangent cone of the quintic  $C_5$  consists of line with multiplicity 4.

(d-1) 
$$C: f(x,y) = (2y^5 + (x+1)y^4 + x^2y^3)^2 + (y^2 + (x+1)y + x^2)^5$$
  
(C,O) ~ B<sub>20,5</sub>,  $\iota = 10, \ \mu = 76.$ 

(d-2) 
$$C: f(x,y) = (y^4 + x^5)^2 + (y^2 + yx)^5, \quad (C,O) \sim (B_{5,4}^2)^{B_{5,2}}, \quad \iota = 9, \ \mu = 68.$$

**Example-Case IV-(e).** We assume that the tangent cone of the quintic  $C_5$  consists of two double lines.

$$C: f(x,y) = (y^5 + y^2 x^2 + x^5)^2 + y^5 x^5, \ (C,O) \sim (B^2_{3,2})^{B_{5,2}} \circ (B^2_{2,3})^{B_{5,2}}, \ \iota = 10, \ \mu = 71.$$

Acknowledgement. I would like to express my deepest gratitude to Professor Mutsuo Oka who has proposed this problem and navigated me during the preparation of this paper.

# References

- B. Audoubert, C. Nguyen, and M. Oka. On alexander polynomials of torus curves. J. Math. Soc. Japan, 57(4):935–957, 2005.
- [2] A. G. Kouchnirenko. Polyèdres de Newton et nombres de Milnor. Invent. Math., 32:1–31, 1976.
- [3] M. Oka. On the bifurcation of the multiplicity and topology of the newton boundary. J. Math. Soc. Japan, 31:435–450, 1979.
- [4] M. Oka. On the resolution of the hypersurface singularities. In Complex analytic singularities, volume 8 of Adv. Stud. Pure Math., pages 405–436. North-Holland, Amsterdam, 1987.
- [5] M. Oka. Non-degenerate complete intersection singularity. Hermann, Paris, 1997.
- M. Oka. Geometry of cuspidal sextics and their dual curves. In Singularities—Sapporo 1998, volume 29 of Adv. Stud. Pure Math., pages 245–277. Kinokuniya, Tokyo, 2000.
- [7] D. T. Pho. Classification of singularities on torus curves of type (2,3). Kodai Math. J., 24(2):259–284, 2001.

DEPARTMENT OF MATHEMATICS TOKYO UNIVERSITY OF SCIENCE WAKAMIYA-CHO 26, SHINJUKU-KU TOKYO 162-0827 E-mail:j1107702@ed.kagu.tus.ac.jp