

AN ALGEBRAIC APPROACH TO THE SET OF INTERVALS.

NICOLAS GOZE, ELISABETH REMM

ABSTRACT. In this paper we present the set of intervals as a normed vector space. We define also a four-dimensional associative algebra whose product gives the product of intervals in any cases. This approach allows to give a notion of divisibility and in some cases an euclidian division.

1. INTERVALS AND GENERALIZED INTERVALS

An interval is a connected closed subset of \mathbb{R} . The classical arithmetic operations on intervals are defined such that the result of the corresponding operation on elements belonging to operand intervals belongs to the resulting interval. That is, if \diamond denotes one of the classical operation $+$, $-$, $*$, we have

$$(1) \quad [x^-, x^+] \diamond [y^-, y^+] = \{x \diamond y / x \in [x^-, x^+], y \in [y^-, y^+]\} \quad .$$

In particular we have

$$\begin{cases} [x^-, x^+] + [y^-, y^+] = [x^- + y^-, x^+ + y^+], \\ [x^-, x^+] - [y^-, y^+] = [x^- - y^+, x^+ - y^-] \end{cases}$$

and

$$[x^-, x^+] - [x^-, x^+] = [x^- - x^+, x^+ - x^-] \neq 0.$$

Let \mathbb{IR} be the set of intervals. It is in one to one correspondance with the half plane of \mathbb{R}^2 :

$$\mathcal{P}_1 = \{(a, b), a \leq b\}.$$

This set is closed for the addition and \mathcal{P}_1 is endowed with a regular semigroup structure. Let \mathcal{P}_2 be the half plane symmetric to \mathcal{P}_1 with respect to the first bisector Δ of equation $y - x = 0$. The substraction on \mathbb{IR} , which is not the symmetric operation of $+$, corresponds to the following operation on \mathcal{P}_1 :

$$(a, b) - (c, d) = (a, b) + s_{\Delta} \circ s_0(c, d),$$

where s_0 is the symmetry with respect to 0, and s_{Δ} with respect to Δ . The multiplication $*$ is not globally defined. Consider the following subset of \mathcal{P}_1 :

$$\begin{cases} \mathcal{P}_{1,1} = \{(a, b) \in \mathcal{P}_1, a \geq 0, b \geq 0\}, \\ \mathcal{P}_{1,2} = \{(a, b) \in \mathcal{P}_1, a \leq 0, b \geq 0\}, \\ \mathcal{P}_{1,3} = \{(a, b) \in \mathcal{P}_1, a \leq 0, b \leq 0\}. \end{cases}$$

We have the following cases:

- 1) If $(a, b), (c, d) \in \mathcal{P}_{1,1}$ the product is written $(a, b) * (c, d) = (ac, bd)$.

This work is supported by 'System'Virtual Prototyping S.A.S, a scientific service company involved in 'complex systems design' and 'Functional Virtual Prototyping' (www.systemvip.com) .

Then if $e_1 = (1, 1)$ and $e_2 = (0, 1)$, these "vectors" generate $\mathcal{P}_{1,1}$:

$$\forall (x, y) \in \mathcal{P}_{1,1} \text{ then } (x, y) = xe_1 + (y - x)e_2, \quad x > 0, \quad y - x > 0.$$

The multiplication corresponds in this case to the following associative commutative algebra:

$$\begin{cases} e_1e_1 = e_1, \\ e_1e_2 = e_2e_1 = e_2e_2 = e_2. \end{cases}$$

2) Assume that $(a, b) \in \mathcal{P}_{1,1}$ and $(c, d) \in \mathcal{P}_{1,2}$ so $c \leq 0$ and $d \geq 0$. Thus we obtain $(a, b) * (c, d) = (bc, bd)$ and this product does not depend of a . Then we obtain the same result for any $a < b$. Let $e_1 = (0, 1)$ and $e_2 = (-1, 0)$. Any interval of $\mathcal{P}_{1,1}$ is written $ae_1 + be_2$ with $b < 0$ and any interval of $\mathcal{P}_{1,2}$, $ce_1 + de_2$ with $c, d > 0$. We have no associative multiplication between (e_1, e_2) which describes the product. We have to add a formal dimension to obtain a 3-dimensional associative algebra and the product appears as the projection in the plane (e_1, e_2) of this associative algebra. Here if we consider the following associative commutative algebra

$$\begin{cases} e_1e_1 = e_1, \quad e_1e_2 = e_2, \quad e_1e_3 = -e_2, \\ e_2e_3 = -e_1, \quad e_2e_2 = e_1, \\ e_3e_3 = e_3. \end{cases}$$

then $(\alpha e_1 + \beta e_2 + \gamma e_3)(\delta e_1 + \epsilon e_2) = \alpha\delta e_1 + \alpha\epsilon e_2$. As $(a, b) = be_1 - ae_2$ and $(c, d) = de_1 - ce_2$, we obtain the expected product.

3) If $(a, b) \in \mathcal{P}_{1,1}$ and $(c, d) \in \mathcal{P}_{1,3}$ then $a \geq 0, b \geq 0$ and $c \leq 0, d \leq 0$ and we have $(a, b) * (c, d) = (bd, ac)$. Let $e_1 = (1, 1)$, $e_2 = (0, 1)$. This product corresponds to the following associative algebra:

$$\begin{cases} e_1e_1 = e_1, \\ e_1e_2 = e_1 - e_2, \\ e_2e_2 = e_1 - e_2. \end{cases}$$

We have similar results for the cases $(\mathcal{P}_{1,2}, \mathcal{P}_{1,2})$, $(\mathcal{P}_{1,2}, \mathcal{P}_{1,3})$ and $(\mathcal{P}_{1,3}, \mathcal{P}_{1,3})$.

All this shows that the set \mathbb{IR} is not algebraically structured. Let us describe a vectorial structure on \mathbb{IR} using the previous geometrical interpretation of \mathbb{IR} with \mathcal{P}_1 . First we extend \mathcal{P}_1 to \mathbb{R}^2 and we obtain an extended set $\overline{\mathbb{IR}}$ which corresponds to the classical interval $[a, b]$ and "generalized intervals" $[a, b]$ with $a > b$. Of course using the addition of \mathbb{R}^2 , we obtain on $\overline{\mathbb{IR}}$ a structure of abelian group and the symmetric of $[a, b] \in \mathbb{IR}$ is $[-a, -b] \in \overline{\mathbb{IR}} \setminus \mathbb{IR}$. In this context $[a, b] + [-a, -b] = 0$. This aspect as been developed in [5].

We have a group homomorphism φ on $\overline{\mathbb{IR}}$ given by

$$\begin{aligned} \varphi : \overline{\mathbb{IR}} &\longrightarrow \overline{\mathbb{IR}} \\ (a, b) &\longrightarrow (b, a). \end{aligned}$$

This map is called dual and we denote by dual (a, b) the generalized interval (b, a) . The corresponding arithmetic has been developed by Kaucher [4] and is naturally called the Kaucher arithmetic. In the following we recall how to complete the semigroup \mathbb{IR} to obtain a natural vectorial structure on $\overline{\mathbb{IR}}$.

2. THE REAL VECTOR SPACE $\overline{\mathbb{IR}}$

2.1. **The semigroup** $(\mathbb{IR}, +)$. Consider $x = [x^-, x^+]$ and $y = [y^-, y^+]$ two elements of \mathbb{IR} . From (1) we get the addition

$$x + y = [x^- + y^-, x^+ + y^+].$$

This operation is commutative, associative and has an unit $[0, 0]$ simply denoted by 0.

Theorem 1. *The semigroup $(\mathbb{IR}, +)$ is commutative and regular.*

Proof. We recall that a semigroup is a nonempty set with an associative unitary operation $+$. It is regular if it satisfies

$$x + z = x + y \implies z = y,$$

for all x, y, z . The semigroup $(\mathbb{IR}, +)$ is regular. In fact

$$x + z = x + y \implies [x^- + z^-, x^+ + z^+] = [x^- + y^-, x^+ + y^+]$$

which gives $z^- = y^-$ and $z^+ = y^+$, that is $z = y$.

2.2. **The group** $(\overline{\mathbb{IR}}, +)$. The goal is to define a subtraction corresponding to an inverse of the addition. For that we build the symmetrized of the semigroup $(\mathbb{IR}, +)$. We consider on the set $\mathbb{IR} \times \mathbb{IR}$ the equivalence relation:

$$(x, y) \sim (z, t) \iff x + t = y + z,$$

for all $x, y, z, t \in \mathbb{IR}$. The quotient set is denoted by $\overline{\mathbb{IR}}$. The addition of intervals is compatible with this equivalence relation:

$$\overline{(x, y)} + \overline{(z, t)} = \overline{(x + z, y + t)}$$

where $\overline{(x, y)}$ is the equivalence class of (x, y) . The unit is $\overline{0} = \{(x, x), x \in \mathbb{IR}\}$ and each element $\overline{(x, y)}$ has an inverse

$$\sphericalangle \overline{(x, y)} = \overline{(y, x)}.$$

Then $(\overline{\mathbb{IR}}, +)$ is a commutative group.

For all $x = [x^-, x^+] \in \mathbb{IR}$, we denote by $l(x)$ his lenght, so $l(x) = x^+ - x^-$, and by $c(x)$ his center, so $c(x) = \frac{x^+ + x^-}{2}$.

Proposition 2. *Let $\mathcal{X} = \overline{(x, y)}$ be in $\overline{\mathbb{IR}}$. Thus*

- *if $l(y) < l(x)$, there is an unique $A \in \mathbb{IR} \setminus \mathbb{R}$ such that $\mathcal{X} = \overline{(A, 0)}$,*
- *if $l(y) > l(x)$, there is an unique $A \in \mathbb{IR} \setminus \mathbb{R}$ such that $\mathcal{X} = \overline{(0, A)} = \sphericalangle \overline{(A, 0)}$,*
- *if $l(y) = l(x)$, there is an unique $A = \alpha \in \mathbb{R}$ such that $\mathcal{X} = \overline{(\alpha, 0)} = \overline{(0, -\alpha)}$.*

Proof. It is based on the following lemmas:

Lemma 3. *Consider $\overline{(x, y)} \in \overline{\mathbb{IR}}$ with $l(x) < l(y)$. Then*

$$(x, y) \sim (0, [y^- - x^-, y^+ - x^+]).$$

Proof. For all $x, y, z, t \in \mathbb{IR}$, we have

$$(x, y) \sim (z, t) \iff \begin{cases} z^- + y^- = x^- + t^-, \\ z^+ + y^+ = x^+ + t^+. \end{cases}$$

If we put $z^- = z^+ = 0$, then

$$\begin{cases} t^- = y^- - x^-, \\ t^+ = y^+ - x^+. \end{cases}$$

with the necessary condition $t^+ > t^-$. We obtain $l(x) < l(y)$. So we have

$$\overline{(x, y)} = \overline{(0, [y^- - x^-, y^+ - x^+])}.$$

Lemma 4. Consider $\overline{(x, y)} \in \overline{\mathbb{IR}}$ with $l(y) < l(x)$, then

$$(x, y) \sim ([x^- - y^-, x^+ - y^+], 0).$$

Proof. For all $x, y, z \in \mathbb{IR}$, we have

$$(x, y) \sim (z, 0) \iff \begin{cases} z^- + y^- = x^- \\ z^+ + y^+ = x^+ \end{cases}.$$

or

$$\begin{cases} z^- = x^- - y^- \\ z^+ = x^+ - y^+ \end{cases}$$

with the condition $z^+ > z^-$ which gives $l(y) < l(x)$. So $\overline{(x, y)} = \overline{([x^- - y^-, x^+ - y^+], 0)}$.

Lemma 5. Consider $\overline{(x, y)} \in \overline{\mathbb{IR}}$ with $l(x) = l(y)$, then

$$(x, y) \sim (\alpha, 0)$$

with $\alpha = x^- - y^-$.

These three lemmas describe the three cases of Proposition 2.

Definition 6. Any element $\mathcal{X} = \overline{(A, 0)}$ with $A \in \mathbb{IR} \setminus \mathbb{R}$ is said positive and we write $\mathcal{X} > 0$. Any element $\mathcal{X} = \overline{(0, A)}$ with $A \in \mathbb{IR} \setminus \mathbb{R}$ is said negative and we write $\mathcal{X} < 0$. We write $\mathcal{X} \geq \mathcal{X}'$ if $\mathcal{X} \setminus \mathcal{X}' \geq 0$.

For example if \mathcal{X} and \mathcal{X}' are positive,

$$\mathcal{X} \geq \mathcal{X}' \iff l(\mathcal{X}) \geq l(\mathcal{X}').$$

The elements $\overline{(\alpha, 0)}$ with $\alpha \in \mathbb{R}^*$ are neither positive nor negative.

Remark. This structure of abelian group on $\overline{\mathbb{IR}}$ has yet been defined by Markov [5]. In his paper he presents the Kaucher arithmetic using the completion of the semigroup \mathbb{IR} . Up to now we have sum up this study. In the following section we will develop a topological vectorial structure. The goal is to define a good differential calculus.

2.3. Vector space structure on $\overline{\mathbb{IR}}$. We are going to construct a real vector space structure on the group $(\overline{\mathbb{IR}}, +)$. We recall that if $A = [a, b] \in \mathbb{IR}$ and $\alpha \in \mathbb{R}^+$, the product αA is the interval $[\alpha a, \alpha b]$. We consider the external multiplication:

$$\cdot : \mathbb{R} \times \overline{\mathbb{IR}} \longrightarrow \overline{\mathbb{IR}}$$

defined, for all $A \in \mathbb{IR}$, by

$$\begin{cases} \alpha \cdot \overline{(A, 0)} = \overline{(\alpha A, 0)}, \\ \alpha \cdot \overline{(0, A)} = \overline{(0, \alpha A)}, \end{cases}$$

for all $\alpha > 0$. If $\alpha < 0$ we put $\beta = -\alpha$. So we take:

$$\begin{cases} \alpha \cdot \overline{(A, 0)} = \overline{(0, \beta A)}, \\ \alpha \cdot \overline{(0, A)} = \overline{(\beta A, 0)}. \end{cases}$$

We denote $\alpha \mathcal{X}$ instead of $\alpha \cdot \mathcal{X}$.

Lemma 7. For any $\alpha \in \mathbb{R}$ and $\mathcal{X} \in \overline{\mathbb{IR}}$ we have:

$$\begin{cases} \alpha(\setminus \mathcal{X}) = \setminus(\alpha \mathcal{X}), \\ (-\alpha)\mathcal{X} = \setminus(\alpha \mathcal{X}). \end{cases}$$

Indeed, if $\alpha \geq 0$ and $\mathcal{X} = \overline{(A, 0)}$ with $A \in \mathbb{IR}$ then $\setminus \mathcal{X} = \overline{(0, A)}$ and

$$\alpha(\setminus \mathcal{X}) = \overline{(0, \alpha A)} = \setminus(\overline{(\alpha A, 0)}) = \setminus(\alpha \mathcal{X}).$$

In the same way $(-\alpha)\mathcal{X} = (-\alpha)\overline{(A, 0)} = \overline{(0, \alpha A)} = \setminus(\alpha \mathcal{X})$. If $\alpha \leq 0$ and $\mathcal{X} = \overline{(A, 0)}$ so $\alpha(\setminus \mathcal{X}) = \alpha\overline{(0, A)} = \overline{(-\alpha A, 0)}$ and $\setminus(\alpha \mathcal{X}) = \setminus(\overline{(0, -\alpha A)}) = \overline{(-\alpha A, 0)}$. The calculus is the same for $\mathcal{X} = \overline{(0, A)}$.

Proposition 8. For all $\alpha, \beta \in \mathbb{R}$, and for all $\mathcal{X}, \mathcal{X}' \in \overline{\mathbb{IR}}$, we have

$$\begin{cases} (\alpha + \beta)\mathcal{X} = \alpha\mathcal{X} + \beta\mathcal{X}, \\ \alpha(\mathcal{X} + \mathcal{X}') = \alpha\mathcal{X} + \alpha\mathcal{X}', \\ (\alpha\beta)\mathcal{X} = \alpha(\beta\mathcal{X}). \end{cases}$$

Proof. We are going to study the differents cases.

1) The result is trivial when $\alpha, \beta > 0$.

2) Suppose $\alpha > 0$, $\beta < 0$ and $\alpha + \beta > 0$. We assume $\mathcal{X} = \overline{(A, 0)}$ and $A \in \mathbb{IR}$. We put $\gamma = -\beta$. So

$$(\alpha + \beta)\mathcal{X} = \overline{((\alpha + \beta)A, 0)}$$

and

$$\alpha\mathcal{X} + \beta\mathcal{X} = \overline{(\alpha A, 0)} + \overline{(0, \gamma A)} = \overline{(\alpha A - \gamma A, 0)} = \overline{((\alpha - \gamma)A, 0)} = \overline{((\alpha + \beta)A, 0)}.$$

The calculus is the same for $\mathcal{X} = \overline{(0, A)}$. So we have $(\alpha + \beta)\mathcal{X} = \alpha\mathcal{X} + \beta\mathcal{X}$ for all $\mathcal{X} \in \overline{\mathbb{IR}}$.

3) If $\alpha > 0$, $\beta < 0$ and $\alpha + \beta < 0$, the proof is the same that the previous case.

4) If $\beta > 0$ and $\alpha < 0$, we refer to the cases 2) et 3).

5) If $\alpha, \beta < 0$, we put $\beta_1 = -\beta$ and $\alpha_1 = -\alpha$. Thus $\alpha_1, \beta_1 > 0$, and we find again the first case.

So we have the result:

Theorem 9. The triplet $(\overline{\mathbb{IR}}, +, \cdot)$ is a real vector space.

2.4. Basis and dimension. We consider the vectors $\mathcal{X}_1 = \overline{([0, 1], 0)}$ and $\mathcal{X}_2 = \overline{([1, 1], 0)}$ of $\overline{\mathbb{IR}}$.

Theorem 10. *The family $\{\mathcal{X}_1, \mathcal{X}_2\}$ is a basis of $\overline{\mathbb{IR}}$. So $\dim_{\mathbb{R}} \overline{\mathbb{IR}} = 2$.*

Proof. We have the following decompositions:

$$\begin{cases} \overline{([a, b], 0)} = (b - a)\mathcal{X}_1 + a\mathcal{X}_2, \\ \overline{(0, [c, d])} = (c - d)\mathcal{X}_1 - c\mathcal{X}_2. \end{cases}$$

The linear map

$$\varphi : \overline{\mathbb{IR}} \longrightarrow \mathbb{R}^2$$

defined by

$$\begin{cases} \varphi(\overline{([a, b], 0)}) = (b - a, a), \\ \varphi(\overline{(0, [c, d])}) = (c - d, -c) \end{cases}$$

is a linear isomorphism and $\overline{\mathbb{IR}}$ is canonically isomorphic to \mathbb{R}^2 .

Remark. Let E be the subspace generated by \mathcal{X}_2 . The vectors of E correspond to the elements which have a non defined sign. Then the relation \leq defined in the paragraph 1.2 gives an order relation on the quotient space $\overline{\mathbb{IR}}/E$.

2.5. A Banach structure on $\overline{\mathbb{IR}}$. Let us begin to define a norm on $\overline{\mathbb{IR}}$. Any element $\mathcal{X} \in \overline{\mathbb{IR}}$ is written $\overline{(A, 0)}$ or $\overline{(0, A)}$. We define its length $l(\mathcal{X})$ as the length of A and its center as $c(A)$ or $-c(A)$ in the second case.

Theorem 11. *The map $\|\cdot\| : \overline{\mathbb{IR}} \longrightarrow \mathbb{R}$ given by*

$$\|\mathcal{X}\| = l(\mathcal{X}) + |c(\mathcal{X})|$$

for any $\mathcal{X} \in \overline{\mathbb{IR}}$ is a norm.

Proof. We have to verify the following axioms:

$$\begin{cases} 1) \|\mathcal{X}\| = 0 \iff \mathcal{X} = 0, \\ 2) \forall \lambda \in \mathbb{R} \|\lambda\mathcal{X}\| = |\lambda|\|\mathcal{X}\|, \\ 3) \|\mathcal{X} + \mathcal{X}'\| \leq \|\mathcal{X}\| + \|\mathcal{X}'\|. \end{cases}$$

1) If $\|\mathcal{X}\| = 0$, then $l(\mathcal{X}) = |c(\mathcal{X})| = 0$ and $\mathcal{X} = 0$.

2) Let $\lambda \in \mathbb{R}$. We have

$$\|\lambda\mathcal{X}\| = l(\lambda\mathcal{X}) + |c(\lambda\mathcal{X})| = |\lambda|l(\mathcal{X}) + |\lambda||c(\mathcal{X})| = |\lambda|\|\mathcal{X}\|.$$

3) We consider that I refers to \mathcal{X} and J refers to \mathcal{X}' thus $\mathcal{X} = \overline{(I, 0)}$ or $\overline{(0, I)}$. We have to study the two different cases:

i) If $\mathcal{X} + \mathcal{X}' = \overline{(I + J, 0)}$ or $\overline{(0, I + J)}$, then

$$\begin{aligned} \|\mathcal{X} + \mathcal{X}'\| &= l(I + J) + |c(I + J)| = l(I) + l(J) + |c(I) + c(J)| \leq l(I) + |c(I)| + l(J) + |c(J)| \\ &= \|\mathcal{X}\| + \|\mathcal{X}'\|. \end{aligned}$$

ii) Let $\mathcal{X} + \mathcal{X}' = \overline{(I, J)}$. If $\overline{(I, J)} = \overline{(K, 0)}$ then $K + J = I$ and

$$\|\mathcal{X} + \mathcal{X}'\| = \|\overline{(K, 0)}\| = l(K) + |c(K)| = l(I) - l(J) + |c(I) - c(J)|$$

that is

$$\|\mathcal{X} + \mathcal{X}'\| \leq l(I) + |c(I)| - l(J) + |c(J)| \leq l(I) + |c(I)| + l(J) + |c(J)| = \|\mathcal{X}\| + \|\mathcal{X}'\|.$$

So we have a norm on $\overline{\mathbb{IR}}$.

Now we shall show that $\overline{\mathbb{IR}}$ is a Banach space that is all Cauchy sequences converge in $\overline{\mathbb{IR}}$.

Theorem 12. *The normed vector space $\overline{\mathbb{IR}}$ is a Banach space.*

Proof. We recall that a sequence $(\mathcal{X}_n)_{n \in \mathbb{N}}$ with $\mathcal{X}_n \in \overline{\mathbb{IR}}$ is a Cauchy sequence if

$$\forall \varepsilon > 0, \exists N \in \mathbb{N}, \forall n, m \geq N, \|\mathcal{X}_n \setminus \mathcal{X}_m\| \leq \varepsilon.$$

We have to verify that all Cauchy sequences in $\overline{\mathbb{IR}}$ converge in $\overline{\mathbb{IR}}$. Let $(\mathcal{X}_n)_{n \in \mathbb{N}}$ be a Cauchy sequence. We suppose that for all $n, m \geq N$ we have $\mathcal{X}_n = \overline{(A_n, 0)}$ and $\mathcal{X}_m = \overline{(A_m, 0)}$. So we have $\mathcal{X}_n \setminus \mathcal{X}_m = \overline{(A_n, A_m)}$ and

$$\|\mathcal{X}_n \setminus \mathcal{X}_m\| = |l(A_n) - l(A_m)| + |c(A_n) - c(A_m)| \leq \varepsilon.$$

Then for $n, m > N$, $|l(A_n) - l(A_m)| \leq \varepsilon$ and $|c(A_n) - c(A_m)| \leq \varepsilon$. The sequences $(l(A_n))_{n \in \mathbb{N}}$ and $(c(A_n))_{n \in \mathbb{N}}$ are Cauchy sequences in \mathbb{R} , which is a complete space, thus the two sequences converge. Let l and c be their limits. It exists an unique L so that $l(L) = l$ and $c(L) = c$. Thus we can thus deduce that $(\mathcal{X}_n)_{n \in \mathbb{N}}$ converges to L . It is the same if all terms of the sequence are of type $\mathcal{X}_n = \overline{(0, A_n)}$. We suppose now that the sequence $(\mathcal{X}_n)_{n \in \mathbb{N}}$ is not of constant sign from a certain rank. So

$$\exists n, m > N \text{ such that } \mathcal{X}_n = \overline{(A_n, 0)} \text{ and } \mathcal{X}_m = \overline{(0, A_m)},$$

and

$$\|\mathcal{X}_n \setminus \mathcal{X}_m\| = \|\overline{(A_n + A_m, 0)}\| = l(A_n) + l(A_m) + |c(A_n) + c(A_m)| \leq \varepsilon. \quad (*)$$

We deduce that $l(A_n) \xrightarrow[n \rightarrow +\infty]{} 0$. Moreover, we can consider the subsequence whose terms are of type $(\overline{(A_n, 0)})_{n \in \mathbb{N}}$. It converges to \mathcal{X} with $l(\mathcal{X}) = 0$ thus $\mathcal{X} = \overline{(\alpha, 0)}$. In the same way we can consider the subsequence of terms are $(\overline{(0, A_n)})$. It converges to $\mathcal{X}' = \overline{(0, \beta)}$ since $l(\mathcal{X}') = 0$. But the Inequation (*) implies $\alpha + \beta \leq \varepsilon$ for all ε thus $\mathcal{X}' = \overline{(0, -\alpha)} = \mathcal{X}$. It follows that $\overline{\mathbb{IR}}$ is complete.

3. A 4-DIMENSIONAL ASSOCIATIVE ALGEBRA ASSOCIATED TO $\overline{\mathbb{IR}}$

3.1. Classical product of intervals. We consider $X, Y \in \mathbb{IR}$. The multiplication of intervals is defined by

$$X \cdot Y = [\min(x^-y^-, x^-y^+, x^+y^-, x^+y^+), \max(x^-y^-, x^-y^+, x^+y^-, x^+y^+)].$$

Let $\mathcal{X} = \overline{(X, 0)}$ and $\mathcal{X}' = \overline{(Y, 0)}$ be in $\overline{\mathbb{IR}}$. We put

$$\mathcal{X}\mathcal{X}' = \overline{(XY, 0)}.$$

For this product we have:

Proposition 13. *For all $\mathcal{X} = \overline{(X, 0)}$ and $\mathcal{X}' = \overline{(Y, 0)}$ in $\overline{\mathbb{IR}}$, we have*

$$\|\mathcal{X}\mathcal{X}'\| \leq \|\mathcal{X}\| \|\mathcal{X}'\|.$$

Proof. In the following table, the boxes represent $\|\mathcal{X}\mathcal{X}'\|$ following the values of $\|\mathcal{X}\|$ and $\|\mathcal{X}'\|$.

If $\mathcal{X} = \overline{([x_1, x_2], 0)}$ then

$$\begin{cases} \|\mathcal{X}\| = \frac{3x_2 - x_1}{2} & \text{if } c(\mathcal{X}) > 0, \\ \|\mathcal{X}\| = \frac{x_2 - 3x_1}{2} & \text{if } c(\mathcal{X}) < 0. \end{cases}$$

Considering the different situations, we obtain

$$\|\mathcal{X}\| \|\mathcal{X}'\| - \|\mathcal{X}\mathcal{X}'\| = \frac{3}{4}l(\mathcal{X})l(\mathcal{X}')$$

or $\frac{1}{2}\|\mathcal{X}\|l(\mathcal{X}')$ or $\frac{1}{2}\|\mathcal{X}'\|l(\mathcal{X})$. These expressions are always positive. We have $\|\mathcal{X}\| \|\mathcal{X}'\| = \|\mathcal{X}\mathcal{X}'\|$ if \mathcal{X} or \mathcal{X}' are reduce to one point.

Proposition 14. *We consider $\mathcal{X} = \overline{(X, 0)}$ and $\mathcal{X}' = \overline{(Y, 0)}$ in $\overline{\mathbb{IR}}$. We have*

$$X \subset Y \Rightarrow \|\mathcal{X}\| \leq \|\mathcal{X}'\|.$$

Proof. Consider $X = [x_1, x_2]$ and $Y = [y_1, y_2]$.

First case: $y_1 \geq 0$. So $2\|\mathcal{X}'\| = 3y_2 - y_1$. As $X \subset Y$, then $2\|\mathcal{X}\| = 3x_2 - x_1$ and $\|\mathcal{X}\| \leq \|\mathcal{X}'\|$.

Second case: $y_1 < 0, y_2 > 0$. If $c(Y) \geq 0$, so $2\|\mathcal{X}'\| = 3y_2 - y_1$. If $c(X) \geq 0$, from the first case $\|\mathcal{X}\| \leq \|\mathcal{X}'\|$. Otherwise $2\|\mathcal{X}\| = x_2 - 3x_1$. Thus $\|\mathcal{X}\| \leq \|\mathcal{X}'\|$ if and only if $3y_2 - y_1 \geq x_2 - 3x_1$, that is $3(y_2 + x_1) \geq x_2 + y_1$ which is true.

If $c(Y) \leq 0$, then $2\|\mathcal{X}'\| = y_2 - 3y_1$. If $c(X) \leq 0$, thus $2\|\mathcal{X}\| = x_2 - 3x_1$ and $\|\mathcal{X}\| \leq \|\mathcal{X}'\|$. If $c(X) \geq 0$, $\|\mathcal{X}\| \leq \|\mathcal{X}'\|$ is equivalent to $y_2 - 3y_1 \geq 3x_2 - x_1$. But $c(Y) \leq 0$ implies $y_1 + y_2 \leq 0$ and $y_2 - 3y_1 \geq 4y_2$. Similarly $3x_2 - x_1 \leq 4x_2$, thus $y_2 - 3y_1 \geq 3x_2 - x_1$ because $x_2 \leq y_2$.

Third case: $y_1 < 0, y_2 < 0$. Similar computations give the result.

Remark. If $\mathcal{X} > 0$, i.e. $\mathcal{X} = \overline{(X, 0)}$, and $\mathcal{X}' < 0$, i.e. $\mathcal{X}' = \overline{(0, Y)}$, so $\setminus\mathcal{X}' > 0$ and if $X \subset Y$ we deduce $\|\mathcal{X}\| \leq \|\setminus\mathcal{X}'\| = \|\mathcal{X}'\|$.

3.2. Definition of \mathcal{A}_4 . Recall that by an algebra we mean a real vector space with an associative ring structure. Consider the 4-dimensional associative algebra whose product in a basis $\{e_1, e_2, e_3, e_4\}$ is given by

	e_1	e_2	e_3	e_4
e_1	e_1	0	0	e_4
e_2	0	e_2	e_3	0
e_3	0	e_3	e_2	0
e_4	e_4	0	0	e_1

The unit is the vector $e_1 + e_2$. This algebra is a direct sum of two ideals: $\mathcal{A}_4 = I_1 + I_2$ where I_1 is generated by e_1 and e_4 and I_2 is generated by e_2 and e_3 . It is not an integral domain, that is, we have divisors of 0. For example $e_1 \cdot e_2 = 0$.

Proposition 15. *The ring \mathcal{A}_4 is principal that is every ideal is generated by one element.*

Proof. In fact we have only two ideals I_1 and I_2 and I_1 is generated by e_4 and I_2 is generated by e_3 . We denote by \mathcal{A}_4^* the group of invertible elements. We compute this group. The cartesian expression of this product is, for $x = (x_1, x_2, x_3, x_4)$ and $y = (y_1, y_2, y_3, y_4)$ in \mathcal{A}_4 :

$$x \cdot y = (x_1y_1 + x_4y_4, x_2y_2 + x_3y_3, x_3y_2 + x_2y_3, x_4y_1 + x_1y_4).$$

We consider the equation

$$x \cdot y = (1, 1, 0, 0).$$

We obtain

$$\begin{cases} x_1y_1 + x_4y_4 = 1, \\ x_2y_2 + x_3y_3 = 1, \\ x_3y_2 + x_2y_3 = 0, \\ x_4y_1 + x_1y_4 = 0. \end{cases}$$

For a given vector x , we obtain a solution y if and only if:

$$(x_1^2 - x_4^2)(x_2^2 - x_3^2) \neq 0.$$

Proposition 16. *The multiplicative group \mathcal{A}_4^* is the set of elements $x = (x_1, x_2, x_3, x_4)$ such that*

$$\begin{cases} x_4 \neq \pm x_1, \\ x_3 \neq \pm x_2. \end{cases}$$

If $x \in \mathcal{A}_4^*$ we have:

$$x^{-1} = \left(\frac{x_1}{x_1^2 - x_4^2}, \frac{x_2}{x_2^2 - x_3^2}, \frac{x_3}{x_2^2 - x_3^2}, \frac{x_4}{x_1^2 - x_4^2} \right).$$

3.3. An embedding of \mathbb{IR} in \mathcal{A}_4 . We define a correspondance between \mathbb{IR} and \mathcal{A}_4 . Let φ be the map

$$\varphi : \mathbb{IR} \longrightarrow \mathcal{A}_4$$

defined by:

$$\varphi(X) = \begin{cases} (x_1, x_2, 0, 0) & \text{if } x_1, x_2 \geq 0, \\ (0, x_2, -x_1, 0) & \text{if } x_1 \leq 0 \text{ and } x_2 \geq 0, \\ (0, 0, -x_1, -x_2) & \text{if } x_1, x_2 \leq 0, \end{cases}$$

for every $X = [x_1, x_2]$ in \mathbb{IR} .

Theorem 17. *Let $X = [x_1, x_2]$ and $Y = [y_1, y_2]$ be in \mathbb{IR} .*

- If $x_1x_2 > 0$ or $y_1y_2 > 0$, then

$$\varphi(XY) = \varphi(X) \cdot \varphi(Y).$$

- If $x_1x_2 < 0$ and $y_1y_2 < 0$, then

$$\varphi^{-1}(\varphi(XY)) \subset \varphi^{-1}(\varphi(X) \cdot \varphi(Y)).$$

Proof. For the first case we have the following table:

	$\varphi(X)$	$\varphi(Y)$	$\varphi(XY)$
$\begin{cases} x_1, x_2 > 0 \\ y_1, y_2 > 0 \end{cases}$	$(x_1, x_2, 0, 0)$	$(y_1, y_2, 0, 0)$	$(x_1y_1, x_2y_2, 0, 0)$
$\begin{cases} x_1, x_2 > 0 \\ y_1 < 0, y_2 > 0 \end{cases}$	$(x_1, x_2, 0, 0)$	$(0, y_2, -y_1, 0)$	$(0, x_2y_2, -x_2y_1, 0)$
$\begin{cases} x_1, x_2 > 0 \\ y_1, y_2 < 0 \end{cases}$	$(x_1, x_2, 0, 0)$	$(0, 0, -y_1, -y_2)$	$(0, 0, -x_2y_1, -x_1y_2)$
$\begin{cases} x_1, x_2 < 0 \\ y_1 < 0, y_2 > 0 \end{cases}$	$(0, 0, -x_1, -x_2)$	$(0, y_2, -y_1, 0)$	$(0, x_1y_1, -x_1y_2, 0)$
$\begin{cases} x_1, x_2 < 0 \\ y_1, y_2 < 0 \end{cases}$	$(0, 0, -x_1, -x_2)$	$(0, 0, -y_1, -y_2)$	$(x_2y_2, x_1y_1, 0, 0)$

and we see that in each case we have $\varphi(XY) = \varphi(X) \cdot \varphi(Y)$. In the second case,

$$\varphi(X) = (0, x_2, -x_1, 0) \text{ and } \varphi(Y) = (0, y_2, -y_1, 0) .$$

Then $\varphi(X) \cdot \varphi(Y) = (0, x_2y_2 + x_1y_1, -x_1y_2 - x_2y_1, 0)$. But XY can be equal to one of the following intervals

$$XY = \begin{cases} [x_1y_2, x_2y_2], \\ [x_2y_1, x_2y_2], \\ [x_1y_2, x_1y_1], \\ [x_2y_1, x_1y_1]. \end{cases}$$

Then

$$\varphi(XY) \in \{(0, x_2y_2, -x_1y_2, 0), (0, x_2y_2, -x_2y_1, 0), (0, x_1y_1, -x_1y_2, 0), (0, x_1y_1, -x_2y_1, 0)\}$$

and

$$\varphi^{-1}(\varphi(X) \cdot \varphi(Y)) = [x_1y_2 + x_2y_1, x_2y_2 + x_1y_1] \supset \varphi^{-1}(\varphi(XY)).$$

Remark. This inclusion is, in some sense, universal because it contains all the cases and it is the minimal expression satisfying this property. For a computational use of this product, this is not a problem because the result $\varphi(X) \cdot \varphi(Y)$ contains the classical product (it is a constraint asked by the arithmetic programming).

3.4. An embedding of $\overline{\mathbb{IR}}$ in \mathcal{A}_4 .

$$\varphi : \mathbb{IR} \longrightarrow \mathcal{A}_4$$

we would like to define $\overline{\varphi} : \overline{\mathbb{IR}} \longrightarrow \mathcal{A}_4$ such that

$$\overline{\varphi}(\overline{K, 0}) = \varphi(K).$$

To define $\overline{\varphi}(\overline{0, K})$ we consider $K = [x_1, x_2]$ and

$$\overline{\varphi}(\overline{0, K}) = \begin{cases} (0, 0, x_1, x_2) \text{ if } x_1 \geq 0, \\ (-x_1, 0, 0, x_2) \text{ if } x_1x_2 \leq 0, \\ (-x_1, -x_2, 0, 0) \text{ if } x_2 \leq 0. \end{cases}$$

Let $\overline{(A, B)}$ be in $\overline{(K, 0)}$. This means that $K = [a_1 - b_1, a_2 - b_2]$. As $\overline{(A, B)} = \overline{(A, 0)} \setminus \overline{(0, B)}$ we have, if $\overline{\varphi}$ is linear

$$\overline{\varphi}(\overline{A, B}) = \overline{\varphi}(\overline{A, 0}) - \overline{\varphi}(\overline{0, B}).$$

Suppose that $a_1, b_1 \geq 0$, and $a_1 - b_1 \geq 0$.

$$\begin{aligned}\overline{\varphi}(A, B) &= (a_1, a_2, 0, 0) - (0, 0, b_1, b_2) \\ &= (a_1, a_2, -b_1, -b_2).\end{aligned}$$

But $\overline{\varphi}(K, 0) = (a_1 - b_1, a_2 - b_2, 0, 0)$ and $Im\overline{\varphi}$ depends on the representants of the equivalent classes. Let us introduce on \mathcal{A}_4 the following equivalence relation

$$(x_1, x_2, x_3, x_4) \sim (y_1, y_2, y_3, y_4) \iff \begin{cases} x_1 - y_1 = -x_3 + y_3, \\ x_2 - y_2 = -x_4 + y_4. \end{cases}$$

We have

$$\begin{cases} a_1 - (a_1 - b_1) = b_1, \\ a_2 - (a_2 - b_2) = b_2, \end{cases}$$

that is $(a_1, a_2, -b_1, -b_2) \simeq (a_1 - b_1, a_2 - b_2, 0, 0)$ and we have to consider

$$\overline{\varphi} : \overline{\mathbb{IR}} \longrightarrow \frac{\mathcal{A}_4}{F} = \overline{\mathcal{A}_4}.$$

There exists always a representant in $\overline{\mathcal{A}_4}$ whose coordinates are positive. In fact

$$\begin{aligned}(x_1, x_2, x_3, x_4) &\simeq (x_1 - x_3, x_2 - x_4, 0, 0) \\ &\simeq (x_1 - x_3, 0, 0, x_4 - x_2) \\ &\simeq (0, x_2 - x_4, -x_3 + x_1, 0) \\ &\simeq (0, 0, -x_3 + x_1, x_4 - x_2),\end{aligned}$$

and for one of them, the coordinates are positive.

Now we consider $Y = \overline{(y_1, y_2, 0, 0)} \in \overline{\mathcal{A}_4}$ with $y_1, y_2 > 0$.

- If $y_1 < y_2$ then $Y = \overline{\varphi}(\overline{[y_1, y_2]}, 0)$.
- If $y_1 > y_2$ then $Y = \overline{\varphi}(0, \overline{[-y_1, -y_2]})$.

If $Y = \overline{(0, y_2, y_1, 0)}$ with $y_1, y_2 > 0$ then $Y = \overline{\varphi}(\overline{[-y_1, y_2]}, 0)$.

If $Y = \overline{(y_1, 0, 0, y_2)}$ with $y_1, y_2 > 0$ then $Y = \overline{\varphi}(0, \overline{[-y_1, y_2]})$.

If $Y = \overline{(0, 0, y_1, y_2)}$ with $y_1, y_2 > 0$ then $Y = \overline{\varphi}(0, \overline{[-y_1, y_2]})$ then

- If $y_1 < y_2$ then $Y = \overline{\varphi}(0, \overline{[y_1, y_2]})$
- If $y_1 > y_2$ then $Y = \overline{\varphi}(\overline{[-y_1, -y_2]}, 0)$.

Thus we define $\psi : \overline{\mathcal{A}_4} \longrightarrow \overline{\mathbb{IR}}$ as follows: let $M = (a, b, c, d)$ be in \mathcal{A}_4 .

- If $a - c < b - d$, then $\psi(\overline{M}) = \overline{([a - c, b - d], 0)}$.
- If $b - d < a - c$, then $\psi(\overline{M}) = \overline{(0, [c - a, d - b])}$.

We obtain a product in $\overline{\mathbb{IR}}$ by taking $\mathcal{X} \bullet \mathcal{X}' = \psi(\overline{\varphi}(\mathcal{X}) \cdot \overline{\varphi}(\mathcal{X}'))$ for all $\mathcal{X}, \mathcal{X}' \in \overline{\mathbb{IR}}$.

3.5. On the monotony of the product. We define on \mathcal{A}_4 a partial order relation by

$$\begin{cases} (x_1, x_2, 0, 0) \leq (y_1, y_2, 0, 0) \iff y_1 \leq x_1 \text{ and } x_2 \leq y_2, \\ (x_1, x_2, 0, 0) \leq (0, y_2, y_3, 0) \iff x_2 \leq y_2, \\ (0, x_2, x_3, 0) \leq (0, y_2, y_3, 0) \iff x_3 \leq y_3 \text{ and } x_2 \leq y_2, \\ (0, 0, x_3, x_4) \leq (0, y_2, y_3, 0) \iff x_3 \leq y_3, \\ (0, 0, x_3, x_4) \leq (0, 0, y_3, y_4) \iff x_3 \leq y_3 \text{ and } y_4 \leq x_4. \end{cases}$$

Proposition 18. Monotony property: Let $\mathcal{X}_1, \mathcal{X}_2 \in \overline{\mathbb{IR}}$. Then

$$\begin{cases} \mathcal{X}_1 \subset \mathcal{X}_2 \implies \mathcal{X}_1 \bullet \mathcal{Z} \subset \mathcal{X}_2 \bullet \mathcal{Z} \text{ for all } \mathcal{Z} \in \overline{\mathbb{IR}}. \\ \overline{\varphi}(\mathcal{X}_1) \leq \overline{\varphi}(\mathcal{X}_2) \implies \overline{\varphi}(\mathcal{X}_1 \bullet \mathcal{Z}) \leq \overline{\varphi}(\mathcal{X}_2 \bullet \mathcal{Z}) \end{cases}$$

Proof. Let us note that the second property is equivalent to the first. It is its translation in $\overline{\mathcal{A}_4}$. For any $\mathcal{X} \in \overline{\mathbb{IR}}$, we denote by $\overline{\mathcal{X}} = \overline{\varphi}(\mathcal{X})$ its image in $\overline{\mathcal{A}_4}$. Let $\mathcal{X}_1, \mathcal{X}_2$ and \mathcal{Z} be in $\overline{\mathbb{IR}}$. We denote by (z_1, z_2, z_3, z_4) the image of \mathcal{Z} in $\overline{\mathcal{A}_4}$, this implies $z_i \geq 0$ and $z_1 z_3 = z_2 z_4 = 0$. We assume that $\overline{\varphi}(\mathcal{X}_1) \leq \overline{\varphi}(\mathcal{X}_2)$.

First case. $\overline{\mathcal{X}}_1 = (x_1, x_2, 0, 0), \overline{\mathcal{X}}_2 = (y_1, y_2, 0, 0)$. We have

$$\begin{cases} \overline{\varphi}(\mathcal{X}_1 \bullet \mathcal{Z}) = (x_1 z_1, x_2 z_2, x_2 z_3, x_1 z_4), \\ \overline{\varphi}(\mathcal{X}_2 \bullet \mathcal{Z}) = (y_1 z_1, y_2 z_2, y_2 z_3, y_1 z_4). \end{cases}$$

As $y_1 z_1 \leq x_1 z_1$ and $x_2 z_2 \leq y_2 z_2$, then $\overline{\varphi}(\mathcal{X}_1 \bullet \mathcal{Z}) \leq \overline{\varphi}(\mathcal{X}_2 \bullet \mathcal{Z})$.

Second case. $\overline{\mathcal{X}}_1 = (x_1, x_2, 0, 0), \overline{\mathcal{X}}_2 = (0, y_2, y_3, 0)$. We have

$$\begin{cases} \overline{\varphi}(\mathcal{X}_1 \bullet \mathcal{Z}) = (x_1 z_1, x_2 z_2, x_2 z_3, x_1 z_4), \\ \overline{\varphi}(\mathcal{X}_2 \bullet \mathcal{Z}) = (0, y_2 z_2 + y_3 z_3, y_2 z_3 + y_3 z_2, 0). \end{cases}$$

If $z_1 \neq 0$ we have $y_2 z_2 + y_3 z_3 \geq x_2 z_2$. If $z_1 = 0$ and $z_2 \neq 0$, we have $y_2 z_3 + y_3 z_2 \geq x_2 z_3$ and $y_2 z_2 + y_3 z_3 \geq x_2 z_2$. If $z_1 = z_2 = 0$, we have $x_2 z_3 \leq y_2 z_3$. This implies that in any case $\overline{\varphi}(\mathcal{X}_1 \bullet \mathcal{Z}) \leq \overline{\varphi}(\mathcal{X}_2 \bullet \mathcal{Z})$.

Third case. $\overline{\mathcal{X}}_1 = (0, x_2, x_3, 0), \overline{\mathcal{X}}_2 = (0, y_2, y_3, 0)$. We have

$$\begin{cases} \overline{\varphi}(\mathcal{X}_1 \bullet \mathcal{Z}) = (0, x_2 z_2 + x_3 z_3, x_2 z_3 + x_3 z_2, 0), \\ \overline{\varphi}(\mathcal{X}_2 \bullet \mathcal{Z}) = (0, y_2 z_2 + y_3 z_3, y_2 z_3 + y_3 z_2, 0). \end{cases}$$

Thus

$$\overline{\varphi}(\mathcal{X}_1 \bullet \mathcal{Z}) \leq \overline{\varphi}(\mathcal{X}_2 \bullet \mathcal{Z}) \iff \begin{cases} (x_2 - y_2)z_2 + (x_3 - y_3)z_3 \leq 0, \\ (x_2 - y_2)z_3 + (x_3 - y_3)z_2 \leq 0. \end{cases}$$

But $(x_2 - y_2), (x_3 - y_3) \leq 0$ and $z_2, z_3 \geq 0$. This implies $\overline{\varphi}(\mathcal{X}_1 \bullet \mathcal{Z}) \leq \overline{\varphi}(\mathcal{X}_2 \bullet \mathcal{Z})$.

Fourth case. $\overline{\mathcal{X}}_1 = (0, 0, x_3, x_4), \overline{\mathcal{X}}_2 = (0, y_2, y_3, 0)$. We have

$$\begin{cases} \overline{\varphi}(\mathcal{X}_1 \bullet \mathcal{Z}) = (x_4 z_4, x_3 z_3, x_3 z_2, x_4 z_1), \\ \overline{\varphi}(\mathcal{X}_2 \bullet \mathcal{Z}) = (0, y_2 z_2 + y_3 z_3, y_2 z_3 + y_3 z_2, 0). \end{cases}$$

If $z_4 \neq 0$, then $z_2 = 0$ and $x_3 z_3 \leq y_3 z_3$. If $z_4 = 0$ and $z_3 \neq 0$, we have $x_3 z_3 \leq y_2 z_2 + y_3 z_3$ and $x_3 z_2 \leq y_2 z_3 + y_3 z_2$. If $z_4 = z_3 = 0$ then $x_3 z_2 \leq x_3 z_2$. This implies that in any case $\overline{\varphi}(\mathcal{X}_1 \bullet \mathcal{Z}) \leq \overline{\varphi}(\mathcal{X}_2 \bullet \mathcal{Z})$.

Fifth case. $\overline{\mathcal{X}}_1 = (0, 0, x_3, x_4), \overline{\mathcal{X}}_2 = (0, 0, y_3, y_4)$. We have

$$\begin{cases} \overline{\varphi}(\mathcal{X}_1 \bullet \mathcal{Z}) = (x_4 z_4, x_3 z_3, x_3 z_2, x_4 z_1), \\ \overline{\varphi}(\mathcal{X}_2 \bullet \mathcal{Z}) = (y_4 z_4, y_3 z_3, y_3 z_2, y_4 z_1). \end{cases}$$

But $\overline{\varphi}(\mathcal{X}_1 \bullet \mathcal{Z}) \leq \overline{\varphi}(\mathcal{X}_2 \bullet \mathcal{Z})$ is equivalent to $\begin{cases} x_3 z_2 \leq y_3 z_2, \\ y_4 z_1 \leq x_4 z_1, \end{cases}$ and this is always satisfied.

3.6. Application. To work in the intervals set, we propose the following program:

1. Translate the problem in \mathcal{A}_4 through $\overline{\mathbb{IR}}$.
2. Solve the problem in \mathcal{A}_4 (which is a normed associative algebra).
3. Return to $\overline{\mathbb{IR}}$ and then to \mathbb{IR} .

The last condition require the use of ψ .

3.7. Remark. In the first version of this paper we have considered an another product in \mathcal{A}_4 but this product was not minimal.

4. DIVISIBILITY AND AN EUCLIDEAN DIVISION

We have computed the inversible elements of \mathcal{A}_4 . If $x = (x_1, x_2, x_3, x_4) \in \mathcal{A}_4$ and if $\Delta = (x_1^2 - x_4^2)(x_2^2 - x_3^2) \neq 0$ then

$$x^{-1} = \left(\frac{x_1}{x_1^2 - x_4^2}, \frac{x_2}{x_2^2 - x_3^2}, \frac{x_3}{x_2^2 - x_3^2}, \frac{x_4}{x_1^2 - x_4^2} \right).$$

The elements associated to $\mathcal{X} = \overline{(K, 0)}$ are of the form

$$\begin{cases} (x_1, x_2, 0, 0) & \text{if } 0 < x_1 < x_2, \\ (0, x_2, -x_1, 0) & \text{if } x_1 < 0 < x_2, \\ (0, 0, -x_1, -x_2) & \text{if } x_1 < x_2 < 0, \end{cases}$$

and to $\mathcal{X} \in \overline{(0, K)}$

$$\begin{cases} (0, 0, x_1, x_2) & \text{if } 0 < x_1 < x_2, \\ (-x_1, 0, 0, x_2) & \text{if } x_1 < 0 < x_2, \\ (-x_1, -x_2, 0, 0) & \text{if } x_1 < x_2 < 0. \end{cases}$$

The inverse of $(x_1, x_2, 0, 0)$ with $0 < x_1 < x_2$ is $\left(\frac{1}{x_1}, \frac{1}{x_2}, 0, 0 \right)$.

The inverse of $(0, 0, -x_1, -x_2)$ with $x_1 < x_2 < 0$ is $\left(0, 0, -\frac{1}{x_1}, -\frac{1}{x_2} \right)$.

The inverse of $(0, 0, x_1, x_2)$ with $0 < x_1 < x_2$ is $\left(0, 0, \frac{1}{x_1}, \frac{1}{x_2} \right)$.

The inverse of $(-x_1, -x_2, 0, 0)$ with $x_1 < x_2 < 0$ is $\left(-\frac{1}{x_1}, -\frac{1}{x_2}, 0, 0 \right)$.

For $\mathcal{X} = (0, x_2, -x_1, 0)$ or $(-x_1, 0, 0, x_2)$ with $x_1 x_2 < 0$, then $\Delta = 0$ and \mathcal{X} is not invertible. Then if $\Delta \neq 0$ the inverse is always represented by an element of $\overline{\mathbb{IR}}$ through ψ .

4.1. Division by an invertible element. We denote by $\overline{\mathbb{IR}}^+$ the subset $\overline{(X, 0)}$ with $X = [x_1, x_2]$ and $0 \leq x_1$.

Proposition 19. *Let $\mathcal{X} = \overline{(X, 0)}$ and $\mathcal{Y} = \overline{(Y, 0)}$ be in $\overline{\mathbb{IR}}^+$ with $X = [x_1, x_2]$, $Y = [y_1, y_2]$. If $\frac{y_2}{y_1} \geq \frac{x_2}{x_1}$ then there exists an unique $\mathcal{Z} = \overline{(Z, 0)} \in \overline{\mathbb{IR}}^+$ such that $\mathcal{Y} = \mathcal{X}\mathcal{Z}$.*

Proof. Let \mathcal{Z} be defined by $c(\mathcal{Z}) = \frac{1}{2} \left(\frac{y_2}{x_2} + \frac{y_1}{x_1} \right)$ and $l(\mathcal{Z}) = \left(\frac{y_2}{x_2} - \frac{y_1}{x_1} \right)$. Then $l(\mathcal{Z}) \geq 0$ if and only if $\frac{y_2}{x_2} \geq \frac{y_1}{x_1}$ that is $\frac{y_2}{y_1} \geq \frac{x_2}{x_1}$. Thus we have $\mathcal{Y} = \mathcal{X}\mathcal{Z}$. In fact

$$(\overline{\varphi(\mathcal{X})})^{-1} = \overline{\left(\frac{1}{x_1}, \frac{1}{x_2}, 0, 0 \right)} = \psi \left(\overline{\left(0, \left[-\frac{1}{x_1}, -\frac{1}{x_2} \right] \right)} \right).$$

Thus

$$\overline{\varphi(\mathcal{Y})} \cdot \overline{\varphi(\mathcal{X})}^{-1} = (y_1, y_2, 0, 0) \cdot \left(\frac{1}{x_1}, \frac{1}{x_2}, 0, 0 \right) = \left(\frac{y_1}{x_1}, \frac{y_2}{x_2}, 0, 0 \right).$$

As $\frac{y_1}{x_1} \leq \frac{y_2}{x_2}$,

$$\psi(\overline{\varphi(\mathcal{Y})} \cdot \overline{\varphi(\mathcal{X})}^{-1}) = \overline{\left(\left[\frac{y_1}{x_1}, \frac{y_2}{x_2} \right], 0 \right)}.$$

We can note also that

$$\overline{\left(0, \left[-\frac{1}{x_1}, -\frac{1}{x_2} \right] \right)} \bullet \overline{\left(\left[y_1, y_2 \right], 0 \right)} = \overline{\left(\left[\frac{y_1}{x_1}, \frac{y_2}{x_2} \right], 0 \right)}.$$

Then the divisibility corresponds to the multiplication by the inverse element.

4.2. Division by a non invertible element. Let $X = [-x_1, x_2]$ with $x_1, x_2 > 0$. We have seen that $\varphi(X) = (0, x_2, x_1, 0)$ is not invertible in \mathcal{A}_4 . For any $M = (y_1, y_2, y_3, y_4) \in \mathcal{A}_4$ we have

$$\varphi(X) \cdot M = (0, x_2 y_2 + x_1 y_3, x_1 y_2 + x_2 y_3, 0)$$

and this point represents a non invertible interval. Thus we can solve the equation $\mathcal{Y} = \mathcal{X} \bullet \mathcal{Z}$ for $\mathcal{X} = \overline{\left([-x_1, x_2], 0 \right)}$, $\mathcal{Y} = \overline{\left([-y_1, y_2], 0 \right)}$ with $x_1, x_2 > 0$ and $y_1, y_2 > 0$. Putting $\varphi(\mathcal{Z}) = (z_1, z_2, z_3, z_4)$, we obtain

$$(0, y_2, y_1, 0) = (0, x_2, x_1, 0) \cdot (z_1, z_2, z_3, z_4),$$

that is

$$\begin{cases} y_2 = x_2 z_2 + x_1 z_3, \\ y_1 = x_2 z_3 + x_1 z_2, \end{cases}$$

or

$$\begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} x_1 & x_2 \\ x_2 & x_1 \end{pmatrix} \begin{pmatrix} z_2 \\ z_3 \end{pmatrix}.$$

If $x_1^2 - x_2^2 \neq 0$,

$$\begin{cases} z_2 = \frac{x_1 y_1 - x_2 y_2}{x_1^2 - x_2^2}, \\ z_3 = \frac{-x_2 y_1 + x_1 y_2}{x_1^2 - x_2^2}. \end{cases}$$

If $x_1^2 - x_2^2 = 0$ then $x_1 = x_2$ and the center of $X = [-x_1, x_1]$ is 0. Let us assume that $x_1 \neq x_2$. If $x_1^2 - x_2^2 < 0$ that is $x_1 < x_2$ then

$$\begin{cases} x_1 y_1 - x_2 y_2 < 0, \\ x_1 y_2 - x_2 y_1 < 0, \end{cases}$$

and $\frac{x_1}{x_2} < \frac{y_2}{y_1}$, $\frac{x_1}{x_2} < \frac{y_1}{y_2}$. If $\alpha = \frac{x_1}{x_2} < 1$ we have $y_2 > \alpha y_1$, $y_1 > \alpha y_2$ then $y_2 > \alpha^2 y_2$ and $1 - \alpha^2 > 0$. This case admits solution.

Proposition 20. *Let $\mathcal{X} = \overline{([-x_1, x_2], 0)}$ with $x_1, x_2 > 0$ and $x_1 < x_2$. Then for any $\mathcal{Y} = \overline{([-y_1, y_2], 0)}$ with $y_1, y_2 > 0$ and $\frac{x_1}{x_2} < \frac{y_2}{y_1}$, $\frac{x_1}{x_2} < \frac{y_1}{y_2}$, there is $\mathcal{Z} = \overline{([-z_1, z_2], 0)}$ such that $\mathcal{Y} = \mathcal{X} \bullet \mathcal{Z}$.*

Suppose now that $x_1^2 - x_2^2 > 0$ that us $x_1 > x_2$. In this case we have

$$\begin{cases} x_1 y_1 - x_2 y_2 > 0, \\ x_1 y_2 - x_2 y_1 > 0, \end{cases}$$

that is $\frac{y_2}{y_1} < \frac{x_1}{x_2}$ and $\frac{y_1}{y_2} < \frac{x_1}{x_2}$.

Proposition 21. *Let $\mathcal{X} = \overline{([-x_1, x_2], 0)}$ with $x_1, x_2 > 0$ and $x_1 > x_2$. For any $\mathcal{Y} = \overline{([-y_1, y_2], 0)}$ with $y_1, y_2 > 0$, $\frac{x_1}{x_2} > \frac{y_2}{y_1}$, $\frac{x_1}{x_2} > \frac{y_1}{y_2}$, there is $\mathcal{Z} = \overline{([-z_1, z_2], 0)}$ such that $\mathcal{Y} = \mathcal{X} \bullet \mathcal{Z}$.*

Example. $\mathcal{X} = \overline{([-4, 2], 0)}$, $\mathcal{Y} = \overline{([-2, 3], 0)}$. We have $\frac{x_2}{x_1} = \frac{1}{2}$, $\frac{x_1}{x_2} = 2$ and $\frac{3}{2} < 2 < 6$. Then \mathcal{Z} exists and it is equal to $\mathcal{Z} = \overline{([- \frac{8}{12}, \frac{2}{12}], 0)}$.

4.3. An euclidean division. Consider $\mathcal{X} = \overline{([x_1, x_2], 0)}$ and $\mathcal{Y} = \overline{([y_1, y_2], 0)}$ in $\overline{\mathbb{IR}^+}$. We have seen that \mathcal{Y} is divisible by \mathcal{X} as soon as $\frac{x_1}{x_2} \geq \frac{y_1}{y_2}$. We suppose now that $\frac{x_1}{x_2} < \frac{y_1}{y_2}$. In this case we have

Theorem 22. *Let \mathcal{X} and \mathcal{Y} be in $\overline{\mathbb{IR}^+}$ with $\frac{x_1}{x_2} < \frac{y_1}{y_2}$. There is a unique pair $(\mathcal{Z}, \mathcal{R})$ unique in $\overline{\mathbb{IR}^+}$ such that*

$$\begin{cases} \mathcal{Y} = \mathcal{X} \bullet \mathcal{Z} + \mathcal{R}, \\ l(\mathcal{R}) = 0 \text{ and } c(\mathcal{R}) \text{ minimal.} \end{cases}$$

This pair is given by

$$\begin{cases} \mathcal{Z} = \frac{y_2 - y_1}{x_2 - x_1} \overline{([1, 1], 0)}, \\ \mathcal{R} = \frac{x_2 y_1 - x_1 y_2}{x_2 - x_1} \overline{([1, 1], 0)}. \end{cases}$$

Proof. We consider $\mathcal{Z} = \overline{([z_1, z_2], 0)}$ with $z_1 > 0$. Then $\mathcal{Y} = \mathcal{X} \bullet \mathcal{Z} + \mathcal{R}$ gives

$$\mathcal{R} = \overline{([y_1, y_2], [z_1 x_1, z_2 x_2])}.$$

We have $\mathcal{R} \in \overline{\mathbb{IR}^+}$ if and only if $0 \leq y_1 - x_1 z_1 \leq y_2 - x_2 z_2$ that is

$$\begin{cases} z_1 \leq \frac{y_1}{x_1}, \\ z_2 \leq \frac{y_2}{x_2}, \\ z_1 \geq \frac{y_1 - y_2 + x_2 z_2}{x_1}. \end{cases}$$

The condition $z_1 \leq z_2$ implies $\frac{y_1 - y_2 + x_2 z_2}{x_1} \leq z_2$ that is $z_2 \leq \frac{y_2 - y_1}{x_2 - x_1}$. Consider the case $z_2 = \frac{y_2 - y_1}{x_2 - x_1}$. Then $z_1 \geq \frac{y_1 - y_2 + x_2 z_2}{x_1} = \frac{y_2 - y_1}{x_2 - x_1} = z_2$ and $z_1 = z_2$. This case corresponds to

$$\begin{cases} \mathcal{Z} = \frac{y_2 - y_1}{x_2 - x_1} \overline{([1, 1], 0)}, \\ \mathcal{R} = \frac{x_2 y_1 - x_1 y_2}{x_2 - x_1} \overline{([1, 1], 0)}. \end{cases}$$

Let us note that $y_1 x_2 - x_1 y_2 > 0$ is equivalent to $\frac{y_1}{y_2} > \frac{x_1}{x_2}$ which is satisfied by hypothesis.

We have also for this solution $l(\mathcal{R}) = 0$ and $c(\mathcal{R}) = \frac{x_2 y_1 - x_1 y_2}{x_2 - x_1}$.

Conversely, if $l(\mathcal{R}) = 0$ then $y_1 - z_1 x_1 = y_2 - z_2 x_2$ and $z_1 = z_2 \frac{x_2}{x_1} + \frac{y_1 - y_2}{x_1}$. As $z_1 > 0$, we obtain $z_2 > \frac{y_1 - y_2}{x_1}$ and $z_1 \leq z_2$ implies

$$\frac{y_2 - y_1}{x_2} \leq z_2 \leq \frac{y_2 - y_1}{x_2 - x_1}.$$

But $c(\mathcal{R}) = y_1 - z_1 x_1 = y_2 - z_2 x_2$. Thus

$$\frac{x_2 y_1 - x_1 y_2}{x_2 - x_1} \leq c(\mathcal{R}) \leq y_1.$$

The norm is minimal when $c(\mathcal{R}) = \frac{x_2 y_1 - x_1 y_2}{x_2 - x_1}$.

Example. Let $\mathcal{X} = \overline{([1, 4], 0)}$ and $\mathcal{Y} = \overline{([1, 3], 0)}$. We have $\frac{x_1}{x_2} = \frac{1}{4} < \frac{y_1}{y_2} = \frac{1}{3}$. Thus $\mathcal{Z} = \frac{2}{3} \overline{([1, 1], 0)}$ and $\mathcal{R} = \frac{1}{3} \overline{([1, 1], 0)}$. The division writes

$$\overline{([1, 3], 0)} = \overline{([1, 4], 0)} \cdot \overline{\left(\frac{2}{3}, \frac{2}{3}\right), 0} + \overline{\left(\frac{1}{3}, \frac{1}{3}\right), 0}.$$

Suppose now that \mathcal{X} and \mathcal{Y} are not invertible, that is $\mathcal{X} = \overline{([-x_1, x_2], 0)}$ and $\mathcal{Y} = \overline{([-y_1, y_2], 0)}$ with x_1, x_2, y_1, y_2 positive. We have seen that \mathcal{Y} is divisible by \mathcal{X} as soon as

$$\begin{cases} \frac{x_1}{x_2} > \frac{y_2}{y_1} \text{ and } \frac{x_1}{x_2} > \frac{y_1}{y_2}, \\ \text{or} \\ \frac{x_1}{x_2} < \frac{y_2}{y_1} \text{ and } \frac{x_1}{x_2} < \frac{y_1}{y_2}. \end{cases}$$

We suppose now that these conditions are not satisfied. For example we assume that

$$\frac{x_1}{x_2} > \frac{y_2}{y_1} \text{ and } \frac{x_1}{x_2} < \frac{y_1}{y_2}$$

(The other case is similar). If $\mathcal{Y} = \mathcal{X} \bullet \mathcal{Z} + \mathcal{R}$ then $\mathcal{R} = \overline{([-r_1, r_2], 0)}$ with $r_1 \geq 0$ and with $r_2 \geq 0$ because $\overline{\varphi}(\mathcal{R}) = (0, r_2, r_1, 0)$. This shows that we can choose \mathcal{Z} such that

$\overline{\varphi}(\mathcal{R}) = (0, z_2, z_3, 0)$ and

$$\begin{cases} z_2 = \frac{x_1(y_1 - r_1) - x_2(y_2 - r_2)}{x_1^2 - x_2^2}, \\ z_3 = \frac{x_1(y_2 - r_2) - x_2(y_1 - r_1)}{x_1^2 - x_2^2}, \end{cases}$$

with the condition $z_2 \geq 0$ and $z_3 \geq 0$. If $x_1 < x_2$ then this is equivalent to

$$\begin{cases} \frac{x_1}{x_2} < \frac{y_2 - r_2}{y_1 - r_1}, \\ \frac{x_1}{x_2} < \frac{y_1 - r_1}{y_2 - r_2}. \end{cases}$$

If we suppose that $\mathcal{R} \leq \mathcal{Y}$, thus $0 < r_2 < y_2$ and $0 < r_1 < y_1$, we obtain

$$r_1 > \frac{x_2}{x_1}r_2 + \frac{-x_2y_2 + x_1y_1}{x_1} < r_1 < \frac{x_1}{x_2}r_2 + \frac{x_2y_1 - x_1y_2}{x_2}.$$

Then length $l(\mathcal{R}) = r_1 + r_2$ is minimal if and only if $r_2 = 0$ and in this case $r_1 = \frac{x_1y_1 - x_2y_2}{x_1}$.

We obtain

$$\begin{cases} z_2 = 0, \\ z_3 = \frac{y_2}{x_1}. \end{cases}$$

Theorem 23. Let $\mathcal{X} = \overline{([-x_1, x_2], 0)}$ with $x_1, x_2 > 0$ and $x_1 > x_2$. If $\mathcal{Y} = \overline{([-y_1, y_2], 0)}$ with $y_1, y_2 > 0$, satisfies $\frac{x_1}{x_2} > \frac{y_2}{y_1}$ and $\frac{x_1}{x_2} < \frac{y_1}{y_2}$, there is a unique pair \mathcal{R}, \mathcal{Z} of non invertible elements such that

$$\begin{cases} l(\mathcal{R}) \text{ minimal,} \\ \mathcal{R} < \mathcal{Y}. \end{cases}$$

This pair is given by

$$\begin{cases} \mathcal{Z} = \overline{([- \frac{y_2}{x_1}, 0], 0)}, \\ \mathcal{R} = \overline{([- \frac{x_1y_1 - x_2y_2}{x_1}, 0], 0)}. \end{cases}$$

5. APPLICATIONS

5.1. **Linear programming on $\overline{\mathbb{IR}}$.** We consider the following linear programm

$$\begin{cases} Ax = b, \\ \max(f(x)), \end{cases}$$

where A is a real $n \times p$ matrix and $f : \mathbb{R}^n \rightarrow \mathbb{R}$ a linear map. We want to translate this problem in the context of intervals. This is easy because the set of intervals is endowed with a vectorial space structure. Then we assume here that each variable x_i belongs to an interval $\mathcal{X}_i = \overline{(X_i, 0)}$. We assume also that the constraints b_i belongs to $Y_i = \overline{(B_i, 0)}$ with $B_i = [b_i^1, b_i^2]$ and $b_i^1 \geq 0$. We can extend the linear $f : \mathbb{R}^n \rightarrow \mathbb{R}$ which is written $f(x_1, \dots, x_n) = \sum a_i x_i$ by a linear map, denoted by \overline{f} , given by $\overline{f}(\mathcal{X}_1, \dots, \mathcal{X}_n) = \sum a_i \mathcal{X}_i$. In $\overline{\mathbb{IR}}$ we claim that

$$\mathcal{X} \geq \mathcal{X}' \iff \mathcal{X} \setminus \mathcal{X}' = \overline{(K, 0)}.$$

We have the following linear programming in terms of intervals:

$$\begin{cases} A\mathcal{X} = \mathcal{B} \text{ with } \mathcal{X} \in \overline{\mathbb{IR}}^n \text{ and } \mathcal{B} \in \overline{\mathbb{IR}}^p, \\ \mathcal{B} = (\mathcal{B}_1, \dots, \mathcal{B}_p) \text{ and } \mathcal{B}_i \geq 0, \\ \max(\overline{f}(\mathcal{X})). \end{cases}$$

To solve this program, we extend the classical simplex algorithm. The simplex method is based on an algorithm using the Gauss elimination. We consider the linear program on the intervals, with $\mathcal{X} = (\mathcal{X}_1, \dots, \mathcal{X}_n) \in \overline{\mathbb{IR}}^n$ and $\mathcal{B} = (\mathcal{B}_1, \dots, \mathcal{B}_p) \in \overline{\mathbb{IR}}^p$. By hypothesis $\mathcal{B}_i = \overline{(Y_i, 0)}$ is a positive vector. We have to define the pivot. The column pivot is defined by the largest positive coefficient of the economic function \overline{f} . In choosing the line, the goal is to keep the second member of positive vectors of $\overline{\mathbb{IR}}^p$. Let k be the number of the column containing the pivot. We note $l(Y_j)$ the length of the interval Y_j and let i be such that

$$\frac{l(Y_i)}{a_{ik}} = \min_j \left\{ \frac{l(Y_j)}{a_{jk}}, a_{jk} > 0 \right\}.$$

We choose a_{ik}^k as pivot. The line l_j is transformed into $a_{ik}l_j - a_{jk}l_i$. In this case the second member becomes $a_{ik}\mathcal{B}_j \setminus a_{jk}\mathcal{B}_i$. Let us compute this value. We assume $a_{jk} > 0$

$$\begin{aligned} a_{ik}\mathcal{B}_j \setminus a_{jk}\mathcal{B}_i &= \overline{(a_{ik}Y_j, a_{jk}Y_i)} \\ &= \overline{(a_{ik}[Y_j^1, Y_j^2], a_{jk}[Y_i^1, Y_i^2])} \\ &= \overline{([a_{ik}Y_j^1, a_{ik}Y_j^2], [a_{jk}Y_i^1, a_{jk}Y_i^2])} \\ &= \overline{([a_{ik}Y_j^1 - a_{jk}Y_i^1, a_{ik}Y_j^2 - a_{jk}Y_i^2], 0)}. \end{aligned}$$

This is well defined. Indeed

$$\begin{aligned} (a_{ik}Y_j^1 - a_{jk}Y_i^1) &< a_{ik}Y_j^2 - a_{jk}Y_i^2 \\ \iff a_{jk}(Y_i^2 - Y_i^1) &< a_{ik}(Y_j^2 - Y_j^1) \\ \iff \frac{l(Y_i)}{a_{ik}} &< \frac{l(Y_j)}{a_{jk}}, \end{aligned}$$

what is assumed by hypothesis. So $a_{ik}\mathcal{B}_j \setminus a_{jk}\mathcal{B}_i$ is positive. In the case where $a_{jk} < 0$, $l(Y_j)$ is transformed into $a_{ik}l(Y_j) - a_{jk}l(Y_i)$ and thus the second member is given by

$$\overline{(a_{ik}Y_j, a_{jk}Y_i)} = \overline{(a_{ik}Y_j, 0)} + \overline{(-a_{jk}Y_i, 0)} = \overline{(a_{ik}Y_j - a_{jk}Y_i, 0)}.$$

This vector is always positive. This process, as in the classical simplex algorithm, gives in terms of intervals the maximum of the economic function.

5.2. Differential calculus on $\overline{\mathbb{IR}}$. As $\overline{\mathbb{IR}}$ is a Banach space, we can describe a notion of differential function on it. Consider $\mathcal{X}_0 = \overline{(X_0, 0)}$ in $\overline{\mathbb{IR}}$. The norm $\|\cdot\|$ defines a topology on $\overline{\mathbb{IR}}$ whose a basis of neighbourhoods is given by the balls $\mathcal{B}(X_0, \varepsilon) = \{X \in \overline{\mathbb{IR}}, \|X \setminus \mathcal{X}_0\| < \varepsilon\}$. Let us characterize the elements of $\mathcal{B}(X_0, \varepsilon)$. $\mathcal{X}_0 = \overline{(X_0, 0)} = \overline{([a, b], 0)}$.

Proposition 24. *Consider $\mathcal{X}_0 = \overline{(X_0, 0)} = \overline{([a, b], 0)}$ in $\overline{\mathbb{IR}}$. Then every element of $\mathcal{B}(X_0, \varepsilon)$ is of type $\mathcal{X} = \overline{(X, 0)}$ and satisfies*

$$l(X) \in B_{\mathbb{R}}(l(X_0), \varepsilon_1) \text{ and } c(X) \in B_{\mathbb{R}}(c(X_0), \varepsilon_2)$$

with $\varepsilon_1, \varepsilon_2 \geq 0$ and $\varepsilon_1 + \varepsilon_2 \leq \varepsilon$, where $B_{\mathbb{R}}(x, a)$ is the canonical open ball in \mathbb{R} of center x and radius a .

Proof. First case : Assume that $\mathcal{X} = \overline{(X, 0)} = \overline{([x, y], 0)}$. We have

$$\begin{aligned} \mathcal{X} \setminus \mathcal{X}_0 &= \overline{(X, X_0)} = \overline{([x, y], [a, b])} \\ &= \begin{cases} \overline{([x-a, y-b], 0)} & \text{if } l(X) \geq l(X_0) \\ \overline{(0, [a-x, b-y])} & \text{if } l(X) \leq l(X_0) \end{cases} \end{aligned}$$

If $l(X) \geq l(X_0)$ we have

$$\begin{aligned} \|\mathcal{X} \setminus \mathcal{X}_0\| &= (y-b) - (x-a) + \left| \frac{y-b+x-a}{2} \right| \\ &= l(X) - l(X_0) + |c(X) - c(X_0)|. \end{aligned}$$

As $l(X) - l(X_0) \geq 0$ and $|c(X) - c(X_0)| \geq 0$, each one of this term is less than ε . If $l(X) \leq l(X_0)$ we have

$$\|\mathcal{X} \setminus \mathcal{X}_0\| = l(X_0) - l(X) + |c(X_0) - c(X)|.$$

and we have the same result.

Second case : Consider $\mathcal{X} = \overline{(0, X)} = \overline{([x, y], 0)}$. We have

$$\mathcal{X} \setminus \mathcal{X}_0 = \overline{(0, X_0 + X)} = \overline{([x+a, y+b])}$$

and

$$\|\mathcal{X} \setminus \mathcal{X}_0\| = l(X_0) + l(X) + |c(X_0) + c(X)|.$$

In this case, we cannot have $\|\mathcal{X} \setminus \mathcal{X}_0\| < \varepsilon$ thus $X \notin \mathcal{B}(X_0, \varepsilon)$.

Definition 25. A function $f : \overline{\mathbb{R}} \rightarrow \overline{\mathbb{R}}$ is continuous at \mathcal{X}_0 if

$$\forall \varepsilon > 0, \exists \eta > 0 \text{ such that } \|\mathcal{X} \setminus \mathcal{X}_0\| < \varepsilon \text{ implies } \|f(\mathcal{X}) \setminus f(\mathcal{X}_0)\| < \varepsilon.$$

Consider $(\mathcal{X}_1, \mathcal{X}_2)$ the basis of $\overline{\mathbb{R}}$ given in section 2. We have

$$f(\mathcal{X}) = f_1(\mathcal{X})\mathcal{X}_1 + f_2(\mathcal{X})\mathcal{X}_2 \text{ with } f_i : \overline{\mathbb{R}} \rightarrow \mathbb{R}.$$

If f is continuous at \mathcal{X}_0 so

$$f(\mathcal{X}) \setminus f(\mathcal{X}_0) = (f_1(\mathcal{X}) - f_1(\mathcal{X}_0))\mathcal{X}_1 + (f_2(\mathcal{X}) - f_2(\mathcal{X}_0))\mathcal{X}_2.$$

To simplify notations let $\alpha = f_1(\mathcal{X}) - f_1(\mathcal{X}_0)$ and $\beta = f_2(\mathcal{X}) - f_2(\mathcal{X}_0)$. If $\|f(\mathcal{X}) \setminus f(\mathcal{X}_0)\| < \varepsilon$, and if we assume $f_1(\mathcal{X}) - f_1(\mathcal{X}_0) > 0$ and $f_2(\mathcal{X}) - f_2(\mathcal{X}_0) > 0$ (other cases are similar), then we have

$$l(\alpha\mathcal{X}_1 + \beta\mathcal{X}_2) = l(\overline{[\beta, \alpha + \beta], 0}) < \varepsilon$$

thus $f_1(\mathcal{X}) - f_1(\mathcal{X}_0) < \varepsilon$. Simillary,

$$c(\alpha\mathcal{X}_1 + \beta\mathcal{X}_2) = c(\overline{[\beta, \alpha + \beta], 0}) = \frac{\alpha}{2} + \beta < \varepsilon$$

and this implies that $f_2(\mathcal{X}) - f_2(\mathcal{X}_0) < \varepsilon$.

Corollary 26. f is continuous at \mathcal{X}_0 if and only if f_1 and f_2 are continuous at \mathcal{X}_0 .

Examples.

- $f(\mathcal{X}) = \mathcal{X}$. This function is continuous at any point.

- $f(\mathcal{X}) = \mathcal{X}^2$. Consider $\mathcal{X}_0 = \overline{(X_0, 0)} = \overline{([a, b], 0)}$ and $\mathcal{X} \in \mathcal{B}(X_0, \varepsilon)$. We have

$$\begin{aligned} \|\mathcal{X}^2 \setminus \mathcal{X}_0^2\| &= \|(\mathcal{X} \setminus \mathcal{X}_0)(\mathcal{X} + \mathcal{X}_0)\| \\ &\leq \|\mathcal{X} \setminus \mathcal{X}_0\| \|\mathcal{X} + \mathcal{X}_0\|. \end{aligned}$$

Given $\varepsilon > 0$, let $\eta = \frac{\varepsilon}{\|\mathcal{X} + \mathcal{X}_0\|}$, thus if $\|\mathcal{X} \setminus \mathcal{X}_0\| < \eta$, we have $\|\mathcal{X}^2 \setminus \mathcal{X}_0^2\| < \varepsilon$ and f is continuous.

- Consider $P = a_0 + a_1X + \dots + a_nX^n \in \mathbb{R}[X]$. We define $f : \overline{\mathbb{R}} \rightarrow \overline{\mathbb{R}}$ with $f(\mathcal{X}) = a_0\mathcal{X}_0 + a_1\mathcal{X} + \dots + a_n\mathcal{X}^n$ where $\mathcal{X}^n = \mathcal{X} \cdot \mathcal{X}^{n-1}$. From the previous example, all monomials are continuous, it implies that f is continuous.

Definition 27. Consider \mathcal{X}_0 in $\overline{\mathbb{R}}$ and $f : \overline{\mathbb{R}} \rightarrow \overline{\mathbb{R}}$ continuous. We say that f is differentiable at \mathcal{X}_0 if there is $g : \overline{\mathbb{R}} \rightarrow \overline{\mathbb{R}}$ linear such as

$$\|f(\mathcal{X}) \setminus f(\mathcal{X}_0) \setminus g(\mathcal{X} \setminus \mathcal{X}_0)\| = o(\|\mathcal{X} \setminus \mathcal{X}_0\|).$$

5.3. Non Linear programming. We consider here the following programm

$$\begin{cases} Ax = b, \\ \max(f(x)), \end{cases}$$

where A is a real $n \times p$ matrix and $f : \mathbb{R}^n \rightarrow \mathbb{R}$ a non linear differentiable map. We translate this programm in terms of intervals:

$$\begin{cases} A\mathcal{X} = \mathcal{B} \text{ with } \mathcal{X} \in \overline{\mathbb{R}}^n \text{ and } \mathcal{B} \in \overline{\mathbb{R}}^p, \\ \mathcal{B} = (\mathcal{B}_1, \dots, \mathcal{B}_p) \text{ and } \mathcal{B}_i \geq 0, \\ \max(\tilde{f}(\mathcal{X})). \end{cases}$$

where $\tilde{f} : \overline{\mathbb{R}}^n \rightarrow \overline{\mathbb{R}}$ is the transfered function. It is differentiable in the previous sense. Assume that the system $A\mathcal{X} = \mathcal{B}$ is indetermined and all the solutions are expressed in term of one parameter \mathcal{X}_0 . In this case, a solution of the non linear programming is given by a root of the equation $f'(\mathcal{X}_0) = 0$.

REFERENCES

- [1] A. Goldsztejn. *Définition et Applications des Extensions des Fonctions Réelles aux Intervalles Généralisés: Nouvelle Formulation de la Théorie des Intervalles Modaux et Nouveaux Résultats*, PhD thesis, University of Nice Sophia Antipolis, 2005.
- [2] M. Goze; N. Goze. *Arithmétique des Intervalles Infiniment Petits*. Preprint Mulhouse, 2008.
- [3] L. Jaulin; M. Kieffer; O. Didrit; E. Walter. *Applied Interval Analysis*. Springer Editors, 2001.
- [4] E. Kaucher. *Interval Analysis in the Extended Interval Space IIR*, Computing Suppl. 2 , pp. 33–49, 1980.
- [5] M. Markov. *Isomorphic Embeddings of Abstract Interval Systems*. Reliable Computing 3: 199–207, 1997.
- [6] N. Revol. *Introduction à l'arithmétique par intervalles*, research report RR 2001-41, LIP, École Normale Supérieure de Lyon and INRIA research report RR-4297, 2001.
- [7] N. Ramdani. *Méthodes Ensemblistes pour l'estimation*, Habilitation à Diriger des Recherches, Université Paris XII, 2005.