# AN EXAMPLE OF A TEICHMÜLLER DISK IN GENUS 4 WITH DEGENERATE KONTSEVICH-ZORICH SPECTRUM

GIOVANNI FORNI AND CARLOS MATHEUS

ABSTRACT. We construct an orientable holomorphic quadratic differential on a Riemann surface of genus 4 whose  $SL(2, \mathbb{R})$ -orbit is closed and has a highly degenerate Kontsevich-Zorich spectrum. This example is related to a previous similar construction in genus 3 by the first author.

## 1. INTRODUCTION

The goal of this note is the construction of an orientable quadratic differential q on a Riemann surface M of genus 4 such that the Kontsevich-Zorich cocycle along the  $SL(2, \mathbb{R})$ -orbit of (M, q) has a *totally degenerate* Kontsevich-Zorich spectrum, in the sense that all non-trivial Lyapunov exponents of the Kontsevich-Zorich cocycle are equall to zero. A similar example in genus 3 appeared in [6]. Our example is given by a non-primitive Veech surface in the stratum  $\mathcal{H}(2, 2, 2)$  of abelian differentials with 3 double zeros. Unpublished work of M. Möller implies that there are very few examples of this kind and only for abelian differentials on surfaces of genus  $g \leq 5$ . It is likely that the Forni's example in genus 3 [6] and the example presented here are the only examples of Veech surfaces with vanishing non-trivial Konstevich-Zorich exponents [11].

In order to explain more precisely our results, let us recall the definition of the Teichmüller flow and the Kontsevich-Zorich cocycle.

Given a surface M, we denote by  $\text{Diff}^+(M)$  the group of orientationpreserving diffeomorphisms of M and  $\text{Diff}_0^+(M)$  the connected component of the identity in  $\text{Diff}^+(M)$  (i.e.,  $\text{Diff}_0^+(M)$  is the subset of diffeomorphisms in  $\text{Diff}^+(M)$  which are isotopic to the identity).

**Definition 1.1.** Let  $Q_g$  be the *Teichmüller* space of holomorphic quadratic differentials on a surface of genus  $g \ge 1$ :

 $Q_g = \{\text{holomorphic quadratic differentials}\} / \text{Diff}_0^+(M).$ Also, let  $\mathcal{M}_g$  be the *moduli* space of holomorphic quadratic differentials on a surface of genus  $g \ge 1$ :

 $\mathcal{M}_g = \{ \text{holomorphic quadratic differentials} \} / \Gamma_g,$ 

Date: September 30, 2008.

where  $\Gamma_q$  is the mapping class group  $\Gamma_q := \operatorname{Diff}^+(M) / \operatorname{Diff}^+_0(M)$ .

2

Moreover, we note that the group  $SL(2, \mathbb{R})$  acts naturally on  $\mathcal{M}_g$  by linear transformations on the pairs of real-valued 1-forms  $(\operatorname{Re}(q^{1/2}), \operatorname{Im}(q^{1/2}))$ . The *Teichmüller* (geodesic) flow  $G_t$  is given by the action of the diagonal subgroup diag $(e^t, e^{-t})$  of  $SL(2, \mathbb{R})$  on  $\mathcal{M}_g$ .

For later reference, we recall some of the main structures of the Teichmüller space  $Q_q$  and the moduli space  $\mathcal{M}_q$ :

- $\mathcal{M}_g$  and  $Q_g$  are stratified into analytic spaces  $\mathcal{M}_{\kappa}$  and  $Q_{\kappa}$  obtained by fixing the multiplicities  $\kappa = (k_1, \ldots, k_{\sigma})$  of the zeroes  $\{p_1, \ldots, p_{\sigma}\}$  of the quadratic differentials (here  $\sum k_i = 4g 4$ );
- the total area function  $A: \mathcal{M}_g \to \mathbb{R}^+$ ,  $A(q) = \int_M |q|$  is  $SL(2, \mathbb{R})$ invariant so that the unit bundle  $\mathcal{M}_g^{(1)} := A^{-1}(1)$  and its strata  $\mathcal{M}_{\kappa}^{(1)} := \mathcal{M}_g \cap \mathcal{M}_g^{(1)}$  are  $SL(2, \mathbb{R})$ -invariant (and, *a fortiori*,  $G_t$ invariant);
- a stratum M<sub>κ</sub> of *orientable quadratic differentials* (that is, quadratic differentials obtained as squares of holomorphic 1-forms) has a locally affine structure modeled on the cohomology H<sup>1</sup>(M, Σ<sub>κ</sub>, ℂ), relative to the zero set Σ<sub>κ</sub> := {p<sub>1</sub>,..., p<sub>σ</sub>}, with local charts are given by the period map q ↦ [q<sup>1/2</sup>] ∈ H<sup>1</sup>(M, Σ<sub>κ</sub>, ℂ);
- the Lebesgue measure on the Euclidean space H<sup>1</sup>(M, Σ<sub>κ</sub>, C), appropriately normalized, induces an absolutely continuous SL(2, R)-invariant measure μ<sub>κ</sub> on M<sub>κ</sub> such that the conditional measure μ<sub>κ</sub><sup>(1)</sup> induced on M<sub>κ</sub><sup>(1)</sup> is SL(2, R)-invariant (and hence G<sub>t</sub>-invariant).

Once we get the existence of a good invariant measure  $\mu_{\kappa}^{(1)}$  for the Teichmüller flow, it is natural to ask whether  $\mu_{\kappa}^{(1)}$  has finite mass and/or  $\mu_{\kappa}^{(1)}$ is ergodic with respect to the Teichmüller dynamics. In this direction, Veech [13] showed that the strata are *not* always connected. More recently, Kontsevich and Zorich [8] (in the orientable case) and Lanneau [9] (in the non-orientable case) gave a complete classification of the connected components of all strata of holomorphic quadratic differentials. Taking this into account, we have the following result:

**Theorem 1.2** (Masur [10], Veech [12]). The total volume of  $\mu_{\kappa}^{(1)}$  is finite and the Teichmüller flow  $G_t = diag(e^t, e^{-t})$  is ergodic on each connected component of  $\mathcal{M}_{\kappa}$  with respect to  $\mu_{\kappa}^{(1)}$ .

In order to analyze the Lyapunov spectrum (i.e., the collection of the Lyapunov exponents) of the Teichmüller flow, Kontsevich and Zorich [7] introduced the following notion: the *Kontsevich-Zorich* cocycle  $G_t^{KZ}$  is the quotient of the trivial cocycle  $G_t \times id : Q_g \times H^1(M, \mathbb{R}) \to Q_g \times H^1(M, \mathbb{R})$  with respect to the action of the mapping class group  $\Gamma_q$ . It is known that

the cocycle  $G_t^{KZ}$  is *symplectic*, so that the Lyapunov spectrum of  $G_t^{KZ}$  with respect to any  $G_t$ -invariant ergodic probability  $\mu$  is *symmetric*:

$$1 = \lambda_1^{\mu} \ge \dots \lambda_g^{\mu} \ge 0 \ge -\lambda_g^{\mu} \ge \dots \ge -\lambda_1^{\mu} = -1.$$

It turns out that the g numbers  $1 = \lambda_1^{\mu} \ge \lambda_2^{\mu} \ge \cdots \ge \lambda_g^{\mu}$  appearing in the non-negative part of the Kontsevich-Zorich spectrum determines the Lyapunov spectrum of the Teichmüller flow (this is one of the motivation for introducing the Kontsevich-Zorich cocycle). Indeed, it is possible to show that, for any ergodic probability measure  $\mu$  on  $\mathcal{M}_g$  supported on a stratum of orientable quadratic differentials with  $\sigma \in \mathbb{N}$  distinct zeros, the Lyapunov spectrum of the Teichmüller flow is

$$2 = (1 + \lambda_1^{\mu}) \ge \dots \ge (1 + \lambda_g^{\mu}) \ge \overbrace{1 = \dots = 1}^{\sigma - 1} \ge (1 - \lambda_g^{\mu}) \ge \dots$$
$$\ge (1 - \lambda_2^{\mu}) \ge 0 \ge -(1 - \lambda_2^{\mu}) \ge \dots \ge -(1 - \lambda_g^{\mu})$$
$$\ge \underbrace{-1 = \dots = -1}_{\sigma - 1} \ge -(1 + \lambda_g^{\mu}) \ge \dots \ge -(1 + \lambda_1^{\mu}) = -2.$$

On the other hand, concerning the Lyapunov spectrum of the Kontsevich-Zorich cocycle, Zorich and Kontsevich conjectured that the Lyapunov exponents of  $G_t^{KZ}$  for the canonical absolutely continuous measure  $\mu_{\kappa}^{(1)}$  on any stratum of orientable quadratic differentials are all non-zero (i.e., nonuniform hyperbolicity) and distinct (i.e., all Lyapunov exponents have multiplicity 1). After the fundamental works of G. Forni [5] (showing the nonuniform hyperbolicity of  $G_t^{KZ}$ ) and Avila, Viana [1] (proving the simplicity of the Lyapunov spectrum), it follows that Zorich-Kontsevich conjecture is true. In other words, the Lyapunov exponents of a  $\mu_{\kappa}^{(1)}$ -generic point are all non-zero and they have multiplicity 1. However, it remains to understand the dynamical behavior under  $G_t^{KZ}$  of the non-generic orbits (with respect to  $\mu_{\kappa}^{(1)}$ ). In particular, one can follow Veech and ask how "degenerate" the Lyapunov spectrum of  $G_t^{KZ}$  can be along a non-typical orbit. This question was first answered by G. Forni [6] who exhibited an example of an orientable holomorphic quadratic differential q on a Riemann surface M of genus q = 3 such that the Kontsevich-Zorich spectrum of the  $SL(2,\mathbb{R})$ -invariant measure  $\mu$  supported on the (closed)  $SL(2,\mathbb{R})$ -orbit of (M,q) verifies  $\lambda_2^{\mu} = \lambda_3^{\mu} = 0.$ 

At this point, we are able to state our main result:

**Theorem 1.3.** There exists an orientable holomorphic quadratic differential q on a Riemman surface M of genus 4 such that the Lyapunov exponents (with respect to  $G_t^{KZ}$ ) of the  $SL(2, \mathbb{R})$ -invariant probability  $\mu$  supported on the  $SL(2,\mathbb{R})$ -orbit of (M,q) verifies

$$\lambda_2^\mu = \lambda_3^\mu = \lambda_4^\mu = 0.$$

The organization of this note is the following: in §2, we will recall Forni's version of the Kontsevich-Zorich formula for the sum of the Lyapunov exponents of  $G_t^{KZ}$ , in §3 we review Forni's method [6] to exlpoit symmetries and, in section §4, we will complete the proof of our theorem.

Acknowledgements: We would like to thank J.-C. Yoccoz, who motivated our work by asking whether the example in [6] could be generalized. We are also very grateful to M. Möller who introduced us to the notion of a cyclic cover, directed us to the reference [2] and suggested the correct algebraic equation of our example.

# 2. THE KONTSEVICH-ZORICH FORMULA REVISITED

Let q be a holomorphic quadratic differential on a Riemann surface M of genus  $g \ge 2$ . Fix z = x + iy a holomorphic local coordinate and write  $q = \phi(z)dz^2$ . It follows that the (degenerate) Riemannian metric  $R_q$  and the area form  $\omega_q$  induced by q are  $R_q = |\phi(z)|^{1/2}(dx^2 + dy^2)^{1/2}$ ,  $\omega_q = |\phi(z)|dx \wedge dy$ . Denote by  $S = \partial/\partial x$  and  $T = \partial/\partial y$  the horizontal and vertical directions. Define  $L_q^2(M) := L^2(M, \omega_q)$  the space of complex-valued square-integrable functions and  $H_q^1(M)$  the Sobolev space of functions  $v \in L_q^2(M)$  such that  $Sv, Tv \in L_q^2(M)$ .

**Lemma 2.1** (Forni [4], prop. 3.2). The Cauchy-Riemann operators  $\partial^{\pm} := \frac{S \pm iT}{2}$  with (dense) domain  $H^1_q(M) \subset L^2_q(M)$  are closed. Morever,  $\partial^{\pm}$  has closed range of finite codimension (equal to the genus g of M). Furthermore, denoting by  $\mathfrak{M}^{\pm}_q \subset L^2_q(M)$  the subspaces of meromorphic (resp. anti-meromorphic) functions, we have the orthogonal decompositions:

$$L^2_q(M) = \operatorname{Ran}(\partial_q^+) \oplus \mathfrak{M}_q^- = \operatorname{Ran}(\partial_q^-) \oplus \mathfrak{M}_q^+$$

Denote by  $\pi_q^{\pm}: L_q^2(M) \to \mathfrak{M}_q^{\pm}$  the orthogonal projection. Let  $H_q$  be the (non-negative definite) Hermitian form on  $\mathfrak{M}_q^+ \subset L_q^2(M)$  given by

$$H_q(m_1^+, m_2^+) := (\pi_q^-(m_1^+), \pi_q^-(m_2^+))_q := \int_M \pi_q^-(m_1^+) \cdot \overline{\pi_q^-(m_2^+)} \omega_q$$

for all  $m_1^+, m_2^+ \in \mathfrak{M}_q$ . Let

$$1 \equiv \Lambda_1(q) \ge \Lambda_2(q) \ge \dots \ge \Lambda_g(q) \ge 0$$

be the eigenvalues of  $H_q$ .

Next, we consider the sets

$$\mathcal{R}_{g}^{(1)}(k) := \{ q \in \mathcal{M}_{g}^{(1)} : \Lambda_{k+1}(q) = \dots = \Lambda_{g}(q) = 0 \}.$$

4

**Definition 2.2.** The set  $\mathcal{R}_g^{(1)}(g-1)$  is called the *determinant locus*.

The relevance of the natural filtration of sets  $\mathcal{R}_g^{(1)}(1) \subset \cdots \subset \mathcal{R}_g^{(1)}(g-1)$  for the study of the Lyapunov spectrum of  $G_t^{KZ}$  becomes evident from the following version of a formula by Kontsevich and Zorich [7] for the sum of Lyapunov exponents:

**Theorem 2.3** (Forni [5], Corollary 5.3). Let  $\mu$  be a  $SL(2, \mathbb{R})$ -invariant ergodic probability measure on  $\mathcal{M}_g^{(1)}$ . Then, the Lyapunov exponents of  $G_t^{KZ}$  with respect to  $\mu$  satisfy the formula:

$$\lambda_1^{\mu} + \dots + \lambda_g^{\mu} = \int_{\mathcal{M}_g^{(1)}} (\Lambda_1(q) + \dots + \Lambda_g(q)) d\mu(q).$$

In particular, since  $\lambda_1^{\mu} = 1 \equiv \Lambda_1(q)$ , we have

$$\lambda_2^{\mu} + \dots + \lambda_g^{\mu} = \int_{\mathcal{M}_g^{(1)}} (\Lambda_2(q) + \dots + \Lambda_g(q)) d\mu(q).$$

A direct consequence of this formula is:

**Corollary 2.4** (Forni [6], Corollary 7.1). Let  $\mu$  be any  $SL(2, \mathbb{R})$ -invariant ergodic probability measure on  $\mathcal{M}_g^{(1)}$  supported on a stratum of orientable quadratic differentials. The measure  $\mu$  is supported on the locus  $\mathcal{R}_g^{(1)}(1)$  if and only if the non-trivial Kontsevich-Zorich spectrum vanishes, that is,

$$\lambda_2^{\mu} = \cdots = \lambda_g^{\mu} = 0$$
.

#### 3. Symmetries

In this section we recall the simple method developed in [6] to derive bounds on the rank of the matrix  $H_q$  from symmetries  $\hat{E}$  of the orientable holomorphic quadratic differential q. Let  $\operatorname{Aut}(M)$  be the goup of holomorphic automorphism of the Riemann surface M. Let  $\operatorname{Aut}(q) \subset \operatorname{Aut}(M)$  is the subgroup formed by automorphisms  $a \in \operatorname{Aut}(M)$  such that  $a^*(q) = q$ . Note that there is a natural unitary action of  $\operatorname{Aut}(q)$  on  $\mathfrak{M}_q^+$  (by pull-back). Given  $a \in \operatorname{Aut}(q)$ , fix  $\{m_1^+(a), \ldots, m_g^+(a)\}$  an orthonormal basis of eigenvectors and denote by  $\{u_1(a), \ldots, u_g(a)\}$  the associated eigenvalues. Let  $B^a(q)$  be the matrix of the operator  $\pi_q^- : \mathfrak{M}_q^+ \to \mathfrak{M}_q^-$  with respect to the basis  $\{m_1^+(a), \ldots, m_g^+(a)\} \subset \mathfrak{M}_q^+$  and  $\{\overline{m_1^+(a)}, \ldots, \overline{m_g^+(a)}\} \subset \mathfrak{M}_q^-$ :

$$B_{ij}^a(q) = \int_M m_i^+(a)m_j^+(a)\omega_q.$$

For any  $I, J \subset \{1, \ldots, g\}$  with #I = #J, denote by det  $B^a_{IJ}$  the minor of the matrix  $B^a(q)$  with entries  $B^a_{ij}(q), i \in I, j \in J$ .

Lemma 3.1 (Forni [6], Lemma 7.2). The following holds:

$$\prod_{i \in I} \prod_{j \in J} u_i(a) u_j(a) \neq 1 \Longrightarrow \det B^a_{IJ}(q) = 0.$$

**Corollary 3.2.** If  $\prod_{i \in I} \prod_{j \in J} u_i(a)u_j(a) \neq 1$  for all  $I, J \subset \{1, \ldots, g\}$  with #I = #J = k, then  $rank(H_q) \leq g - k$ , hence  $q \in \mathcal{R}_g^{(1)}(k)$ .

*Proof.* Since the matrix H(q) of  $H_q$  with respect to  $\{m_1^+(a), \ldots, m_g^+(a)\}$  satisfies  $H(q) = B^a(q)^*B^a(q) = \overline{B^a(q)}B^a(q)$  (see equation (44) of [6]), the statement follows immediately from the previous lemma.

At this point, we are ready to present our genus 4 example and prove Theorem 1.3.

## 4. The example

Given an integer N > 1 and a 4-tuple of integers  $\mathbf{a} := (a_1, \ldots, a_4)$ with  $0 < a_{\mu} < N$ ,  $gcd(N, a_1, \ldots, a_4) = 1$  and  $\sum_{\mu=1}^{4} a_{\mu} \equiv 0 \pmod{N}$ . Let  $x_1, x_2, x_3, x_4 \in \mathbb{C}$  be 4 distinct points and let  $M := M_N(\mathbf{a})$  be the connected, non-singular Riemann surface determined by the algebraic equation:

$$w^N = (z - x_1)^{a_1} (z - x_2)^{a_2} (z - x_3)^{a_3} (z - x_4)^{a_4}.$$

The surface M is a cyclic cover of the Riemann sphere  $\mathbb{P}^1(\mathbb{C})$  branched over the points  $x_1, \ldots, x_4 \in \mathbb{C} \equiv \mathbb{P}^1(\mathbb{C}) \setminus \{\infty\}$ . In fact, every such cyclic cover (up to isomorphisms) can be written as above (see Example 4.2 in [2]). The surface M has genus

$$g = N + 1 - \frac{1}{2} \sum_{\mu=1}^{4} \gcd(a_{\mu}, N)$$
.

The surface M is called a cyclic cover since its automorphism group is cyclic. In fact, it is generated by the automorphism  $T: M \to M$  given by

$$T(z,w) = (z,\varepsilon w),$$

where  $\varepsilon^N = 1$  is a primitive *N*th root of unity.

For  $i \in \{1, ..., N-1\}$ , let  $L_i$  denote the eigenspace with eigenvalue  $\epsilon^i$  for the action of T on holomorphic abelian differentials:

$$L_i := \{ \omega \in H^1_{\mathrm{dR}}(M, \mathbb{C}) : T^* \omega = \varepsilon^i \cdot \omega \}.$$

Lemma 4.1 (Bouw [2], Lemma 4.3). The following formulas hold:

$$\dim_{\mathbb{C}} L_i = \sum_{\mu=1}^4 \langle \frac{i \cdot a_\mu}{N} \rangle - 1$$

(here  $\langle \cdot \rangle$  denotes the fractional part).

We recall that the genus 3 example in [6] was given by the  $SL(2, \mathbb{R})$  orbit of orientable quadratic differentials (or, equivalently, of abelian differentials) constructed from cyclic covers of type  $M_4(1, 1, 1, 1)$ .

*Proof of Theorem* 1.3. Our example is given by an orientable holomorphic quadratic differential constructed from cyclic covers of type  $M_6(1, 1, 1, 3)$ .

Let M be any branched cover of type  $M_6(1, 1, 1, 3)$ . Observe that M has genus 4. Lemma 4.1 yields:

(4.1)  
$$\dim_{\mathbb{C}} L_1 = \dim_{\mathbb{C}} L_2 = 0;$$
$$\dim_{\mathbb{C}} L_3 = \dim_{\mathbb{C}} L_4 = 1;$$
$$\dim_{\mathbb{C}} L_5 = 2.$$

Thus, we can fix  $\theta_1 \in L_3$ ,  $\theta_2 \in L_4$  and  $\theta_3, \theta_4 \in L_5$  so that  $\{\theta_1, \ldots, \theta_4\}$  is a basis of the space of holomorphic differentials on M and the action of  $T^*$  on this basis is diagonal with eigenvalues  $-1 = \varepsilon^3$  (with multiplicity 1),  $\varepsilon^4$  (with multiplicity 1) and  $\varepsilon^5$  (with multiplicity 2). Let  $q = \theta_1^2$  be the orientable quadratic differential obtained as pull-back to M of the unique holomorphic quadratic differential on  $\mathbb{P}(\mathbb{C})$  with simple poles at the branching points  $\{x_1, \ldots, x_4\}$ . It follows that the spectrum of the action of  $T \in \operatorname{Aut}(q)$  on the space  $\mathfrak{M}_q^+ \subset L_q^2(M)$  is

$$u_1(T) = 1$$
,  $u_2(T) = -\varepsilon^4$ ,  $u_3(T) = -\varepsilon^5$ ,  $u_4(T) = -\varepsilon^5$ .

Consequently,  $q \in \mathcal{R}_4^{(1)}(1)$ . In fact, a direct application of Corollary 3.2 shows that the form  $H_q$  has rank 1, so that  $q \in \mathcal{R}_4^{(1)}(1)$ .

Let  $\mathcal{V}$  be the set of all orientable quadratic differentials on a cyclic cover of type  $M_6(1, 1, 1, 3)$  branched over 4 points  $\{x_1, \ldots, x_4\} \in \mathbb{P}(\mathbb{C})$  obtained as pull-back of the unique (non-orientable) holomorphic quadratic differential on  $\mathbb{P}(\mathbb{C})$ , of total area equal to 1, with simple poles at  $x_1, \ldots, x_4$ . Since the stratum of all quadratic differentials on the Riemann sphere with 4 simple poles is  $SL(2, \mathbb{R})$ -invariant and consists of a single  $SL(2, \mathbb{R})$  orbits, the set  $\mathcal{V}$  consists of a single  $SL(2, \mathbb{R})$  orbit in the stratum of orientable quadratic differentials with 3 distinct zeros of order 4 (equivalently, of the stratum of squares of abelian differentials with 3 distinct double zeros). The above computation shows that  $\mathcal{V} \subset \mathcal{R}_4^{(1)}(1)$ , hence (in view of Corollary 2.4) the proof of Theorem 1.3 is completed.  $\Box$  We conclude by the remark that our exploration of other cyclic covers has failed to provide additional example of totally degenerate Kontsevich-Zorich spectra. We recall that according to Möller [11], there are no more examples of totally degenerate Kontsevich-Zorich spectrum among Veech surfaces in any genus.

#### REFERENCES

- [1] A. Avila and M. Viana, Simplicity of Lyapunov Spectra: Proof of the Zorich-Kontsevich conjecture, Acta Math., v. 146, no. 2, 295–344, 1997.
- [2] I. Bouw, *The p-rank of ramified covers of curves*, Compositio Math., v. 126, 295–322, 2001.
- [3] I. Bouw and M. Möller, *Teichmüller curves, triangle groups and Lyapunov exponents*, to appear in Ann. of Math.
- [4] G. Forni, Solutions of the cohomological equation for area-preserving flows on compact surfaces of higher genus, Ann. of Math., v. 146, no. 2, 295–344, 1997.
- [5] \_\_\_\_\_, Deviation of ergodic averages for area-preserving flows on surfaces of higher genus, Annals of Math., v.155, no. 1, 1–103, 2002.
- [6] \_\_\_\_\_, On the Lyapunov exponents of the Kontsevich-Zorich cocycle, Handbook of Dynamical Systems v. 1B, B. Hasselblatt and A. Katok, eds., Elsevier, 2006, 549-580.
- [7] M. Kontsevich, Lyapunov exponents and Hodge theory, in 'The mathematical beauty of physics', Saclay, 1996. Adv. Ser. Math. Phys. v. 24,318–332, World Scientific, River Edge, NJ, 1997.
- [8] M. Kontsevich and A. Zorich, Connected components of the moduli spaces of Abelian differentials with prescribed singularities, Invent. Math. v. 153, no. 3, 631-678, 2003.
- [9] E. Lanneau, Connected components of the strata of the moduli spaces of quadratic differentials, An. Sc. de l'ENS, v. 41, 1-56, 2008.
- [10] H. Masur, Interval exchange transformations and measured foliations, Annals of Mathematics, v. 115, 169-200, 1982.
- [11] M. Möller, personal communication, August 2008.
- [12] W. Veech, *Teichmüller Geodesic Flow*, Annals of Mathematics, v. 124, 441-530, 1986.
- [13] \_\_\_\_\_, Moduli spaces of quadratic differentials, J. Anal. Math., v. 55, 117–171, 1990.

GIOVANNI FORNI: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK, MD 20742-4015, USA

*E-mail address*: gforni@math.umd.edu.

CARLOS MATHEUS: COLLÈGE DE FRANCE, 3, RUE D'ULM, PARIS, CEDEX 05, FRANCE

*E-mail address*: matheus@impa.br.