

Return of the quantum cosmic censor

Shahar Hod

The Ruppin Academic Center, Emeq Hefer 40250, Israel
and

The Hadassah Institute, Jerusalem 91010, Israel

(Dated: November 17, 2021)

The influential theorems of Hawking and Penrose demonstrate that spacetime singularities are ubiquitous features of general relativity, Einstein's theory of gravity. The utility of classical general relativity in describing gravitational phenomena is maintained by the cosmic censorship principle. This conjecture, whose validity is still one of the most important open questions in general relativity, asserts that the undesirable spacetime singularities are always hidden inside of black holes. In this Letter we reanalyze extreme situations which have been considered as counterexamples to the cosmic censorship hypothesis. In particular, we consider the absorption of *fermion* particles by a spinning black hole. Ignoring quantum effects may lead one to conclude that an incident fermion wave may over spin the black hole, thereby exposing its inner singularity to distant observers. However, we show that when quantum effects are properly taken into account, the integrity of the black-hole event horizon is irrefutable. This observation suggests that the cosmic censorship principle is intrinsically a quantum phenomena.

Spacetime singularities that arise in gravitational collapse are always hidden inside of black holes, invisible to distant observers. This is the essence of the weak cosmic censorship conjecture (WCCC), put forward by Penrose forty years ago [1, 2, 3, 4]. The validity of this hypothesis is essential for preserving the predictability of Einstein's theory of gravity. The conjecture is based on the common wisdom that singularities are not pervasive [4] and it has become one of the cornerstones of general relativity. Moreover, it is being envisaged as a basic principle of nature. However, despite the flurry of research over the years, the validity of this conjecture is still an open question (see e.g. [5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23] and references therein).

The destruction of a black-hole event horizon is ruled out by this principle because it would expose the inner singularities to distant observers. Moreover, the horizon area of a black hole, A , is associated with an entropy $S_{BH} = A/4$ [24] (we use natural units in which $G = c = \hbar = 1$). Therefore, without any obvious physical mechanism to compensate for the loss of the black-hole enormous entropy, the destruction of the black-hole event horizon would violate the generalized second law (GSL) of thermodynamics [24]. For these two reasons, any process which seems, at first sight, to remove the black-hole horizon is expected to be unphysical. For the advocates of the cosmic censorship principle the task remains to find out how such candidate processes eventually fail to remove the horizon.

According to the uniqueness theorems [25, 26, 27, 28, 29], all stationary solutions of the Einstein-Maxwell equations are uniquely described by the Kerr-Newman metric which is characterized by three conserved parameters: the gravitational mass M , the angular momentum J , and the electric charge Q . A black-hole solution must satisfy

the relation

$$M^2 - Q^2 - a^2 \geq 0, \quad (1)$$

where $a \equiv J/M$ is the specific angular momentum of the black hole. Extreme black holes are the ones which saturate the relation (1). As is well known, the Kerr-Newman metric with $M^2 - Q^2 - a^2 < 0$ does not contain an event horizon, and it is therefore associated with a naked singularity rather than a black hole.

One may try to "over spin" a black hole by injecting into it particles with small energy and large angular momentum. In this work we inquire into the physical mechanism which protects the black-hole horizon from being eliminated by the absorption of waves which may "super-saturate" the extremality condition, Eq. (1). In order to analyze such processes one should study the propagation and scattering of various fields in the black-hole spacetime.

The dynamics of a wave field Ψ in the rotating Kerr-Newman spacetime is governed by the Teukolsky equation [30, 31]. One may decompose the field as

$$\Psi_{lm}(t, r, \theta, \phi) = e^{im\phi} {}_s S_{lm}(\theta; a\omega) \psi_{lm}(r; a\omega) e^{-i\omega t}, \quad (2)$$

where (t, r, θ, ϕ) are the Boyer-Lindquist coordinates, ω is the (conserved) frequency of the mode, l is the spheroidal harmonic index, and m is the azimuthal harmonic index with $-l \leq m \leq l$. The parameter s is called the spin weight of the field, and is given by $s = \pm 2$ for gravitational perturbations, $s = \pm 1$ for electromagnetic perturbations, $s = \pm \frac{1}{2}$ for massless neutrino perturbations, and $s = 0$ for scalar perturbations. (We shall henceforth omit the indices l, m , and s for brevity.) With the decomposition (2), ψ and S obey radial and angular equations, both of confluent Heun type [32, 33], coupled by a separation constant $A(a\omega)$ [34].

For the scattering problem one should impose physical boundary conditions of purely ingoing waves at the

black-hole horizon and a mixture of both ingoing and outgoing waves at infinity (these correspond to incident and scattered waves, respectively). Namely,

$$\psi \sim \begin{cases} e^{-i\omega y} + \mathcal{R}(\omega)e^{i\omega y} & \text{as } r \rightarrow \infty \text{ (} y \rightarrow \infty \text{)} ; \\ \mathcal{T}(\omega)e^{-i(\omega-m\Omega)y} & \text{as } r \rightarrow r_+ \text{ (} y \rightarrow -\infty \text{)} , \end{cases} \quad (3)$$

where the ‘‘tortoise’’ radial coordinate y is defined by $dy = [(r^2 + a^2)/\Delta]dr$, with $\Delta \equiv r^2 - 2Mr + Q^2 + a^2$. [The zeroes of Δ , $r_{\pm} = M \pm (M^2 - Q^2 - a^2)^{1/2}$, are the black hole (event and inner) horizons.] Here $\Omega = a/(r_+^2 + a^2)$ is the angular velocity of the black hole. The coefficients $\mathcal{T}(\omega)$ and $\mathcal{R}(\omega)$ are the transmission and reflection amplitudes for a wave incident from infinity.

The transmission and reflection amplitudes satisfy the usual probability conservation equation $|\mathcal{T}(\omega)|^2 + |\mathcal{R}(\omega)|^2 = 1$. The calculation of these scattering amplitudes in the low frequency limit, $M\omega \ll 1$, is a common practice in the physics of black holes, see e.g. [35, 36] and references therein. For *boson* fields ($s = 0, \pm 1, \pm 2$) one finds [22, 36]

$$|\mathcal{T}(\omega)|^2 \sim \frac{\omega - m\Omega}{\pi T_{BH}} (AT_{BH}\omega)^{2l+1} , \quad (4)$$

for the transmission probability, where $T_{BH} = (r_+ - r_-)/A$ is the Bekenstein-Hawking temperature of the black hole, and $A = 4\pi(r_+^2 + a^2)$ is its surface area.

The transmission probability, Eq. (4), implies that those modes for which the frequency ω and the azimuthal quantum number m are related by $\omega < m\Omega$ have negative transmission probabilities. These modes are actually amplified rather than absorbed. This is the well-known black-hole superradiance phenomena [37, 38]. One therefore finds that only modes for which

$$\omega > m\Omega , \quad (5)$$

can be absorbed by the black hole. Thus, it is impossible to increase the black-hole spin without increasing its mass simultaneously. This fact guarantees that the black-hole condition, Eq. (1), would still be respected after the absorption of the mode [22, 39]. One therefore concludes that the incident mode cannot remove the black-hole horizon. Cosmic censorship is therefore respected!

We have just learned that, thanks to the superradiance phenomena the black-hole is protected from being over-spun by an incident *bosonic* mode. It should be emphasized, however, that the same cannot be said if the incident mode is of a *fermion* type. It is a well-known fact that there is no superradiance effect for fermion fields [23, 40, 41, 45, 46, 47]. That is, the superradiant term $\omega - m\Omega$ is absent from the expression of the transmission probability of fermion fields. For fermion fields one simply finds [23, 36]

$$|\mathcal{T}(\omega)|^2 \sim (AT_{BH}\omega)^{2l+1} . \quad (6)$$

It may therefore seem, at first sight, that fermion particles of low-energy and high angular momentum are not hindered from entering the black hole. This has led Richartz and Saa [23] to conclude that an incident fermion wave may over spin the black hole, thereby exposing its inner singularity to distant observers.

Everything in our past experience in physics tells us that a black hole should defeat any attempt of destroying its event horizon. It seems every time we think we have finally found a sophisticated way of violating the cosmic censorship conjecture, nature still has the final word. The cosmic censorship principle asserts that nature should always provide a black hole with some physical mechanism which would protect its integrity, thereby preventing one from exposing the black-hole inner singularity.

Where should we look for the physical mechanism which may protect the cosmic censorship principle in the current case? The absence of superradiance for fermion fields is a direct consequence the Pauli exclusion principle (for fermion fields there can only be one quantum per state). Being a quantum principle, it suggests that there may be some *quantum* phenomena which protects the integrity of the black-hole horizon in the present gedanken experiment.

Indeed, it has been shown by Unruh [40, 41] that the intrinsic parity non-conservation of neutrinos (i.e., massless neutrinos have only one helicity) would lead to vacuum polarization effects about a spinning black hole. What is the physical cause for this polarization effect? In the rotating Kerr spacetime there is a gravitational spin-orbit coupling between orbiting particles and the black-hole spin. Near the black hole (more precisely, inside the ergosphere), the spin-orbit coupling becomes strong enough to create negative energy orbits (as seen from infinity) [42]. The phenomena of vacuum polarization in the Kerr spacetime is a direct consequence of the possibility of decay into such negative-energy orbits [40, 41].

The vacuum polarization effect is the wave analog of the Penrose process [43], in which rotational energy can be extracted from a rotating black hole. The process utilizes the existence of retrograde particle orbits in the ergosphere of rotating black holes, for which the energy, as it would have been measured by an observer at infinity, is negative [44]. Such orbits can not come out to infinity. However, the negative-energy particles can induce changes in the energies of other particles, which do come out to infinity.

The spontaneous polarization of the vacuum around rotating black holes involves the creation of two modes. One of these modes is co-rotating with the black hole and is characterized by a positive energy as measured at infinity, while the other one is counter rotating, having negative energy. The positive-energy particle can escape to infinity, while the negative-energy particle must fall into the black hole. An observer at infinity may detect the emitted positive-energy particle. He also measures a

decrease in the rotational energy of the black hole, caused by the infall of the retrograde, negative-energy particle into it. He therefore concludes that by the spontaneous polarization process rotational energy was extracted from the black hole [43, 44].

It is worth noting that, the fact that the energy-momentum tensor of neutrinos can have negative values inside the ergosphere [40, 41, 45, 46, 47] is in immediate contradiction with the positive energy condition. This energy criterion is commonly assumed to be valid for classical matter distributions. Thus, the spontaneous creation of particles in the Kerr spacetime is obviously a *quantum* phenomena.

It should be emphasized that the quantum polarization effect of neutrinos in the Kerr spacetime exists even for extremal, zero-temperature black holes. In fact, taking cognizance of Hawking's expression for the expected number of particles in each fermion mode of the black-hole radiation [48]

$$\langle N_{lms\omega} \rangle = |\mathcal{T}_{lms}(\omega)|^2 \{ \exp[(\omega - m\Omega)/T_{BH}] + 1 \}^{-1}, \quad (7)$$

and substituting $T_{BH} = 0$ for the extremal limit, one finds

$$\langle N_{lms\omega} \rangle = |\mathcal{T}_{lms}(\omega)|^2 \Theta(m\Omega - \omega), \quad (8)$$

where $\Theta(x)$ is the Heaviside step function.

The result Eq. (8) implies that even cold ($T_{BH} = 0$) spinning black holes emit fermion particles with frequencies below $m\Omega$. As indicated by Unruh [40, 41], this vacuum polarization effect serves to constantly decrease both the mass and the angular momentum of the black hole. It should be emphasized that only fermion modes for which $\omega < m\Omega$ are being spontaneously emitted by the extremal spinning black hole. This implies that the black hole loses angular momentum more rapidly than it loses its (squared) mass. One therefore concludes that the black hole is constantly pushing itself away from the extremal limit [that is, the dimensionless ratio $(r_+ - r_-)/r_+$ increases with time due to the spontaneous emission of neutrinos].

The important point to realize is that the Pauli exclusion principle would prevent one from beaming low-energy neutrinos (characterized by $\omega < m\Omega$) on the rotating black hole any more rapidly than they are spontaneously emitted [49]. Thus, the exclusion principle implies that one can at best suppress the constant increase of the ratio $(r_+ - r_-)/r_+$. Cosmic censorship is therefore respected!

A closely related problem is that of a rotating black hole immersed in a thermal radiation bath [50]. Taking into account the quantum character of the black hole (namely, the spontaneous polarization of the vacuum around the black hole) one finds that the probability $p(0|1)$ that one fermion mode is incident upon the black

hole with no reflection is given by [50]

$$p(0|1) = |\mathcal{T}(\omega)|^2 (1 + e^{-x})^{-1}, \quad (9)$$

where

$$x \equiv (\omega - m\Omega)/T_{BH}, \quad (10)$$

and $|\mathcal{T}(\omega)|^2$ is given by Eq. (6).

The absorption probability (9) implies that the claim of Ref. [23] that the probability to over spin a black hole is given by $|\mathcal{T}(\omega)|^2$ [as defined in Eq. (6)] is actually erroneous. The correct probability, given by Eq. (9), is actually suppressed by a factor of $(1 + e^{-x})^{-1}$. It is clear that in order to over spin the black hole one must consider an incident mode characterized by $\omega < m\Omega$ (that is, a mode with $x < 0$). For extremal black holes (characterized by $T_{BH} = 0$) this implies $x \rightarrow -\infty$, and one therefore finds from Eq. (9) that $p(0|1) = 0$. The incident mode is therefore reflected with probability 1. In particular, it fails to over spin the extremal black hole.

Suppose we have a near extremal black hole with mass M and angular momentum $J = M^2 - 1$ (this is the "nearest extreme" black hole considered in [23]). Of course, our analysis is meaningful only in the semiclassical regime, where $M \gg 1$. In this limit, the temperature of the near-extremal black hole is given by $T_{BH} \simeq \sqrt{2}/4\pi M^2$. Consider an incident mode with small frequency and $l = m = 3/2$. In this case $x = -3\pi M/\sqrt{2}$. Taking cognizance of Eq. (9) one finds that the probability of such a 'dangerous' mode to be absorbed by the black hole is

$$p(0|1) = |\mathcal{T}(\omega)|^2 e^{-3\pi M/\sqrt{2}}. \quad (11)$$

It is obvious that the absorption probability is extremely small in the semiclassical regime $M \gg 1$ considered here. For example, for a black hole of mass $M = 1\text{gr}$, the suppression factor $(1 + e^{-x})^{-1}$ is $\sim 10^{-133009}$ [51]. The number 10^{133009} is much larger than the total number of particles in the whole universe. It is therefore clear that the absorption probability of a dangerous mode is practically zero.

Moreover, such dangerous absorptions (with extremely low probability) are not cumulative—most of the incident modes will merely be scattered off the black hole. In being scattered they will always radiate into the black hole some gravitational waves [38] which will push the black hole away from the extremal limit.

In summary, extreme situations which have been considered as counterexamples to the weak cosmic censorship conjecture were reexamined. In particular, we have re-analyzed gedanken experiments in which *fermion* waves are beamed from far away towards a near-extremal rotating black hole. The unique feature which characterizes fermion fields is the absence of the superradiance phenomena. It therefore seems, at first sight, that

fermion modes of low energy and large angular momentum can be absorbed by the black hole. One may thus give in to the temptation of concluding that the black hole can acquire enough angular momentum to over spin, $M^2 - Q^2 - a^2 < 0$. Previous analyzes [23] indeed claimed that this process may provide a counterexample to the WCCC. However, we have demonstrated that a more complete analysis of the gedanken experiment (in which *quantum* effects are properly taken into account) reveals that it does *not* violate the weak cosmic censorship conjecture.

The physical mechanism which protects the integrity of the black-hole horizon in this gedanken experiment is the spontaneous emission of low-energy ($\omega < m\Omega$) fermions by the rotating black hole (quantum instability of the vacuum around a spinning object). This vacuum polarization effect was discovered by Unruh in the 70's after performing a detailed analysis of the second quantization of fields in the rotating Kerr spacetime [40, 41]. It is interesting to note that, historically one could have *predicted* the existence of such a spontaneous quantum emission already in the 60's, following the formulation of the cosmic censorship principle. We thus conclude that the cosmic censor must be cognizant of both general relativity and quantum physics.

ACKNOWLEDGMENTS

This research is supported by the Meltzer Science Foundation. I thank Jacob D. Bekenstein, Tsvi Piran and Liran Shimshi for stimulated discussions. I also thank Yael Oren for helpful discussions.

[1] S. W. Hawking and R. Penrose, Proc. R. Soc. London A **314**, 529 (1970).
 [2] R. Penrose, Riv. Nuovo Cimento **1**, 252 (1969); in *General Relativity, an Einstein Centenary Survey*, edited by S. W. Hawking and W. Israel (Cambridge University Press, Cambridge, England, 1979).
 [3] S. W. Hawking, Phys. Rev. D **14**, 2460 (1976).
 [4] P. R. Brady, I. G. Moss, and R. C. Myers, Phys. Rev. Lett. **80**, 3432 (1998).
 [5] R. M. Wald, "Gravitational Collapse and Cosmic Censorship", e-print gr-qc/9710068.
 [6] T. P. Singh, "Gravitational Collapse, Black Holes and Naked Singularities", e-print gr-qc/9805066.
 [7] C. J. S. Clarke, Classical and Quantum Gravity **11**, 1375 (1994).
 [8] C. V. Vishveshwara, Phys. Rev. D **1**, 2870 (1970).
 [9] R. Price, Phys. Rev. D **5**, 2419 (1972); **5**, 2439 (1972).
 [10] R. Wald, Ann. Phys. (N.Y.) **83**, 548 (1974).
 [11] W. A. Hiscock, Ann. Phys. (N.Y.) **131**, 245 (1981).
 [12] B. S. Kay and R. M. Wald, Classical and Quantum Gravity **4**, 893 (1987).
 [13] J. D. Bekenstein and C. Rosenzweig, Phys. Rev. D **50**, 7239 (1994).

[14] V. E. Hubeny, Phys. Rev. D **59**, 064013 (1999).
 [15] T. C. Quinn and R. M. Wald, Phys. Rev. D **60**, 064009 (1999).
 [16] S. Hod, Phys. Rev. D **60**, 104031 (1999).
 [17] S. Hod, e-print gr-qc/9908004; S. Hod and T. Piran, Gen. Relativ. Gravit. **32**, 2333 (2000).
 [18] S. Hod, Phys. Rev. D **66**, 024016 (2002).
 [19] L. H. Ford and T. A. Roman, Phys. Rev. D **41**, 3662 (1990).
 [20] L. H. Ford and T. A. Roman, Phys. Rev. D **46**, 1328 (1992).
 [21] G. E. A. Matsas and A. R. R. da Silva, Phys. Rev. Lett. **99**, 181301 (2007).
 [22] S. Hod, Phys. Rev. Lett. **100**, 121101 (2008).
 [23] M. Richartz and A. Saa, e-print arXiv:0804.3921.
 [24] J. D. Bekenstein, Phys. Rev. D **9**, 3292 (1974).
 [25] W. Israel, Phys. Rev. **164**, 1776 (1967); Commun. Math. Phys. **8**, 245 (1968).
 [26] B. Carter, Phys. Rev. Lett. **26**, 331 (1971).
 [27] S. W. Hawking, Commun. Math. Phys. **25**, 152 (1972).
 [28] D. C. Robinson, Phys. Rev. D **10**, 458 (1974); Phys. Rev. Lett. **34**, 905 (1975).
 [29] J. Ispert, Phys. Rev. Lett. **27**, 529 (1971).
 [30] S. A. Teukolsky, Phys. Rev. Lett. **29**, 1114 (1972); Astrophys. J. **185**, 635 (1973).
 [31] A. L. Dudley and J. D. Finley, J. Math. Phys. **20**, 311 (1979).
 [32] A. Ronveaux, *Heun's differential equations*. (Oxford University Press, Oxford, UK, 1995).
 [33] C. Flammer, *Spheroidal Wave Functions* (Stanford University Press, Stanford, 1957).
 [34] The functions $S(\theta; a\omega)$ are the spheroidal wave functions [30, 33]. In the $a\omega \ll 1$ limit they become the familiar spin-weighted spherical harmonics with the corresponding angular eigenvalues ${}_sA_{lm} = l(l+1) - s(s+1) + O(a^2\omega^2)$.
 [35] S. Chandrasekhar, *The mathematical theory of black holes* (Oxford University Press, Oxford, UK, 1985).
 [36] D. N. Page, Phys. Rev. D **13**, 198 (1976).
 [37] W. H. Press and S. A. Teukolsky, Nature **238**, 211 (1972).
 [38] J. D. Bekenstein, Phys. Rev. D **7**, 949 (1973).
 [39] It has been claimed [21] that this process may nevertheless push a near-extremal, charged (Reissner-Nördstrom) black hole over the extremal limit. This claim was based on the fact that the "bare" black hole has no angular momentum ($\Omega = 0$). This may suggest that one can inject waves of low energy and large angular momentum into the black hole. The authors of [21] therefore concluded that such processes may serve as counterexamples to the WCCC.
 It should be stressed, however, that previous analyzes [21] considered only the zeroth-order interaction between the black hole and the incident wave. That is, the wave was assumed to propagate on a *fixed* (unperturbed) Reissner-Nördstrom background. It is important to realize that while the field spirals into the black hole it interacts with the black hole, so the horizon generators start to rotate. This implies that, even if the initial ("bare") black hole was a non-rotating one (as assumed in [21]), the field would "ignite" its rotation, such that the propagation of the field itself is actually taking place on a slowly rotating perturbed spacetime.
 However small, these higher-order backreaction effects turn out to be a crucial ingredient of the analysis. In par-

ticular, we have demonstrated [22] that a more complete analysis of the gedanken experiments (in which backreaction effects are properly taken into account) reveals that they do *not* violate the WCCC.

- [40] W. Unruh, Phys. Rev. Lett. **31**, 1265 (1973).
- [41] W. Unruh, Phys. Rev. D **10**, 3194 (1974).
- [42] B. Carter, Phys. Rev. **174**, 1559 (1968).
- [43] R. Penrose, Riv. Nuovo Cimento **1**, 252 (1969).
- [44] T. Piran and J. Shaham, Phys. Rev. D **16**, 1615 (1977).
- [45] R. Güven, Phys. Rev. D **16**, 1706 (1977).
- [46] S. M. Wagh and N. Dadhich, Phys. Rev. D **32**, 1863 (1985).
- [47] A. Gould, Phys. Rev. D **35**, 449 (1987).
- [48] S. W. Hawking, Commun. Math. Phys. **43**, 199 (1975).
- [49] Note that one can still send high-energy neutrinos (characterized by $\omega > m\Omega$) into the black hole. However, such high-energy modes would certainly push the black hole away from the extremal limit. [That is, they will increase the ratio $(r_+ - r_-)/r_+$.] Thus, high-energy particles pose no challenge to the cosmic censorship conjecture.
- [50] J. D. Bekenstein, Phys. Rev. D **15**, 2775 (1977).
- [51] It is worth emphasizing that the pre-factor $|\mathcal{T}(\omega)|^2$ is also very small, see Eq. (6).