Planar aggregation and the coalescing Brownian flow

James Norris ¹ & Amanda Turner ²

November 3, 2018

Abstract

We study a scaling limit associated to a model of planar aggregation. The model is obtained by composing certain independent random conformal maps. The evolution of harmonic measure on the boundary of the cluster is shown to converge to the coalescing Brownian flow.

1 Introduction

A simplified model of aggregation in two dimensions may be formulated as follows. Let K_0 be the closed disc of radius 1, with centre at the origin, and let P_1 be another closed disc, of diameter $\delta > 0$ say, tangent to K_0 at a point chosen uniformly at random. Set $D_0 = (\mathbb{C} \cup \{\infty\}) \setminus K_0$ and $D_1 = (\mathbb{C} \cup \{\infty\}) \setminus K_1$, where $K_1 = K_0 \cup P_1$. Write F_1 for the unique conformal isomorphism $D_0 \to D_1$ such that $F_1(\infty) = \infty$ and $F'_1(\infty) > 0$. Suppose then that F_2, F_3, \ldots are independent and identically distributed copies of F_1 , and define

$$D_n = F_1 \circ \cdots \circ F_n(D_0).$$

The complementary sets $K_n = \mathbb{C} \setminus D_n$ form an increasing family and may be considered as a model of aggregation, where, for each $n \ge 0$, we add at time n + 1 a new particle $P_{n+1} = K_{n+1} \setminus K_n$. Note that P_{n+1} is the image under the conformal map $F_1 \circ \cdots \circ F_n$ of a disc distributed as P_1 and independent of K_n . Since harmonic measure is invariant under conformal maps, conditional on K_n , the random point at which P_{n+1} is attached to K_n is distributed on the boundary of K_n according to the normalized harmonic measure from infinity. In this respect, the model is appropriate for the aggregation of diffusive particles 'coming in from infinity'. We note however that the added particle is distorted in a way depending on the current cluster K_n .

¹Statistical Laboratory, Centre for Mathematical Sciences, Wilberforce Road, Cambridge, CB3 0WB, UK

²Department of Mathematics and Statistics, Fylde College, Lancaster University, Lancaster, LA1 4YF, UK

In this paper, we identify a scaling limit for the restrictions to the boundary of D_0 of the maps $(F_m \circ \cdots \circ F_n)^{-1}$, for $m \leq n$, as $m, n \to \infty$ and $\delta \to 0$ in a suitable way. This can be thought of as identifying the time-evolution of the total harmonic measure carried by the various 'fingers' of the growing cluster. The limit is not sensitive to the shape of the first added particle: although we specified discs in the description above, the same sort of limit applies more generally. The limit object is the flow of coalescing Brownian motions on the circle, also known as the Brownian web.

A brief review of related work is given next, followed by some illustrations of typical clusters, for certain cases of the model. From Section 4 on, we confine our attention to the restrictions of maps to the boundary of D_0 , generalizing at the same time to a natural class of Lévy random flows on the circle. We first show a weak convergence result for the time-evolutions of finitely many points. In Section 5 we describe a new canonical space for the coalescing Brownian flow. Then, in Section 6, a larger flow-space, of Skorokhod type, is introduced. This is a complete separable metric space suitable for the formulation of weak convergence of stochastic flows which are not necessarily continuous in time or space. The convergence of Lévy flows to the coalescing Brownian flow, is shown in Section 7. Time reversal of flows is discussed in Section 8. Then, in Section 9, convergence results for the aggregation model are deduced as corollaries of the convergence of Lévy flows. Some technical details are left to an Appendix, where we discuss, in particular, the relation between our formulation of the coalescing Brownian flow and the Brownian webs of Fontes et al.

2 Review of related work

2.1 Coalescing Brownian motions

It is straightforward to define a finite family of (standard) Brownian motions in one dimension, with given space-time starting points, independent until they collide, and coalescing on collision. The possibility to extend such a model to a space-time continuum of starting points was shown by Arratia in 1979 in his PhD thesis [1], where the model was considered as a limit object for coalescing random walks on the integers. Subsequent work has been carried out by many people including Harris [8], who was interested in general coalescing stochastic flows, and Piterbarg [14], who showed that Arratia's flow arises as a weak limit of rescaled isotropic stochastic flows. Further properties were developed by Tóth and Werner [16] in 1998, who used the flow to contruct an object which they called the 'continuous true self-repelling motion'. Tsirelson [17] studied Arratia's flow, formulated in terms of L^2 -spaces, rather than pathwise, as an example in the general theory of stochastic flows. Fontes, Isopi, Newman and Ravishankar [6, 7] introduced the name 'Brownian web' in 2004, to describe a number of new formulations of Arratia's flow, and gave further characterization and convergence results. The paper [6] characterizes the Brownian web as a random element of a space of compact collections of paths with specified starting points.

In this paper we follow most closely the viewpoint of Tóth and Werner but lay greater stress on the almost sure flow-type properties, formulating Arratia's flow as a random variable in a certain complete separable metric space of continuous weak flows. We show that there is a unique Borel probability measure on this space with respect to which all the *n*-point motions are coalescing Brownian motions, and that this measure is invariant under time reversal. Moreover, we show that any sequence of random such flows, all of whose *n*-point motions converge to coalescing Brownian motions, converges to the coalescing Brownian flow. The exact correspondence between our work and that in [6] is discussed in the Appendix.

2.2 Planar random growth

Our motivation for looking at the coalescing Brownian flow arises from a surprising connection with planar random growth processes.

The simplest sorts of planar growth process to formulate take values in finite subsets of \mathbb{Z}^2 , starting from a singleton at the origin, and grow by the successive addition of sites adjacent to the present cluster, which are chosen according to some distribution determined by the present cluster. In 1961, Eden [5] introduced one such process, where the added site is simply chosen uniformly from all adjacent sites. This has been considered as a model for the growth of bacterial cells or tissue cultures of cells that are constrained from moving. In 1981, Witten and Sander [18] put forward another such process, known as diffusion-limited aggregation or DLA. Here, particles perform random walks 'starting from infinity' until they reach a site adjacent to the cluster. This site is then added to the cluster. DLA is considered as a model for the formation of aggregates by deposition, such as soot particles. A family of further processes, indexed by a parameter $\eta \in [0, 1]$ and called dielectric-breakdown models, were discussed by Niemeyer et al. [13]. In these processes, the random walk hitting probabilities on sites adjacent to the cluster are interpreted as an electric field E; the probabilities of attachment for new sites are then chosen proportional to E^{η} . Thus the case $\eta = 0$ is the Eden model and the case $\eta = 1$ is DLA.

The primary interest in these and other related processes is in the asymptotic behaviour of large clusters. Computational investigations reveal the emergence of complex structures, of fractal type, which sometimes appear a good match for observed phenomena. However, such investigations are highly sensitive to model variations and there are few notable mathematical results, with the exception of Kesten's 1987 growth estimate [11] for DLA. Moreover, the simulations appear to show that clusters depend on the fine lattice structure at large scales. This suggests that lattice-based processes may never properly model continuum phenomena.

In 1998, Hastings and Levitov [9] formulated a family of continuum growth models in terms of sequences of iterated conformal maps, indexed by a parameter $\alpha \in [0, 2]$. As in the

model described in the Introduction, they identify the cluster after the arrival of n particles with a conformal map $\Phi_n = F_1 \circ \cdots \circ F_n$ on D_0 , where, conditional on Φ_n , the map F_{n+1} corresponds to a particle attached at a random point z_{n+1} on the unit circle, but now of radius $\delta_{n+1} = \delta_0 |\Phi'_n(z_{n+1})|^{-\alpha}$. Hastings and Levitov argue, by comparing local growth rates, that their model can be related to lattice dielectric-breakdown by setting $\eta = \alpha - 1$. Further exploration of this relation is discussed in the survey paper by Bazant and Crowdy [2].

Carleson and Makarov [4], in 2001, obtained a growth estimate for a deterministic analogue of the DLA model. In 2005, Rohde and Zinsmeister [15] considered the case $\alpha = 0$ in the Hastings-Levitov family. They established a scaling limit, in the case where the particle size is small (but fixed), as the number of particles tends to infinity, and showed that the limit sets were one dimensional. They also gave estimates for the dimension of the limit sets in the case of general α , and discussed limits of deterministic variants.

We also consider the case $\alpha = 0$ but in the limiting regime where the particle diameter δ becomes small and the number of particles is of the order δ^{-3} . We then show that the resulting flow map, restricted to points on the unit circle, converges to the coalescing Brownian flow.

3 Some illustrations

In the Introduction we described the construction of a sequence of iterated conformal maps that can be interpreted as a model of the aggregation of diffusive particles 'coming in from infinity'. We chose the basic particle P_1 to be a closed disc of diameter $\delta > 0$. However our analysis turns out to apply to a wide range of basic particle shapes, as we shall see in Section 9. Recall that D_0 denotes the complement of the closed unit disc K_0 in $\mathbb{C} \cup \{\infty\}$ and D_1 denotes the complement of K_1 in $\mathbb{C} \cup \{\infty\}$, where $K_1 = K_0 \cup P_1$. Write F_1 for the conformal isomorphism $D_0 \to D_1$ fixing ∞ , with $F'_1(\infty) > 0$ and write G_1 for the inverse isomorphism $D_1 \to D_0$. Denote the upper half-plane by H.

We now discuss briefly the form of G_1 for two particular choices of P_1 . The first of these, corresponding to the slit $[1, 1 + \delta]$, was used to generate some realizations of the cluster for various values of δ , which are presented in Figure 1.

For the slit, we can obtain the map G_1 by composing the sequence of maps

$$D_1 \to H \setminus (0, i\sqrt{t}] \to H \to H \to D_0,$$

given by

$$g_1(z) = i\frac{z-1}{z+1}, \quad g_2(z) = \sqrt{z^2+t}, \quad g_3(z) = \lambda z, \quad g_4(z) = \frac{i+z}{i-z}$$

Here, we choose $t = \delta^2/(2+\delta)^2$ so that $g_2 \circ g_1(1+\delta) = g_2(i\delta/(2+\delta)) = 0$ and choose $\lambda = 1/\sqrt{1-t}$ so that $g_3 \circ g_2 \circ g_1(\infty) = g_3 \circ g_2(i) = \lambda i \sqrt{1-t} = i$. Then $G_1(1+\delta) = 1$ and $G_1(\infty) = \infty$, as required. The maps extend to the boundaries of their domains in obvious



(a) The cluster after a few arrivals with $\delta=1.$



(c) The cluster after 800 arrivals with $\delta=0.1.$



(e) The cluster after 20000 arrivals with $\delta = 0.02$.



(b) The cluster after 100 arrivals with $\delta = 1$.



(d) The cluster after 5000 arrivals with $\delta = 0.04$.



(f) The stochastic flow $(X_{t0})_{t \in [0,1]}$ with $\delta = 0.02$.

Figure 1: The slit model case of simplified Hastings-Levitov DLA

ways. In particular the boundary of the slit maps under G_1 to a boundary arc $[-\theta_{\delta}, \theta_{\delta}]$ of the unit circle. Note that $G_1(1) = g_4(\pm \lambda \sqrt{t})$ and $g'_4(0) = -2i$, from which we see that

$$\frac{\theta_{\delta}}{\delta} \to 1 \quad \text{as} \quad \delta \to 0.$$
 (1)

We observe in Figure 1(b), when $\delta = 1$, that incoming particles are markedly distorted and, in particular, that particles arriving later tend to be larger. This effect is diminished when we examine smaller values of δ . In Figure 1(e), the cluster is a rough ball, but with some sort of internal structure. The colours label arrivals in different epochs, showing that growth over time is uniform and there is a close relationship between time of arrival and the distance from the origin at which the particle sticks. Figure 1(f) focuses on the motion of points on the original circular boundary, which is the aspect of these simulations examined theoretically in this paper. The coalescing motion displayed agrees with our theoretical predictions. The size of the gaps between the flow lines gives the amount of harmonic measure carried on fingers of the cluster based between the chosen points on the unit circle.

In the case where P_1 is the lune $L = \{z \in \mathbb{C} : |z - 1| \leq \delta, |z| > 1\}$, the map G_1 is given by

$$G_1(z) = \frac{(e^{i\theta}z - 1)^a - (e^{-i\theta}z - 1)^a}{(z - e^{-i\theta})^a - (z - e^{i\theta})^a},$$

where

$$\theta = 2 \tan^{-1} \delta \sqrt{1 - \delta^2/4}, \quad a = \frac{\pi}{\pi - \cos^{-1}(\delta/2)}$$

We shall use in Section 9 the following estimate of the logarithmic capacity of K_1 for this case. As $\delta \to 0$,

$$\operatorname{cap}(K_1) = -\log \lim_{z \to \infty} \frac{G_1(z)}{z} = \log \frac{a \sin \theta}{\sin(a\theta)} = 2\delta^2 + o(\delta^2).$$
(2)

4 A class of Lévy flows on the circle

We introduce a class of random flows on the circle, whose distributions are invariant under rotations of the circle, and, under which, each point on the circle performs a Lévy process of mean 0 with no diffusive part. The flow maps are in general not continuous on the circle, but have a non-crossing property. In a certain asymptotic regime, the motion of the flow from a countable family of starting points is shown to converge weakly to a family of coalescing Brownian motions. We specify a particular flow by the choice of a non-decreasing, right-continuous function $f^+ : \mathbb{R} \to \mathbb{R}$ with the following *degree* 1 property³

$$f^+(x+n) = f^+(x) + n, \quad x \in \mathbb{R}, \quad n \in \mathbb{Z}.$$
(3)

Denote the set of such functions by \mathcal{R} and write \mathcal{L} for the analogous set of left-continuous functions. Each $f^+ \in \mathcal{R}$ has a left-continuous modification $f^- \in \mathcal{L}$, given by $f^-(x) = f(x)$. Write \mathcal{D} for the set of all pairs $f = \{f^-, f^+\}$. When f^+ is continuous, we shall also write $f = f^+$ and, generally, we write f in place of f^{\pm} in expressions where the choice of left or rightcontinuous modification makes no difference to the value. For $f^+ \in \mathcal{R}$ and $z = \tilde{z} + \mathbb{Z} \in \mathbb{R}/\mathbb{Z}$, define $(\tau_z f)^+ \in \mathcal{R}$ by

$$(\tau_z f)^+(x) = \tilde{z} + f^+(x - \tilde{z}), \quad x \in \mathbb{R}.$$

The degree 1 property ensures that there is no dependence on the choice of representative \tilde{z} . The sets \mathcal{R} and \mathcal{L} are closed under composition, but \mathcal{D} is not. In fact, if $f_1, f_2 \in \mathcal{D}$, then $f_2^- \circ f_1^-$ is the left-continuous modification of $f_2^+ \circ f_1^+$ if and only if f_1 sends no interval of positive length to a point of discontinuity of f_2 . We say in this case that $f_2 \circ f_1 \in \mathcal{D}$, denoting by $f_2 \circ f_1$ the pair $\{f_2^- \circ f_1^-, f_2^+ \circ f_1^+\}$. Define $\operatorname{id}(x) = x$ and set $\mathcal{D}^* = \mathcal{D} \setminus \{\operatorname{id}\}$. We assume throughout that our basic map $f \in \mathcal{D}^*$. Write \tilde{f}^\pm for the periodic functions $\tilde{f}^\pm(x) = f^\pm(x) - x$. Define constants $\rho = \rho(f) > 0$ and $\beta = \beta(f) \in \mathbb{R}$ by

$$\rho \int_0^1 \tilde{f}(x)^2 dx = 1, \quad \beta = \rho \int_0^1 \tilde{f}(x) dx.$$
(4)

We now define a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Take Ω to be the set of those integer-valued Borel measures on $\mathbb{R} \times (\mathbb{R}/\mathbb{Z})$ which are finite on bounded sets, and also have the property that $\omega(\{t\} \times (\mathbb{R}/\mathbb{Z})) \in \{0, 1\}$ for all $t \in \mathbb{R}$. We interpret the first factor \mathbb{R} of the underlying space as time. Write \mathcal{F}^o for the σ -algebra on Ω generated by evaluations on Borel sets. For intervals $I \subseteq \mathbb{R}$, write \mathcal{F}^o_I for the σ -algebra on Ω generated by evaluations on Borel subsets of $I \times (\mathbb{R}/\mathbb{Z})$. There is a unique probability measure $\mathbb{P} = \mathbb{P}^\rho$ on (Ω, \mathcal{F}^o) which makes the coordinate map $\mu(\omega, dt, dz) = \omega(dt, dz)$ into a Poisson random measure on $\mathbb{R} \times (\mathbb{R}/\mathbb{Z})$ with intensity $\nu(dt, dz) = \rho dt dz$. Here dz denotes Lebesgue measure on \mathbb{R}/\mathbb{Z} : we shall use, without further comment, in defining integrals, the obvious identification of \mathbb{R}/\mathbb{Z} and [0, 1). Write \mathcal{F} for the completion of \mathcal{F}^o with respect to \mathbb{P} , extending \mathbb{P} to \mathcal{F} as usual.

We first construct the flow in the case where $\beta = 0$. Given $\omega \in \Omega$, for each $t \in \mathbb{R}$ define $F_t = \{F_t^-, F_t^+\} \in \mathcal{D}$ by

$$F_t^{\pm} = \begin{cases} (\tau_z f)^{\pm}, & \text{if } \omega \text{ has an atom at } (t, z), \\ \text{id}, & \text{otherwise.} \end{cases}$$

³These functions can be considered as liftings of maps from the circle \mathbb{R}/\mathbb{Z} to itself having an orderpreserving property. In practice, the circle map will be a perturbation of the identity map and our basic map f^+ will be the unique lifting which is close to the identity map on \mathbb{R} .

For each finite interval $I \subseteq \mathbb{R}$, define $X_I^+ \in \mathcal{R}$ and $X_I^- \in \mathcal{L}$ by

$$X_I^{\pm} = F_{T_n}^{\pm} \circ \dots \circ F_{T_1}^{\pm},$$

where $T_1 < \cdots < T_n$ are the times of the atoms of ω in $I \times (\mathbb{R}/\mathbb{Z})$. We take $X_I^{\pm} = \text{id}$ if there are no such atoms.

In the case where $\beta \neq 0$, we replace ω in the preceding construction by ω^{β} , given by

$$\omega^{\beta}(dt, dz) = \omega(dt, d(z + \beta t)),$$

to obtain $X_I^{\beta,\pm}$, and then set

$$X_I^{\pm}(x) = X_I^{\beta,\pm}(x+\beta s) - \beta t,$$

where $s = \inf I$ and $t = \sup I$.

Since f can have at most countably many points of discontinuity and intervals of constancy, and since, under \mathbb{P} , the positions of the atoms of ω are distributed uniformly on \mathbb{R}/\mathbb{Z} , we have, almost surely, $X_I = \{X_I^-, X_I^+\} \in \mathcal{D}$ for all intervals I. Write $I = I_1 \oplus I_2$ if I_1, I_2 are disjoint intervals with sup $I_1 = \inf I_2$ and $I = I_1 \cup I_2$. Note that $(X_I : I \subseteq \mathbb{R})$ has the following properties:

$$X_I^+(x)$$
 and $X_I^-(x)$ are random variables for all finite intervals I and all $x \in \mathbb{R}$, (5)

$$X_I^+ = X_{I_2}^+ \circ X_{I_1}^+ \text{ and } X_I^- = X_{I_2}^- \circ X_{I_1}^- \text{ whenever } I = I_1 \oplus I_2,$$
 (6)

$$X^{+}_{(s,t)}(x) = X^{-}_{(s,t)}(x) = x - \beta(t-s) \text{ for all } x \in \mathbb{R}, \text{ eventually as } s \uparrow t \text{ or } t \downarrow s.$$
(7)

Fix $e = (s, x) \in \mathbb{R}^2$, and define two processes $X_t^{e,-}$ and $X_t^{e,+}$, both starting from e, by setting $X_t^{e,\pm} = X_{(s,t]}^{\pm}(x)$ for $t \ge s$. Then $X^{e,-}$ and $X^{e,+}$ are both piecewise continuous, cadlag, and satisfy the integral equations

$$X_t^{e,\pm} = x + \int_{(s,t]\times[0,1)} \tilde{f}^{\pm} (X_{r-}^{e,\pm} - z)(\mu - \nu)(dr, dz), \quad t \ge s$$

Under \mathbb{P} , we have $X_t^{e,-} = X_t^{e,+}$ for all $t \ge s$, almost surely. We shall therefore drop the \pm from now on and write simply X^e . Write μ_e^f for the distribution of X^e under \mathbb{P} on the Skorokhod space $D_e = D_x([s,\infty),\mathbb{R})$ of cadlag paths starting from x at time s. Write μ_e for the distribution on D_e of a standard Brownian motion starting from e.

In the context of planar aggregation f describes the action, on the boundary of the unit disc, of the conformal map corresponding to the arrival of a particle. This action is described precisely in Section 9. Where particles are symmetric around the axis through their attachment point, $\beta = 0$. Asymmetric particles can give rise to non-zero values of β and in these cases a drift is induced for which we have compensated. **Proposition 4.1.** We have $\mu_e^f \to \mu_e$ weakly on D_e , uniformly in $f \in \mathcal{D}^*$ as $\rho(f) \to \infty$.

Proof. We drop the superscript e within the proof to lighten the notation. Under \mathbb{P} , the process $(X_t)_{t \ge s}$ is a martingale. In fact, it is a Lévy process, with characteristic exponent χ given by

$$\chi(\theta) = \rho \int_0^1 \left\{ e^{i\theta \tilde{f}(z)} - 1 - i\theta \tilde{f}(z) \right\} dz, \quad \theta \in \mathbb{R}.$$

In particular, we have $\mathbb{E}(|X_{t_1} - X_{t_2}|^2) = |t_1 - t_2|$ for all $t_1, t_2 \ge s$. A standard criterion (see for example [3, page 143] or [10, page 355]) allows us to deduce that the family of laws $(\mu_e^f : f \in \mathcal{D}^*)$ is tight in D_e . We note that the process $(X_t^2 - t)_{t\ge s}$ is also a martingale, and that the jumps of $(X_t)_{t\ge s}$ are bounded in absolute value by $\|\tilde{f}\| = \sup_{x\in\mathbb{R}} |\tilde{f}(x)|$. Let μ be any weak limit law for the limit $\rho \to \infty$. Write $(Z_t)_{t\ge s}$ for the coordinate process on D_e . Under μ , by standard arguments, both $(Z_t)_{t\ge s}$ and $(Z_t^2 - t)_{t\ge s}$ are local martingales in the natural filtration of $(Z_t)_{t\ge s}$. By using the fact that f is non-decreasing, we can obtain for all $\rho \ge 1$ the estimate

$$\|\tilde{f}\| \leqslant 2\rho^{-1/3}.\tag{8}$$

Hence μ is supported on continuous paths and must therefore be μ_e by Lévy's characterization of Brownian motion.

We next fix a sequence $E = (e_k : k \in \mathbb{N})$ in \mathbb{R}^2 , where $e_k = (s_k, x_k)$ say, and consider the sequence of processes $X^E = (X^k : k \in \mathbb{N})$, where $X^k = X^{e_k}$. Then X^E is a random variable in the complete separable metric space $D_E = \prod_{k=1}^{\infty} D_{e_k}$, where we define the metric d_E on D_E by

$$d_E(\xi,\xi') = \sum_{k=1}^{\infty} 2^{-k} \{ d(\xi^k, {\xi'}^k) \land 1 \},\$$

and where d denotes appropriate instances of the Skorokhod metric. Write μ_E^f for the distribution of X^E on D_E under \mathbb{P} .

We consider now a limit in which the basic map f is an increasingly well-localized perturbation of the identity, where we quantify this property in terms of the smallest constant $\lambda = \lambda(f) \in (0, 1]$ such that

$$\rho \int_0^1 |\tilde{f}(x+a)\tilde{f}(x)| dx \leqslant \lambda, \quad a \in [\lambda, 1-\lambda].$$

Denote by $(Z_t^k)_{t \ge s_k}$ the *k*th coordinate process on D_E , given by $Z_t^k(\xi) = \xi_t^k$, and consider the filtration $(\mathcal{Z}_t)_{t \in \mathbb{R}}$ on D_E , where \mathcal{Z}_t is the σ -algebra generated by $(Z_s^k : s_k < s \le t \lor s_k, k \in \mathbb{N})$. Write C_E for the (measurable) subset of D_E where each coordinate path is continuous. Define also

$$T^{jk} = \inf\{t \ge s_j \lor s_k : Z_t^j - Z_t^k \in \mathbb{Z}\}.$$

We are thinking of the paths $(Z_t^k)_{t \ge s_k}$ as liftings of paths in the circle. Thus T^{jk} is the collision time of these circle-valued paths. The following is a convenient reformulation of a result of Arratia [1].

Proposition 4.2. There exists a unique Borel probability measure μ_E on D_E under which, for all $j, k \in \mathbb{N}$, the processes $(Z_t^k)_{t \ge s_k}$ and $(Z_t^j Z_t^k - (t - T^{jk})^+)_{t \ge s_j \lor s_k}$ are both continuous local martingales in the filtration $(\mathcal{Z}_t)_{t \in \mathbb{R}}$.

Of course μ_E is supported on C_E and may naturally be considered as a measure defined there. We sketch a proof. For existence, one can take independent Brownian motions from each of the given time-space starting points and then impose a rule of coagulation on collision, deleting the path of lower index. The law of the resulting process has the desired properties. On the other hand, given a probability measure such as described in the proposition, on some larger probability space, one can use a supply of independent Brownian motions to resurrect the paths deleted at each collision. Then Lévy's characterization can be used to see that one has recovered the set-up used for existence. This gives uniqueness.

Proposition 4.3. We have $\mu_E^f \to \mu_E$ weakly on D_E , uniformly in $f \in \mathcal{D}^*$, as $\rho(f) \to \infty$ and $\lambda(f) \to 0$.

Proof. The families of marginal laws $(\mu_{e_k}^f : f \in \mathcal{D}^*)$ are all tight, as shown in Proposition 4.1. Hence the family of laws $(\mu_E^f : f \in \mathcal{D}^*)$ is also tight. Let μ be any weak limit law for the limits $\rho \to \infty$ and $\lambda \to 0$. Under \mathbb{P} , for j, k distinct, the process

$$X_t^j X_t^k - \int_{s_j \lor s_k}^t b(X_s^j, X_s^k) ds, \quad t \ge s_j \lor s_k,$$

is a martingale, where

$$b(x,x') = \rho \int_0^1 \tilde{f}(x-z)\tilde{f}(x'-z)dz.$$

We have $|b(x, x')| \leq \lambda$ whenever $\lambda \leq |x - x'| \leq 1 - \lambda$. Hence, by standard arguments, under μ , the process $(Z_t^j Z_t^k : s_j \lor s_k \leq t < T^{jk})$ is a local martingale. We know from the proof of Proposition 4.1 that, under μ , the processes $(Z_t^j : t \geq s_j)$, $((Z_t^j)^2 - t : t \geq s_j)$ and $(Z_t^k : t \geq s_k)$ are continuous local martingales. But μ inherits from the laws μ_E^f the property that, almost surely, for all $n \in \mathbb{Z}$, the process $(Z_t^j - Z_t^k + n : t \geq s_j \lor s_k)$ does not change sign. Hence, by an optional stopping argument, $Z_t^j - Z_t^k$ is constant for $t \geq T^{jk}$. It follows that $(Z_t^j Z_t^k - (t - T^{jk})^+)_{t \geq s_j \lor s_k}$ is a continuous local martingale. Hence $\mu = \mu_E$, by Proposition 4.2.

We note that the characterizing property of μ_E is invariant under a permutation of the sequence $(e_k : k \in \mathbb{N})$, as is the topology of the metric space D_E . Hence we can define for any

countable set $E \subseteq \mathbb{R}^2$, a unique Borel probability measure on μ_E on $D_E = \prod_{e \in E} D_e$, having the given property. Then, by Proposition 4.3, the distribution μ_E^f of $X^E = (X^e : e \in E)$ on D_E converges weakly to μ_E as $\rho(f) \to \infty$ and $\lambda(f) \to 0$.

5 A new state-space for the coalescing Brownian flow

The preceding statement is unsatisfactory in that it expresses convergence of our Lévy flows only for a given countable set of time-space starting points. To remedy this, we must first formulate a suitable limit object. This object is known as Arratia's flow, or the Brownian web, and has been studied in some depth. However, we have found it convenient to introduce a new state-space of flows, which we now describe.

We begin by defining a metric on \mathcal{D} . Let \mathcal{S} denote the set of all periodic contractions on \mathbb{R} having period 1. Each $f \in \mathcal{D}$ can be identified with some $f^{\times} \in \mathcal{S}$ by drawing new axes at an angle $\pi/4$ with the old, and scaling appropriately. See Figure 2. More formally, since



Figure 2: The map f^{\times} obtained from f by rotating the axes by $\frac{\pi}{4}$.

 $x + f^+(x)$ is strictly increasing in x, there is for each $t \in \mathbb{R}$ a unique $x \in \mathbb{R}$ such that

$$\frac{x+f^{-}(x)}{2} \leqslant t \leqslant \frac{x+f^{+}(x)}{2}$$

Define $f^{\times}(t) = t - x$. Note that $\mathrm{id}^{\times} = 0$. Then the map $f \mapsto f^{\times} : \mathcal{D} \to \mathcal{S}$ is a bijection, so we can define a metric $d_{\mathcal{D}}$ on \mathcal{D} by

$$d_{\mathcal{D}}(f,g) = \|f^{\times} - g^{\times}\| = \sup_{t \in [0,1)} |f^{\times}(t) - g^{\times}(t)|$$

A proof of the italicized assertion is given in the Appendix. The same is true for some further technical assertions which will be made below, written also in italics. The metric space $(\mathcal{S}, \| \dots \|)$ is complete and locally compact, so the same is true for $(\mathcal{D}, d_{\mathcal{D}})$. An alternative characterization⁴ of the metric $d_{\mathcal{D}}$ is as follows: for $f, g \in \mathcal{D}$ and $\varepsilon > 0$, we have

$$d_{\mathcal{D}}(f,g) \leqslant \varepsilon \iff f^-(x-\varepsilon) - \varepsilon \leqslant g^-(x) \leqslant g^+(x) \leqslant f^+(x+\varepsilon) + \varepsilon \text{ for all } x \in \mathbb{R}.$$

We deduce that, for $f, g \in \mathcal{D}$,

$$d_{\mathcal{D}}(f,g) \leqslant ||f-g||, \quad 2d_{\mathcal{D}}(f,\mathrm{id}) = ||f-\mathrm{id}||,$$

and

$$d_{\mathcal{D}}(f, g \circ f) \leq \|g - \operatorname{id}\|$$
 when $g \circ f \in \mathcal{D}$, $d_{\mathcal{D}}(f, f \circ g) \leq \|g - \operatorname{id}\|$ when $f \circ g \in \mathcal{D}$.

Moreover, for any sequence $(f_n : n \in \mathbb{N})$ in \mathcal{D} ,

$$f_n \to f \iff f_n(x) \to f(x)$$
 at every point x where f is continuous

Here and below, we write $f_n \to f$ to mean convergence in the metric $d_{\mathcal{D}}$.

We now define our space of flows. We call them weak flows to emphasise that the usual flow property may fail at points of spatial discontinuity. Consider $\phi = (\phi_{ts} : s, t \in \mathbb{R}, s < t)$, with $\phi_{ts} \in \mathcal{D}$ for all s, t. Say that ϕ is a *weak flow* if

$$\phi_{ut}^- \circ \phi_{ts}^- \leqslant \phi_{us}^- \leqslant \phi_{us}^+ \leqslant \phi_{ut}^+ \circ \phi_{ts}^+, \quad s < t < u.$$

Say that ϕ is *continuous* if, for all $t \in \mathbb{R}$,

$$\phi_{ts} \to \mathrm{id} \ \mathrm{as} \ s \uparrow t, \quad \phi_{ut} \to \mathrm{id} \ \mathrm{as} \ u \downarrow t.$$

Write $C^{\circ}(\mathbb{R}, \mathcal{D})$ for the set of all continuous weak flows⁵. It will be convenient sometimes to extend a continuous weak flow ϕ to the diagonal, which we do by setting $\phi_{ss} = \text{id}$ for all $s \in \mathbb{R}$. Then, for any $\phi \in C^{\circ}(\mathbb{R}, \mathcal{D})$, the map

$$(s,t) \mapsto \phi_{ts} : \{(s,t) : s \leqslant t\} \to \mathcal{D} \tag{9}$$

⁴Thus, $d_{\mathcal{D}}$ is a close relative of the Lévy metric sometimes used on the set of distribution functions for real random variables. The relationships of such a metric to the operations of composition and inversion in \mathcal{D} , which are significant for us, do not appear to have been studied.

⁵In Section 9 we shall work with a modified space of flows $C^{\circ}((0,\infty),\mathcal{D}_0)$. Here we restrict to intervals $I \subseteq (0,\infty)$, and we take $\phi_I \in \mathcal{D}_0$, where \mathcal{D}_0 is the set of circle maps whose liftings are in \mathcal{D} . Given the continuity of the map (9) and the requirement $\phi_{ss} = \text{id}$, there is an obvious identification of this space with $C^{\circ}((0,\infty),\mathcal{D})$

is continuous.

Define, for $\phi, \psi \in C^{\circ}(\mathbb{R}, \mathcal{D})$,

$$d_C(\phi, \psi) = \sum_{n=1}^{\infty} 2^{-n} \{ d_C^{(n)}(\phi, \psi) \land 1 \},\$$

where

$$d_C^{(n)}(\phi,\psi) = \sup_{s,t \in (-n,n), s < t} d_{\mathcal{D}}(\phi_{ts},\psi_{ts}).$$

Then d_C is a metric on $C^{\circ}(\mathbb{R}, \mathcal{D})$, under which this space is complete and separable. For this metric, for all $s, t \in \mathbb{R}$ with s < t, and all $x \in \mathbb{R}$, the evaluation map

$$\phi \mapsto \phi_{ts}^+(x) : C^\circ(\mathbb{R}, \mathcal{D}) \to \mathbb{R}$$

is continuous. Moreover the Borel σ -algebra on $C^{\circ}(\mathbb{R}, \mathcal{D})$ is generated by the set of all such evaluation maps with s,t and x rational.

For $e = (s, x) \in \mathbb{R}^2$ and $\phi \in C^{\circ}(\mathbb{R}, \mathcal{D})$, the map

$$t \mapsto \phi_{ts}^+(x) : [s, \infty) \to \mathbb{R}$$

is continuous. Hence we can define a measurable map $Z^{e,+}: C^{\circ}(\mathbb{R}, \mathcal{D}) \to C_e = C_x([s, \infty), \mathbb{R})$ by

$$Z^{e,+}(\phi) = (\phi_{ts}^+(x) : t \ge s).$$

Given a countable set $E \subseteq \mathbb{R}^2$, we then define a measurable map $Z^{E,+} : C^{\circ}(\mathbb{R}, \mathcal{D}) \to C_E = \prod_{e \in E} C_e$ by

$$Z^{E,+}(\phi)^e = Z^{e,+}(\phi).$$

Set $C_E^{\circ,+} = \{Z^{E,+}(\phi) : \phi \in C^{\circ}(\mathbb{R}, \mathcal{D})\}$. Define similarly $Z^{e,-}, Z^{E,-}$ and $C_E^{\circ,-}$ and set $C_E^{\circ} = C_E^{\circ,+} \cap C_E^{\circ,-}$.

The following result translates into the language of continuous weak flows a result of Tóth and Werner [16, Theorem 2.1], which itself was a variant of a result of Arratia [1]. We shall give a complete proof, in part because we need most components of the proof also for our main convergence result, and in part because our framework leads to some simplifications, for example in the probabilistic underpinnings contained in Proposition 10.10. The formulation in terms of continuous weak flows has advantages, in leading to a unique object, with a natural time-reversal invariance, and for the derivation of weak limits. Recall that, for a countable set $F \subseteq \mathbb{R}^2$, we denote by μ_F the law on C_F of a family of coalescing Brownian motions starting from F, as discussed at the end of the preceding section. **Theorem 5.1.** There exists a unique Borel probability measure μ_A on $C^{\circ}(\mathbb{R}, \mathcal{D})$ such that, for any finite set $F \subseteq \mathbb{R}^2$, we have

$$\mu_A \circ (Z^{F,+})^{-1} = \mu_F. \tag{10}$$

Moreover, for all $e \in \mathbb{R}^2$, we have, μ_A -almost surely, $Z^{e,+} = Z^{e,-}$.

Proof. Fix a countable subset E of \mathbb{R}^2 containing \mathbb{Q}^2 . Consider $\phi, \psi \in C^{\circ}(\mathbb{R}, \mathcal{D})$ and suppose that $Z^{E,+}(\phi) = Z^{E,+}(\psi)$. Then $\phi_{ts}^+(x) = \psi_{ts}^+(x)$ for all $s, x \in \mathbb{Q}$ and all t > s. This extends to all $x \in \mathbb{R}$ because ϕ_{ts}^+ and ψ_{ts}^+ are both right continuous, and then to all $s \in \mathbb{R}$ using (9), so $\phi = \psi$. Hence the map $Z^{E,+} : C^{\circ}(\mathbb{R}, \mathcal{D}) \to C_E^{\circ,+}$ is a bijection. Write $\Phi^{E,+}$ for the inverse bijection $C_E^{\circ,+} \to C^{\circ}(\mathbb{R}, \mathcal{D})$. Similarly, write $\Phi^{E,-}$ for the inverse of the bijection $Z^{E,-} : C^{\circ}(\mathbb{R}, \mathcal{D}) \to C_E^{\circ,-}$. Then $C_E^{\circ,+}$ and $\Phi^{E,+}$ are measurable, we have $\mu_E(C_E^{\circ}) = 1$, and $\Phi^{E,+} = \Phi^{E,-}$ on C_E° . So we can define a Borel probability measure μ_A on $C^{\circ}(\mathbb{R}, \mathcal{D})$ by

$$\mu_A = \mu_E \circ (\Phi^{E,+})^{-1}$$

Then

$$\iota_E = \mu_A \circ (Z^{E,+})^{-1}.$$

For $F \subseteq E$, we have $Z^{F,+} = \pi_{E,F} \circ Z^{E,+}$, and $\mu_F = \mu_E \circ \pi_{E,F}^{-1}$, where $\pi_{E,F} : C_E \to C_F$ is the obvious projection. So

$$\mu_F = \mu_A \circ (Z^{E,+})^{-1} \circ \pi_{E,F}^{-1} = \mu_A \circ (Z^{F,+})^{-1}.$$

On the other hand, for $E \subseteq E'$, we have $Z^E = \pi_{E',E} \circ Z^{E'}$, and $\mu_E = \mu_{E'} \circ \pi_{E',E}^{-1}$. Then $\pi_{E',E}$ restricts to a bijection $C_{E'}^{\circ,+} \to C_E^{\circ,+}$, so we obtain that

$$\mu_A = \mu_{E'} \circ \pi_{E',E}^{-1} \circ \Phi_E^{-1} = \mu_{E'} \circ \Phi_{E'}^{-1}$$

This shows that μ_A does not depend on E, and so (10) holds for all finite sets F. This property then characterizes μ_A by a standard π -system argument. Finally, μ_A -almost surely, $Z^{E,+} \in C_E^{\circ}$, so $Z^{E,+} = Z^{E,-}$, so $Z^{e,+} = Z^{e,-}$ for all $e \in E$.

We call any $C^{\circ}(\mathbb{R}, \mathcal{D})$ -valued random variable with law μ_A a coalescing Brownian flow. The relationship of this space and measure to the Brownian web of Fontes et al. is explored in the Appendix.

6 A Skorokhod-type space of non-decreasing flows on the circle

Since our Lévy flows are not continuous in time, it will be necessary to introduce a larger flow space to accommodate them. Consider now $\phi = (\phi_I : I \subseteq \mathbb{R})$, where $\phi_I \in \mathcal{D}$ and I ranges over

all non-empty finite intervals. Recall that we write $I = I_1 \oplus I_2$ if I_1, I_2 are disjoint intervals with $\sup I_1 = \inf I_2$ and $I = I_1 \cup I_2$. Say that ϕ is a *weak flow* if,

$$\phi_{I_2}^- \circ \phi_{I_1}^- \leqslant \phi_I^- \leqslant \phi_I^+ \leqslant \phi_{I_2}^+ \circ \phi_{I_1}^+, \quad I = I_1 \oplus I_2.$$
(11)

Say that ϕ is $cadlag^6$ if, for all $t \in \mathbb{R}$,

$$\phi_{(s,t)} \to \mathrm{id}$$
 as $s \uparrow t$, $\phi_{(t,u)} \to \mathrm{id}$ as $u \downarrow t$.

Write $D^{\circ}(\mathbb{R}, \mathcal{D})$ for the set of cadlag weak flows. It will be convenient to extend a cadlag weak flow ϕ to the empty interval by setting $\phi_{\emptyset} = \text{id}$. Given a finite interval I and a sequence of finite intervals $(I_n : n \in \mathbb{N})$, write $I_n \to I$ if

$$I = \bigcup_{n} \bigcap_{m \ge n} I_m = \bigcap_{n} \bigcup_{m \ge n} I_m$$

For any $\phi \in D^{\circ}(\mathbb{R}, \mathcal{D})$, we have

$$\phi_{I_n} \to \phi_I \quad as \ I_n \to I.$$
 (12)

Let ϕ be a cadlag weak flow and suppose that $\phi_{\{t\}} = \text{id}$ for all $t \in \mathbb{R}$. Then, using (11), we have $\phi_{(s,t)} = \phi_{(s,t)} = \phi_{[s,t]}$ for all s < t and, denoting all these functions by ϕ_{ts} , the family $(\phi_{ts} : s, t \in \mathbb{R}, s < t)$ is a continuous weak flow in the sense of the preceding section.

For $\phi, \psi \in D^{\circ}(\mathbb{R}, \mathcal{D})$ and $n \ge 1$, define

$$d_D^{(n)}(\phi,\psi) = \inf_{\lambda} \left\{ \gamma(\lambda) \lor \sup_{I \subseteq \mathbb{R}} \|\chi_n(I)\phi_I^{\times} - \chi_n(\lambda(I))\psi_{\lambda(I)}^{\times}\| \right\},\,$$

where the infimum is taken over the set of increasing homeomorphisms λ of \mathbb{R} , where

$$\gamma(\lambda) = \sup_{t \in \mathbb{R}} |\lambda(t) - t| \lor \sup_{s, t \in \mathbb{R}, s < t} \left| \log \left(\frac{\lambda(t) - \lambda(s)}{t - s} \right) \right|,$$

and where χ_n is the cutoff function⁷ given by

$$\chi_n(I) = 0 \lor (n+1-R) \land 1, \quad R = \sup I \lor (-\inf I).$$

$$\|\chi_n(I)\phi_I^{\times} - \chi_n(\lambda(I))\psi_{\lambda(I)}^{\times}\| = d_{\mathcal{D}}(\phi_I, \psi_{\lambda(I)}).$$

Also, for all intervals I, we have $|\chi_n(\lambda(I)) - \chi_n(I)| \leq \gamma(\lambda)$ and

$$\|\chi_n(I)\phi_I^{\times} - \chi_n(\lambda(I))\psi_{\lambda(I)}^{\times}\| \leq \chi_n(I)d_{\mathcal{D}}(\phi_I,\psi_{\lambda(I)}) + |\chi_n(\lambda(I)) - \chi_n(I)| \|\psi_{\lambda(I)}^{\times}\|$$

 $^{^{6}\}mathrm{This}$ definition is more symmetrical in time than is usual for 'cadlag': a more accurate acronym would be laglad

⁷As in the case of the standard Skorokhod topology, localization in time sits awkwardly with the stretching of time introduced via the homeomorphisms λ . There is no fundamental obstacle, just some messiness at the edges. We note that, when $I \cup \lambda(I) \subseteq [-n, n]$, we have

Then define

$$d_D(\phi, \psi) = \sum_{n=1}^{\infty} 2^{-n} \{ d_D^{(n)}(\phi, \psi) \land 1 \}.$$
 (13)

Then d_D is a metric on $D^{\circ}(\mathbb{R}, \mathcal{D})$ under which $D^{\circ}(\mathbb{R}, \mathcal{D})$ is complete and separable. Moreover the metrics d_C and d_D generate the same topology on $C^{\circ}(\mathbb{R}, \mathcal{D})$. For the metric d_D , for all finite intervals I and all $x \in \mathbb{R}$, the evaluation map

$$\phi \mapsto \phi_I^+(x) : D^\circ(\mathbb{R}, \mathcal{D}) \to \mathbb{R}$$

is Borel measurable. Moreover the Borel σ -algebra on $D^{\circ}(\mathbb{R}, \mathcal{D})$ is generated by the set of all such evaluation maps with I = (s, t] and s, t and x rational.

For $e = (s, x) \in \mathbb{R}^2$ and $\phi \in D^{\circ}(\mathbb{R}, \mathcal{D})$, the map

$$t \mapsto \phi^+_{(s,t]}(x) : [s,\infty) \to \mathbb{R}$$

is cadlag. Hence we can extend the map $Z^{e,+}$, which we defined on $C^{\circ}(\mathbb{R}, \mathcal{D})$ in the preceding section, to a measurable map $Z^{e,+}: D^{\circ}(\mathbb{R}, \mathcal{D}) \to D_e = D_x([s, \infty), \mathbb{R})$ by setting

$$Z^{e,+}(\phi) = (\phi^+_{(s,t]}(x) : t \ge s).$$

Given a countable set $E \subseteq \mathbb{R}^2$, we then define a measurable map $Z^{E,+}: D^{\circ}(\mathbb{R}, \mathcal{D}) \to D_E$ by

$$Z^{E,+}(\phi)^e = Z^{e,+}(\phi)$$

and set

$$D_E^{\circ,+} = \{ Z^{E,+}(\phi) : \phi \in D^\circ(\mathbb{R}, \mathcal{D}) \}.$$

Define similarly $Z^{e,-}$, $Z^{E,-}$ and $D_E^{\circ,-}$, and set $D_E^{\circ} = D_E^{\circ,+} \cap D_E^{\circ,-}$.

7 Convergence of Lévy flows to the coalescing Brownian flow

We defined in Section 4, for a given basic map $f \in \mathcal{D}^*$, on the canonical probability space $(\Omega, \mathcal{F}, \mathbb{P})$ of a Poisson random measure, a certain random flow $(X_I : I \subseteq \mathbb{R})$, with $X_I \in \mathcal{D}$ for all finite intervals I. The properties (5), (6) and (7), noted above, imply that $X(\omega) \in D^{\circ}(\mathbb{R}, \mathcal{D})$ for \mathbb{P} -almost all $\omega \in \Omega$, and that the function $X : \Omega \to D^{\circ}(\mathbb{R}, \mathcal{D})$ is measurable for the Borel σ -algebra of $D^{\circ}(\mathbb{R}, \mathcal{D})$. Moreover, we know that

$$Z^{e,+}(X) = Z^{e,-}(X) \quad \text{almost surely, for all } e \in \mathbb{R}^2.$$
(14)

We denote by μ_A^f the law of X on the Borel σ -algebra of $D^{\circ}(\mathbb{R}, \mathcal{D})$.

Theorem 7.1. We have $\mu_A^f \to \mu_A$ weakly on $D^{\circ}(\mathbb{R}, \mathcal{D})$, uniformly in $f \in \mathcal{D}^*$, as $\rho(f) \to \infty$ and $\lambda(f) \to 0$.

Proof. Take $E = \mathbb{Q}^2$. Consider $\phi, \psi \in D^{\circ}(\mathbb{R}, \mathcal{D})$ and suppose that $Z^{E,+}(\phi) = Z^{E,+}(\psi)$. Thus $\phi_{(s,t]}^+(x) = \psi_{(s,t]}^+(x)$ for all $s, x \in \mathbb{Q}$ and all t > s. This extends to all $x \in \mathbb{R}$ because $\phi_{(s,t]}^+$ and $\psi_{(s,t]}^+$ are both right continuous. Then, using (12), we obtain $\phi_I(x) = \psi_I(x)$ for all finite intervals I. Hence the map $Z^{E,+} : D^{\circ}(\mathbb{R}, \mathcal{D}) \to D_E^{\circ,+}$ is a bijection. Write $\Phi^{E,+}$ for the inverse bijection $D_E^{\circ,+} \to D^{\circ}(\mathbb{R}, \mathcal{D})$. Similarly, write $\Phi^{E,-}$ for the inverse of the bijection $Z^{E,-} : D^{\circ}(\mathbb{R}, \mathcal{D}) \to D_E^{\circ,+}$ and $\Phi^{E,+}$ are measurable, and we have $\Phi^{E,+} = \Phi^{E,-}$ on D_E° . Write Φ^E for the common restriction of these functions to D_E° . Then Φ^E is continuous at z for all $z \in C_E^{\circ}$. We see from (14) that $\mu_E^f(D_E^{\circ}) = 1$ for all $f \in \mathcal{D}^*$, and we recall from the proof of Theorem 5.1 that $\mu_E(C_E^{\circ}) = 1$.

Consider the limit where $\rho(f) \to \infty$ and $\lambda(f) \to 0$. By Proposition 4.3, we have $\mu_E^f \to \mu_E$ weakly on D_E . Then, by a standard weak convergence argument, see for example [3], we obtain $\mu_A^f = \mu_E^f \circ (\Phi^E)^{-1} \to \mu_E \circ (\Phi^E)^{-1} = \mu_A$ weakly on $D^\circ(\mathbb{R}, \mathcal{D})$.

8 Time reversal

For $f^+ \in \mathcal{R}$ and $f^- \in \mathcal{L}$, we define a *left-continuous inverse* $(f^+)^{-1} \in \mathcal{L}$ and a *right-continuous inverse* $(f^-)^{-1} \in \mathcal{R}$ by

$$(f^+)^{-1}(y) = \inf\{x \in \mathbb{R} : f^+(x) > y\}, \quad (f^-)^{-1}(y) = \sup\{x \in \mathbb{R} : f^-(x) < y\}.$$

The map $f^+ \mapsto (f^+)^{-1} : \mathcal{R} \to \mathcal{L}$ is a bijection, with $((f^+)^{-1})^{-1} = f^+$ and

$$(f_1^+ \circ f_2^+)^{-1} = (f_2^+)^{-1} \circ (f_1^+)^{-1}, \quad f_1, f_2 \in \mathcal{R}.$$

We have $f^+ \circ (f^+)^{-1} = \text{id}$ if and only if f^+ is a homeomorphism. Define for $f = \{f^-, f^+\} \in \mathcal{D}$ the *inverse* $f^{-1} = \{(f^+)^{-1}, (f^-)^{-1}\} \in \mathcal{D}$. Note that $(f^{-1})^{\times} = -f^{\times}$, so the map $f \mapsto f^{-1} : \mathcal{D} \to \mathcal{D}$ is an isometry.

Define the time-reversal map $\phi \mapsto \hat{\phi}$ on $D^{\circ}(\mathbb{R}, \mathcal{D})$ by

$$\hat{\phi}_I = \phi_{-I}^{-1},$$

where $-I = \{-x : x \in I\}$. It is straightforward to check that this is a well-defined isometry of $D^{\circ}(\mathbb{R}, \mathcal{D})$, which restricts to an isometry of $C^{\circ}(\mathbb{R}, \mathcal{D})$. Write $\hat{\mu}_A^f$ for the law of \widehat{X}^f , where $X^f \sim \mu_A^f$ is the Lévy flow with basic map $f \in \mathcal{D}^*$.

Proposition 8.1. We have $\hat{\mu}_A^f = \mu_A^{f^{-1}}$ for all $f \in \mathcal{D}^*$.

Proof. Fix $f \in \mathcal{D}^*$. Set $g = f^{-1}$ and

$$\Delta = \{(x, y) \in \mathbb{R}^2 : y < f(x)\} = \{(x, y) \in \mathbb{R}^2 : x > g(y)\}, \quad \Delta_0 = \{(x, y) \in \mathbb{R}^2 : y < x\}.$$

Then, by Fubini's theorem,

$$\int_{0}^{1} \tilde{f}(x)dx = \int_{0}^{1} \int_{\mathbb{R}} (1_{\Delta} - 1_{\Delta_{0}})(x, y)dxdy = -\int_{0}^{1} \tilde{g}(y)dy$$

and

$$\int_0^1 \tilde{f}(x)^2 dx = \int_0^1 \int_{\mathbb{R}} 2(y-x)(1_\Delta - 1_{\Delta_0})(x,y) dx dy = \int_0^1 \tilde{g}(y)^2 dy.$$

So $\rho(g) = \rho(f)$ and $\beta(g) = -\beta(f)$. Given $\omega \in \Omega$, define $\hat{\omega} \in \Omega$ by $\hat{\omega}(I \times B) = \omega((-I) \times B)$, for intervals $I \subseteq \mathbb{R}$ and Borel sets $B \subseteq (\mathbb{R}/\mathbb{Z})$. The map $\omega \mapsto \hat{\omega}$ is a measure preserving transformation of $(\Omega, \mathcal{F}, \mathbb{P})$. We have also $(\hat{\omega})^{\beta} = \widehat{\omega^{-\beta}}$. Now, suppressing the \pm notation, for any finite interval I,

$$X_{-I}^{f}(\hat{\omega}, x) = \hat{F}_{\hat{T}_{n}} \circ \dots \circ \hat{F}_{\hat{T}_{n}}(x - \beta t) + \beta s,$$

where $\hat{T}_1 < \cdots < \hat{T}_n$ are the times of the atoms of $(\hat{\omega})^{\beta}$ in -I, and where

$$\hat{F}_t = \begin{cases} \tau_z f, & \text{if } (\hat{\omega})^\beta \text{ has an atom at } (t, z), \\ \text{id}, & \text{otherwise.} \end{cases}$$

Set

$$G_t(y) = \begin{cases} \tau_z g, & \text{if } \omega^{-\beta} \text{ has an atom at } (t, z), \\ \text{id}, & \text{otherwise.} \end{cases}$$

Note that $\hat{T}_k = -T_{n-k+1}$, where $T_1 < \cdots < T_n$ are the times of the atoms of $\omega^{-\beta}$ in I, and $(\tau_z f)^{-1} = \tau_z g$, so $(\hat{F}_{\hat{T}_k})^{-1} = G_{T_{n-k+1}}$. Then

$$\widehat{X_I^f}(\hat{\omega}, y) = (X_{-I}^f)^{-1}(\hat{\omega}, y) = G_{T_n} \circ \cdots \circ G_{T_1}(y - \beta s) + \beta t = X_I^g(\omega, y).$$

Hence X^g and $\widehat{X^f}$ have the same distribution.

We get as a corollary the reversibility of the limit, which is already known in various guises. See, for example, [1],[7] and [16].

Corollary 8.2. The law μ_A of the coalescing Brownian flow is invariant under time-reversal.

Proof. Write $\hat{\mu}_A$ for the image measure of μ_A under time reversal. Fix r > 0 and define $f \in \mathcal{D}$ by

$$f^+(n+x) = n + (x \lor r), \quad n \in \mathbb{Z}, \quad x \in [0,1).$$

Then $\tilde{f}^+(x) = (r-x) \vee 0$ for $x \in [0,1)$, so $\rho(f) = 3/r^3$ and

$$\int_0^1 \tilde{f}(x)\tilde{f}(x+a)dx = 0, \quad r \leqslant a \leqslant 1-r,$$

so $\lambda(f) \leq r$. Moreover $\rho(f^{-1}) = \rho(f)$ and $\lambda(f^{-1}) = \lambda(f)$. Consider the limit $r \to 0$. By Theorem 7.1, we know that $\mu_A^f \to \mu_A$ and $\mu_A^{f^{-1}} \to \mu_A$, weakly on $D^{\circ}(\mathbb{R}, \mathcal{D})$. Since the time-reversal map $\phi \mapsto \hat{\phi}$ is an isometry, it follows, using the preceding proposition, that $\mu_A^{f^{-1}} = \hat{\mu}_A^f \to \hat{\mu}_A$, weakly on $D^{\circ}(\mathbb{R}, \mathcal{D})$. Hence $\mu_A = \hat{\mu}_A$.

9 Scaling limit of the aggregation model

We now return to the aggregation model described in the Introduction and deduce from Theorem 4.3 that, as the particle diameter $\delta \to 0$ and time is suitably speeded up, then the evolution of harmonic measure on the fingers of the cluster converges to the coalescing Brownian flow. In fact, this scaling limit applies more generally and depends on the shape of the attached particles through only a single scalar parameter, which determines the speed of the associated flow. We therefore begin this section by giving a more general formulation of the aggregation model than that in the Introduction.

Let K_0 denote the closed unit ball in \mathbb{C} with centre at 0. Set $D_0 = (\mathbb{C} \cup \{\infty\}) \setminus K_0$. Let P be a closed, connected, simply connected subset of $\overline{D_0}$ of diameter $\delta \in (0, 1]$ such that $P \cap K_0 = \{1\}$. Set $D = (\mathbb{C} \cup \{\infty\}) \setminus (K_0 \cup P)$. The set P models an incoming particle, which is attached to K_0 at 1. Write F for the unique conformal isomorphism $F : D_0 \to D$ such that $F(\infty) = \infty$ and $F'(\infty) > 0$. Let R_1, R_2, \ldots be a sequence of random rotations of the plane about the origin and set $F_n = R_n \circ F \circ R_n^{-1}$. Now construct the aggregation model from the sequence F_1, F_2, \ldots as in the Introduction.

Write G for the inverse isomorphism $D \to D_0$. There exists a unique $g \in \mathcal{D}$ such that g restricts to a continuous map from the interval (0, 1) to itself, and such that

$$G(e^{2\pi ix}) = e^{2\pi ig(x)}, \quad x \in (0,1).$$

Define $\rho(P) = \rho(g) > 0$ and $\beta(P) = \beta(g) \in \mathbb{R}$ as at (4). Note that, if P is symmetric in the real axis, then g is an odd function, so $\beta(P) = 0$.

Set $\Gamma_n = G_n \circ \cdots \circ G_1$, where $G_n = F_n^{-1}$, so that $\Gamma_n : D_n \to D_0$. The restriction of Γ_n to the boundary $\partial K_n = \partial D_n$, gives a natural parametrization of the boundary of the *n*th cluster by the unit circle. It has the property that, for $\xi, \eta \in \partial K_n$, the normalized harmonic

measure h (from ∞) of the positively oriented boundary segment from ξ to η is given by $\Gamma_n(\eta)/\Gamma_n(\xi) = e^{2\pi i h}$. For $m, n \in \mathbb{N}$ with m < n, set

$$\Gamma_{nm} = G_n \circ \cdots \circ G_{m+1}|_{\partial K_0}.$$

Set $\Gamma_{nn} = id$. The circle maps Γ_{nm} belong to \mathcal{D}_0 (see Footnote 5) and have the flow property

$$\Gamma_{nm} \circ \Gamma_{mk} = \Gamma_{nk}, \quad k \leqslant m \leqslant n.$$

The map Γ_{nm} expresses how the harmonic measure on ∂K_m is transformed by the arrival of new particles up to time n. The following result identifies the asymptotic behaviour of this family of maps, for symmetric particles P, in the limit as the particle diameter δ becomes small.⁸ We embed in continuous time by defining, for an interval $I \subseteq (0, \infty)$, $\Gamma_I = \Gamma_{nm}$ where m and n are the smallest and largest integers, respectively, in $\rho(P)I$. Then $(\Gamma_I : I \subseteq (0, \infty))$ is a random variable in $D^{\circ}((0, \infty), \mathcal{D}_0)$. We denote its law by μ_A^P . Write here μ_A for the law of the coalescing Brownian flow on $C^{\circ}((0, \infty), \mathcal{D}_0)$.

Theorem 9.1. We have $\mu_A^P \to \mu_A$ weakly on $D^{\circ}((0,\infty), \mathcal{D}_0)$ as $\delta(P) \to 0$.

Proof. By a standard argument, it will suffice to prove that $\tilde{\mu}_A^P \to \mu_A$ weakly on $D^{\circ}((0, \infty), \mathcal{D}_0)$ as $\delta(P) \to 0$, where we obtain $\tilde{\mu}_A^P$ by a Poissonized embedding, taking $\tilde{\Gamma}_{(s,t]} = \Gamma_{N_tN_s}$, with N a Poisson process of rate $\rho(P)$. In the light of Proposition 9.3 below, the theorem is then a straightforward translation of Theorem 7.1.

Corollary 9.2. Let x_1, \ldots, x_n be a positively oriented set of points in \mathbb{R}/\mathbb{Z} and set $x_0 = x_n$. Set $K_t = K_{\lfloor \rho(P)t \rfloor}$. For $k = 1, \ldots, n$, write H_t^k for the harmonic measure in K_t of the boundary segment of all fingers in K_t attached between x_{k-1} and x_k . Let $(B_t^1, \ldots, B_t^n)_{t \ge 0}$ be a family of coalescing Brownian motions in \mathbb{R}/\mathbb{Z} starting from (x_1, \ldots, x_n) . Then, in the limit $\delta(P) \to 0$, $(H_t^1, \ldots, H_t^n)_{t \ge 0}$ converges weakly in $D([0, \infty), [0, 1]^n)$ to $(B_t^1 - B_t^0, \ldots, B_t^n - B_t^{n-1})_{t \ge 0}$.

Proposition 9.3. There is a universal constant $C < \infty$ such that $\delta^{-3}/C \leq \rho(P) \leq C\delta^{-3}$ and $\lambda(P) \leq C\delta^{1/4}$.

Proof. It is shown in Lawler [12] that there exists some universal constant $c < \infty$ such that if $c\delta \leq x \leq 1 - c\delta$, then

$$|\tilde{g}(x)| \le \frac{3 \operatorname{cap}(P \cup K_0)}{2\pi |1 - e^{2\pi i x}|} = \frac{3 \operatorname{cap}(P \cup K_0)}{4\pi \sin(\pi x)},$$

$$\Gamma_I(e^{2\pi ix}) = e^{-2\pi i\beta t} \Gamma_{nm}(e^{2\pi i(x+\beta s)}),$$

where $s = \inf I$ and $t = \sup I$. The cluster exhibits in this case a first order spinning at speed $\beta = \beta(P)$, with second order fluctuations given by the coalescing Brownian flow.

⁸If P is not symmetric, we obtain the same result once the definition of Γ_I is modified to

where $\operatorname{cap}(P \cup K_0) = -\log G'(\infty)$ is the logarithmic capacity of $P \cup K_0$. If $K_1 \subseteq K_2$, then $\operatorname{cap}(K_1) \leq \operatorname{cap}(K_2)$ (see, for example, [12]), and hence $\operatorname{cap}(P \cup K_0) \leq \operatorname{cap}(L \cup K_0) \leq c'\delta^2$ for some constant c' > 0, by the estimate (2).

Now suppose $x \in (-c\delta, c\delta)$. Then $g(-c\delta) \leq g(x) \leq g(c\delta)$ and hence, for some c'' > 0,

$$|\tilde{g}(x)| \leq |\tilde{g}(c\delta)| \vee |\tilde{g}(-c\delta)| + 2c\delta \leq \frac{3c'\delta^2}{4\pi\sin(\pi c\delta)} + 2c\delta \leq c''\delta.$$

Hence, we have

$$\rho(g)^{-1} = \int_0^1 |\tilde{g}(x)|^2 dx \leqslant 2c\delta(c''\delta)^2 + \int_{c\delta}^{1-c\delta} \frac{9c'^2\delta^4}{16\pi^2 \sin^2(\pi x)} dx$$
$$= 2cc''^2\delta^3 + \frac{9c'^2\delta^4}{8\pi^3} \cot(c\pi\delta) = O(\delta^3).$$

To show the other side of the inequality, recall the estimate (8), which implies that $\rho(g) \leq 8 \|\tilde{g}\|^{-3}$. So it is enough to show that there exists some $x \in [0, 1)$ with $|\tilde{g}(x)| \geq c'' \delta$ for some c'' > 0. Define

$$\tau_A = \inf\{t > 0 : B_t \in A\},\$$

where B_t is a Brownian motion starting from infinity. Since Brownian motion is invariant under conformal transformations,

$$g^+(0) - g^-(0) = \mathbb{P}(\tau_P \le \tau_{K_0})$$

Since P has diameter δ , there exist two points, $w, z \in \overline{P}$, with $w \in \overline{P} \cap K_0$ and $|w - z| \ge \delta/2$. Since P is connected there is a path, $l \subseteq \overline{P}$ joining w and z. Let r be the line joining 0 to z. Reflect l in r and call the reflection l' and the reflection of point w, w'. Now

$$2\mathbb{P}(\tau_l \le \tau_{K_0}) \ge \mathbb{P}(\tau_{l \cup l'} \le \tau_{K_0})$$

and

$$\mathbb{P}(\tau_{l\cup l'} \le \tau_{K_0}) \ge \mathbb{P}(\tau_r \le \tau_{K_0}) \ge 8c'''(|z|-1),$$

by the limit (1), but also

$$\mathbb{P}(\tau_{l\cup l'} \le \tau_{K_0}) \ge \mathbb{P}(B_{\tau_{K_0}} \in \operatorname{arc}(w - w')) \ge 8c''' |w - w'|$$

for some constant c''' > 0. Either (|z| - 1) or $|w - w'| \ge \delta/4$ which finishes the argument.

Finally, to show that $\lambda(g) \leq C\delta^{1/4}$ observe that for δ sufficiently small, if $a \in [\delta^{1/4}, 1-\delta^{1/4}]$, then at most one of x and x + a lie in $\bigcup_{n \in \mathbb{Z}} [n - c\delta, n + c\delta]$. Hence

$$\rho \int_0^1 |\tilde{g}(x+a)\tilde{g}(x)| dx \leqslant 2\rho c'' \delta \int_{c\delta}^{1-c\delta} \frac{3c'\delta^2}{4\pi\sin(\pi x)} dx = \frac{3\rho c'c''\delta^3}{\pi^2} \log\tan(\pi c\delta) = O(\delta).$$

Hence there exists some C > 0 such that, for all $a \in [C\delta^{1/4}, 1 - C\delta^{1/4}]$, we have

$$\rho \int_0^1 |\tilde{g}(x+a)\tilde{g}(x)| dx < C\delta^{1/4}.$$

10 Appendix

10.1 Some properties of the space \mathcal{D} of non-decreasing functions of degree 1

We give proofs in this subsection of a number of assertions made in Section 5.

Proposition 10.1. The map $f \mapsto f^{\times} : \mathcal{D} \to \mathcal{S}$ is a well-defined bijection, with inverse given by

$$f^{-}(x) = \inf\{t + f^{\times}(t) : t \in \mathbb{R}, x = t - f^{\times}(t)\}, \quad f^{+}(x) = \sup\{t + f^{\times}(t) : t \in \mathbb{R}, x = t - f^{\times}(t)\}$$

Proof. Recall that $f^{\times}(t) = t - x$, where x is the unique point such that $f^{-}(x) \leq 2t - x \leq f^{+}(x)$. The periodicity of f^{\times} is an easy consequence of the degree 1 condition. We now show that f^{\times} is a contraction. Fix $s, t \in \mathbb{R}$ and suppose that $f^{\times}(s) = s - y$. Switching the roles of s and t if necessary, we may assume without loss that $x \geq y$. If x = y, then $f^{\times}(s) - f^{\times}(t) = s - t$. On the other hand, if x > y, then $2s - y \leq f^{+}(y) \leq f^{-}(x) \leq 2t - x$, so

$$-(t-s) \leqslant -(t-s) + (2t-x) - (2s-y) = f^{\times}(t) - f^{\times}(s) = (t-s) - (x-y) < t-s.$$

In both cases, we see that $|f^{\times}(t) - f^{\times}(s)| \leq |t - s|$. Hence $f^{\times} \in \mathcal{S}$.

Suppose now that $g \in \mathcal{S}$. Consider, for each $x \in \mathbb{R}$, the set

$$I_x = \{t + g(t) : t \in \mathbb{R}, x = t - g(t)\}.$$

Since g is a contraction, these sets are all intervals, and, since g is bounded, they cover \mathbb{R} . For $x, y \in \mathbb{R}$ with x > y, and for $s, t \in \mathbb{R}$ with x = t - g(t), y = s - g(s), we have t - s - (g(t) - g(s)) = x - y > 0, so $s \leq t$, and so

$$t + g(t) - (s + g(s)) = t - s + (g(t) - g(s)) \ge 0$$

Define $h^+(y) = \sup I_y$ and $h^-(x) = \inf I_x$. We have shown that $h^+(y) \leq h^-(x)$. Moreover, since the intervals I_x cover \mathbb{R} , the functions h^{\pm} must be the left-continuous and rightcontinuous versions of a non-decreasing function h, which then has the degree 1 property, because g is periodic. Thus $h \in \mathcal{D}$. For each $t \in \mathbb{R}$, we have $h^{\times}(t) = t - x$, where $2t - x \in I_x$, and so 2t - x = s + g(s) for some $s \in \mathbb{R}$ with x = s + g(s). Then s = t and so $h^{\times}(t) = g(t)$. Hence $h^{\times} = g$. On the other hand, if we take $g = f^{\times}$ and if x is a point of continuity of f, then we find $I_x = \{f(x)\}$, so $h^+(x) = h^-(x) = f(x)$. Hence h = f. We have now shown that $f \mapsto f^{\times} : \mathcal{D} \to \mathcal{S}$ is a bijection, and that its inverse has the claimed form.

Proposition 10.2. For $f, g \in \mathcal{D}$ and $\varepsilon > 0$,

$$d_{\mathcal{D}}(f,g) \leqslant \varepsilon \iff f^{-}(x-\varepsilon) - \varepsilon \leqslant g^{-}(x) \leqslant g^{+}(x) \leqslant f^{+}(x+\varepsilon) + \varepsilon \text{ for all } x \in \mathbb{R}.$$

Moreover, for any sequence $(f_n : n \in \mathbb{N})$ in \mathcal{D} ,

 $f_n \to f \text{ in } \mathcal{D} \quad \iff \quad f_n^+(x) \to f(x) \text{ at all points } x \in \mathbb{R} \text{ where } f \text{ is continuous.}$

Proof. Suppose that $d_{\mathcal{D}}(f,g) \leq \varepsilon$ and that x is a continuity point of g. Then $g(x) = t + g^{\times}(t)$ for some $t \in \mathbb{R}$ with $x = t - g^{\times}(t)$. We must have $x + \varepsilon \geq t - f^{\times}(t)$ and $g(x) \leq t + f^{\times}(t) + \varepsilon$, so $f^+(x + \varepsilon) + \varepsilon \geq t + f^{\times}(t) + \varepsilon \geq g(x)$. Similarly $f^-(x - \varepsilon) - \varepsilon \leq g(x)$. These inequalities extend to all $x \in \mathbb{R}$ by taking left and right limits along continuity points.

Conversely, suppose that $t \in \mathbb{R}$ is such that $|f^{\times}(t) - g^{\times}(t)| = d_{\mathcal{D}}(f,g)$ and let $x = t - g^{\times}(t)$ and $y = t - f^{\times}(t)$. Then x is the unique point with $g^{-}(x) + x \leq 2t \leq g^{+}(x) + x$ and y is the unique point such that $f^{-}(y) + y \leq 2t \leq f^{+}(y) + y$. Hence $f^{-}(x - \varepsilon) - \varepsilon \leq g^{-}(x) \leq g^{+}(x) \leq f^{+}(x + \varepsilon) + \varepsilon$ implies $y \in [x - \varepsilon, x + \varepsilon]$ and so $d_{\mathcal{D}}(f, g) = |y - x| \leq \varepsilon$.

It follows directly that for any sequence $(f_n : n \in \mathbb{N})$ in \mathcal{D} , if $d_{\mathcal{D}}(f_n, f) \to 0$ as $n \to \infty$, then $f_n^+(x) \to f(x)$ at all points $x \in \mathbb{R}$ where f is continuous.

Now suppose $f_n^+(x) \to f(x)$ at all points $x \in \mathbb{R}$ where f is continuous. By equicontinuity, it will suffice to show that $f_n^{\times}(t) \to f^{\times}(t)$ for each $t \in \mathbb{R}$. Set $x = t - f^{\times}(t)$ and $x_n = t - f_n^{\times}(t)$. Given $\varepsilon > 0$, choose $y_1 \in (x - \varepsilon, x)$ and $y_2 \in (x, x + \varepsilon)$, both points of continuity of f. Now $f(y_1) + y_1 < 2t < f(y_2) + y_2$, so there exists $N \in \mathbb{N}$ such that for all $n \ge N$, we have $f_n^+(y_1) + y_1 < 2t < f_n^+(y_2) + y_2$, which implies $x_n \in [y_1, y_2]$ and hence $|f_n^{\times}(t) - f^{\times}(t)| < \varepsilon$, as required.

Proposition 10.3. Suppose $f_n \to f, g_n \to g, h_n \to h$ in \mathcal{D} with $h_n^+ \leq f_n^+ \circ g_n^+$ for all n. Then $h^+ \leq f^+ \circ g^+$.

Proof. It will suffice to establish the inequality at all points x where g and h are both continuous. Given $\varepsilon > 0$, since f^+ is right-continuous, there exists a point y > g(x) where f is continuous and such that $f(y) < f^+(g(x)) + \varepsilon$. Then $f_n^+(y) < f^+(g(x)) + \varepsilon$ and $g_n^+(x) \leq y$ eventually, so

$$h_n^+(x) \leqslant f_n^+(g_n^+(x)) \leqslant f_n^+(y) < f^+(g(x)) + \varepsilon$$

eventually. Hence $h^+(x) = \lim_{n \to \infty} h_n^+(x) \leq f^+(g^+(x))$, as required.

23

10.2 Some properties of the the continuous flow-space $C^{\circ}(\mathbb{R}, D)$ and cadlag flow-space $D^{\circ}(\mathbb{R}, D)$

We give proofs in this subsection of a number of assertions made in Sections 5 and 6.

Proposition 10.4. For $(s, x) \in \mathbb{R}^2$ and $\phi \in D^{\circ}(\mathbb{R}, \mathcal{D})$, the map

$$t \mapsto \phi^+_{(s\,t]}(x) : [s,\infty) \to \mathbb{R}$$

is cadlag, and is moreover continuous whenever $\phi \in C^{\circ}(\mathbb{R}, \mathcal{D})$.

Proof. Given $t \ge s$ and $\varepsilon > 0$, we can choose $\delta > 0$ so that $d_{\mathcal{D}}(\phi_{(t,u]}, \mathrm{id}) < \varepsilon/2$ for all $u \in (t, t + \delta]$. For such u and for x a point of continuity of $\phi_{(s,t]}$, we have

$$\phi_{(s,t]}^+(x) - \varepsilon = \phi_{(s,t]}^-(x) - \varepsilon \le \phi_{(t,u]}^- \circ \phi_{(s,t]}^-(x) \le \phi_{(s,u]}^-(x) \le \phi_{(s,u]}^+(x) \le \phi_{(t,u]}^+ \circ \phi_{(s,t]}^+(x) \le \phi_{(s,t]}^+(x) + \varepsilon,$$

so $|\phi_{(s,u]}^+(x) - \phi_{(s,t]}^+(x)| \leq \varepsilon$. The final estimate extends to all x by right-continuity. Hence the map is right continuous. A similar argument shows that, for $u \in (s,t)$, we have $|\phi_{(s,u]}^+(x) - \phi_{(s,t)}^+(x)| \to 0$ as $u \to t$, so that the map has a left limit at t given by $\phi_{(s,t)}^+(x)$. Finally, if $\phi \in C^{\circ}(\mathbb{R}, \mathcal{D})$, then $\phi_{(s,t)} = \phi_{(s,t]}$, so the map is continuous.

Proposition 10.5. For all $\phi \in C^{\circ}(\mathbb{R}, \mathcal{D})$, the map $(s, t) \mapsto \phi_{ts} : \{(s, t) : s \leq t\} \to \mathcal{D}$ is continuous. Moreover, for all $\phi \in D^{\circ}(\mathbb{R}, \mathcal{D})$ and for any sequence of finite intervals $I_n \to I$, we have $\phi_{I_n} \to \phi_I$.

Proof. The first assertion follows from the second: given $\phi \in C^{\circ}(\mathbb{R}, \mathcal{D})$ and sequences $s_n \to s$ and $t_n \to t$, then, passing to a subsequence if necessary, we can assume that $(s_n, t_n] \to I$ for some interval I with $\inf I = s$ and $\sup I = t$. Then, by the second assertion, we have $\phi_{t_n s_n} \to \phi_I = \phi_{ts}$, as required.

So, let us fix $\phi \in D^{\circ}(\mathbb{R}, \mathcal{D})$ and a sequence of finite intervals $I_n \to I$. By combining the cadlag and weak flow properties, we can show the following variant of the cadlag property: for all $t \in \mathbb{R}$, we have

$$\phi_{[s,t)} \to \mathrm{id} \quad \mathrm{as} \ s \uparrow t, \quad \phi_{(t,u]} \to \mathrm{id} \quad \mathrm{as} \ u \downarrow t.$$
 (15)

For each n, there exist two disjoint intervals J_n and J'_n , possibly empty, such that $I \triangle I_n = J_n \cup J'_n$. For any such J_n and J'_n , using the weak flow property, we obtain

$$d_{\mathcal{D}}(\phi_I, \phi_{I_n}) \leqslant \|\phi_{J_n} - \operatorname{id}\| + \|\phi_{J'_n} - \operatorname{id}\|$$

Set $s = \inf I$, $s_n = \inf I_n$, $t = \sup I$ and $t_n = \sup I_n$. Then $s_n \to s$, $t_n \to t$, and

if $s \in I$ then $s \in I_n$ eventually, if $s \notin I$ then $s \notin I_n$ eventually,

if $t \in I$ then $t \in I_n$ eventually, if $t \notin I$ then $t \notin I_n$ eventually.

Hence, using the cadlag property or (15), or both, we find that $\phi_{J_n} \to \text{id}$ and $\phi_{J'_n} \to \text{id}$, which proves the proposition.

Proposition 10.6. The metrics d_C and d_D generate the same topology on $C^{\circ}(\mathbb{R}, \mathcal{D})$.

Proof. On comparing the definitions of d_n^C and d_n^D for each $n \in \mathbb{N}$, and considering the choice $\lambda = \operatorname{id}$, we see that $d_D \leq d_C$. Hence, it will suffice to show, given $\phi \in C^{\circ}(\mathbb{R}, \mathcal{D})$, $n \in \mathbb{N}$ and $\varepsilon > 0$, that there exists $\varepsilon' > 0$ such that, for all $\psi \in C^{\circ}(\mathbb{R}, \mathcal{D})$, we have $d_C^{(n)}(\phi, \psi) < \varepsilon$ whenever $d_{n+1}^D(\phi, \psi) < \varepsilon'$. By the preceding proposition, there exists a $\delta \in (0, 1]$ such that $d_D(\phi_{ts}, \phi_{t's'}) < \varepsilon/2$ whenever $|s - s'|, |t - t'| \leq \delta$ and $s, t \in (-n, n)$. Set $\varepsilon' = \delta \wedge (\varepsilon/2)$ and suppose that $d_{n+1}^D(\phi, \psi) < \varepsilon'$. Then there exists an increasing homeomorphism λ of \mathbb{R} , with $|\lambda(t) - t| \leq \delta$ for all t, such that, for all intervals I, we have $\|\chi_{n+1}(I)\psi_I^{\times} - \chi_{n+1}(\lambda(I))\phi_{\lambda(I)}^{\times}\| < \varepsilon/2$. Given $s, t \in (-n, n)$ with s < t, take I = (s, t]. Then $\chi_{n+1}(I) = \chi_{n+1}(\lambda(I)) = 1$, so $d_D(\phi_{\lambda(t)\lambda(s)}, \psi_{ts}) = \|\psi_I^{\times} - \phi_{\lambda(I)}^{\times}\| < \varepsilon/2$. But then, for all such s, t, we have

$$d_{\mathcal{D}}(\phi_{ts},\psi_{ts}) \leqslant d_{\mathcal{D}}(\phi_{ts},\phi_{\lambda(t)\lambda(s)}) + d_{\mathcal{D}}(\phi_{\lambda(t)\lambda(s)},\psi_{ts}) < \varepsilon,$$

so $d_C^{(n)}(\phi,\psi) < \varepsilon$, as required.

Proposition 10.7. The metric spaces $(C^{\circ}(\mathbb{R}, \mathcal{D}), d_C)$ and $(D^{\circ}(\mathbb{R}, \mathcal{D}), d_D)$ are complete and separable.

Proof. The argument for completeness is a variant of the corresponding argument for the usual Skorokhod space $D(\mathbb{R}, S)$ of cadlag paths in complete separable metric space S, as found for example in [3]. Suppose then that $(\psi^n)_{n\geq 1}$ is a Cauchy sequence in $D^{\circ}(\mathbb{R}, \mathcal{D})$. There exists a subsequence $\phi^k = \psi^{n_k}$ such that $d_D^{(n)}(\phi^n, \phi^{n+1}) < 2^{-n}$ for all $n \geq 1$. It will suffice to find a limit in $D^{\circ}(\mathbb{R}, \mathcal{D})$ for $(\phi^n)_{n\geq 1}$. There exist increasing homeomorphisms μ_n of \mathbb{R} for which $\gamma(\mu_n) < 2^{-n}$ and

$$d_{\mathcal{D}}(\phi_I^n, \phi_{\mu_n(I)}^{n+1}) < 2^{-n}, \quad I \cup \mu_n(I) \subseteq (-n, n).$$

For each $n \ge 1$, the sequence $(\mu_{n+m} \circ \cdots \circ \mu_n)_{m \ge 1}$ converges uniformly on \mathbb{R} to an increasing homeomorphism, λ_n say, with $\gamma(\lambda_n) < 2^{-n+1}$. Then $\mu_n \circ \lambda_n^{-1} = \lambda_{n+1}^{-1}$, so

$$d_{\mathcal{D}}(\phi_{\lambda_n^{-1}(I)}^n, \phi_{\lambda_{n+1}^{-1}(I)}^{n+1}) < 2^{-n}, \quad I \subseteq (-n+1, n-1).$$

So, for all $m \ge n$,

$$d_{\mathcal{D}}(\phi_{\lambda_n^{-1}(I)}^n, \phi_{\lambda_{n+m}^{-1}(I)}^{n+m}) < 2^{-n+1}, \quad I \subseteq (-n+1, n-1).$$
(16)

Hence, for all finite intervals $I \subseteq \mathbb{R}$, $(\phi_{\lambda_n^{-1}(I)}^n)_{n\geq 1}$ is a Cauchy sequence in \mathcal{D} , which, since \mathcal{D} is complete, has a limit $\phi_I \in \mathcal{D}$. On letting $m \to \infty$ in (16), we obtain

$$d_{\mathcal{D}}(\phi_{\lambda_n^{-1}(I)}^n, \phi_I) < 2^{-n+1}, \quad I \subseteq (-n+1, n-1).$$

By Proposition 10.3, $\phi = (\phi_I : I \subseteq \mathbb{R})$ has the weak flow property. To see that ϕ is cadlag, suppose given $\varepsilon > 0$ and $t \in \mathbb{R}$. Choose *n* such that $2^{-n+1} \leq \varepsilon/3$ and $|t| \leq n-2$. Then choose $\delta \in (0, 1]$ such that

$$d_{\mathcal{D}}(\phi_{\lambda_n^{-1}(s,t)}^n, \mathrm{id}) < \varepsilon/3, \quad d_{\mathcal{D}}(\phi_{\lambda_n^{-1}(t,u)}^n, \mathrm{id}) < \varepsilon/3$$

whenever $s \in (t - \delta, t)$ and $u \in (t, t + \delta)$. For such s and u, we then have

$$d_{\mathcal{D}}(\phi_{(s,t)}, \mathrm{id}) < \varepsilon, \quad d_{\mathcal{D}}(\phi_{(t,u)}, \mathrm{id}) < \varepsilon.$$

Hence $\phi \in D^{\circ}(\mathbb{R}, \mathcal{D})$. For $m \leq n-3$, we have

$$d_D^{(m)}(\phi^n, \phi) \leqslant \gamma(\lambda_n) \lor \sup_{I \subseteq (-m-2, m+2)} \|\chi_m(\lambda_n^{-1}(I))\phi_{\lambda_n^{-1}(I)}^{n \times} - \chi_m(I)\phi_I^{\times}\|$$

$$\leqslant \gamma(\lambda_n) \lor \sup_{I \subseteq (-m-2, m+2)} \left\{ d_{\mathcal{D}}(\phi_{\lambda_n^{-1}(I)}^n, \phi_I) + \gamma(\lambda_n) \|\phi_I^{\times}\| \right\}$$

$$\leqslant 2^{-n+1} (1 + \sup_{I \subseteq (-m-2, m+2)} \|\phi_I^{\times}\|).$$

Hence $d_D(\phi^n, \phi) \to 0$ as $n \to \infty$. We have shown that $D^{\circ}(\mathbb{R}, \mathcal{D})$ is complete. If the sequence $(\phi^n)_{n\geq 1}$ in fact lies in $C^{\circ}(\mathbb{R}, \mathcal{D})$, then, by an obvious variation of the argument for the cadlag property, the limit ϕ also lies in $C^{\circ}(\mathbb{R}, \mathcal{D})$. Hence $C^{\circ}(\mathbb{R}, \mathcal{D})$ is also complete. In particular, $C^{\circ}(\mathbb{R}, \mathcal{D})$ is a closed subspace in $D^{\circ}(\mathbb{R}, \mathcal{D})$.

We turn to the question of separability. Let us write D_N for the set of those $\phi \in D^{\circ}(\mathbb{R}, \mathcal{D})$ such that:

- (i) for some $n \in \mathbb{N}$ and some rationals $t_1 < \cdots < t_n$, we have $\phi_J = \text{id}$ for all time intervals J which do not intersect the set $\{t_1, \ldots, t_n\}$;
- (ii) for all other time intervals I, the maps ϕ_I and ϕ_I^{-1} on \mathbb{R} are constant on all space intervals which do not intersect $2^{-N}\mathbb{Z}$.

Note that each $\phi \in D_N$ is determined by the maps $\phi_{(t_k,t_m]}$, for integers $0 \leq k < m \leq n$, where $t_0 < t_1$, and for each of these maps there are only countably many possibilities (finitely many if we insist that $\phi(0) \in [0,1)$). Hence D_N is countable and so is $D_* = \bigcup_{N \geq 1} D_N$. We shall show that D_* is also dense in $D^{\circ}(\mathbb{R}, \mathcal{D})$.

Fix $\phi \in D^{\circ}(\mathbb{R}, \mathcal{D})$ and $n_0 \geq 1$. It will suffice to find, for a given $\varepsilon > 0$, a $\psi \in D_*$ with $d_D^{(n_0)}(\phi, \psi) < \varepsilon$. By the cadlag property and compactness, there exist $n \in \mathbb{N}$ and reals $s_1 < \cdots < s_n$ in $I_0 = (-n_0 - 1, n_0 + 1)$ such that $d_{\mathcal{D}}(\phi_I, \mathrm{id}) < \varepsilon/4$ for every subinterval I of I_0 which does not intersect $\{s_1, \ldots, s_n\}$. Then we can find rationals $t_1 < \cdots < t_n$ in I_0 and an increasing homeomorphism λ of \mathbb{R} , with $\lambda(t) = t$ for $t \notin I_0$, with $\gamma(\lambda) \sup_{I \subseteq I_0} ||\phi_I^{\times}|| < \varepsilon/4$, and such that $\lambda(t_m) = s_m$ for all m. Set $s_0 = t_0 = -n_0 - 1$.

For $f \in \mathcal{D}$, write $\Delta(f)$ for the set of points where f is not continuous. Define, for $m = 0, 1, \ldots, n$,

$$\Delta_m = \bigcup_{k=0}^{m-1} \Delta(\phi_{(s_k,s_m]}^{-1}) \cup \bigcup_{k=m+1}^n \Delta(\phi_{(s_m,s_k]}).$$

Then Δ_m is countable, so we can choose $N \ge 1$ with $16.2^{-N} \le \varepsilon$ and choose $\varepsilon_m \in \mathbb{R}$ with $|\varepsilon_m| \le 2^{-N}$ such that

$$\tau_m(\Delta_m) \cap 2^{-N} \mathbb{Z} = \emptyset, \quad m = 0, 1, \dots, n_q$$

where $\tau_m(x) = x + \varepsilon_m$. Set

$$\delta^{-}(x) = 2^{N} \lceil 2^{-N} x \rceil, \quad \delta^{+}(x) = 2^{N} \lfloor 2^{-N} x \rfloor + 1.$$

Note that $\delta = {\delta^-, \delta^+} \in \mathcal{D}$. Define for $0 \leq k < m \leq n$

$$\psi_{(t_k,t_m]}^- = (\delta^{-1})^- \circ (\tau_m)^{-1} \circ \phi_{(s_k,s_m]}^- \circ \tau_k \circ \delta^-, \quad \psi_{(t_k,t_m]}^+ = (\delta^{-1})^+ \circ (\tau_m)^{-1} \circ \phi_{(s_k,s_m]}^+ \circ \tau_k \circ \delta^+.$$

Then $\psi_{(t_k,t_m]} = \{\psi_{(t_k,t_m]}^-, \psi_{(t_k,t_m]}^+\} \in \mathcal{D}$ by our choice of ε_k and ε_m . Moreover $\delta^+ \circ (\delta^{-1})^+ \ge \mathrm{id}$ and $\delta^- \circ (\delta^{-1})^- \le \mathrm{id}$ so, for $0 \le m < m' < m'' \le n$, we obtain the inequalities

$$\psi_{(t_{m'},t_{m''}]}^{-} \circ \psi_{(t_m,t_{m'}]}^{-} \leqslant \psi_{(t_m,t_{m''}]}^{-} \leqslant \psi_{(t_m,t_{m''}]}^{+} \leqslant \psi_{(t_{m'},t_{m''}]}^{+} \circ \psi_{(t_m,t_{m'}]}^{+}$$

from the corresponding inequalities for ϕ . We use the equations $\|\delta - \mathrm{id}\| = 2^{-N}$ and $\|\tau_m - \mathrm{id}\| = |\varepsilon_m|$ to see that

$$d_{\mathcal{D}}(\phi_{(s_k, s_m]}, \psi_{(t_k, t_m]}) \leqslant 4.2^{-N}, \quad 0 \leqslant k < m \leqslant n$$

Define $\psi_J = \psi_{(t_k,t_m]}$ for all intervals J such that $J \cap \{t_1,\ldots,t_n\} = \{t_{k+1},\ldots,t_m\}$. For such intervals J, with $J \subseteq I_0$, we have $d_{\mathcal{D}}(\phi_{(s_k,s_m]\setminus\lambda(J)}, \mathrm{id}) < \varepsilon/4$ and $d_{\mathcal{D}}(\phi_{\lambda(J)\setminus(s_k,s_m]}, \mathrm{id}) < \varepsilon/4$ so, using the weak flow property for ϕ ,

$$d_{\mathcal{D}}(\psi_J, \phi_{\lambda(J)}) \leqslant d_{\mathcal{D}}(\psi_{(t_k, t_m]}, \phi_{(s_k, s_m]}) + d_{\mathcal{D}}(\phi_{(s_k, s_m]}, \phi_{\lambda(J)}) \leqslant 4.2^{-N} + 2\varepsilon/4 < 3\varepsilon/4.$$

Define $\psi_J = \text{id for all intervals } J$ which do not intersect $\{t_1, \ldots, t_n\}$. For such intervals J with $J \subseteq I_0$, we have $d_{\mathcal{D}}(\psi_J, \phi_{\lambda(J)}) \leq d_{\mathcal{D}}(\text{id}, \phi_{\lambda(J)}) \leq \varepsilon/4$. Now $\psi \in D_N$ and

$$d_D^{(n_0)}(\phi,\psi) \leqslant \gamma(\lambda) \lor \sup_{J \subseteq I_0} \{ d_{\mathcal{D}}(\psi_J,\phi_{\lambda(J)}) + \gamma(\lambda) \| \phi_J^{\times} \| \} < \varepsilon,$$

as required. This proves that $D^{\circ}(\mathbb{R}, \mathcal{D})$ is separable and, since $C^{\circ}(\mathbb{R}, \mathcal{D})$ is a closed subspace of $D^{\circ}(\mathbb{R}, \mathcal{D})$, it follows that $C^{\circ}(\mathbb{R}, \mathcal{D})$ is also separable.

Proposition 10.8. For all $s, t \in \mathbb{R}$ with s < t, and all $x \in \mathbb{R}$, the map $\phi \mapsto \phi_{ts}^+(x)$ on $C^{\circ}(\mathbb{R}, \mathcal{D})$ is continuous. Moreover the Borel σ -algebra on $C^{\circ}(\mathbb{R}, \mathcal{D})$ is generated by the set of all such maps with s, t and x rational.

For all finite intervals $I \subseteq \mathbb{R}$ and all $x \in \mathbb{R}$, the map $\phi \mapsto \phi_I^+(x)$ on $D^{\circ}(\mathbb{R}, \mathcal{D})$ is Borel measurable, and the Borel σ -algebra on $D^{\circ}(\mathbb{R}, \mathcal{D})$ is generated by the set of all such maps where I = (s, t] with s, t and x rational.

Proof. The results for $C^{\circ}(\mathbb{R}, \mathcal{D})$ can be proved more simply than those for $D^{\circ}(\mathbb{R}, \mathcal{D})$. We omit details of the former, but note that these follow also from the latter, by general measure theoretic arguments, given what we already know about the two spaces.

The proof for $D^{\circ}(\mathbb{R}, \mathcal{D})$ is an adaptation of the analogous result for the classical Skorokhod space, see for example [10, page 335]. We prove first the Borel measurability of the evaluation maps. Given a finite interval I and $x \in \mathbb{R}$, we can find $s_n, t_n \in \mathbb{R}$ such that $(s_n, t_n] \to I$ as $n \to \infty$. Then $\phi_I^+(x) = \lim_{m \to \infty} \lim_{n \to \infty} \phi_{(s_n, t_n]}^+(x + 1/m)$, by Proposition 10.5. Hence, it will suffice to consider intervals I of the form (s, t]. Fix s, t and x and define for each $m, n \in \mathbb{N}$ a function $F_{m,n}$ on $D^{\circ}(\mathbb{R}, \mathcal{D})$ by

$$F_{m,n}(\phi) = \int_{s}^{s+1/n} \int_{t}^{t+1/n} \int_{x}^{x+1/m} \phi^{+}_{(s',t']}(x') dx' dt' ds'.$$

Suppose $\phi^k \to \phi$ in $D^{\circ}(\mathbb{R}, \mathcal{D})$. We can choose increasing homeomorphisms λ_k of \mathbb{R} such that, $\gamma(\lambda_k) \to 0$ and, uniformly in $r \in [s-1, s+1]$ and $u \in [t-1, t+1]$, we have

$$d_{\mathcal{D}}(\phi_{\lambda_k(r,u]}^k, \phi_{(r,u]}) \to 0.$$

Define

$$f(r,u) = \int_{x}^{x+1/m} \phi_{\lambda(r,u]}(x')dx', \quad f_k(r,u) = \int_{x}^{x+1/m} \phi_{\lambda_k(r,u]}^k(x')dx'.$$

Then $f_k(r, u) \to f(r, u)$, uniformly in $r \in [s - 1, s + 1]$ and $u \in [t - 1, t + 1]$. Set $\mu_k = \lambda_k^{-1}$. Then

$$F_{m,n}(\phi^k) = \int_{\mu_k(s)}^{\mu_k(s+1/n)} \int_{\mu_k(t)}^{\mu_k(t+1/n)} f_k(r, u) d\lambda_k(u) d\lambda_k(r) \to \int_s^{s+1/n} \int_t^{t+1/n} f(r, u) du dr = F_{m,n}(\phi),$$

so $F_{m,n}$ is continuous on $D^{\circ}(\mathbb{R}, \mathcal{D})$. By Proposition 10.5, we have

$$\phi_{(s,t]}^+(x) = \lim_{m \to \infty} \lim_{n \to \infty} \frac{1}{mn^2} F_{m,n}(\phi).$$

Hence $\phi \mapsto \phi^+_{(s,t]}(x)$ is Borel measurable, as required.

Write now \mathcal{E} for the σ -algebra on $D^{\circ}(\mathbb{R}, \mathcal{D})$ generated by all maps of this form with s, t and x rational. It remains to show that \mathcal{E} contains the Borel σ -algebra of $D^{\circ}(\mathbb{R}, \mathcal{D})$. Write $\{(I_k, z_k) : k \in \mathbb{N}\}$ for an enumeration of the set $\{(s, t] : s, t \in \mathbb{Q}, s < t\} \times \mathbb{Q}$. It is straightforward to show that, for all k, the map $\phi \mapsto \phi_{I_k}^{\times}(z_k)$ is \mathcal{E} -measurable. Fix $n \in \mathbb{N}$, $\phi^0 \in D^{\circ}(\mathbb{R}, \mathcal{D}), r \in (0, \infty)$ and $k \in \mathbb{N}$, and consider the set

$$A(k,r) = \{ \phi \in D^{\circ}(\mathbb{R}, \mathcal{D}) : (\chi_n(I_1)\phi_{I_1}^{\times}(z_1), \dots, \chi_n(I_k)\phi_{I_k}^{\times}(z_k)) \in B(k,r) \},\$$

where

$$B(k,r) = \bigcup_{\lambda} \{ (y_1, \dots, y_k) \in \mathbb{R}^k : \max_{j \leq k} |y_j - \chi_n(\lambda(I_j))\phi_{\lambda(I_j)}^{0\times}(z_j)| < r \},\$$

where the union is taken over all increasing homeomorphisms λ of \mathbb{R} with $\gamma(\lambda) < r$. Note that B(k,r) is an open set in \mathbb{R}^k , so $A(k,r) \in \mathcal{E}$, so $A = \bigcup_{m \in \mathbb{N}} \bigcap_{k \in \mathbb{N}} A(k,r-1/m) \in \mathcal{E}$.

Consider the set

$$C = \{ \phi \in D^{\circ}(\mathbb{R}, \mathcal{D}) : d_D^{(n)}(\phi, \phi^0) < r \}.$$

It is straightforward to check from the definition of $d_D^{(n)}$, that $C \subseteq A$. Suppose that $\phi \in A$. We shall show that $\phi \in C$. Then C = A, so $C \in \mathcal{E}$, and since sets of this form generate the Borel σ -algebra, we are done.

We can find an $m \in \mathbb{N}$ and, for each $k \in \mathbb{N}$, a λ_k with $\gamma(\lambda_k) < r - 1/m$ such that

$$|\chi_n(I_j)\phi_{I_j}^{\times}(z_j) - \chi_n(\lambda_k(I_j))\phi_{\lambda_k(I_j)}^{0\times}(z_j)| < r - 1/m, \quad j = 1, \dots, k.$$

Without loss of generality, we may assume that the sequence $(\lambda_k : k \in \mathbb{N})$ converges uniformly on compacts, and that its limit, λ say, satisfies $\gamma(\lambda) \leq r - 1/m$. By Proposition 10.5, for each j, there is an interval \hat{I}_j , having the same endpoints as I_j such that $\phi_{\lambda(\hat{I}_j)}$ is a limit point in \mathcal{D} of the sequence $(\phi_{\lambda_k(I_j)} : k \in \mathbb{N})$, so $\phi_{\lambda(\hat{I}_j)}^{\times}$ is a limit point in \mathcal{S} of the sequence $(\phi_{\lambda_k(I_j)}^{\times} : k \in \mathbb{N})$. Then

$$|\chi_n(I_j)\phi_{I_j}^{\times}(z_j) - \chi_n(\lambda(\hat{I}_j))\phi_{\lambda(\hat{I}_j)}^{0\times}(z_j)| \leqslant r - 1/m,$$

for all j. For all finite intervals I and all $z \in \mathbb{R}$, we can find a sequence $(j_p : p \in \mathbb{N})$ such that $I_{j_p} \to I$, $\hat{I}_{j_p} \to I$ and $z_{j_p} \to z$. So we obtain

$$|\chi_n(I)\phi_I^{\times}(z) - \chi_n(\lambda(I))\phi_{\lambda(I)}^{0\times}(z)| \leq r - 1/m.$$

Hence $d_D^{(n)}(\phi, \phi^0) \leq r - 1/m$ and $\phi \in C$, as we claimed.

Recall that we define, for $e = (s, x) \in \mathbb{R}^2$ and $\phi \in D^{\circ}(\mathbb{R}, \mathcal{D})$,

$$Z^{e,+}(\phi) = (\phi^+_{(s,t]}(x) : t \ge s)$$

and for $E \subseteq \mathbb{R}^2$, we define $Z^{E,+}: D^{\circ}(\mathbb{R}, \mathcal{D}) \to D_E$ by $Z^{E,+}(\phi)^e = Z^{e,+}(\phi)$. Recall also that we set

$$C_{E}^{\circ,+} = \{ Z^{E,+}(\phi) : \phi \in C^{\circ}(\mathbb{R}, \mathcal{D}) \}, \quad D_{E}^{\circ,+} = \{ Z^{E,+}(\phi) : \phi \in D^{\circ}(\mathbb{R}, \mathcal{D}) \},$$

and that we define analogously $Z^{E,-}$, $C_E^{\circ,-}$ and $D_E^{\circ,-}$, and we set $C_E^{\circ} = C_E^{\circ,+} \cap C_E^{\circ,-}$ and $D_E^{\circ} = D_E^{\circ,+} \cap D_E^{\circ,-}.$

Proposition 10.9. Let E be a countable subset of \mathbb{R}^2 containing \mathbb{Q}^2 . Then $Z^{E,+} : C^{\circ}(\mathbb{R}, \mathcal{D}) \to \mathbb{R}$ $C_E^{\circ,+}$ is a bijection, $C_E^{\circ,+}$ is a measurable subset of C_E , and the inverse bijection $\Phi^{E,+}: C_E^{\circ,+} \to C_E^{\circ,+}$ $C^{\circ}(\mathbb{R}, \mathcal{D})$ is a measurable map.

Moreover $Z^{E,+}: D^{\circ}(\mathbb{R}, \mathcal{D}) \to D_{E}^{\circ,+}$ is also a bijection, $D_{E}^{\circ,+}$ is a measurable subset of D_{E} and the inverse bijection $\Phi^{E,+}: D_{E}^{\circ,+} \to D^{\circ}(\mathbb{R}, \mathcal{D})$ is also a measurable map. Moreover the same statements hold with + replaced by -, and we have $\Phi^{E,+} = \Phi^{E,-}$ on

 D_E° .

Proof. We discuss only the cadlag case. The usual comments apply concerning the relationship of this case with the continuous case. The simple argument that $Z^{E,+}$ is injective on $D_E^{\circ,+}$ was given in the proof of Theorem 7.1. Consider for $z \in D_E$ the conditions

$$z_t^{(s,x+n)} = z_t^{(s,x)} + n, \quad s, t, x \in \mathbb{Q}, \quad s < t, \quad n \in \mathbb{Z}$$

$$(17)$$

and

$$z_t^{(s,x)} = \inf_{y \in \mathbb{Q}, y > x} z_t^{(s,y)}, \quad (s,x) \in E, \quad t \in \mathbb{Q}, t > s.$$
(18)

Under these conditions, define for $s, t \in \mathbb{Q}$ with s < t and for $x \in \mathbb{R}$,

$$\Phi_{(s,t]}^{-}(x) = \sup_{y \in \mathbb{Q}, y < x} z_t^{(s,y)}, \quad \Phi_{(s,t]}^{+}(x) = \inf_{y \in \mathbb{Q}, y > x} z_t^{(s,y)}.$$

Then $\Phi_{(s,t]} = \{\Phi_{(s,t]}^{-}, \Phi_{(s,t]}^{+}\} \in \mathcal{D}$ and

$$\Phi^+_{(s,t]}(x) = z_t^{(s,x)}, \quad s, t, x \in \mathbb{Q}, \quad s < t.$$

Now consider the following additional conditions on z:

$$\Phi_{(t,u]}^{-} \circ \Phi_{(s,t]}^{-} \leqslant \Phi_{(s,u]}^{-} \leqslant \Phi_{(s,u]}^{+} \leqslant \Phi_{(t,u]}^{+} \circ \Phi_{(s,t]}^{+}, \quad s, t, u \in \mathbb{Q}, \quad s < t < u;$$
(19)

and

for all $\varepsilon > 0$ and all $n \in \mathbb{N}$, there exist $\delta > 0$, $m \in \mathbb{Z}^+$ and $u_1, \ldots, u_m \in (-n, n)$ such that

$$\|\Phi_{(s,t]} - \operatorname{id}\| < \varepsilon \tag{20}$$

whenever $s, t \in \mathbb{Q} \cap (-n, n)$ with $0 < t - s < \delta$ and $(s, t] \cap \{u_1, \dots, u_m\} = \emptyset$.

Note that the inequalities between functions required in (19) hold whenever the same inequalities hold between their restrictions to \mathbb{Q} , by left and right continuity. Note also that condition (20) is equivalent to the following condition involving quantifiers only over countable sets:

for all rationals $\varepsilon > 0$ and all $n \in \mathbb{N}$, there exist a rational $\delta > 0$ and an $m \in \mathbb{Z}^+$ such that, for all rationals $\eta > 0$, there exist rationals $s_1, t_1, \ldots, s_m, t_m \in (-n, n)$, with $s_i < t_i$ for all *i* and with $\sum_{i=1}^m (t_i - s_i) < \eta$, such that

$$\|\Phi_{(s,t]} - \operatorname{id}\| < \varepsilon$$

whenever $s, t \in \mathbb{Q} \cap (-n, n)$ with $0 < t - s < \delta$ and $(s, t] \cap ((s_1, t_1] \cup \cdots \cup (s_m, t_m]) = \emptyset$.

Denote by $D_E^{*,+}$ the set of those $z \in D_E$ where conditions (17), (18), (19) and (20) all hold. Then $D_E^{*,+}$ is a measurable subset of D_E . Fix $z \in D_E^{*,+}$. Given finite interval I, we can find sequences of rationals s_n and t_n such that $(s_n, t_n] \to I$ as $n \to \infty$. Then, by conditions (19) and (20),

$$d_{\mathcal{D}}(\Phi_{(s_n,t_n]},\Phi_{(s_m,t_m]}) \leqslant ||\Phi_{(s_n,s_m]} - \mathrm{id} || + ||\Phi_{(t_n,t_m]} - \mathrm{id} || \to 0$$

as $n, m \to \infty$. So the sequence $\Phi_{(s_n, t_n]}$ converges in \mathcal{D} , with limit Φ_I , say, and Φ_I does not depend on the approximating sequences of rationals. In the case where $I = I_1 \oplus I_2$, there exists another sequence of rationals u_n such that $(s_n, u_n] \to I_1$ and $(u_n, t_n] \to I_2$ as $n \to \infty$. Hence $\Phi = (\Phi_I : I \subseteq \mathbb{R})$ has the weak flow property, by Proposition 10.3. It is straightforward to deduce from (20) that Φ is moreover cadlag, so $\Phi = \Phi(z) \in D^{\circ}(\mathbb{R}, \mathcal{D})$. It follows from its construction, and the preceding proposition, that the map $z \mapsto \Phi(z) : D_E^{*,+} \to D^{\circ}(\mathbb{R}, \mathcal{D})$ is measurable.

Now, for all $z \in D_E^{*,+}$, we have $Z^{E,+}(\Phi(z)) = z$ and for all $\phi \in D^{\circ}(\mathbb{R}, \mathcal{D})$, we have $Z^{E,+}(\phi) \in D_E^*$ and $\Phi(Z^{E,+}(\phi)) = \phi$. Hence $D_E^{\circ,+} = D_E^{*,+}$ and $\Phi^{E,+} = \Phi$, showing that these are measurable, as claimed.

Consider now for $z \in D_E$ the condition

$$z_t^{(s,x)} = \sup_{y \in \mathbb{Q}, y < x} z_t^{(s,y)}, \quad (s,x) \in E, \quad t \in \mathbb{Q}, t > s.$$
(21)

Denote by $D_E^{*,-}$ the set of those $z \in D_E$ where conditions (17), (19), (20) and (21) all hold, and define Φ on $D_E^{*,-}$ exactly as on $D_E^{*,+}$. Then, by a similar argument, $D_E^{\circ,-} = D_E^{*,-}$ and $\Phi = \Phi^{E,-}$ on $D_E^{\circ,-}$. In particular $\Phi^{E,+} = \Phi^{E,-}$ on D_E° , as claimed.

Proposition 10.10. Let E be a countable subset of \mathbb{R}^2 containing \mathbb{Q}^2 . Then $\mu_E(C_E^\circ) = 1$.

Proof. We use an identification of C_E° analogous to that implied for D_E° by the preceding proof. The same five conditions (17), (18), (19), (20) and (21) characterize C_E° inside C_E , except that, in (20), only the case m = 0 is allowed. Recall that, under μ_E , for time-space starting points e = (s, x) and e' = (s', x'), the coordinate processes Z^e and $Z^{e'}$ behave as independent Brownian motions up to

$$T^{ee'} = \inf\{t \ge s \lor s' : Z_t^e - Z_t^{e'} \in \mathbb{Z}\},\$$

after which they continue to move as Brownian motions, but now with a constant separation. In particular, if s = s' and x' = x + n for some $n \in \mathbb{Z}$, then $T^{ee'} = 0$, so $Z_t^{e'} = Z_t^e + n$ for all $t \ge s$, so (17) holds almost surely.

Let $(s, x) \in E$ and $t, u \in \mathbb{Q}$, with $s \leq t < u$. Consider the event

$$A = \left\{ \sup_{y \in \mathbb{Q}, y < Z_t^{(s,x)}} Z_u^{(t,y)} = Z_u^{(s,x)} = \inf_{y' \in \mathbb{Q}, y' > Z_t^{(s,x)}} Z_u^{(t,y')} \right\}.$$

Fix $n \in \mathbb{N}$ and set $Y = n^{-1} \lfloor nZ_t^{(s,x)} \rfloor$ and Y' = Y + 1/n. Then Y and Y' are \mathcal{F}_t -measurable, Q-valued random variables. Now $\mathbb{P}(Y < Z_t^{(s,x)} < Y') = 1$ and

$$\{Y < Z_t^{(s,x)} < Y'\} \cap \{T^{(t,Y)(t,Y')} \leqslant u\} \subseteq A.$$

By the Markov property of Brownian motion, almost surely,

$$\mathbb{P}(T^{(t,Y)(t,Y')} \leqslant u | \mathcal{F}_t) = 2\Phi\left(\frac{1}{n\sqrt{2(u-t)}}\right),$$

and the right-hand side tends to 1 as $n \to \infty$. So, by bounded convergence, we obtain $\mathbb{P}(A) = 1$. On taking a countable intersection of such sets A over the possible values of s, x, t and u, we deduce that conditions (18), (19) and (21) hold almost surely.

It remains to establish the continuity condition (20). For a standard Brownian motion B starting from 0, we have, for $n \ge 4$,

$$\mathbb{P}\left(\sup_{t\leqslant 1}|B_t|>n\right)\leqslant e^{-n^2/2}.$$

Define, for $\delta > 0$ and $e = (s, x) \in E$,

$$V^e(\delta) = \sup_{s \leqslant t \leqslant s + \delta^2} |Z_t^e - x|$$

Then, by scaling,

$$\mathbb{P}(V^e(\delta) > n\delta) \leqslant e^{-n^2/2}.$$

Consider, for each $n \in \mathbb{N}$ the set

$$E_n = \{ (j2^{-2n}, k2^{-n}) : j \in \frac{1}{2}\mathbb{Z} \cap [-2^{2n}, 2^{2n}), k = 0, 1, \dots, 2^n - 1 \}$$

and the event

$$A_n = \bigcup_{e \in E_n} \{ V^e(2^{-n}) > n2^{-n} \}.$$

Then $\mathbb{P}(A_n) \leq |E_n|e^{-n^2/2}$, so $\sum_n \mathbb{P}(A_n) < \infty$, so by Borel–Cantelli, almost surely, there is a random $N < \infty$ such that $V^e(2^{-n}) \leq n2^{-n}$ for all $e \in E_n$, for all $n \geq N$.

Given $\varepsilon > 0$, choose $n \ge N$ such that $(4n+2)2^{-n} \le \varepsilon$ and set $\delta = 2^{-2n-1}$. Then, for all rationals $s, t \in (-n, n)$ with $0 < t - s < \delta$ and all rationals $x \in [0, 1]$, there exist $e^{\pm} = (r, y^{\pm}) \in E_n$ such that

$$r \leq s < t \leq r + 2^{-2n}, \quad x + n2^{-n} < y^+ \leq x + (n+1)2^{-n}, \quad x - (n+1)2^{-n} \leq y^- < x - n2^{-n},$$

Then, $Z_s^{e^-} < x < Z_s^{e^+}$, so

$$x - \varepsilon \leqslant Z_t^{e^-} \leqslant Z_t^{(s,x)} \leqslant Z_t^{e^+} \leqslant x + \varepsilon.$$

Hence $\|\Phi_{(s,t]} - \operatorname{id}\| \leq \varepsilon$, as required.

Recall that Φ^E denotes the common restriction of $\Phi^{E,+}$ and $\Phi^{E,-}$ to D_E° .

Proposition 10.11. Let $E = \mathbb{Q}^2$. Then Φ^E is continuous at z for all $z \in C_E^{\circ}$.

Proof. Consider a sequence $(z_k : k \in \mathbb{N})$ in D_E° and suppose that $z_k \to z$ in D_E , with $z \in C_E^\circ$. Set $\phi^k = \Phi^E(z_k)$ and $\phi = \Phi^E(z)$. It will suffice to show that, for all $n \in \mathbb{N}$, we have $d_C^{(n)}(\phi^k, \phi) \to 0$ as $k \to \infty$. Given $\varepsilon > 0$, choose $\varepsilon' > 0$ and $\eta > 0$ so that $\varepsilon' + 2\eta < \varepsilon$. Then choose $m \in \mathbb{N}$ so that $\varepsilon' + 2\eta + 1/m < \varepsilon$ and so that $\|\phi_{ts} - \operatorname{id}\| < \eta$ for all $s, t \in (-n, n)$ with 0 < t - s < 1/m. Consider the finite set

$$F = (m^{-1}\mathbb{Z} \cap [-n, n]) \times (m^{-1}\mathbb{Z} \cap [0, 1]).$$

There exists a $K < \infty$ such that, for all $k \ge K$, all $(s_0, x_0) \in F$, and all $t \in (s_0, n]$,

$$|\phi_{(s_0,t]}^{k,+}(x_0) - \phi_{ts}^+(x_0)| = |\phi_{(s_0,t]}^{k,-}(x_0) - \phi_{ts}^-(x_0)| < \varepsilon'.$$

For all $s \in [-n, n)$ and all $x \in [0, 1)$, there exists $(s_0, x_0) \in F$ such that

$$s_0 \leq s < s_0 + 1/m, \quad x_0 \leq x + \varepsilon' + \eta + 1/m < x_0 + 1/m.$$

Then

$$\phi_{(s_0,s]}^{k,+}(x_0) \ge \phi_{ss_0}^+(x_0) - \varepsilon' \ge x_0 - \varepsilon' - \eta > x,$$

 \mathbf{SO}

$$\phi_{(s_0,t]}^{k,+}(x_0) \geqslant \phi_{(s,t]}^{k,+}(x), \quad t \geqslant s.$$

Also, we have

$$\phi_{ss_0}^+(x_0) \leqslant x_0 + \eta \leqslant x + \varepsilon' + 2\eta + 1/m < x + \varepsilon,$$

so

$$\phi_{ts_0}^+(x_0) \leqslant \phi_{ts}^+(x+\varepsilon), \quad t \ge s.$$

Now, for all $t \in (s, n]$,

$$\phi_{(s_0,t]}^{k,+}(x_0) \leqslant \phi_{ts_0}^+(x_0) + \varepsilon_s$$

so

 $\phi_{(s,t]}^{k,+}(x) \leqslant \phi_{ts}^+(x+\varepsilon) + \varepsilon.$

By a similar argument, for all $t \in (s, n]$,

$$\phi_{(s,t]}^{k,-}(x) \ge \phi_{ts}^{-}(x-\varepsilon) - \varepsilon,$$

so $d_{\mathcal{D}}(\phi_{(s,t]}^k, \phi_{(s,t]}) \leq \varepsilon$. Hence $d_C^{(n)}(\phi^k, \phi) \to 0$ as $k \to \infty$, as required.

10.3 From the coalescing Brownian flow to the Brownian webs

The coalescing Brownian flow, which provides the limit object for our main result is a refinement, in certain respects, of Arratia's flow of coalescing Brownian motions, via the work of Tóth and Werner. In a series of works, beginning with [6], Fontes, Newman and others have already provided another such refinement, in fact several, which they call Brownian webs. They direction taken by Fontes et al. emphasises path properties: the Brownian web is conceived as a random element of a space \mathcal{H} of compact collections of \mathbb{R} -valued paths with specified starting points. In our formulation, one does not see so clearly the possible varieties of path, but we find the state space $C^{\circ}(\mathbb{R}, \mathcal{D})$ a convenient one, with a natural time-reversal map, and just one candidate for a probability measure corresponding to Arratia's flow. We now attempt to clarify the relationship between the coalescing Brownian flow and one of the Brownian webs.

In [6], the authors make the comment that there is more than one natural distribution of an \mathcal{H} -valued⁹ random variable \mathcal{W} that satisfies the following two conditions.

- (i) From any deterministic point (x, t) in space-time, there is almost surely a unique path $W_{x,t}$ in \mathcal{W} starting from (x, t).
- (ii) For any deterministic n and $(x_1, t_1), \ldots, (x_n, t_n)$, the joint distribution of $W_{x_1,t_1}, \ldots, W_{x_n,t_n}$ is that of coalescing Brownian motions (with unit diffusion constant).

 $^{^{9}}$ We refer to [6] for precise definitions and attempt here only to give a flavour of their results.

The standard Brownian web satisfies (i) and (ii) together with a certain minimality condition. On the other hand, the forward full Brownian web, introduced in [7], satisfies (i) and (ii) together with a certain maximality condition, subject to a non-crossing condition. Also in [7], the authors characterize a third object, the full Brownian web, which is a random variable on the space \mathcal{H}^F of compact collections of paths from $\mathbb{R} \to \mathbb{R}$, and explain how this is naturally related to the other Brownian webs.

We now describe in detail the space \mathcal{H}^F of the full Brownian web and give its characterization. We then show that there is a natural way to realise the full Brownian web (and hence also the standard Brownian web and forward full Brownian web) as a random variable on $C^{\circ}(\mathbb{R}, \mathcal{D})$. Define the function $\Phi : [-\infty, \infty] \times \mathbb{R} \to [0, 1]$ by

$$\Phi(x,t) = \frac{\tanh(x)}{1+|t|}.$$

Now construct the two metric spaces (Π^F, d) and $(\mathcal{H}^F, d_{\mathcal{H}^F})$ as follows. Let Π^F denote the set of functions $f : \mathbb{R} \to [-\infty, \infty]$ such that $\Phi(f(t), t)$ is continuous, and define

$$d^{F}(f_{1}, f_{2}) = \sup_{t \in \mathbb{R}} |\Phi(f_{1}(t), t) - \Phi(f_{2}(t), t)|.$$

Then (Π^F, d^F) is a complete separable metric space. Now let \mathcal{H}^F denote the set of compact subsets of (Π^F, d^F) , with $d_{\mathcal{H}^F}$ the induced Hausdorff metric

$$d_{\mathcal{H}^F}(K_1, K_2) = \sup_{g_1 \in K_1} \inf_{g_2 \in K_2} d^F(g_1, g_2) \lor \sup_{g_2 \in K_2} \inf_{g_1 \in K_1} d^F(g_1, g_2).$$

The space $(\mathcal{H}^F, d_{\mathcal{H}^F})$ is also complete and separable. Let $\mathcal{F}_{\mathcal{H}^F}$ be the Borel σ -algebra on \mathcal{H}^F . The full Brownian web is defined in [7] as follows.

Definition 10.12. A full Brownian web $\overline{\mathcal{W}}^F$ is any $(\mathcal{H}^F, \mathcal{F}_{\mathcal{H}^F})$ -valued random variable whose distribution has the following properties.

- (a) Almost surely the paths of $\bar{\mathcal{W}}^F$ are noncrossing (although they may touch, including coalescing and bifurcating).
- (b₁) From any deterministic point $(x,t) \in \mathbb{R}^2$, there is almost surely a unique path $W_{x,t}^F$ passing through x at time t.
- (b₂) For any deterministic m, $\{(x_1, t_1), \ldots, (x_m, t_m)\}$, the joint distribution of the semipaths $\{W_{x_j,t_j}^F(t), t \ge t_j, j = 1, \ldots, m\}$ is that of a flow of coalescing Brownian motions (with unit diffusion constant).

They show that any two full Brownian webs have the same distribution. We make a variation of this definition, to tie up with our work, in requiring in (b_2) that the semipaths have the law of coalescing Brownian motions on the circle.

Fix E, a countable subset of \mathbb{R}^2 containing \mathbb{Q}^2 . We define a map $\theta_F : C^{\circ}(\mathbb{R}, \mathcal{D}) \to \mathcal{H}^F$ as follows. Let

$$\Pi = \bigcup_{t \in \mathbb{R}} C([t, \infty), [-\infty, \infty]).$$

Let $\overline{\mathcal{H}}$ be the set of all subsets A of $\Pi \cup \Pi^F$ having the following noncrossing property

for all $f, g \in A$ and all $s, t \in \text{dom}(f) \cap \text{dom}(g), f(s) < g(s)$ implies $f(t) \leq g(t)$.

We say that a set $U \in \overline{\mathcal{H}}$ is maximal if, for any $f \in \Pi \cup \Pi^F$, $U \cup \{f\} \in \overline{\mathcal{H}}$ implies $f \in U$. For $\phi \in C^{\circ}(\mathbb{R}, \mathcal{D})$, define $\theta_0(\phi) = \{Z^{e,+}(\phi) : e \in E\}$. Note that, since E is dense in \mathbb{R}^2 , if $f, g \in \Pi \cup \Pi^F$ and $\theta_0(\phi) \cup \{f\}, \theta_0(\phi) \cup \{g\} \in \overline{\mathcal{H}}$, then also $\theta_0(\phi) \cup \{f, g\} \in \overline{\mathcal{H}}$. Hence there exists a unique maximal set $\theta_{FF}(\phi) \in \overline{\mathcal{H}}$ containing $\theta_0(\phi)$. Furthermore, this set is independent of the choice of E. Define $\theta_F(\phi) = \theta_{FF}(\phi) \cap \Pi^F$.

This can be viewed as a random variable on $C^{\circ}(\mathbb{R}, \mathcal{D})$ as a consequence of Propositions 10.13 and 10.15 below. We note that, under the measure μ_W , the conditions to be a full Brownian web are almost surely satisfied: (a) is immediate from the construction; (b_1) follows from Proposition 10.14, where condition (i) holds almost surely by Proposition 10.10, and condition (ii) depends on a countable number of almost sure conditions; (b_2) follows from Theorem 5.1. In fact θ_{FF} defined above is a forward full Brownian web, and the standard Brownian web can be constructed similarly, but we omit the details here.

Proposition 10.13. The set $\theta_F(\phi) \subset \Pi^F$ is compact.

Proof. Suppose that $f_1, f_2, \ldots \in \theta_F(\phi)$. Since paths in $\theta_F(\phi)$ are noncrossing, there exists a subsequence n_r such that $f_{n_r}(t)$ is monotone for all $t \in \mathbb{R}$ and so there exists some $f : \mathbb{R} \to [-\infty, \infty]$ such that $f_{n_r} \to f$ pointwise.

Since $\phi_{ts}(\cdot)$ is periodic with period 1, if $g \in \theta_F(\phi)$, then $g + m \in \theta_F(\phi)$ for all $m \in \mathbb{Z}$. Suppose $f(s) = \infty$ for some $s \in \mathbb{R}$. By restricting to a further subsequence if necessary, we may assume $f_{n_r}(s) \ge f_{n_{r-1}}(s) + 1$ for all r. But then, by noncrossing, $f_{n_r}(t) \ge f_{n_{r-1}}(t) + 1$ for all $t \in \mathbb{R}$. Therefore $f(t) = \infty$ for all $t \in \mathbb{R}$ and $f_{n_r} \to f$ in (Π^F, d^F) . A similar argument applies if $f(s) = -\infty$ for some $s \in \mathbb{R}$. Hence we may assume $|f(t)| < \infty$ for all $t \in \mathbb{R}$.

Note that $d^F(f,g) \leq d^{(n)}(f,g) \vee 2/(n+1)$ where

$$d^{(n)}(f,g) = \sup_{-n < t < n} |f(t) - g(t)|.$$

Given $\varepsilon > 0$, there exist $n > 2\varepsilon^{-1}$, and $-n = a_0 < \cdots < a_M = n$ such that $\|\phi_{sa_k}^+ - \operatorname{id}\| < \frac{\varepsilon}{4}$ for all $a_k \leq s \leq a_{k+1}$, $k = 0, \ldots, M - 1$. Pick N sufficiently large that if $n_r > N$,

 $|f_{n_r}(a_k) - f(a_k)| < \frac{\varepsilon}{4}$ for $k = 1, \ldots, M - 1$. Then for $a_k \leq s < a_{k+1}$,

$$|f_{n_r}(s) - f(s)| \le |\phi_{sa_k}^+(f(a_k) + \frac{\varepsilon}{4}) - \phi_{sa_k}^+(f(a_k) - \frac{\varepsilon}{4})| < \varepsilon.$$

Hence, $f_{n_r} \to f$ in (Π^F, d^F) . Therefore, f is continuous and $\theta_{FF}(\phi) \cup \{f\}$ is noncrossing, and so $f \in \theta_F(\phi)$ proving compactness.

Proposition 10.14. For every $(x, s) \in \mathbb{R}^2$, there exists some $f \in \theta_F(\phi)$ with f(s) = x. Moreover, f is unique if the following two conditions hold.

- (i) $\phi_{ts}^+(x) = \phi_{ts}^+(x)$ for all $t \ge s$;
- (ii) $\phi_{st}^+(y) \neq x$ for all $(y,t) \in \mathbb{Q}^2$ with t < s.

Proof. Define a function $f : \mathbb{R} \to \mathbb{R}$ by

$$f(t) = \begin{cases} \inf\{y : \phi_{st}^+(y) > x\} & \text{if } t < s \\ \phi_{ts}^+(x) & \text{if } t \ge s. \end{cases}$$

We first show that f is continuous. Clearly this is the case on $[s, \infty)$. Suppose $t < u \leq s$. Pick a sequence $y_n \downarrow f(t)$. By the weak flow property of ϕ , $\phi_{su}^+(\phi_{ut}^+(y_n)) \geq \phi_{st}^+(y_n) > x$ and so $\phi_{ut}^+(y_n) \geq f(u)$ for all n. Letting $n \to \infty$ gives $\phi_{ut}^+(f(t)) \geq f(u)$. Similarly, using $\inf\{y: \phi_{st}^+(y) > x\} = \sup\{y: \phi_{st}^-(y) \leq x\}, \phi_{ut}^-(f(t)) \leq f(u)$. Now given $\varepsilon > 0$, there exists some $\delta > 0$ such that $\|\phi_{ut} - \operatorname{id}\| < \varepsilon$ for all $0 < u - t < \delta$. But then $|f(t) - f(u)| \leq |f(t) - \phi_{ut}^-(f(t))| < |f(t) - \phi_{ut}^+(f(t))| < \varepsilon$, proving continuity. Also, by construction, $\theta_0(\phi) \cup \{f\} \in \overline{\mathcal{H}}$ and so $f \in \theta_F(\phi)$ as required.

For uniqueness, suppose $f_1, f_2 \in \theta_F(\phi)$ with $f_1(s) = f_2(s) = x$. By the noncrossing property, for all $t \ge s$, $\phi_{ts}^-(x) \le f_i(t) \le \phi_{ts}^+(x)$. Condition (i) therefore implies $f_1(t) = f_2(t)$ for all $t \ge s$. If t < s, then $\sup\{y : \phi_{st}^-(y) < x\} \le f_i(t) \le \inf\{y : \phi_{st}^+(y) > x\}$. If $\inf\{y : \phi_{st}^+(y) > x\} - \sup\{y : \phi_{st}^-(y) < x\} > 0$, then by continuity, there exist rationals (y, u)with $\phi_{st}^+(y) = x$. Hence (ii) ensures $f_1(t) = f_2(t)$ for all t < s.

Proposition 10.15. The function $\theta_F : C^{\circ}(\mathbb{R}, \mathcal{D}) \to \mathcal{H}^F$ is continuous.

Proof. A sequence $\phi^m \to \phi$ in $(C^{\circ}(\mathbb{R}, \mathcal{D}), d_C)$ if and only if $d_C^{(n)}(\phi^m, \phi) \to 0$ for all $n \in \mathbb{N}$. Since $d^F(f,g) \leq d^{(n)}(f,g) \vee 2/(n+1)$, where $d^{(n)}$ is defined in the previous proposition, in order to show that θ_F is continuous, it is enough to show that for every n, $d_{\mathcal{H}^F}^{(n)}(\theta_F(\phi^1), \theta_F(\phi^2)) \leq d_C^{(n)}(\phi^1, \phi^2)$ for all $\phi^1, \phi^2 \in C^{\circ}(\mathbb{R}, \mathcal{D})$, where

$$d_{\mathcal{H}^F}^{(n)}(K_1, K_2) = \sup_{g_1 \in K_1} \inf_{g_2 \in K_2} d^{(n)}(g_1, g_2) \lor \sup_{g_2 \in K_2} \inf_{g_1 \in K_1} d^{(n)}(g_1, g_2).$$

Suppose $\phi^1, \phi^2 \in C^{\circ}(\mathbb{R}, \mathcal{D})$. By compactness, there exist $f_1 \in \theta_F(\phi^1)$, $f_2 \in \theta_F(\phi^2)$ such that $d^{(n)}(f_1, f_2) = d^{(n)}_{\mathcal{H}^F}(\theta_F(\phi^1), \theta_F(\phi^2))$. Without loss of generality, suppose that f_2 is chosen so that $d^{(n)}(f_1, f_2) = \inf_{g \in \theta_F(\phi^2)} d^{(n)}(f_1, g) = f_2(s) - f_1(s)$ for some $s \in \mathbb{R}$. Let $f_1(s) = x$ and $f_2(s) = y$. By continuity, there exists some t > s such that $f_1(t) - \phi^-_{ts}(y) = d^{(n)}(f_1, f_2)$, otherwise it would be possible to pick some $g \in \theta_F(\phi^2)$ with $d^{(n)}(f_1, g) < d^{(n)}(f_1, f_2)$. Since $f_1(t) \in [\phi^{1,-}_{ts}(x), \phi^{1,+}_{ts}(x)]$,

$$\phi_{ts}^{2,-}(y) - \phi_{ts}^{1,+}(x) \le x - y \le \phi_{ts}^{2,+}(y) - \phi_{ts}^{1,-}(x).$$

Hence there exists some

$$u \in \left[\frac{1}{2}(x + \phi_{ts}^{1,-}(x)), \frac{1}{2}(x + \phi_{ts}^{1,+}(x))\right] \cap \left[\frac{1}{2}(y + \phi_{ts}^{2,-}(y)), \frac{1}{2}(y + \phi_{ts}^{2,+}(y))\right].$$

Therefore,

$$d_{\mathcal{H}^{F}}(\theta_{F}(\phi^{1}), \theta_{F}(\phi^{2})) = |x - y|$$

= $|\phi_{ts}^{1\times}(u) - \phi_{ts}^{2\times}(u)|$
 $\leq ||\phi_{ts}^{1\times} - \phi_{ts}^{2\times}||$
 $\leq d_{C}^{(n)}(\phi^{1}, \phi^{2}),$

as required.

References

- Richard A. Arratia. Coalescing Brownian motions on the line. PhD thesis, University of Wisconsin, Madison, 1979.
- [2] Martin Z. Bazant and Darren Crowdy. Conformal mapping methods for interfacial dynamics. arXiv:cond-mat/0409439, 2005.
- [3] Patrick Billingsley. Convergence of probability measures. Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley & Sons Inc., New York, 1999.
- [4] L. Carleson and N. Makarov. Aggregation in the plane and Loewner's equation. Comm. Math. Phys., 216(3):583–607, 2001.
- [5] Murray Eden. A two-dimensional growth process. In Proc. 4th Berkeley Sympos. Math. Statist. and Prob., Vol. IV, pages 223–239. Univ. California Press, Berkeley, Calif., 1961.

- [6] L. R. G. Fontes, M. Isopi, C. M. Newman, and K. Ravishankar. The Brownian web: characterization and convergence. *Ann. Probab.*, 32(4):2857–2883, 2004.
- [7] Luiz Renato Fontes and Charles M. Newman. The full Brownian web as scaling limit of stochastic flows. Stoch. Dyn., 6(2):213–228, 2006.
- [8] Theodore E. Harris. Coalescing and noncoalescing stochastic flows in R¹. Stochastic Process. Appl., 17(2):187–210, 1984.
- [9] M. B. Hastings and L. S. Levitov. Laplacian growth as one-dimensional turbulence. *Physica D*, 116 (1-2):244, 1998.
- [10] Jean Jacod and Albert N. Shiryaev. Limit theorems for stochastic processes, volume 288 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 2003.
- [11] Harry Kesten. Hitting probabilities of random walks on \mathbb{Z}^d . Stochastic Process. Appl., 25(2):165–184, 1987.
- [12] Gregory F. Lawler. Conformally invariant processes in the plane, volume 114 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2005.
- [13] L. Niemeyer, L. Pietronero, and H. J. Wiesmann. Fractal dimension of dielectric breakdown. *Phys. Rev. Lett.*, 52:1033–1036, 1984.
- [14] Vladimir V. Piterbarg. Expansions and contractions of isotropic stochastic flows of homeomorphisms. Ann. Probab., 26(2):479–499, 1998.
- [15] Steffen Rohde and Michel Zinsmeister. Some remarks on Laplacian growth. *Topology* Appl., 152(1-2):26–43, 2005.
- [16] Bálint Tóth and Wendelin Werner. The true self-repelling motion. Probab. Theory Related Fields, 111(3):375–452, 1998.
- [17] Boris Tsirelson. Nonclassical stochastic flows and continuous products. Probab. Surv., 1:173–298 (electronic), 2004.
- [18] T. A. Witten and L. M. Sander. Diffusion-limited aggregation, a kinetic critical phenomenon. Phys. Rev. Lett., 47(19):1400–1403, 1981.