

Quantum switching at a mean-field instability of a Bose-Einstein condensate in an optical lattice

V. S. Shchesnovich¹ and V. V. Konotop²

¹*Centro de Ciências Naturais e Humanas,*

Universidade Federal do ABC, Santo André, SP, 09210-170 Brazil,

²*Centro de Física Teórica e Computacional,*

Universidade de Lisboa, Complexo Interdisciplinar,

Avenida Professor Gama Pinto 2, Lisboa 1649-003,

Portugal; Departamento de Física, Faculdade de Ciências,

Universidade de Lisboa, Campo Grande,

Ed. C8, Piso 6, Lisboa 1749-016, Portugal

Abstract

It is shown that bifurcations of the mean-field dynamics of a Bose-Einstein condensate can be related with the quantum phase transitions of the original many-body system. As an example we explore the inter-band tunneling in the two-dimensional optical lattice. Such a system allows for easy control by the lattice depth as well as for macroscopic visualization of the phase transition. The system manifests switching between two self-trapping states or from a self-trapping state to a superposition of the macroscopically populated self-trapping states with the step-like variation of the control parameter about the bifurcation point. We have also observed the magnification of the microscopic difference between the even and odd number of atoms to a macroscopically distinguishable dynamics of the system.

PACS numbers: 03.75.Lm; 03.75.Nt

Since the very beginning of the quantum mechanics its relation to the classical dynamics constitutes one of the central questions of the theory. Dependence of the distribution of energy levels of a quantum system on the type of dynamics of the corresponding classical system [1], in general, and its response to variation of the bifurcation parameters controlling the qualitative changes of the classical behavior [2] are among the major issues [3]. One of the main tools in studies of the quantum-classical correspondence is the WKB approximation, where, loosely speaking, the Planck constant \hbar is regarded as a small parameter.

On the other hand, it is known that for an N -boson system the limit $N \rightarrow \infty$ at a constant density, leading to the mean-field approximation, can also be understood as a semiclassical limit. This latter approach has received a great deal of attention during the last decade [4], due its high relevance to the theory of Bose-Einstein condensates (BECs), many properties of whose dynamics are remarkably well described within the framework of the mean-field models [5]. More recently, it was shown [6, 7] that the mean-field description of a few-mode N -boson system can be recast in a form similar to the WKB approximation for a discrete Schrödinger equation [8], emergent for the coefficients of the wavefunction expansion in the associated Fock space, where $1/N$ plays the role similar to that of the Planck constant in the conventional WKB approximation.

The mean-field equations of a system of interacting bosons are nonlinear, hence, they naturally manifest many common features of the nonlinear dynamics, including bifurcations of the stationary solutions caused by variation of the system parameters. One of the well studied examples is a boson-Josephson junction [9], which can show either equally populated (symmetric) or strongly asymmetric states, characterized by population of only one of the sites (the well known phenomenon of self-trapping [10]). Now, exploring parallels between the semiclassical approach and the mean-field approximation one can pose the natural question: *what changes occur in a manybody system when a control parameter crosses an instability (e.g. bifurcation) point of the limiting mean-field system?*

In the present Letter we aim to give at least partial answer, showing that one of the possibilities is the occurrence of the quantum phase transition of the second type, associated with the switching of the wave-function in the Fock space between the “coherent” and “Bogoliubov” states, possessing distinct features. Considering a flexible (time-dependent) control parameter, we have also found a strong sensitivity of the system to the parity of the total number of atoms N , showing parity-dependent structure of the energy levels and the

macroscopically different dynamics for different parity of N . Observation of the discussed phenomena is feasible in the experimental setting available nowadays.

We consider the nonlinearity-induced intra-band tunneling of BEC within the Brillouin zone $(-q_B, q_B) \times (-q_B, q_B)$ of a two dimensional square optical lattice (OL), specifically, between the two high-symmetry X -points, i.e. with the quasimomenta $\mathbf{q}_1 = (q_B, 0)$ and $\mathbf{q}_2 = (0, q_B)$ and equal energies. This leads to the two mode boson Hamiltonian, where the Bloch waves $\varphi_{\mathbf{q}_j}(\mathbf{r})$ [6] are the basis:

$$\hat{H} = \frac{1}{2N^2} \left\{ n_1^2 + n_2^2 + \Lambda \left[4n_1n_2 + (b_1^\dagger b_2)^2 + (b_2^\dagger b_1)^2 \right] \right\}. \quad (1)$$

Hereafter, b_j and b_j^\dagger are the annihilation and creation operators of the state $\varphi_{\mathbf{q}_j}(\mathbf{r})$, $n_j = b_j^\dagger b_j$, and $\Lambda = \chi_{11}/\chi_{12}$ ($0 \leq \Lambda \leq 1$), where $\chi_{ij} = \int d^3\mathbf{r}(\varphi_{\mathbf{q}_i}^*)^2(\varphi_{\mathbf{q}_j})^2$. The lattice parameter Λ is easily controlled by variation of the lattice depth. The Schrödinger equation for the BEC in a state $|\Psi\rangle$ reads $ih\partial_\tau|\Psi\rangle = \hat{H}|\Psi\rangle$, where $\tau = 2gN\chi_{11}t/\hbar$ is the scaled time, $g = 4\pi\hbar^2 a_s/m$, and $h = 2/N$.

The N -factor in the Hamiltonian \hat{H} becomes clear when the Schrödinger equation is written in the Fock basis, $|k, N-k\rangle = \frac{(b_1^\dagger)^k (b_2^\dagger)^{N-k}}{\sqrt{k!(N-k)!}}|0\rangle$, where only the relative populations k/N and $(N-k)/N$ appear, while h serves as an effective ‘‘Planck constant’’ (the scaled time τ is N -independent for a constant density, i.e. in the thermodynamic limit $N \rightarrow \infty$).

Hamiltonian (1) represents a nonlinear version of the well-known boson-Josephson model (see, e.g. [9, 11]), where the quantum modes are coupled by the exchange of *atom pairs*, what is a fairly common situation for systems with four-wave-mixing, provided by the two-body interactions involving four bosons. The exchange of the bosons by pairs results in the coupling of the states with the same parity of the population and is reflected in the double degeneracy of all $(N+1)/2$ energy levels for odd N , evident from the symmetry relation $\langle 2k, N-2k|\Psi_1\rangle = \langle N-2k, 2k|\Psi_2\rangle$ valid for odd N . For even N the energy levels show quasi degeneracy (see below).

Next we recall, that in the mean-field limit, the system (1) is described by the classical Hamiltonian [6]

$$\mathcal{H} = x(1-x)[2\Lambda - 1 + \Lambda \cos \phi] + \frac{1}{2}, \quad (2)$$

where $x = \langle n_1 \rangle / N$ is the population density and $\phi = \arg\langle (b_2^\dagger)^2 b_1^2 \rangle$ is the relative phase. Formally \mathcal{H} is obtained by replacing the boson operators b_j in (1) by the c-numbers $b_1 \rightarrow \sqrt{Nx}e^{i\phi/4}$ and $b_2 \rightarrow \sqrt{N(1-x)}e^{-i\phi/4}$.

Hamiltonian (2) possesses two stationary points describing equally populated X -states: the classical energy maximum $P_1 = (x = \frac{1}{2}, \phi = 0)$ and minimum $P_2 = (x = \frac{1}{2}, \phi = \pi)$. P_1 is dynamically stable in the domain $\Lambda > \Lambda_c = \frac{1}{3}$. For $\Lambda < \Lambda_c$ it loses its stability, and another set of stationary points $x = 1$ (S_1) and $x = 0$ (S_2) appears. These solutions describe the symmetry breaking leading to self-trapping, i.e. when all atoms populate just one X -state.

The spectrum of the Hamiltonian (1) in the vicinity of the critical value Λ_c allows for an analytical analysis. To emphasize the role of the control parameter we rewrite \hat{H} in terms of the operators $a_{1,2} = (b_1 \mp ib_2)/\sqrt{2}$

$$\hat{H} = \hat{H}_0 + \left(1 - \frac{\Lambda}{\Lambda_c}\right) \hat{V} + \mathcal{E}(\Lambda), \quad \mathcal{E}(\Lambda) = \frac{\Lambda + 1}{4} + \frac{\Lambda}{2N} \quad (3)$$

where $\hat{H}_0 = \frac{2\Lambda}{N^2} a_1^\dagger a_1 a_2^\dagger a_2$ and $\hat{V} = \frac{1}{4N^2} (a_1^\dagger a_2 + a_2^\dagger a_1)^2$. At the critical point the energy spectrum is determined by \hat{H}_0 : $E_m = \frac{2\Lambda_c}{N^2} m(N - m) + \mathcal{E}(\Lambda_c)$, where m is the occupation number corresponding to the operator $a_1^\dagger a_1$. The spectrum of \hat{H}_0 is doubly degenerate (except for the top level for even N) due to the symmetry $m \rightarrow N - m$. The ground state energy is $\mathcal{E}_{min}(\Lambda_c) = E_0 = E_N$, while the top energy level has $m = N/2$ for even N and $m = (N \pm 1)/2$ for odd N . Restricting ourselves to even number of bosons we get $\mathcal{E}_{max}(\Lambda_c) = \frac{1}{2} + \frac{\Lambda_c}{2N}$.

Now consider small deviations of Λ from the bifurcation point Λ_c . To this end, for a fixed N , one can use the basis consisting of the degenerate eigenstates of \hat{H}_0 : $|E_m, j\rangle = \frac{(a_j^\dagger)^m (a_{3-j}^\dagger)^{N-m}}{\sqrt{m!(N-m)!}} |0\rangle$, $j = 1, 2$, $m = 0, \dots, \frac{N}{2}$. The conditions for \hat{V} to be treated as a perturbation depend on m as is seen, for instance, from the diagonal matrix elements:

$$\langle E_m, j | \hat{V} | E_m, j \rangle = \frac{1}{4N} + \frac{m}{2N} \left(1 - \frac{m}{N}\right). \quad (4)$$

At the lower levels ($m \ll N/2$) the energy gaps between the degenerate subspaces and the perturbation both scale as $\Delta E \sim N^{-1}$, hence the condition of applicability is $|\Lambda - \Lambda_c| \ll 1$ and the lower energy subspaces acquire simple shifts. At the upper energy levels ($m \sim N/2$) the above energy gaps behave as $\Delta E \sim N^{-2}$. Since $\langle \hat{V} \rangle \sim 1$ in this case, the perturbation theory is applicable only in an interval of Λ of the size on the order of N^{-2} . There is a dramatic transition in the energy levels, e.g. Fig. 1(b) shows the exchange of the double degeneracy of the top levels for even N in this N^{-2} -small interval of Λ . Finally, by considering the phase of

$$\langle E_m, j | (b_2^\dagger b_1)^2 | E_m, j \rangle = -\frac{N^2}{4} + \frac{N}{4} + \frac{3}{2} m(N - m), \quad (5)$$

one sees that the upper and lower eigenstates correspond to the mean-field stationary points P_1 with $\phi = 0$, and P_2 with $\phi = \pi$.




FIG. 1: (a) The energy levels of \hat{H} for $N = 200$ and (b) a detailed picture in the vicinity of Λ_c . The classical energy lines of the mean-field fixed points P_1 and S_2 are visibly formed. The top energy levels for sufficiently large $|\Lambda - \Lambda_c|$ are quasi-degenerate with the inter-level distances indistinguishable on the scale of the figure (see the discussion in the text below).

For $\Lambda_c^{-1} - \Lambda^{-1} \gg N^{-2}$ the quantum states corresponding to P_1 can be obtained by reduction of the Schrödinger equation in Fock space (similar to the case of boson-Josephson model [12]) or quantizing the local classical Hamiltonian (2) by expanding it with respect to $\xi = x - 1/2$ and ϕ (on this way one loses the term of order $1/N$ in \mathcal{E}_{max}). One assumes the “wave function” $\psi(x) = \sqrt{N}C_k \equiv \sqrt{N}\langle k, N - k|\psi\rangle$, where $x = k/N$, to be localized about $x = 1/2$ and obtains:

$$\left[-\frac{\Lambda}{8}\hat{p}_\xi^2 + U(\Lambda)(1 - 4\xi^2) \right] \psi_E = E\psi_E, \quad (6)$$

with $\hat{p} = -i\hbar\partial_\xi$ and $U(\Lambda) = \frac{3\Lambda-1}{4}$. Eq. (6) is the quantum oscillator problem with the frequency $\omega^2 = 8(3 - \frac{1}{\Lambda})$. The respective descending energy levels read $E_n^{(top)} = \mathcal{E}_{max} + \frac{1}{4}\left(\frac{\Lambda}{\Lambda_c} - 1\right) - \frac{\hbar\Lambda\omega}{4}\left(n + \frac{1}{2}\right)$. Since the eigenfunctions are localized in Fock space, for instance, the $n = 0$ eigenfunction is $\psi_0(x) = C \exp\left[-\frac{\omega}{2\hbar}\left(x - \frac{1}{2}\right)^2\right]$, returning to the discrete Fock

basis $x = k/N$ we see that there are even and odd eigenstates C_{2k} and C_{2k-1} related by the approximate symmetry $C_n \approx C_{n+1}$, hence the energy levels are quasi doubly degenerate [c.f. Fig. 1(b)].

The local approximation becomes invalid as $\Lambda_c^{-1} - \Lambda^{-1} \sim N^{-2}$ (the wave-function delocalizes). Correspondingly, another set of the stationary points, $S_{1,2}$, becomes stable for $\Lambda < \Lambda_c$ in the mean-field case. In this case, however, the phase ϕ is undefined. Let us first consider the full quantum case, i.e. the limit $\langle n_j \rangle \ll N$, say for $j = 1$ (i.e. the point S_2). The resulting reduced Hamiltonian can be easily derived in the Fock basis, it also obtains by formally setting $b_2 = N$ and retaining the lowest-order terms in b_1 and b_1^\dagger :

$$\hat{H} \approx \hat{H}_{S_2} = \frac{1}{2} + \frac{(2\Lambda - 1)}{N} b_1^\dagger b_1 + \frac{\Lambda}{2N} [(b_1^\dagger)^2 + b_1^2]. \quad (7)$$

Hamiltonian (7) can be diagonalized by the Bogoliubov transformation $c = \cosh(\theta)b_1 - \sinh(\theta)b_1^\dagger$, where $\theta = \theta(\Lambda) > 0$ is determined from $\tanh(2\theta) = \Lambda/(1 - 2\Lambda)$. We get

$$\hat{H}_{S_2} = -\frac{\Lambda}{N \sinh(2\theta)} c^\dagger c + \frac{\Lambda \tanh \theta}{2N} + \frac{1}{2}. \quad (8)$$

Thus $c^\dagger c$ gives the number of negative-energy quasi-particles over the Bogoliubov (squeezed) vacuum solving $c|\text{vac}\rangle = 0$. In the atom-number basis $|\text{vac}\rangle$ is a superposition of the Fock states with

$$C_{2k}^{(\text{vac})} = \frac{\tanh^k(\theta)}{2^k} \frac{\sqrt{(2k)!}}{k!} C_0, \quad C_{2k-1}^{(\text{vac})} = 0, \quad (9)$$

where C_0 is obtained by the normalization.

The parameter θ determines the average number of atoms in the Bogoliubov vacuum $\langle \hat{n} \rangle = \sinh^2(\theta)$ and its fluctuations $\Delta \hat{n} = \sinh(2\theta)/\sqrt{2}$. Thus the validity condition of the approximation (7), given by $n, \Delta n \ll N$, can be rewritten in the form $\tanh^{-2}(2\theta) \gg 1 + N^{-2}$, what is the same as $\Lambda^{-1} - \Lambda_c^{-1} \gg N^{-2}$. In this case, the Bogoliubov states, in particular the squeezed vacuum $|\text{vac}\rangle$, are well-localized in the atom-number Fock space, i.e. the coefficients C_{2k} decay fast enough. The above condition coincides with the quantum perturbation theory result, i.e. that the transition between the coherent states and the Bogoliubov states occurs on the interval of Λ of order of N^{-2} . The convergence of the eigenstates of \hat{H}_{S_2} to that of the full Hamiltonian (1) turns out to be remarkably fast as it is shown in Fig. 2(a). In Fig. 2b the dramatic deformation of the top energy eigenstate of \hat{H} (corresponding to the S_2 - P_1 transition) about the critical Λ_c is shown. Finally, we note that for even N the quasi

double degeneracy of the energy levels for $\Lambda^{-1} - \Lambda_c^{-1} \gg N^{-2}$ (c.f. Fig. 1(b)) is due to the exchange symmetry between S_1 and S_2 resulting in equal energy levels of the Hamiltonians \hat{H}_{S_1} and \hat{H}_{S_2} .




FIG. 2: (a) Convergence of the four upper eigenstates of the Hamiltonian (1) to the eigenstates of \hat{H}_{S_2} (shown by dots) for $N = 100$ (solid lines) and $N = 1000$ (dashed lines), for $\Lambda = \Lambda_c - 0.1$. (b) The contour plot of the state corresponding to the top energy level in the vicinity of Λ_c for $N = 200$.

Returning to the mean-field description of the stationary point S_2 , the associated Hamiltonian is defined by replacing the boson operators in Eq. (7) by the c-numbers $b_1 = \sqrt{N}\alpha$ and $b_2 = \beta$. Using $|\alpha|^2 + |\beta|^2 = 1$ and fixing the irrelevant common phase by setting β real we get the dynamical variables α and α^* and the classical Hamiltonian in the form $\mathcal{H}_{S_2} = \frac{1}{2} + \frac{1}{2}(1 - |\alpha|^2)\{2(2\Lambda - 1)|\alpha|^2 + \Lambda[\alpha^2 + (\alpha^*)^2]\}$, from which the stability of the point S_2 ($\alpha = 0$) for $\Lambda < \Lambda_c$ follows since the quadratic form $2(2\Lambda - 1)|\alpha|^2 + \Lambda[\alpha^2 + (\alpha^*)^2]$ [c.f. Eq. (7)] is positive.

Thus, the passage through the bifurcation point Λ_c of the mean-field model, manifested in the exchange of the stable stationary points, corresponds to the quantum phase transition of the original many-body system reflected in the deformation of the spectrum and dramatic change of the system wave-function in the Fock space. The described change of the system is related to the change of the symmetry of the particle distribution, and thus it is the second order phase transition.

In our case this scenario corresponds to loss of stability of the boundary solution S_1 and S_2 and to appearance of the stable stationary point P_1 . In the quantum description this happens by a set of avoided crossings of the top energy levels (and splitting of the quasi-degenerate energy levels for even N) as the parameter Λ sweeps the small interval on the order of N^{-2} about the critical value Λ_c (see Fig. 1). For lower energy levels the avoided crossings (and splitting of the energy levels for even N) also occur but along the two straight lines approximating the classical energies of the two involved stationary points: $\mathcal{H}(P_1) = \frac{1}{2} + \frac{(3\Lambda-1)}{4}$ (for $\Lambda < \Lambda_c$) and $\mathcal{E}_{max} = \frac{1}{2}$ ($\Lambda > \Lambda_c$), see Fig. 1.

The obtained results have strong implications for time-dependent variation of Λ (achievable experimentally by varying the intensity of the lasers creating the OL). The states with either $x = 0$ or $x = 1$ correspond to the occupation of just one of the high-symmetry X -points of the Brillouin zone. Such an initial condition can be experimentally created by switching on a moving lattice with $\Lambda < \Lambda_c$ (see e.g. [7]). As the lattice parameter $\Lambda(\tau)$ passes the critical value from below, the Bogoliubov states describing the self-trapping at one X -point are replaced by the coherent states (in Fock space) with comparable average occupations of the two X -points.

A more interesting dynamics is observed when $\Lambda(t)$ is a step-like function, which we chose in the form

$$\Lambda = \Lambda_1 + \frac{\Lambda_2 - \Lambda_1}{2} \left[\tanh\left(\frac{\tau - \tau_1}{T}\right) - \tanh\left(\frac{\tau - \tau_2}{T}\right) \right]. \quad (10)$$

Then the system dynamics and the resulting state dramatically depends not only on the macroscopic parameters, Λ_1 , Λ_2 , T and $\tau_2 - \tau_1$, but also on *parity* of the number of atoms, as shown in Fig. 3.

More specifically, we are able to identify two distinct scenarios which can be described as a switching dynamics between the self-trapping states at the two X -points or a dynamic creation of the superposition of the two macroscopically populated self-trapping states, see Fig. 3. For instance, for fixed $\Lambda_{1,2}$ and T we have observed that the system dynamics crucially depends on the hold time $\tau_2 - \tau_1$. Moreover, the formation of the superposition of macroscopically populated states shows anomalous dependence on parity of the number of atoms (i.e. giving the same behavior for two macroscopically different N of the same parity, while a macroscopically distinct for $N \pm 1$) see Fig. 3(c). We also note that the corresponding classical (i.e. mean-field) dynamics is close to the quantum one in the switching case, while




FIG. 3: The average population densities $\langle n_1 \rangle / N$, (a) and (c), and the atom-number probabilities $|C_k|^2$, (b) and (d), for $\Lambda(\tau)$ given by (10). The corresponding classical dynamics is shown by the dash-dot lines in (a) and (c). Here $\Lambda_1 = 0.25$, $\Lambda_2 = 0.5$, $T = 1$, $\tau_1 = 50$ and $\tau_2 = 85$ (a) with $N = 500$ and 501 (indistinguishable), while in (c) $\tau_2 = 135$ with $N = 500$ and 400 (the blue lines) and $N = 501$ and 401 (the red lines). The initial state is $|\text{vac}\rangle$ at S_1 , but using $\frac{(b_1^\dagger)^N}{\sqrt{N!}}|0\rangle$ gives a similar picture.

it is dramatically different in the superposition case.

In conclusion, we have demonstrated that behind the mean-field instability in the inter-band tunneling of BEC in an optical lattice is a quantum phase transition between macroscopically distinct states, giving a macroscopic magnification of the microscopic quantum features of the system. A spectacular demonstration of this is a dynamic formation of the superposition of two macroscopically distinct self-trapping states which shows an anomalous dependence on parity of BEC atoms reflecting distinct energy level structure for even and odd number of atoms.

The work of VSS was supported by the FAPESP of Brazil.

[1] I. C. Percival. J. Phys. B **6**, L229 (1973); M. V. Berry and M. Tabor, Proc. R. Soc. Lond. A **356**, 375 (1977).

- [2] P. Pechukas, Phys. Rev. Lett. **51**, 943 (1983); T. Yukawa, Phys. Rev. Lett. **54**, 1883 (1985).
- [3] see e.g. K. Nakamura, *Quantum Chaos* (Cambridge University Press, 1993); F. Haake, *Quantum Signatures of Chaos* (Springer-Verlag Berlin Heidelberg, 2001)
- [4] L. Pitaevskii and S. Stringari, *Bose-Einstein Condensation* (Clarendon Press, Oxford, 2003)
- [5] C. W. Gardiner, Phys. Rev. A **56**, 1414 (1997); Y. Castin and R. Dum, Phys. Rev. A **57**, 3008 (1998).
- [6] V. S. Shchesnovich and V. V. Konotop, Phys. Rev. A **75**, 063628 (2007).
- [7] V. S. Shchesnovich and V. V. Konotop, Phys. Rev. A **77**, 013614 (2008).
- [8] P. A. Braun, Rev. Mod. Phys. **65**, 115 (1993).
- [9] A. J. Leggett, Rev. Mod. Phys. **73**, 307 (2001).
- [10] V. M. Kenkre and D. K. Campbell, Phys. Rev. B **34**, 4959 (1986).
- [11] R. Gati and M. K. Oberthaler, J. Phys. B: At. Mol. Opt. Phys. **40**, R61 (2007).
- [12] V. S. Shchesnovich and M. Trippenbach, arXiv: cond-mat/08040234; Phys. Rev. A **78**, 023611 (2008).