Comment on "Electron screening and excitonic condensation in double-layer graphene systems"

Recently we[1, 2] predicted that a clean[3] doublelayer graphene system with a small layer separation and nested electron and hole Fermi spheres induced by external gates will exhibit pair-condensation[2] at temperatures ~ $0.1\epsilon_F[1]$, where ϵ_F is the Fermi energy of the gate-induced electron gases. A very different conclusion was reached in a recent preprint[4] by M. Y. Kharitonov and K. Efetov who conclude that the maximum paircondensation temperature is ~ $10^{-7}\epsilon_F$. The stark contrast between these estimates is a combined consequence of the different approximations used for the effective interlayer interaction and the extreme sensitivity of the condensation temperature to the strength of this interaction when it is weak.

Kharitonov and Efetov's estimate for the critical temperature is based on using a mean-field-theory linearized T_c equation combined with Thomas-Fermi screened interlayer interactions. As already discussed in our original paper, we agree with Kharitonov and Efetov that low transition temperature estimates follow from this procedure: "Screening and other beyond-mean-field inducedinteraction effects are difficult to describe. In the case of weakly interacting atomic gases induced interaction effects can[18] either increase or decrease T_c , depending on the number of fermion flavors g. For the present Coulomb interaction case, a static Thomas-Fermi screening approximation with normal state screening wavevectors reduces interaction strengths very substantially when spin and valley degeneracies (q = 4) are included. \cdots On the other hand, when the screening wavevectors are evaluated in the condensed state there is little influence on T_{KT} at small $k_{\rm F}d$ both because the large gap weakens screening and because T_{KT} is proportional to the Fermi energy and not to the interaction strength in this limit. All this leads us to suspect that at low-temperatures there is a first-order phase transition as a function of layer separation d between condensed and electron-hole plasma states, similar to the transitions studied experimentally[19] in quantum Hall exciton condensates and theoretically[20] in parabolic band bilayers." References [18-20] in Ref.[1] are listed here as References [5-7]. We elaborate on our views below.

Corrections to mean-field-theory estimates of T_c can be treated rigorously only in the limit of weak shortrange interactions[5]. For the circumstance[8] treated in Ref. [5], induced interactions corrections reduce T_c for g = 2, but enhance T_c for g = 4. When interlayer interactions are strong, the static screening approximation used by Kharitonov and Efetov does not represent a systematic improvement on mean-field theory even if the number of fermion flavors is large. Nevertheless, our assessment of the importance of screening for spontaneous bilayer coherence follows from the observation that two distinct solutions emerge when mean-field theory is consistently combined with Thomas-Fermi screening theory, the low T_c solution discussed by Kharitonov and Efetov and a high T_c solution. In our opinion the low T_c solution is unlikely to be physically realistic at any layer separation; we predict instead that there is a first-order phase transition as a function of layer separation between a state without spontaneous coherence and a high- T_c state as discussed in Ref. [1].

The influence of screening on the interlayer and intralayer interactions in bilayer systems was considered long ago in the context of semiconductor based bilayers. In the absence of interlayer coherence it is common practice to estimate the screened potentials using the random phase approximation. However, in the condensed state it is more difficult to consistently account for dynamical screening and we resort to the Thomas Fermi approximation. Within that approximation a straightforward calculation for the screened potentials of a balanced bilayer yields

$$V^{sc} = \frac{1}{2} \frac{V_S + V_D}{1 + \Pi_+ (V_S + V_D)} \pm \frac{1}{2} \frac{V_S - V_D}{1 + \Pi_- (V_S - V_D)}$$
(1)

where the plus sign corresponds to the screened intralayer potential and the minus sign to the screened interlayer potential. Here $V_S = 2\pi e^2/\varepsilon q$ is the 2D Coulomb interaction between electrons in the same layer, $V_D = \exp(-qd)V_S$ is the interlayer interaction with d being the distance between layers, and

$$\Pi_{\pm} = \Pi_S \pm \Pi_D. \tag{2}$$

In Eq.(2), Π_S and Π_D are respectively the static in-

tralayer and interlayer polarization operators that relate a shift in the chemical potential (relative to the electrostatic potential) in one layer to the corresponding density in the same-layer (S) or in the opposite layer (D=different). We note that Π_D differs from zero only in the condensed state. Only then does the density in one layer depend on the chemical potential of the other layer.

In the normal state Π_D vanishes and $\Pi_S = g\nu_0$ where

 ν_0 is the density-of-states at the Fermi energy per valley and spin in either layer and g = 4. It then follows that the screened interlayer interaction is

$$V_D^{sc} = \frac{1}{\nu_0 g} \frac{q_{\rm TF} \exp(-qd)}{q + 2q_{\rm TF} + q_{\rm TF}^2 \left[1 - \exp(-2qd)\right]/q} \le \frac{1}{2\nu_0 g}$$
(3)

where $q_{\rm TF} = 2\pi e^2 g\nu_0/\varepsilon$ is the Thomas-Fermi wave vector. We have calculated the mean field critical temperature with this interaction and found that $T_c \sim 6 \cdot 10^{-5} \epsilon_F$. Although not in complete agreement with the estimate of Kharitonov and Efetov, we fully agree that this analysis predicts a small value of T_c rendering the pairing phenomena improbable in realistic disordered bilayer systems[3].

Quite a different conclusion is reached if Thomas-Fermi screening theory is applied consistently in the condensed state. While Π_{-} retains it normal state value $g\nu_0$,

$$\Pi_{+} = g \sum_{\mathbf{k}} \partial_{E_{k}} n_{\mathrm{F}} \left(E_{k} \right) \tag{4}$$

is considerably reduced. Here E_k is the energy of a quasiparticle in the condensed state. In fact, due to the excitation gap Π_+ vanishes at zero temperature. Surprisingly at low temperatures $V_D^{sc} \approx (V_S + V_D)/2$ and hence, the mean-field theory gaps tend to be even larger than when screening is neglected. Although we do not take this argument for enhanced interactions particularly seriously, we conclude that interlayer phase coherence is completely consistent with naive Thomas Fermi screening considerations.

At high temperatures $\Pi_+ \sim g\nu_0$ as well, screening substantially diminishes the order parameter and the system undergoes a first order phase transition into the normal phase. It follows from Eq.(4) that the transition to the normal state occurs when $\Delta \sim T$. As a consequence, the mean field T_c overestimates the real critical temperature. Nevertheless, mean field theory is adequate for determining the Kosterlitz-Thouless temperature, $T_{\rm KT}$, as long as $\Delta(T_{\rm KT}) \gtrsim T_{\rm KT}$. We expect this condition to be satisfied in the strongly interacting regime where $\Delta(T_{\rm KT}) \sim \Delta(0)$.

In the absence of experimental studies of graphene double-layer systems, we can attempt to take guidance from two other partially related systems in which electron-pair condensation has been achieved. A combination of experiment, quantum Monte-Carlo, and various approximate calculations suggests that cold-atom fermions with strongly attractive interactions are reasonably well described at low-temperatures by including collective fluctuations around the BCS mean-field theory state. Moreover, at unitarity numerical calculations find that $T_c \approx 0.2E_{\rm F}$ [9].

A closer comparison is possible between graphenebilayers and semi-conductor based bilayers in the quantum Hall regime. In this case the Kharitonov and Efetov estimate would yield $T_c \rightarrow 0$ because the Thomas-Fermi screening wavevector diverges in concert with the zerowidth of 2D Landau bands. Although experiments are not yet able to extract unambiguous signals of a finite-T phase transition, it is nevertheless clear that spontaneous coherence anomalies begin to emerge below the Kosterlitz-Thouless temperature estimated by a T = 0mean-field-theory phase stiffness calculation. Moreover it is known[10] that spontaneous interlayer phase coherence occurs at small layer separations irrespective of the number of fermion flavors.

Admittedly, an accurate estimate for the critical temperature in the strongly interacting regime $k_{\rm F}d \lesssim 1$ does not exist at present[11]. Nevertheless we feel that the attractive features of graphene bilayers: high carrier densities, high electron and hole Fermi energies, and nearly-perfect particle-hole symmetry suggest that interesting and novel phenomena are likely in this system. We hope that the intriguing thermodynamic and transport aspects of bilayer graphene as well as its potential use in applications encourage experimentalists to construct this system and explore its electrical properties in different geometries[12].

R. Bistritzer, H. Min, J. J. Su, and A.H. MacDonald Department of Physics,The University of Texas at Austin,Austin Texas 78712

- H. Min, R. Bistritzer, J.-J. Su, and A.H. MacDonald, Phys. Rev. B 78, 121401(R) (2008).
- [2] Closely related predictions were made recently by C.-H. Zhang and Yogesh N. Joglekar, Phys. Rev. B 77, 233405 (2008).
- [3] R. Bistritizer and A.H. MacDonald, arXiv:0808.1310.
- [4] M. Y. Kharitonov, and K. Efetov arXiv:0808.2164.
- [5] H. Heiselberg, C. J. Pethick, H. Smith, and L. Viverit, Phys. Rev. Lett. 85, 2418 (2000).
- [6] A. R. Champagne, J. P. Eisenstein, L. N. Pfeiffer, and K. W. West, Phys. Rev. Lett. **100**, 096801 (2008).
- [7] S. De Palo, F. Rapisarda, and G. Senatore, Phys. Rev. Lett. 88, 206401 (2002).
- [8] The model studied in this paper has attractive interactions between all particles, whereas a bilayer has attractive interlayer electron-hole interactions and repulsive intralayer electron-electron interactions.
- [9] V. K. Akkineni, D. M. Ceperley and N. Trivedi, Phys. Rev. B 76, 165116 (2007)
- [10] K. Yang, S. Das Sarma and A. H. MacDonald, Phys. Rev. B 74, 075423 (2006)
- [11] S. Hands and C. Strouthos, arXiv:0800.2720. Perhaps the Monte-Carlo techniques which have been used to explore quantum phase transitions in single-layer graphene can be turned in this direction.
- [12] J.-J. Su and A.H. MacDonald, arXiv:0801.3694 (to appear in Nat. Phys.).