FOUR-DIMENSIONAL RICCI-FLAT SPACE DEFINED BY THE KP EQUATION

Valery Dryuma

Institute of Mathematics and Informatics AS Moldova, Kishinev

E-mail: valery@dryuma.com; cainar@mail.md

Abstract

Four-dimensional affinely connected Ricci-flat space dependent from solutions of the Kadomtsev-Petviashvili equation are constructed. Conditions of metrizabilty of corresponding connection are investigated. Their properties are discussed.

1 The Ricci-flat 8-dim metrics

An examples of the eight-dimensional Ricci-flat metrics of Riemann extensions of affinely-connected four-dimensional spaces A^4 dependent from solutions of the KP-equation were obtained recently by author ([8]).

Here we give the example of four-dimensional Ricci-flat affinely connected space E^4 defined by solutions of the KP-equation which has Ricci-flat the Riemann extension also dependent from solutions of the KP-equation.

Proposition 1 Eight-dimensional Riemann space in local coordinates (x, y, z, t, P, Q, U, V) equipped with the metric

$$ds^{2} = \left(-2\,P\mathit{H11}(y,z,t) - 2\,P\frac{\partial}{\partial t}F(y,z,t) - 2\,\mathit{H12}(y,z,t)Q - 2\,\Gamma_{11}^{\beta}(y,z,t)\,U + 2\,\mathit{H22}(y,z,t)\,V\right)dx^{2} + 2\,\mathit{H22}(y,z,t)\,U + 2\,\mathit{H22}(y,z,t)\,V$$

$$+2\,\left(2\,\mathit{H11}(y,z,t)Q-2\,\mathit{H21}(y,z,t)V\right)\,\mathit{dx}\,\mathit{dy}+2\,\left(-2\,\left(\frac{\partial}{\partial z}F(y,z,t)\right)V+2\,\left(\frac{\partial}{\partial t}F(y,z,t)\right)U\right)\mathit{dx}\,\mathit{dz}+2\,\left(2\,\mathit{H11}(y,z,t)Q-2\,\mathit{H21}(y,z,t)V\right)\,\mathit{dx}\,\mathit{dy}+2\,\left(-2\,\left(\frac{\partial}{\partial z}F(y,z,t)\right)V+2\,\left(\frac{\partial}{\partial t}F(y,z,t)\right)U\right)\,\mathit{dx}\,\mathit{dz}+2\,\left(-2\,\left(\frac{\partial}{\partial z}F(y,z,t)\right)V+2\,\left(\frac{\partial}{\partial t}F(y,z,t)\right)U\right)\,\mathit{dx}\,\mathit{dz}+2\,\left(-2\,\left(\frac{\partial}{\partial z}F(y,z,t)\right)U\right)\,\mathit{dx}\,\mathit{dz}+2\,\left(-2\,\left(\frac{\partial}{\partial z}F(y,z,t)\right)U\right)\,\mathit{dz}+2\,\left(-2\,\left(\frac{\partial}{\partial z}F(y,z,t)\right)U\right)\,\mathit{dz}+2$$

$$+2 dx dP + (2 H11(y, z, t)U - 2 H31(y, z, t)V) dy^{2} + 2 dy dQ + 2 dz dU + 2 dt dV$$
 (1)

is a Ricci-flat

$$R_{ij} = 0$$

if the conditions on the coefficients Hij(y, z, t)

$$\begin{split} &\frac{\partial}{\partial y}H12\left(y,z,t\right)-\frac{\partial}{\partial t}H22\left(y,z,t\right)=0\\ &-\frac{\partial}{\partial y}H11\left(y,z,t\right)+\frac{\partial}{\partial t}H21\left(y,z,t\right)=0\\ &-\frac{\partial}{\partial z}H11\left(y,z,t\right)+\frac{\partial}{\partial t}H31\left(y,z,t\right)=0 \end{split} \tag{2}$$

are valid.

An arbitrary functions $\Gamma_{11}^3(y,z,t)$ and F(y,z,t) in so doing satisfy the relation

$$\frac{\partial}{\partial z} \Gamma_{11}^{3}(y,z,t) = 2 \left(\frac{\partial}{\partial t} F(y,z,t) \right)^{2} + 2 H11(y,z,t) \frac{\partial}{\partial t} F(y,z,t) + 2 \left(H11(y,z,t) \right)^{2}$$

or

$$\Gamma_{11}^{3}(y,z,t) = \int 2 \left(\frac{\partial}{\partial t} F(y,z,t) \right)^{2} + 2 H11(y,z,t) \frac{\partial}{\partial t} F(y,z,t) + 2 (H11(y,z,t))^{2} dz + \mathcal{F}1(y,t). \tag{3}$$

The metric (1) is an example of the Riemann extension of affinely-connected four-dimensional space with symmetrical connection $\Gamma^i_{jk}(x^l) = \Gamma^i_{kj}(x^l)$ depending from the local coordinates x^i . In general case it is defined by the expression

$$ds^2 = -2\Gamma^i_{jk}\xi_i dx^j dx^k + 2d\xi_k dx^k, \tag{4}$$

where ξ_k -are an additional coordinates ([1]).

After the substitutions of the form

$$H11 (y, z, t) = -1/2 u (y, z, t), \quad H12 (y, z, t) = -1/3 v (y, z, t),$$

$$H21 (y, z, t) = -2/3 v (y, z, t) - 1/2 \frac{\partial}{\partial t} u (y, z, t),$$

$$H31 (y, z, t) = -3/4 w (y, z, t) + 3/8 (u (y, z, t))^{2} - \frac{\partial}{\partial t} v (y, z, t) - 1/2 \frac{\partial^{2}}{\partial t^{2}} u (y, z, t),$$

$$H22 (y, z, t) = -1/2 w (y, z, t) + 1/2 (u (y, z, t))^{2} - \frac{\partial}{\partial t} v (y, z, t) + 1/2 \frac{\partial}{\partial t} u (y, z, t)$$

$$(5)$$

from the conditions (2) the famous KP-equation is followed

$$\frac{\partial^{2}}{\partial t \partial z} u\left(y,z,t\right) - 3/2 \, \left(\frac{\partial}{\partial t} u\left(y,z,t\right)\right)^{2} - 3/2 \, u\left(y,z,t\right) \, \frac{\partial^{2}}{\partial t^{2}} u\left(y,z,t\right) - 1/4 \, \frac{\partial^{4}}{\partial t^{4}} u\left(y,z,t\right) - 3/4 \, \frac{\partial^{2}}{\partial y^{2}} u\left(y,z,t\right) = 0$$

$$\frac{\partial}{\partial t}\left(\frac{\partial u\left(y,z,t\right)}{\partial z}-3/2\,u\left(y,z,t\right)\frac{\partial}{\partial t}u\left(y,z,t\right)-1/4\frac{\partial^{3}}{\partial t^{3}}u\left(y,z,t\right)\right)=3/4\frac{\partial^{2}}{\partial y^{2}}u\left(y,z,t\right)$$

Remark 1 A notion of the Riemann extensions of affinely-connected spaces ([1]) was used by author for the studying of geometrical problems in theory nonlinear dynamical systems and in General Relativity ([2]-[8]).

Remark 2 About presentation of the KP-equation in form of the system (2) see ([9]).

2 Four-dimensional subspace

Eight-dimensional metrics (4) is closely related with a properties of four-dimensional spaces in local coordinates x^k .

Let us consider an example.

The full system of geodesic of the metric (4) decomposes into two parts.

The first part has the form of the linear system of equations for the coordinates $(\xi_k = P, Q, U, V)$

$$\frac{d^2\xi_k}{ds^2} + A(x^i)\frac{d\xi_k}{ds} + B(x^i)\xi_k = 0$$

and the second part for local coordinates $x^i = (x, y, z, t)$ is defined by the system of equations

$$\frac{d^2x^k}{ds^2} + \Gamma^k_{ij}\frac{dx^i}{ds}\frac{dx^j}{ds} = 0.$$

In our case it takes the form

$$\begin{split} \frac{d^2}{ds^2}x(s) + \left(\frac{d}{ds}x(s)\right)^2 H11(y,z,t) + \left(\frac{d}{ds}x(s)\right)^2 \frac{\partial}{\partial t}F(y,z,t) &= 0, \\ \frac{d^2}{ds^2}y(s) + H12(y,z,t) \left(\frac{d}{ds}x(s)\right)^2 - 2 H11(y,z,t) \left(\frac{d}{ds}x(s)\right) \frac{d}{ds}y(s) &= 0, \\ \frac{d^2}{ds^2}z(s) + \Gamma_{11}^3(y,z,t) \left(\frac{d}{ds}x(s)\right)^2 - 2 \left(\frac{\partial}{\partial t}F(y,z,t)\right) \left(\frac{d}{ds}x(s)\right) \frac{d}{ds}z(s) - H11(y,z,t) \left(\frac{d}{ds}y(s)\right)^2 &= 0, \\ \frac{d^2}{ds^2}t(s) - H22(y,z,t) \left(\frac{d}{ds}x(s)\right)^2 + 2 H21(y,z,t) \left(\frac{d}{ds}x(s)\right) \frac{d}{ds}y(s) + 2 \left(\frac{\partial}{\partial z}F(y,z,t)\right) \left(\frac{d}{ds}x(s)\right) \frac{d}{ds}z(s) + \\ + H31(y,z,t) \left(\frac{d}{ds}y(s)\right)^2 &= 0. \end{split}$$

From these relations we find the coefficients of affine connection Γ^i_{jk} of the 4-dimensional subspace in local coordinates $(x^i = x, y, z, t)$

$$\Gamma_{11}^{1} = H11 + \frac{\partial F}{\partial t}, \quad \Gamma_{11}^{2} = H12, \quad \Gamma_{12}^{2} = -H11,$$

$$\Gamma_{11}^{3} = \Gamma_{11}^{3}, \quad \Gamma_{13}^{3} = -\frac{\partial F}{\partial t}, \quad \Gamma_{22}^{3} = -H11,$$

$$\Gamma_{11}^{4} = -H22, \quad \Gamma_{12}^{4} = H21, \quad \Gamma_{13}^{4} = \frac{\partial F}{\partial z}, \quad \Gamma_{22}^{4} = H31.$$
(6)

Corresponding the Ricci tensor

$$R_{ij} = \partial_k \Gamma^k_{ij} - \partial_i \Gamma^k_{kj} + \Gamma^k_{kl} \Gamma^l_{ij} - \Gamma^k_{im} \Gamma^m_{kj}$$

is symmetrical

$$R_{ij} = R_{ji}$$

and it is equal to zero $R_{ik} = 0$ if the conditions (2), (3) are hold.

As the example of geodesic equation connected with an additional coordinates we refer to the expression on the coordinate V

$$\left(\left(\frac{d}{ds} x(s) \right)^2 \frac{\partial}{\partial t} H11(y,z,t) + \left(\frac{d}{ds} x(s) \right)^2 \frac{\partial^2}{\partial t^2} F(y,z,t) \right) P +$$

$$+ \left(-2 \left(\frac{d}{ds} x(s) \right) \left(\frac{d}{ds} y(s) \right) \frac{\partial}{\partial t} H11(y,z,t) + \left(\frac{d}{ds} x(s) \right)^2 \frac{\partial}{\partial t} H12(y,z,t) \right) Q +$$

$$+ \frac{d^2}{ds^2} V(s) - \left(\frac{d}{ds} x(s) \right)^2 \left(\frac{\partial}{\partial t} H22(y,z,t) \right) V - \left(\frac{d}{ds} y(s) \right)^2 \left(\frac{\partial}{\partial t} H11(y,z,t) \right) U +$$

$$+ 2 \left(\frac{d}{ds} x(s) \right) \left(\frac{d}{ds} y(s) \right) \left(\frac{\partial}{\partial t} H21(y,z,t) \right) V + \left(\frac{d}{ds} x(s) \right)^2 \left(\frac{\partial}{\partial t} \Gamma_{11}^3(y,z,t) \right) U -$$

$$- 2 \left(\frac{d}{ds} x(s) \right) \left(\frac{d}{ds} z(s) \right) \left(\frac{\partial^2}{\partial t^2} F(y,z,t) \right) U + \left(\frac{d}{ds} y(s) \right)^2 \left(\frac{\partial}{\partial t} H31(y,z,t) \right) V +$$

$$+2\left(\frac{d}{ds}x(s)\right)\left(\frac{d}{ds}z(s)\right)\left(\frac{\partial^2}{\partial t\partial z}F(y,z,t)\right)V=0.$$

Corresponding equations for the coordinates U, P, Q have more cumbersome form and may be omitted here.

Remark 3 As distinct from the example of eight-dimensional the Ricci-flat Riemann space related with the KP-equation considered in the article ([8]) the metric (1) has a Ricci-flat affinely connected four-dimensional subspace.

In this connection it should be noted that problem of metrizability of the connection Γ_{jk}^i defined by the expressions (6) has of great importance.

It is known that in order to obtain the metric tensor g_{ij} correspondent the connection Γ^i_{jk} we have to solve the system of equations

$$\nabla_i g_{jk} = \partial_i g_{jk} - \Gamma^l_{ij} g_{lk} - \Gamma^l_{ik} g_{jl} = 0.$$

A necessary and sufficient condition of integrability of this system implies in the studying of the properties of the set of equations

$$R_{ijl}^k g_{km} + R_{ijm}^k g_{kl} = 0,$$

$$\nabla_s R_{ijl}^k g_{km} + \nabla_s R_{ijm}^k g_{kl} = 0,$$

where R_{ijl}^k are the components of Riemann tensor of the connection (6).

In the case being considered the number of components of the Riemann tensor is thirty-six and affine connection (6) obeys the condition $\Gamma_{ki}^i = 0$.

On this basis and taking in consideration the relation $\Gamma_{ki}^i = \frac{1}{2g} \frac{\partial g}{\partial x^k}$ there is followed that in the case of metrizability of the connection (6) determinant of corresponding metric tensor must to be constant.

3 Acknowledgements

The research of author was partially supported by Grant CSTD AS Moldova and Grant RFBR.

References

- [1] Paterson E.M. and Walker A.G., Riemann extensions, Quart. J. Math. Oxford, 1952, V.3, 19–28.
- [2] Dryuma V., The Riemann and Einstein-Weyl geometries in the theory of ODE's, their applications and all that, New Trends in Integrability and Partial Solvability, 115-156, (eds.A.B.Shabat et al.) 2004, Kluwer Academic Publishers, arXiv: nlin: SI/0303023, 11 March, 2003, 1–37.
- [3] Dryuma V., On the Riemann Extension of the Gödel space -time metric, *Buletinul A.S.R.M*, *Matematica*, 2005, No.3(49), p.43–62.
- [4] Dryuma V., On solutions of a Heavenly equations and their generalizations, arXiv:math.DG/0510526 v1, 31 Oct 2006, p.1–13.
- [5] Dryuma V., The Riemann Extension of the Schwarzshild space-time, *Buletinul A.S.R.M*, *Matematica*, 2003, No. 3(43), p.92–103.
- [6] Dryuma V., The Riemann Extension of space-time with vanishing curvature invariants, http://www.math.kth.se/4ecm/abstract/14.27.pdf, Sweden, Stockholm, June 27-July 2, ECM'2004.
- [7] Dryuma V., Riemann Extension in theory of the first order systems of differential equations, $arXiv:math.DG/0510526\ v1,\ 25\ Oct\ 2005,\ p.1-21.$

REFERENCES 5

[8] Dryuma V., Eight-dimensional the Ricci-flat Riemann space related with the KP-equation, $arXiv: 0810.0346\ v1,\ 2\ Oct\ 2008,\ p.1-5.$

[9] B.G.Konopelchenko, Quantum deformations of associative algebras and integrable systems, arXiv: 0802.3022 v2 [nlin. SI] 28 Apr, 2008, p.1–24.