

A MAP FROM LAWSON HOMOLOGY TO DELIGNE COHOMOLOGY

WENCHUAN HU

ABSTRACT. A natural map from Lawson homology to Deligne cohomology groups for smooth complex projective varieties is constructed by using the Harvey-Lawson “spark” complexes.

CONTENTS

1. Introduction	1
2. Sparks and differential characters	2
3. The construction of the map	4
References	6

1. INTRODUCTION

Let X be a complex projective variety of dimension n . The **Lawson homology** $L_p H_k(X)$ of p -cycles is defined by

$$L_p H_k(X) := \pi_{k-2p}(\mathcal{Z}_p(X)) \text{ for } k \geq 2p \geq 0,$$

where $\mathcal{Z}_p(X)$ is provided with a natural topology so that it is an abelian topological group (cf. [F], [L1] and [L2]). For general background, the reader is referred to [L2].

Friedlander and Mazur [FM] showed that there are natural maps, called **cycle class maps** $\Phi_{p,k} : L_p H_k(X) \rightarrow H_k(X)$ from Lawson homology to singular homology. Denote by $L_p H_k(X)_{hom}$ the kernel of $\Phi_{p,k}$, i.e., $L_p H_k(X)_{hom} = \ker\{\Phi_{p,k} : L_p H_k(X) \rightarrow H_k(X)\}$.

Temporarily suppose X is a complex manifold. Let Ω_X^k the sheaf of holomorphic k -form on X . The **Deligne complex of level p** is the complex of sheaves

$$\underline{\mathbb{Z}}_{\mathcal{D}}(p) : 0 \rightarrow \mathbb{Z} \xrightarrow{(2i\pi)^p} \Omega_X^0 \rightarrow \Omega_X^1 \rightarrow \Omega_X^2 \rightarrow \cdots \rightarrow \Omega_X^{p-1} \rightarrow 0$$

Date: September 13, 2008.

The **Deligne cohomology** of X in level p we mean the hypercohomology of this complex:

$$H_{\mathcal{D}}^*(X, \mathbb{Z}(p)) := \mathbf{H}^*(X, \underline{\mathbb{Z}}_{\mathcal{D}}(p)).$$

For more details on Deligne cohomology, the reader is referred to [EV].

We define a natural map from Lawson homology to the corresponding Deligne cohomology using the theory of differential characters introduced by Cheeger in [C] and the theory of D-bar sparks developed by Harvey and Lawson in [HL1] and [HL2]. This theory of differential characters has been developed in [CS], [GS], [Ha] and systematically generalized by Harvey, Lawson and Zweck in [HLZ]. This map is a generalization of the Abel-Jacobi map for Lawson homology defined by the author in [Hu].

The main result in this paper is the following

Theorem 1.1. *Let X be a smooth complex projective variety of dimension n . We have a well-defined homomorphism*

$$\hat{a} : L_p H_{k+2p}(X) \rightarrow H_{\mathcal{D}}^{2(n-p)-k}(X, \mathbb{Z}(n-p-k-1)),$$

given by

$$\hat{a}([f]) = \hat{a}_f$$

which coincides with the generalized Abel-Jacobi map defined in [Hu] when \hat{a} is restricted on $L_p H_k(X)_{hom}$ and the projection of the image of \hat{a} under δ_2 is the natural map $\Phi_{p,k}$.

The notations \hat{a}_f and δ_2 will be defined below.

2. SPARKS AND DIFFERENTIAL CHARACTERS

It might necessary to sketch the background materials of sparks and differential characters for our construction in section 3. For details, see [HLZ], [HL1] and [HL2]. In this section X denotes smooth manifold unless otherwise noted.

Definition 2.1. *A **homological spark complex** is a triple of cochain complexes (F^*, E^*, I^*) together with morphisms*

$$\Psi : I^* \rightarrow F^* \supset E^*$$

such that:

- (1) $\Psi(I^k) \cap E^k = \{0\}$ for $k > 0$,
- (2) $H^*(E) \cong H^*(F)$, and
- (3) $\Psi(I^0) \rightarrow F^0$ is injective.

Definition 2.2. In a given spark complex (F^*, E^*, I^*) a **spark of degree k** is a pair

$$(a, r) \in F^k \oplus I^{k+1}$$

which satisfies the spark equation

- (1) $da = e - \Psi(r)$ for some $e \in E^{k+1}$, and
- (2) $dr = 0$.

The group of sparks of degree k is denoted by $\mathcal{S}^k = \mathcal{S}^k(F^*, E^*, I^*)$.

Definition 2.3. Two sparks $(a, r), (a', r') \in \mathcal{S}^k(F^*, E^*, I^*)$ are equivalent if there exists a pair

$$(b, s) \in F^{k-1} \oplus I^k$$

such that

- (1) $a - a' = db + \Psi(s)$, and
- (2) $r - r' = -ds$.

The set of equivalence classes is called the **group of spark classes of degree k** associated to the given spark complex and will be denoted by $\widehat{\mathbf{H}}^k(F^*, E^*, I^*)$ or simply $\widehat{\mathbf{H}}^k(F)$.

As usual, let $Z^k(E) = \{e \in E^k \mid de = 0\}$ and set

$$Z_I^k(E) := \{e \in Z^k(E) \mid [e] = \Psi_*(\rho) \text{ for some } \rho \in H^k(I)\}$$

where $[e]$ denotes the class of e in $H^k(E)$ (note that $de = 0$). The following lemma was proved in [HLZ]:

Lemma 2.1. *There exist well-defined surjective homomorphisms:*

$$\delta_1 : \widehat{\mathbf{H}}^k(F) \rightarrow Z_I^k(E) \quad \text{and} \quad \delta_2 : \widehat{\mathbf{H}}^k(F) \rightarrow H^{k+1}(I)$$

given on any representing spark $(a, r) \in \mathcal{S}^k$ by

$$\delta_1(a, r) = e \quad \text{and} \quad \delta_2(a, r) = [r]$$

where $da = e - \Psi(r)$ as in Definition 2.2.

The following concrete example is the main object which will be dealt with in the next section.

Example 2.2. Now let X be a smooth complex projective variety of dimension n . Set

$$F^m = \mathcal{D}^m(X, q) := \bigoplus_{r+s=m, r < q} \mathcal{D}^{r,s}(X) \quad \text{and} \quad \bar{d} = \Psi \circ d$$

where

$$\Psi : \mathcal{D}^m(X) \rightarrow \mathcal{D}^m(X, q)$$

is the projection $\Psi(a) = a^{0,m} + a^{1,m-1} + \dots + a^{q-1,m-q+1}$.

$$E^m = \mathcal{E}^m(X, q) := \bigoplus_{r+s=m, r < q} \mathcal{E}^{r,s}(X) \quad \text{and} \quad \bar{d} = \Psi \circ d$$

and

$$I^m = \mathcal{I}^m(X)$$

It has been shown in [HL2] that the above triple (F^*, E^*, I^*) is a homological spark complex. The group of associated spark classes in degree m will be denoted by $\widehat{\mathbf{H}}^m(X, q)$. From this homological spark complex, one has

$$\ker(\delta_1) = H_{\mathcal{D}}^{m+1}(X, \mathbb{Z}(q)).$$

3. THE CONSTRUCTION OF THE MAP

In this section, X denotes a smooth projective variety. Let $\mathcal{Z}_p(X)$ be the space of algebraic p -cycles with a natural topology and a base point, i.e., the ‘null’ p -cycle (which we denote by 0) and let $\Omega^k \mathcal{Z}_p(X)$ be the loop space with the given base point. Explicitly, suppose $S^k = \mathbb{R}^k \cup \infty$. We have

$$\Omega^k \mathcal{Z}_p(X) = \{f : S^k \rightarrow \mathcal{Z}_p(X) \mid f \text{ is continuous with } f(\infty) = 0\}.$$

Given such a continuous map $f : S^k \rightarrow \mathcal{C}_p(X)$, we can find a map $g : S^k \rightarrow \mathcal{Z}_p(X)$ such that g is homotopic to f and g is piecewise linear with regard to a triangulation of $\mathcal{C}_p(X)$ and hence one can define a current c_g over X and this current c_g is a cycle. Moreover c_g depends only on the homotopy class of g . For the detail of the construction of c_g from the map $g : S^k \rightarrow \mathcal{Z}_p(X)$, see [Hu].

Consider Example 2.2 with $m = 2(n-p) - k - 1$, $q = n - p - k - 1$. All the following argument will focus on this homological spark complex.

Definition 3.1. Set $a_f := -\Psi(d^*G(c_f))$. Then (a_f, c_f) is called the Hodge spark of the map $f : S^k \rightarrow \mathcal{Z}_p(X)$. Let $\widehat{a}_f \in \widehat{\mathbf{H}}^{2(n-p)-k-1}(X, n-p-k-1)$ be the differential character corresponding to the Hodge spark (a_f, c_f) .

Lemma 3.1. If $f \in \Omega^k \mathcal{Z}_p(X)$, then $\delta_1(\widehat{a}_f) = 0 \in \widehat{\mathbf{H}}^{2(n-p)-k-1}(X, n-p-k-1)$.

Proof. Let $H(c_f)$ be the harmonic part of c_f . Note that $H(c_f)$ and c_f are of the same $(*, *)$ -type. Moreover,

$$c_f \in \bigoplus_{r+s=k, |r-s| \leq k} \mathcal{D}'_{p+r, p+s}(X).$$

By the type reason, the projection of $H(c_f)$ on

$$\bigoplus_{r+s=k, r \geq k+1} \mathcal{D}'_{p+r, p+s}(X)$$

is zero. This is exactly the image of \widehat{a}_f under δ_1 . \square

Hence we get an element $\hat{a}_f \in \ker(\delta_1) = H_{\mathcal{D}}^{2(n-p)-k}(X, \mathbb{Z}(n-p-k-1))$. Therefore we get a map

$$\hat{a} : \Omega^k \mathcal{Z}_p(X) \rightarrow H_{\mathcal{D}}^{2(n-p)-k}(X, \mathbb{Z}(n-p-k-1))$$

which is defined by $\hat{a}(f) = \hat{a}_f$.

If the map f is contractible, then we have the following:

Lemma 3.2. *If $f : S^k \rightarrow \mathcal{Z}_p(X)$ is contractible, then $\hat{a}(f) = 0$.*

Proof. Let $F : D^{k+1} \rightarrow \mathcal{Z}_p(X)$ be the extension of f , i.e., $F|_{\partial(D^{k+1})} = f$. Let c_F be the current over X defined by F . As showed in Lemma 3.1, c_F is a cycle. Moreover, it is a boundary, i.e., $c_F = \partial(c_f)$. By Corollary 12.11 in [HLZ], we have $\hat{a}_f = \widehat{H(\Psi(c_F))}$. Since

$$c_F \in \bigoplus_{r+s=k+1, |r-s| \leq k+1} \mathcal{D}'_{p+r, p+s}(X),$$

the projection of $H(c_F)$ under Ψ on

$$\bigoplus_{r+s=k, r > k+1} \mathcal{D}'_{p+r, p+s}(X)$$

is zero. Note that Ψ commutes with the Laplace operator, we have $\hat{a}_f = 0$. \square

By the Lemma 3.1 and Lemma 3.2, we have a well-defined map

$$\hat{a} : L_p H_k(X) \rightarrow H_{\mathcal{D}}^{2(n-p)-k}(X, \mathbb{Z}(n-p-k-1)),$$

given by

$$\hat{a}([f]) = \hat{a}_f.$$

Recall that the Deligne cohomology can be written as the middle part of an short exact sequence

$$0 \rightarrow \frac{H^{2(n-p)-k-1}(X, n-p-k-1)}{H^{2(n-p)-k-1}(X, \mathbb{Z})} \rightarrow H_{\mathcal{D}}^{2(n-p)-k}(X, \mathbb{Z}(n-p-k-1)) \rightarrow \ker(\Psi_*) \rightarrow 0,$$

where

$$H^{2(n-p)-k-1}(X, n-p-k-1) = \left\{ \bigoplus_{r+s=2(n-p)-k-1, r < n-p-k-1} H^{r,s}(X) \right\}$$

and

$$\ker(\Psi_*) = H^{2(n-p)-k}(X, \mathbb{Z}) \cap \left\{ \bigoplus_{r+s=2(n-p)-k, |r-s| \leq k+2} H^{r,s}(X) \right\}$$

Proposition 3.3. *The restriction of the above map to $L_p H_k(X)_{\text{hom}} := \ker\{\Phi_{p,k} : L_p H_k(X) \rightarrow H_k(X)\}$ is the generalized Abel-Jacobi map defined in [Hu] if we identify the $H^{r,s}(X)$ with $\{H^{n-r, n-s}(X)\}^*$ for all $0 \leq r, s \leq n$ and $H^q(X, \mathbb{Z})$ with $H_{2n-q}(X, \mathbb{Z})$.*

Proof. Recall the definition that

$$\Phi : L_p H_k(X)_{hom} \rightarrow \left\{ \bigoplus_{r>k+1, r+s=k+1} H^{p+r, p+s}(X) \right\}^* / H_{2p+k+1}(X, \mathbb{Z})$$

is given by $\Phi([f]) = \Phi_f$, where $\Phi_f(\omega) = \int_{\tilde{c}} \omega \pmod{H_{2p+k}(X, \mathbb{Z})}$ with $\partial(\tilde{c}) = c_f$. The Lemma 12.10 in [HLZ] implies that the two constructions coincide. \square

Remark 3.4. *It is easy to see from the definition of Ψ_* that the image of $\Phi_{p,k}$ is in $\ker(\Psi_*)$. Hence the natural map $\Phi_{p,k} : L_p H_k(X) \rightarrow H_k(X)$ factors through \hat{a} and the map δ_2 in the first 3×3 grid given in [HL2].*

In summary, we have the following

Theorem 3.5. *Let X be a smooth complex projective variety of dimension n . We have a well-defined map*

$$\hat{a} : L_p H_{k+2p}(X) \rightarrow H_{\mathcal{D}}^{2(n-p)-k}(X, \mathbb{Z}(n-p-k-1)),$$

given by

$$\hat{a}([f]) = \hat{a}_f$$

which coincides with the generalized Abel-Jacobi map defined in [Hu] when \hat{a} is restricted on $L_p H_k(X)_{hom}$; and the projection of the image of \hat{a} under δ_2 is the natural map $\Phi_{p,k}$.

Proof. The first statement follows from Lemma 3.1 and 3.2 that \hat{a} is well-defined. The second one follows from Proposition 3.3 and the the third one follows from 3.4. \square

Remark 3.6. *In general, the map \hat{a} is a nontrivial even if restricted on $L_p H_k(X)_{hom}$. This has been shown by Griffiths [G] for $k = 2p$ and by the author [Hu] for $k > 2p$.*

Acknowledge

I would like to express my gratitude to my former advisor, Blaine Lawson, for all his help.

REFERENCES

- [C] J. Cheeger, *Multiplication of differential characters*. Symposia Mathematica, Vol. XI (Convegno di Geometria, INDAM, Rome, 1972), pp. 441–445. Academic Press, London, 1973.
- [CS] J. Cheeger; J. Simons *Differential characters and geometric invariants*. Geometry and topology (College Park, Md., 1983/84), 50–80, Lecture Notes in Math., 1167, Springer, Berlin, 1985.

- [EV] H. Esnault; E. Viehweg, *Deligne-Beilinson cohomology*. Beilinson's conjectures on special values of L -functions, 43–91, *Perspect. Math.*, 4, Academic Press, Boston, MA, 1988.
- [F] E. Friedlander, *Algebraic cycles, Chow varieties, and Lawson homology*. *Compositio Math.* 77 (1991), no. 1, 55–93.
- [FM] E. Friedlander and B. Mazur, *Filtrations on the homology of algebraic varieties. With an appendix by Daniel Quillen*. *Mem. Amer. Math. Soc.* 110 (1994), no. 529, x+110 pp.
- [GS] H. Gillet and C. Soulé *Arithmetic Chow groups and differential characters*. *Algebraic K-theory: connections with geometry and topology* (Lake Louise, AB, 1987), 29–68, *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, 279, Kluwer Acad. Publ., Dordrecht, 1989.
- [G] P. Griffiths, *On the periods of certain rational integrals I, II*. *Ann. of Math.* (2) 90(1969), 460-495; *ibid.* (2) 90(1969) 496–541.
- [Ha] B. Harris, *Differential characters and the Abel-Jacobi map*. *Algebraic K-theory: connections with geometry and topology* (Lake Louise, AB, 1987), 69–86, *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, 279, Kluwer Acad. Publ., Dordrecht, 1989.
- [HL1] R. Harvey and B. Lawson *From Sparks to Grundles - Differential Characters* [arXiv.org:math.DG/0306193](https://arxiv.org/math.DG/0306193)
- [HL2] R. Harvey and B. Lawson, *D-bar sparks, I*. [arXiv.org:math.DG/0512247](https://arxiv.org/math.DG/0512247)
- [HLZ] R. Harvey; B. Lawson and J. Zweck, *The de Rham-Federer theory of differential characters and character duality*. *Amer. J. Math.* 125 (2003), no. 4, 791–847
- [Hu] W. Hu, *Generalized Abel-Jacobi map on Lawson Homology*. [arXiv.org:math.AG/0608294](https://arxiv.org/math.AG/0608294)
- [L1] B. Lawson, *Algebraic cycles and homotopy theory*. *Ann. of Math.* **129**(1989), 253-291.
- [L2] B. Lawson, *Spaces of algebraic cycles*. pp. 137-213 in *Surveys in Differential Geometry*, 1995 vol.2, International Press, 1995.

DEPARTMENT OF MATHEMATICS, MIT, ROOM 2-363B, 77 MASSACHUSETTS AVENUE, CAMBRIDGE, MA 02139, USA

E-mail address: wenchuan@math.mit.edu