

# The structure of $\mathcal{N} = 2$ multi-instanton and string-loop corrections in toroidal orbifold models

P. G. Cámara

*Centre de Physique Théorique, Ecole Polytechnique, CNRS, F-91128 Palaiseau, France*

We summarize the structure of  $\mathcal{N} = 2$  multi-instanton and string loop corrections to the Kähler potential and the gauge kinetic function in type I toroidal orbifold models. For that aim, we exploit perturbative calculations of the physical gauge couplings on the heterotic dual, whenever such a perturbative description is available.

## 1. INTRODUCTION

The low energy limit of String Theory compactified to four dimensions is often described in terms of an effective supergravity which generically develops classical flat directions. Understanding the leading corrections from string loops and non-perturbative effects is therefore crucial for addressing questions such as moduli stabilization, supersymmetry breaking or cosmological evolution in these models. Although a non-perturbative description of String Theory is still far from being at our disposal, the rich network of dualities allows, already at this stage, to make some explicit computations in simple models. Precisely, type I/heterotic S-duality [1] maps stringy instanton effects due to euclidean 1-branes (E1-branes), into string loop effects in a dual heterotic model. Whereas in ten dimensions S-duality is a strong-weak coupling duality, in four dimensions it is manifested as an electric-magnetic duality, making the mapping between both effective theories straightforward.

In this contribution we mainly summarize some of the results of [2], where the structure of multi-instanton and string loop corrections in a simple class of models were obtained. Concretely, we consider type I string theory compactified on toroidal orbifolds, and use type I/heterotic S-duality to extract the corrections to the Kähler potential and the gauge kinetic function, coming from string loop and multi-instanton effects associated with  $\mathcal{N} = 2$  subsectors of the theory. Additional related material and a more detailed exposition can be found in [2, 3].

A simple way to extract corrections to the Kähler potential,  $K$ , and the gauge kinetic function,  $f_a$ , of the effective theory is to look at the one-loop physical gauge couplings. In a heterotic compactification, these read,

$$4\pi^2 g_a^{-2}(\mu^2)|_{1\text{-loop}} = \frac{k_a}{\ell} + \frac{b_a}{4} \log \frac{M_s^2}{\mu^2} + \frac{\Delta_a(M, \bar{M})}{4}, \quad (1)$$

with  $\ell$  the linear multiplet associated with the dilaton,  $M_s$  the string scale,  $M$  the moduli of the compactification and  $k_a$  the normalization of the gauge group generators, determined by the level of the corresponding Kac-Moody algebra. The  $\beta$ -function coefficient,  $b_a$ , is given in terms of the quadratic Casimir invariants of the gauge group,  $b_a = \sum_r n_r T_a(r) - 3T_a(\text{adj}_a)$ , with  $n_r$  the number of matter multiplets in the representation  $r$ .

On the other hand, in field theory the physical gauge couplings are given as [4, 5],

$$16\pi^2 g_a^{-2}(\mu^2)|_{1\text{-loop}} = 4\text{Re } f_a(M)|_{1\text{-loop}} + b_a \log \frac{M_{\text{Pl}}^2 \mu^{-2}}{S + \bar{S}} + c_a \hat{K}(M, \bar{M}) - 2 \sum_r T_a(r) \log \det Z_r(M, \bar{M}), \quad (2)$$

where  $\det Z_r$  is the determinant of the tree-level Kähler metric associated with the matter multiplets in the representation  $r$ ,  $\hat{K}(M, \bar{M})$  the tree-level Kähler potential for the moduli  $M$  and  $c_a = \sum_r n_r T_a(r) - T(\text{adj}_a)$ .

In order to compare (1) and (2), we recall the relation between the usual complex axion-dilaton  $S$  and the linear multiplet  $\ell$ ,

$$\frac{1}{\ell} = \text{Re } S - \frac{1}{4} \Delta_{\text{univ}}. \quad (3)$$

with  $\Delta_{\text{univ}}$  a gauge group independent (“universal”) function which, for convenience, we split in to its harmonic and non-harmonic parts,

$$\Delta_{\text{univ}}(M, \bar{M}) = V_{(1)}(M, \bar{M}) + H(M) + H^*(\bar{M}). \quad (4)$$

The Kähler potential and the gauge-kinetic function of the effective theory then are given to one-loop by [5, 6, 7]

$$K|_{1\text{-loop}} = -\log\left(S + \bar{S} - \frac{1}{2}V_{(1)}(M, \bar{M})\right) + \hat{K}(M, \bar{M}) , \quad (5)$$

$$\text{Re } f_a|_{1\text{-loop}} = k_a \text{Re } S + \frac{1}{4} \left( \Delta_a(M, \bar{M}) - V_{(1)}(M, \bar{M}) - c_a \hat{K}(M, \bar{M}) - 2 \sum_r T_a(r) \log \det Z_r(M, \bar{M}) \right) . \quad (6)$$

## 2. THE MODELS

We focus on type I toroidal orbifold compactifications which admit a weakly coupled perturbative description in terms of heterotic string theory. More precisely, we consider models where the orbifold action,  $\mathbb{G}$ , contains some subgroup,  $\mathbb{G}_i$ , which leaves unrotated some complex plane, and therefore there is a  $\mathcal{N} = 2$  sector of the theory associated to each of such subgroups. This includes  $\mathbb{Z}_2$ ,  $\mathbb{Z}_2 \times \mathbb{Z}_2$ ,  $\mathbb{Z}_6$ ,  $\mathbb{Z}'_6$ ,  $\mathbb{Z}_8$  or  $\mathbb{Z}_{12}$  orbifold models.

The gauge threshold corrections can be computed in the heterotic side along the lines of [8, 9, 10, 11] and take the expression,

$$\Delta_a = -b_a \log \frac{M_s^2}{\mu^2} - \frac{1}{2} \sum_i \int_{\mathcal{F}} \frac{d^2\tau}{\tau_2} \hat{Z}_i(T_i, U_i) \hat{\mathcal{A}}_{f,i}^a , \quad (7)$$

where the sum runs over the disjoint union of  $\mathcal{N} = 2$  subsectors, each leaving invariant a single complex plane with Kähler and complex structure moduli  $T_i$  and  $U_i$ , and the gauge group is given by a product  $G = \prod_a G_a$ . The explicit expression for the toroidal lattice sums,  $\hat{Z}_i$ , can be found in [2]. The index  $\hat{\mathcal{A}}_{f,i}^a$  is an almost holomorphic modular function which is constrained by modular invariance and  $\mathcal{N} = 2$  supersymmetry to have the structure,

$$\hat{\mathcal{A}}_{f,i}^a(\tau) = 2b_i^a + \frac{\gamma_i}{20\eta^{24}} [D_{10}E_{10} - 528\eta^{24}] , \quad (8)$$

with  $b_i^a$  the  $\beta$ -function coefficient of the  $\mathcal{N} = 2$  gauge theory associated to a would-be  $T^6/\mathbb{G}^i$  orbifold and  $\gamma_i$  a gauge group independent coefficient. Definitions for the Eisenstein series,  $E_k$ , and the modular covariant derivative,  $D_k$ , can be found e.g. in the appendix of [3].

For the sake of clarity, let us briefly consider the simplest case of a single  $\mathcal{N} = 2$  sector. This corresponds to the well known case of type I string theory compactified on  $T^2 \times K3$ . In order to cancel anomalies, 24 instantons are required. In the  $T^4/\mathbb{Z}_2$  orbifold limit of the  $K3$  [12, 13] these are manifest as 8 D5-branes and 16  $U(1)$  bundles hidden at the 16 orbifold singularities. The D5-branes together with the 16 D9-branes also present, lead to a maximal gauge group  $U(16) \times U(16)$ .

In the Coulomb branch of this setup, where a half D5-brane is located at each of the 16 fixed points, only the  $U(16)$  gauge group from the D9-branes remains massless because of the Green-Schwarz mechanism. The spectrum is given by four hypermultiplets, containing the moduli of the  $K3$ , three vector multiplets containing the axion-dilaton and the moduli of the  $T^2$ , a  $\mathbf{120} + \overline{\mathbf{120}}$ , and sixteen  $\mathbf{16}$  from the D5-D9 modes. In this particular point of the moduli space the weakly coupled  $SO(32)$  heterotic dual therefore does not contain  $NS5$ -branes, and can be worked out explicitly [14, 15]. This is given by a  $T^2 \times T^4/\mathbb{Z}_2$  orbifold with shift vector  $V = \frac{1}{4}(1, \dots, 1, -3)$ , corresponding to  $(b_{U(16)}, \gamma) = (12, -1)$  in eq.(8).

More generically, the condition  $\gamma = -1$  is related to points in the moduli space where a dual description can be given in terms of perturbative  $E_8 \times E_8$  heterotic string compactified on a  $T^6$ , with no Wilson lines, and instanton numbers  $(12 + n, 12 - n)$ , with  $n = 2 + \frac{b_a}{3}$  [16]. In the above Coulomb branch, we hence obtain instanton numbers  $(18, 6)$ , and we are sitting at the point where the moduli spaces of perturbative  $SO(32)$  and  $E_8 \times E_8$  heterotic strings intersect.

### 3. PERTURBATIVE AND NON-PERTURBATIVE CORRECTIONS

The integral (7) can be evaluated using the methods of [9], which consists of dividing  $\hat{Z}_1$  into orbits under the modular group and evaluating the resulting integral for each orbit representative in a suitable unfolded region of the upper complex half-plane. Putting all pieces together and making use of eqs.(5) and (6), we obtain the main result of this note [2],

$$K = -\log(S + \bar{S}) - \sum_i \left\{ \log[(T_i + \bar{T}_i)(U_i + \bar{U}_i)] + \frac{1}{2} \frac{V_{1-loop}^i + V_{E1}^i}{S + \bar{S}} \right\} + \dots, \quad (9)$$

$$V_{1-loop}^i = \frac{4\pi\gamma_i}{3} \frac{E(iU_i, 2)}{T_i + \bar{T}_i}, \quad (10)$$

$$V_{E1}^i = \frac{\gamma_i}{\pi} \sum_{k>j\geq 0, p>0} \frac{e^{-2\pi k p T_i}}{(kp)^2} \left[ \frac{\hat{A}_K(\mathcal{U}_i)}{T_i + \bar{T}_i} - \frac{2ikp}{\mathcal{U}_i - \bar{\mathcal{U}}_i} \frac{E_{10}(\mathcal{U}_i)}{\eta^{24}(\mathcal{U}_i)} \right] + \text{c.c.}, \quad (11)$$

$$f_a = S + \sum_i \left\{ \frac{\pi(b_i^a + 6\gamma_i)T_i}{12} - b_i^a \log \eta(iU_i) - \frac{1}{2} \sum_{k>j\geq 0, p>0} \frac{e^{-2\pi k p T_i}}{kp} \mathcal{A}_{f,i}^a(\mathcal{U}_i) \right\} + \dots, \quad (12)$$

where the dots refer to possible additional corrections from other sectors. Here,  $E(U, k)$  are the non-holomorphic Eisenstein series of order  $k$ , whereas  $\mathcal{A}_{f,i}^a$  is defined as in (8), but replacing the regularized Eisenstein series  $\hat{E}_2$ , appearing inside  $D_{10}E_{10}$ , by the holomorphic one,  $E_2$ . The argument is defined as  $\mathcal{U}_i = (j + ipU_i)/k$ , corresponding to the complex structure induced in the worldvolume of the E1-instantons [17, 18], as will be clear below.

Several comments for the  $T^4/\mathbb{Z}_2$  orbifold limit of K3, where the type I/SO(32) heterotic duality is explicit, are in order:

- *One-loop  $\alpha'$  corrections to the Kähler potential and the gauge kinetic function.* These are given by eq.(10) and the logarithmic term in eq.(12). These expressions were also obtained by direct computation in the type I side in [19, 20]. In our context, these corrections come respectively from non-holomorphic and holomorphic terms in contributions of degenerate orbits under the modular group.
- *Tree-level contribution to the gauge kinetic function.* Because D9-branes are fractional, the gauge kinetic function receives an extra contribution proportional to,

$$\sim \sum \sqrt{\det(P[G + F_2])} T, \quad (13)$$

where the sum runs over the  $\mathbb{C}^2/\mathbb{Z}_2$  singularities and  $P[\dots]$  is the pull-back to the collapsed 2-cycle of the singularity. In the orbifold limit, the volume of the 2-cycle is zero and the contribution from the metric vanishes. However, the U(1) instantons hidden at the singularities [14] lead to a non-trivial linear term in  $T$  in the gauge kinetic function of the D9-branes.

- *Non-perturbative E1 multi-instanton corrections to the Kähler potential and the gauge kinetic function.* They appear as the Polyakov action of an instanton wrapping  $N = kp$  times the  $i$ -th 2-torus, weighted by a modular function of  $\mathcal{U}_i$  and summed over  $N$  and all the possible wrappings.

In order for the instantons to contribute to the gauge kinetic function, only four fermionic neutral zero modes should be massless (corresponding to the “goldstinos”) [21]. A U(1) instanton on top of a singularity behaves as a “gauge” instanton for the U(1) gauge theory inside the corresponding half D5-brane. These instantons are analogous to the ones discussed in [22], with the extra fermionic zero modes lifted by couplings with the D5-branes (see also [23]). Therefore they should be responsible for the 1-instanton contribution. Notice however that in this case there is also a Higgs branch which consists of moving the instanton out of the singularity, leading to a SO(1) instanton (plus its image under the orbifold). In this limit, the instanton has too many zero modes and does not correct the gauge kinetic function.

Similarly, for the  $N$ -instanton contribution, in a generic point of the moduli space the instanton gauge group is  $SO(1)^N$ , and has too many zero modes. Only in the special locus on which all the components of the multi-instanton are on top of the singularity and the gauge group is enhanced to  $U(N)$ , four zero modes survive, with the extra zero modes presumably lifted by interactions with the D5-branes. Explicit examples where instantons only contribute in a given locus of their moduli space have been discussed in detail in the recent literature (see e.g. [24, 25, 26]).

## 4. OUTLOOK

Understanding the leading corrections to the low energy effective theory in generic compactifications would constitute an important input for addressing phenomenological and cosmological questions within String Theory. It would be therefore very interesting to extend these results to more involved and phenomenologically relevant String Theory setups. For that, a better understanding of the blow-up modes or the NS5-brane in heterotic string theory are certainly desirable. We hope that the effort along those lines will continue to be pursued in the near future.

## Acknowledgments

I thank my collaborator E. Dudas, for illuminating discussions and ideas during the elaboration of [2] and afterwards. Also I thank M. Lennek and M. Trapletti for useful comments, and the organizers of the ICHEP'08 conference, in particular those of the Formal Theory session, where this talk was presented. This work is supported by an Individual Marie-Curie IEF. Additional support comes from the contracts ANR-05-BLAN-0079-02, MRTN-CT-2004-005104, MRTN-CT-2004-503369, MEXT-CT-2003-509661 and CNRS PICS # 2530, 3059, 3747.

## References

- [1] E. Witten, Nucl. Phys. B **443**, 85 (1995) [hep-th/9503124]; J. Polchinski and E. Witten, Nucl. Phys. B **460**, 525 (1996) [hep-th/9510169]; C. Angelantonj et al., Phys. Lett. B **385**, 96 (1996) [hep-th/9606169].
- [2] P. G. Camara and E. Dudas, JHEP **0808**, 069 (2008) [0806.3102 [hep-th]].
- [3] P. G. Camara, E. Dudas, T. Maillard and G. Pradisi, Nucl. Phys. B **795** (2008) 453 [0710.3080 [hep-th]].
- [4] V. Kaplunovsky, J. Louis, Nucl. Phys. B **422** (1994) 57 [hep-th/9402005]
- [5] V. Kaplunovsky and J. Louis, Nucl. Phys. B **444** (1995) 191 [hep-th/9502077].
- [6] J. P. Derendinger, S. Ferrara, C. Kounnas and F. Zwirner, Nucl. Phys. B **372**, 145 (1992).
- [7] H. P. Nilles and S. Stieberger, Nucl. Phys. B **499** (1997) 3 [hep-th/9702110].
- [8] V. S. Kaplunovsky, Nucl. Phys. B **307** (1988) 145 [Erratum-ibid. B **382** (1992) 436] [hep-th/9205068, 9205070]
- [9] L. J. Dixon, V. Kaplunovsky and J. Louis, Nucl. Phys. B **355** (1991) 649.
- [10] I. Antoniadis, K. S. Narain and T. R. Taylor, Phys. Lett. B **267** (1991) 37.
- [11] I. Antoniadis, E. Gava and K. S. Narain, Nucl. Phys. B **383** (1992) 93 [hep-th/9204030].
- [12] M. Bianchi and A. Sagnotti, Nucl. Phys. B **361** (1991) 519.
- [13] E. G. Gimon and J. Polchinski, Phys. Rev. D **54** (1996) 1667 [hep-th/9601038].
- [14] M. Berkooz et al., Nucl. Phys. B **475**, 115 (1996) [hep-th/9605184].
- [15] G. Aldazabal, A. Font, L. Ibanez, A. Uranga and G. Violero, Nucl. Phys. B **519**, 239 (1998) [hep-th/9706158].
- [16] S. Stieberger, Nucl. Phys. B **541**, 109 (1999) [hep-th/9807124].
- [17] E. Kiritsis and N. A. Obers, JHEP **9710** (1997) 004 [hep-th/9709058]
- [18] C. Bachas, C. Fabre, E. Kiritsis, N. A. Obers and P. Vanhove, Nucl. Phys. B **509** (1998) 33 [hep-th/9707126]
- [19] I. Antoniadis, C. Bachas, C. Fabre, H. Partouche and T. R. Taylor, Nucl. Phys. B **489** (1997) 160 [hep-th/9608012].

- [20] M. Berg, M. Haack and B. Kors, *JHEP* **0511**, 030 (2005) [hep-th/0508043].
- [21] N. Akerblom, R. Blumenhagen, D. Lust and M. Schmidt-Sommerfeld, *JHEP* **0708** (2007) 044 [0705.2366 [hep-th]].
- [22] C. Petersson, *JHEP* **0805** (2008) 078 [0711.1837 [hep-th]].
- [23] M. Billo et al., *JHEP* **0710** (2007) 091 [0708.3806 [hep-th]]; *JHEP* **0611** (2006) 012 [hep-th/0606013]
- [24] R. Blumenhagen, M. Cvetič, R. Richter and T. Weigand, *JHEP* **0710**, 098 (2007) [0708.0403 [hep-th]];
- [25] I. Garcia-Etxebarria and A. M. Uranga, *JHEP* **0801** (2008) 033 [0711.1430 [hep-th]]
- [26] I. Garcia-Etxebarria, F. Marchesano and A. M. Uranga, *JHEP* **0807** (2008) 028 [0805.0713 [hep-th]].