

# THERMODYNAMICAL CHARACTERISTICS OF "CRYSTAL LATTICE" OF MANY INTERACTING KERR BLACK HOLES IN TOUCHING LIMIT

Vladan Panković<sup>\*,‡</sup>, Simo Ciganović<sup>‡</sup>, Jovan Ivanović<sup>‡</sup>

<sup>\*</sup>Department of Physics, Faculty of Sciences, 21000 Novi Sad,

Trg Dositeja Obradovića 4. , Serbia, vdpan@neobee.net

<sup>‡</sup>Gimnazija, 22320 Indjija, Trg Slobode 2a, Serbia

PACS number : 04.70.Dy

## Abstract

In this work, starting by simple, approximate (quasi-classical) methods presented in our previous works, we reproduce effectively and generalize final results of Herdeiro and Rebelo on the basic thermodynamical characteristics (entropy and temperature) of two interacting Kerr black holes (in touching limit) obtained recently by accurate analysis. Like as it has been done in our previous works, we simply suppose that circumference of the horizon of total black hole (that includes two or, generally, a "crystal lattice" of many interacting Kerr black holes in touching limit, without angular momentum) holds integer number of reduced Compton wave lengths corresponding to mass spectrum of a small quantum system captured at horizon. (Obviously it is conceptually analogous to Bohr quantization postulate interpreted by de Broglie relation in Old, Bohr-Sommerfeld, quantum theory.) It, by simple mathematical methods, first neighbour approximation of the black holes interaction and first thermodynamical law, implies mentioned basic thermodynamical characteristic of the total black hole as well as any its part, i.e. single black hole. Especially, it is shown that, in limit of increasing number of the black holes, entropy and horizon surface of the total black hole stand observables of the discrete spectrum while entropy and horizon surface of the single black hole tends toward observables of the continuous spectrum.

Recently, Herdeiro and Rebelo [1] did an interesting, accurate analysis of two interacting Kerr black holes. In touching limit (corresponding to two horizons touching and when both, opposite angular momentums vanish) there are the following final results

$$M_1 = M_2 = M \quad (1)$$

where  $M_1$  represents the mass of the first one and  $M_2$  mass of the second one black hole,

$$A_1 = A_2 = 2(16\pi M^2) = 2(4\pi R^2) = 2A \quad (2)$$

where  $A_1$  represents the surface of the first one and  $A_2$  surface of the second one black hole,

$$S_1 = S_2 = 2(4\pi M^2) = 2\pi R^2 = 2\frac{A}{4} = 2S \quad (3)$$

where  $S_1$  represents the entropy of the first one and  $S_2$  surface of the second one black hole,

$$T_1 = T_2 = \frac{1}{2} \frac{1}{8\pi M} = \frac{1}{2} T \quad (4)$$

where  $T_1$  represents temperature of the first one and  $T_2$  surface of the second one black hole, and where  $M, R = 2M, A = 4\pi R^2 = 16\pi M^2, S = \pi R^2 = 4\pi M^2$  and  $T = \frac{1}{8\pi M}$  represent the mass, horizon radius, surface, entropy and temperature of a Schwarzschild black hole.

In this work, starting by simple, approximate (quasi-classical) methods presented in our previous works [2]-[5], we shall reproduce effectively and generalize final results of Herdeiro and Rebelo on the basic thermodynamical characteristics (entropy and temperature) of two interacting Kerr black holes (in touching limit). Like as it has been done in our previous works, we simply suppose that circumference of the horizon of total black hole (that includes two or, generally, a "crystal lattice" of many interacting Kerr black holes in touching limit, without angular momentum) holds integer number of reduced Compton wave lengths corresponding to mass (energy) spectrum of a small quantum system captured at horizon. (Obviously it is conceptually analogous to Bohr quantization postulate interpreted by de Broglie relation in Old, Bohr-Sommerfeld, quantum theory.) It, by simple mathematical methods, first neighbour approximation of the black holes interaction and first thermodynamical law, implies mentioned basic thermodynamical characteristic of the total black hole as well as any its part, i.e. single black hole. Especially, it will be shown that, in limit of increasing number of the black holes, entropy and horizon surface of the total black hole stand observables of the discrete spectrum while entropy and horizon surface of the single black hole tends toward observables of the continuous spectrum.

Suppose that there is a series of  $2N$  Kerr black holes distributed along z-axis (representing a simple generalization of two interacting Kerr black holes described by Herdeiro and Rebelo [1]), where  $N$  represents a natural number.

Suppose that initially, without touching limit, first black hole holds the angular momentum  $J_1 = \frac{J}{2}$ , second -  $J_2 = -J$ , third -  $J_3 = J$ , ...,  $2N - 1$ -th -  $J_{2N-1} = J$ , and  $2N$ -th -  $J_{2N} = -\frac{J}{2}$ .

Then, in the first neighbour approximation of the interaction between black holes (and according to translation z-axis symmetry characteristics for Herdeiro and Rebelo dynamics [1]), angular momentum of any black hole slows down toward zero by interaction with two closest neighbouring black holes. Simultaneously, distance between any two neighbouring black holes tends toward touching limit (when corresponding two horizons touching).

In touching limit, masses of individual black holes,  $M_n$  for  $n = 1, 2, \dots, 2N - 1, 2N$ , are mutually equivalent and equal

$$M_1 = M_2 = \dots = M_{2N-1} = M_{2N} = M \quad (5)$$

where  $M$  represents the mass of a Schwarzschild black hole. Then mass of total black hole, that includes all single black hole, equals

$$M_{tot} = 2NM. \quad (6)$$

Suppose the following condition

$$m_{tot} n 2N(2\pi R) = n \quad \text{for} \quad m_{tot} n \ll M_{tot} \quad \text{and} \quad n = 1, 2, \dots \quad (7)$$

where  $m_{tot\ n}$  for  $m_{tot\ n} \ll M_{tot}$  and  $n = 1, 2$ , represent the masses, i.e. mass (energy) spectrum of given small quantum system. It corresponds to expression

$$2N(2\pi R) = n \frac{1}{m_n} = n \lambda_{tot\ r\ n} \quad \text{for } m_{tot\ n} \ll M_{tot} \quad \text{and } n = 1, 2, \dots \quad (8)$$

Here  $2N(2\pi R) = 2N(4\pi M) = 4\pi 2NM = 4\pi M_{tot}$  represents the length of a "N times 8"-like twisted but smooth trajectory of given small quantum system at the total black hole horizon equivalent to  $2N$  times circumference of the single black hole horizon, while

$$\lambda_{tot\ r\ n} = \frac{1}{m_{tot\ n}} \quad (9)$$

represents  $n$ -th reduced Compton wavelength of mentioned small quantum system with mass  $m_{tot\ n}$  for  $n = 1, 2, \dots$ . Expression (8) simply means that *length of a "N times 8"-like twisted but smooth trajectory on the total black hole horizon holds exactly n corresponding n -th reduced Compton wave lengths of a small quantum system with mass  $m_{tot\ n}$  captured at the total black hole horizon, for  $n = 1, 2, \dots$* . Obviously, it is essentially analogous to well-known Bohr's angular momentum quantization postulate interpreted via de Broglie relation. However, there is a principal difference with respect to Bohr's atomic model. Namely, in Bohr's atomic model different quantum numbers  $n = 1, 2, \dots$ , correspond to different circular orbits (with circumferences proportional to  $n^2 = 1^2, 2^2, \dots$ ). Here any quantum number  $n = 1, 2, \dots$ , corresponds to the same "N times 8"-like twisted but smooth orbit (with length  $2N(2\pi R)$ ).

According to (7) it follows

$$m_{tot\ n} = n \frac{1}{2N(2\pi R)} = n \frac{1}{8\pi M} = n \frac{1}{4\pi M_{tot}} \equiv n m_{tot1} \quad \text{for } m_{tot\ n} \ll M_{tot} \quad \text{and } n = 1, 2, \dots \quad (10)$$

where  $m_{tot1}$  represents the minimal small quantum system mass

$$m_{tot1} = \frac{1}{2N(2\pi R)} = \frac{1}{4\pi M_{tot}}. \quad (11)$$

Obviously,  $m_{tot1}$  depends of  $M_{tot}$  so that  $m_{tot1}$  decreases when  $M_{tot}$  increases and vice versa. For a "macroscopic" black hole, i.e. for  $M_{tot} \gg 1$  it follows  $m_{tot1} \ll 1 \ll M_{tot}$ .

Now, we shall suppose that quotient of  $M_{tot}$  and  $m_{tot1}$  represents Bekenstein-Hawking total black hole entropy, i.e.

$$S_{tot} = \frac{M_{tot}}{m_{tot1}} = 4\pi M_{tot}^2 = (2N)^2 4\pi M^2 = (2N)^2 S, \quad (12)$$

where, according to Bekenstein postulate,

$$A_{tot} = 4S_{tot} = 16\pi M_{tot}^2 = (2N)^2 16\pi M^2 = (2N)^2 4S \quad (13)$$

represents the total black hole surface area.

Differentiation of (12) yields

$$dS_{tot} = 8\pi M_{tot} dM_{tot} \quad (14)$$

or, after simple transformation,

$$dM_{tot} = \frac{1}{8\pi M_{tot}} dS_{tot}. \quad (15)$$

Expression (15), representing the first thermodynamical law, implies that term

$$T_{tot} = \frac{1}{8\pi M_{tot}} = \frac{m_{tot1}}{2} = \frac{1}{2N} \frac{1}{8\pi M} = \frac{T}{2N} \quad (16)$$

represents Hawking temperature of the total black hole, while  $T = \frac{1}{8\pi M}$ , as it has been previously discussed, represents Hawking temperature of a Schwarzschild black hole.

Further, since entropy represents an additive variable and since all  $2n$  single black holes are physically identical, it follows, according to (29),

$$S_1 = S_2 = \dots = S_{2N-1} = S_{2N} = \frac{S_{tot}}{2N} = (2N)4\pi M^2 = (2N)S. \quad (17)$$

Finally, since according to Bekenstein postulate horizon surface proportional to entropy represents an additive variable too, it follows, according to (12),

$$A_1 = A_2 = \dots = A_{2N-1} = A_{2N} = \frac{A_{tot}}{2N} = (2N)16\pi M^2 = (2N)A. \quad (18)$$

It is not hard to see that our expressions (5), (17) and (18) represent consequent generalization of Herdeiro and Rebelo expressions (1), (3) and (2) respectively. Especially, for  $N = 1$ , i.e.  $2N = 2$ , or for two interacting Kerr black holes in touching limit, our expressions (5), (17) and (18) are identical to Herdeiro and Rebelo expressions (1), (3) and (2) respectively.

Finally, it can be observed the following. By approximate changing of the differentials by finite differences, (14) turns out approximately, in

$$\Delta S_{tot} = 8\pi M_{tot} \Delta M_{tot} \quad \text{for} \quad \Delta M_{tot} \ll M_{tot} \quad \text{and} \quad n = 1, 2, \dots \quad (19)$$

Now, assume

$$\Delta M_{tot} = nm_{tot1} \quad \text{for} \quad n = 1, 2, \dots \quad (20)$$

which, according to (11), after substituting in (19), yields

$$\Delta S_{tot} = 2n \quad \text{for} \quad n = 1, 2, \dots \quad (21)$$

or, according to (13),

$$\Delta A_{tot} = (2n)4 = (2n)2^2 \quad \text{for} \quad n = 1, 2, \dots \quad (22)$$

Obviously, expression (18) and (19) represent Bekenstein quantization of the entropy and horizon surface of the total black hole.

Since entropy and horizon surface represent additive variables it follows that Bekenstein entropy and surface quantization for single black hole, according to (21), (22), are

$$\Delta S_1 = \Delta S_2 = \dots = \Delta S_{2N-1} = \Delta S_{2N} = \frac{\Delta S_{tot}}{2N} = \frac{2n}{2N} \quad \text{for} \quad n = 1, 2, \dots \quad (23)$$

$$\Delta A_1 = \Delta A_2 = \dots = \Delta A_{2N-1} = \Delta A_{2N} = \frac{\Delta A_{tot}}{2N} = 4 \frac{2n}{2N} \quad \text{for} \quad n = 1, 2, \dots \quad (24)$$

It represents an interesting result. Namely, when number of the black holes,  $2N$ , increases, entropy and horizon surface of the total black hole, according to (12), (13), increase too, while

quants of entropy and horizon surface of the total black hole stand the same and discrete according to Bekenstein quantization rules (21), (22). In other words, in limit of increasing  $2N$ , entropy and horizon surface of the total black hole stand observables of the discrete spectrum. Simultaneously, when number of the black holes,  $2N$ , increases, entropy and horizon surface of the single black hole increase too according to (17), (18), while quants of entropy and horizon surface of the single black hole decrease according to (23), (24). In other words, in limit of increasing  $2N$ , entropy and horizon surface of the total black hole stand observables of the discrete spectrum while entropy and horizon surface of the single black hole tends toward observables of the continuous spectrum.

Thus, in conclusion it can be repeated and pointed out the following. In this work we reproduce effectively and generalize recent results of Herdeiro and Rebelo on the basic thermodynamical characteristics (entropy and temperature) of two interacting Kerr black holes (in touching limit) obtained recently by accurate analysis. We started by simple supposition that circumference of the horizon of total black hole (that includes two or, generally, a "crystal lattice" of many interacting Kerr black holes in touching limit) holds integer number of reduced Compton wave lengths corresponding to mass (energy) spectrum of a small quantum systems captured at horizon (that is conceptually analogous to Bohr quantization postulate interpreted by de Broglie relation). It, by simple mathematical methods, first neighbor approximation of the black holes interaction and first thermodynamical law, implies mentioned basic thermodynamic characteristic of the total black hole as well as any its part, i.e. single black hole. Finally, we showed that, in limit of increasing number of the black holes, entropy and horizon surface of the total black hole stand observables of the discrete spectrum while entropy and horizon surface of the single black hole tends toward observables of the continuous spectrum.

## 1 References

- [1 ] C. A. R. Herdeiro, C. Rebelo, *On the Interaction between Two Kerr Black Holes*, gr-qc/0808.3941
- [2 ] V. Pankovic, M. Predojevic, P. Grujic, *A Bohr's Semiclassical Model of the Black Hole Thermodynamics*, gr-qc/0709.1812
- [3 ] V. Pankovic, J. Ivanovic, M. Predojevic, A.-M. Radakovic, *The Simplest Determination of the Thermodynamical Characteristics of Schwarzschild, Kerr and Reisner-Nordström black hole*, gr-qc/0803.0620
- [4 ] V. Pankovic, S. Ciganovic, R. Glavatovic, *The Simplest Determination of the Thermodynamical Characteristics of Kerr-Newman Black Hole*, gr-qc/0804.2327
- [5 ] V. Pankovic, R. Glavatovic, S. Ciganovic, D-H. Petkovic, L. Martinovic, *Single Horizon Black Hole "Laser" and a Solution of the Information Loss Paradox*, gr-qc/0807.1840