

More non-abelian loop Toda solitons

Kh. S. Nirov* and A. V. Razumov†

Fachbereich C-Physik, Bergische Universität Wuppertal
D-42097 Wuppertal, Germany

Abstract

We find soliton-like solutions in a particular case of non-abelian loop Toda equations associated with general linear groups. We use the formalism of rational dressing based on appropriate block-matrix representation suggested by the \mathbb{Z} -gradation under consideration. We present solutions in a form of a direct matrix generalization of the Hirota's soliton constructions already well-known for the case of abelian loop Toda systems.

Mathematics Subject Classification (2000). 37K10, 37K15, 35Q51, 35Q58

Keywords. Non-abelian loop Toda systems, rational dressing method, soliton solutions

1 Introduction: Toda equations

The loop Toda systems associated with complex classical Lie groups, when the corresponding loop Lie algebras are endowed with integrable \mathbb{Z} -gradations with finite-dimensional grading subspaces, can be given by the equation [1, 2, 3, 4, 5] (see also [6, 7, 8, 9, 10] and references therein),

$$\partial_+(\gamma^{-1}\partial_-\gamma) = [c_-, \gamma^{-1}c_+\gamma], \quad (1)$$

supplied by the conditions

$$\partial_+c_- = 0, \quad \partial_-c_+ = 0. \quad (2)$$

Here, γ is a mapping of \mathbb{R}^2 to a complex classical Lie group G having a block-diagonal form

$$\gamma = \begin{pmatrix} \Gamma_1 & & & \\ & \Gamma_2 & & \\ & & \ddots & \\ & & & \Gamma_p \end{pmatrix},$$

*On leave of absence from the *Institute for Nuclear Research of the Russian Academy of Sciences, 60th October Ave 7a, 117312 Moscow, Russia*

†On leave of absence from the *Institute for High Energy Physics, 142281 Protvino, Moscow Region, Russia*

so that for each $\alpha = 1, \dots, p$ the mapping Γ_α is a mapping of \mathbb{R}^2 to the Lie group $GL_{n_\alpha}(\mathbb{C})$, or its certain subgroup. The p positive integers n_α , defining the block-matrix structure, are suggested by the \mathbb{Z} -gradation under consideration. Further, c_+ and c_- are mappings of \mathbb{R}^2 to the Lie algebra \mathfrak{g} of the Lie group G . The mapping c_+ has the block-matrix structure

$$c_+ = \begin{pmatrix} 0 & C_{+1} & & & \\ & 0 & \ddots & & \\ & & \ddots & \ddots & \\ & & & 0 & C_{+(p-1)} \\ C_{+0} & & & & 0 \end{pmatrix},$$

where for each $\alpha = 1, \dots, p-1$ the mapping $C_{+\alpha}$ is a mapping of \mathbb{R}^2 to the space of $n_\alpha \times n_{\alpha+1}$ complex matrices, and C_{+0} is a mapping of \mathbb{R}^2 to the space of $n_p \times n_1$ complex matrices. The mapping c_- has a similar block-matrix structure,

$$c_- = \begin{pmatrix} 0 & & & & C_{-0} \\ C_{-1} & 0 & & & \\ & \ddots & \ddots & & \\ & & \ddots & 0 & \\ & & & C_{-(p-1)} & 0 \end{pmatrix},$$

where for each $\alpha = 1, \dots, p-1$ the mapping $C_{-\alpha}$ is a mapping of \mathbb{R}^2 to the space of $n_{\alpha+1} \times n_\alpha$ complex matrices, and C_{-0} is a mapping of \mathbb{R}^2 to the space of $n_1 \times n_p$ complex matrices. The notation ∂_+ and ∂_- is used for the derivatives over the standard coordinates in \mathbb{R}^2 which we denote by z^+ and z^- .

In the present paper we consider a particular case of non-abelian loop Toda equations associated with the complex general linear group $GL_n(\mathbb{C})$, where the \mathbb{Z} -gradation of the corresponding loop Lie algebra is generated by an inner automorphism of the initial complex Lie algebra $\mathfrak{gl}_n(\mathbb{C})$, leading to an equal-size block-matrix partition of the algebra and group elements, that is $n_\alpha = n/p = n_*$ for all $\alpha = 1, \dots, p$. Moreover, we assume for simplicity that all nonzero entries of the block-matrix representation of c_+ and c_- are unit matrices, $C_{\pm\alpha} = I_{n_\alpha}$.

Now, the Toda equation (1) can be written as an infinite periodic system,

$$\partial_+(\Gamma_\alpha^{-1}\partial_-\Gamma_\alpha) + \Gamma_\alpha^{-1}\Gamma_{\alpha+1} - \Gamma_{\alpha-1}^{-1}\Gamma_\alpha = 0, \quad (3)$$

with Γ_α subject to the condition $\Gamma_{\alpha+p} = \Gamma_\alpha$. This particular case of Toda systems was introduced in [11]. The case of $p = 2$ and $n_* = 2$, with certain unitarity constraints, was also treated in [12] as a simplest matrix generalization of the sin-Gordon equation. Later on, it was considered in [13] as an example of nonlinear integrable systems for which a combination of the notion of quasi-determinants and a Marchenko lemma would allow for constructing certain soliton-like solutions. We will apply to the equations (3) the rational dressing method of [11] in order to construct soliton solutions for this particular case of non-abelian loop Toda equations. Specifically, we follow our recent consideration presented in [14, 15].

2 Rational dressing

We see that the constant matrices c_- and c_+ commute. Hence, it is obvious that

$$\gamma = I_n, \quad (4)$$

where I_n is the $n \times n$ unit matrix, is a solution to the Toda equation (1). For the formalism of rational dressing it is crucial that the Toda equation (1), (2) can be represented as the zero-curvature condition for a flat connection in a trivial principal fiber bundle, satisfying the grading and gauge-fixing conditions, see, for example, the books [1, 2]. Actually, having such a connection we have a solution to the Toda equations (1). In our case, the corresponding base manifold is the Euclidean plane \mathbb{R}^2 and the fiber coincides with the untwisted loop group $\mathcal{L}_{a,p}(\mathrm{GL}_n(\mathbb{C}))$, where a denotes the inner automorphism of $\mathrm{GL}_n(\mathbb{C})$ of order p acting on an element $g \in \mathrm{GL}_n(\mathbb{C})$ in accordance with the equality

$$a(g) = hgh^{-1}.$$

Here h is a block-diagonal matrix defined by the relation

$$h_{\alpha\beta} = \epsilon_p^{p-\alpha+1} I_{n_*} \delta_{\alpha\beta}, \quad \alpha, \beta = 1, \dots, p, \quad (5)$$

where $\epsilon_p = e^{2\pi i/p}$ is the p th principal root of unity.

Using the exponential law [16, 17], it is convenient to identify the mapping generating a flat connection under consideration with a smooth mapping of $\mathbb{R}^2 \times S^1$ to $\mathrm{GL}_n(\mathbb{C})$ and the connection components with smooth mappings of $\mathbb{R}^2 \times S^1$ to $\mathfrak{gl}_n(\mathbb{C})$. Below we think of the circle S^1 as consisting of complex numbers of modulus one.

Denote the mapping generating the connection corresponding to the solution (4) by φ . For the case under consideration, the rational dressing method consists in finding a mapping ψ of $\mathbb{R}^2 \times S^1$ to $\mathrm{GL}_n(\mathbb{C})$, such that the grading and gauge-fixing conditions for the components of the flat connection generated by the mapping $\varphi\psi$ are satisfied. We assume that the analytic extension of the mapping ψ from S^1 to the whole Riemann sphere is a rational mapping given by the expression

$$\psi = \left(I_n + \sum_{i=1}^r \sum_{k=1}^p \frac{\lambda}{\lambda - \epsilon_p^k \mu_i} h^k P_i h^{-k} \right) \psi_0. \quad (6)$$

Here λ is the standard coordinate in \mathbb{C} , ψ_0 is a mapping of \mathbb{R}^2 to the Lie subgroup of $\mathrm{GL}_n(\mathbb{C})$ formed by the elements $g \in \mathrm{GL}_n(\mathbb{C})$ subject to the equality $hgh^{-1} = g$, and P_i are some smooth mappings of \mathbb{R}^2 to the algebra $\mathrm{Mat}_n(\mathbb{C})$ of $n \times n$ complex matrices. It is also assumed here that $\mu_i \neq 0$, $\mu_i^p \neq \mu_j^p$ for all $i \neq j$. The expression (6) is obtained from an initial rational mapping by averaging over the action of the inner automorphism a , where it was used that $a^p = \mathrm{id}_{\mathrm{GL}(\mathbb{C})}$. Further, we suppose that the analytic extension of the corresponding inverse mapping to the whole Riemann sphere has a similar structure,

$$\psi^{-1} = \psi_0^{-1} \left(I_n + \sum_{i=1}^r \sum_{k=1}^p \frac{\lambda}{\lambda - \epsilon_p^k v_i} h^k Q_i h^{-k} \right),$$

with the pole positions satisfying the conditions $v_i \neq 0$, $v_i^p \neq v_j^p$ for all $i \neq j$, and additionally $v_i^p \neq \mu_j^p$ for any i and j . Note that the mappings $\psi(\lambda)$ and $\psi^{-1}(\lambda)$ are regular at the points $\lambda = 0$ and $\lambda = \infty$.

By definition, the equality

$$\psi^{-1}\psi = I_n$$

is valid on S^1 . Since ψ and ψ^{-1} are rational mappings, this equality is valid on the whole Riemann sphere. Therefore, the residues of $\psi^{-1}\psi$ at the points μ_i and ν_i must vanish. This leads to certain relations to be satisfied by the mappings P_i and Q_i . Further, for the components of the flat connection generated by the mapping $\varphi\psi$ we find the expressions

$$\begin{aligned}\omega_- &= \psi^{-1}\partial_-\psi + \lambda^{-1}\psi^{-1}c_-\psi, \\ \omega_+ &= \psi^{-1}\partial_+\psi + \lambda\psi^{-1}c_+\psi.\end{aligned}$$

We see that ω_- is a rational mapping having simple poles at μ_i, ν_i and zero. Similarly, ω_+ is a rational mapping having simple poles at μ_i, ν_i and infinity. We need a connection satisfying the grading and gauge-fixing conditions. The grading condition in our case means that for each point of \mathbb{R}^2 the component ω_- is rational and has the only simple pole at zero, and the component ω_+ is rational and has the only simple pole at infinity. Therefore, we require that the residues of ω_- and ω_+ at the points μ_i and ν_i must vanish. And this requirement imposes additional conditions on the mappings P_i and Q_i . Having these and the above mentioned relations fulfilled by P_i and Q_i , and besides assuming that $\psi_0 = I_n$ that resolves the gauge-fixing constraints, we can see that the mapping $\gamma = \psi(\infty)$, with fixed $2r$ complex numbers μ_i, ν_i , satisfies the Toda equation (1). In the block-matrix form we have the expressions

$$\gamma_{\alpha\beta} = \delta_{\alpha\beta} \left(I_{n_*} + p \sum_{i=1}^r (P_i)_{\alpha\alpha} \right), \quad \gamma_{\alpha\beta}^{-1} = \delta_{\alpha\beta} \left(I_{n_*} + p \sum_{i=1}^r (Q_i)_{\alpha\alpha} \right), \quad (7)$$

where $(P_i)_{\alpha\beta}$ and $(Q_i)_{\alpha\beta}$ are $n_* \times n_*$ complex matrices satisfying certain conditions as briefly mentioned above, see [14, 15].

To make the solution (7) explicit we should specify the matrix-valued functions P_i and Q_i . We first note that, if we suppose that the functions P_i and Q_i take values in the space of the matrices of maximum rank, then we come to the trivial solution (4). Hence, we assume that the values of the functions P_i and Q_i are not matrices of maximum rank. The case given by matrices of rank one was elaborated in [15]. Now we consider another interesting case, where the functions P_i and Q_i take values in the space of $n \times n$ matrices of rank n_* . Such functions can be represented as

$$P_i = u_i {}^t w_i, \quad Q_i = x_i {}^t y_i, \quad (8)$$

where u, w, x and y are functions on \mathbb{R}^2 taking values in the space of $n \times n_*$ complex matrices of rank n_* . The used \mathbb{Z} -gradation suggests the most convenient block-matrix representation for the matrices of the form (8) as

$$(P_i)_{\alpha\beta} = u_{i,\alpha} {}^t w_{i,\beta}, \quad (Q_i)_{\alpha\beta} = x_{i,\alpha} {}^t y_{i,\beta},$$

where the standard matrix multiplication of the $n_* \times n_*$ matrix-valued functions $u_{i,\alpha}, x_{i,\alpha}$ by the $n_* \times n_*$ matrix-valued functions ${}^t w_{i,\beta}, {}^t y_{i,\beta}$ is implied.

Then, within the formalism of rational dressing we see that the functions w and x can be expressed via the functions y and u , and we find the expressions

$$(P_i)_{\alpha\beta} = -\frac{1}{p} u_{i,\alpha} \sum_{j=1}^r (R_\beta^{-1})_{ij} {}^t y_{j,\beta}, \quad (Q_i)_{\alpha\beta} = \frac{1}{p} \sum_{j=1}^r u_{j,\alpha} \frac{1}{\mu_j} (R_{\alpha+1}^{-1})_{ji} \nu_i {}^t y_{i,\beta},$$

with $n_* r \times n_* r$ matrix-valued functions R_α defined through its $n_* \times n_*$ blocks as

$$(R_\alpha)_{ij} = \frac{1}{v_i^p - \mu_j^p} \sum_{\beta=1}^p v_i^{p-|\beta-\alpha|_p} \mu_j^{|\beta-\alpha|_p} {}^t y_{i,\beta} u_{j,\beta},$$

where $|x|_p$ denotes the residue of division of x by p . Note that, unlike the abelian and non-abelian cases considered in [14, 15], here $(R_\alpha)_{ij}$ is an $n_* \times n_*$ matrix-valued function for any i, j .

It is convenient to use quantities defined as $\tilde{u}_{i,\alpha} = u_{i,\alpha} \mu_i^\alpha$, $\tilde{y}_{i,\alpha} = y_{i,\alpha} v_i^{-\alpha}$ and $(\tilde{R}_\alpha)_{ij} = v_i^{-\alpha} (R_\alpha)_{ij} \mu_j^\alpha$. For the matrices \tilde{R}_α we have explicitly the relation

$$(\tilde{R}_\alpha)_{ij} = \frac{1}{v_i^p - \mu_j^p} \left(\mu_j^{\alpha-1} \sum_{\beta=1}^{\alpha-1} \tilde{y}_{i,\beta} \tilde{u}_{j,\beta} + v_i^p \sum_{\beta=\alpha}^p \tilde{y}_{i,\beta} \tilde{u}_{j,\beta} \right).$$

Then, in terms of these quantities, the $n_* \times n_*$ matrix-valued functions Γ_α can be written as

$$\Gamma_\alpha = I_{n_\alpha} - \sum_{i,j=1}^r \tilde{u}_{i,\alpha} (\tilde{R}_\alpha^{-1})_{ij} \tilde{y}_{j,\alpha}.$$

Similarly, for the corresponding inverse mappings we obtain the expression

$$\Gamma_\alpha^{-1} = I_{n_\alpha} + \sum_{i,j=1}^r \tilde{u}_{i,\alpha} (\tilde{R}_{\alpha+1}^{-1})_{ij} \tilde{y}_{j,\alpha},$$

which can be useful for verifying the Toda equations.

To finally satisfy the conditions imposed earlier on the matrix-valued functions P_i and Q_i , we also demand the validity of the equations

$$\partial_- u_i = -\mu_i^{-1} c_- u_i, \quad \partial_+ u_i = -\mu_i c_+ u_i \quad (9)$$

$$\partial_- y_i = v_i^{-1} {}^t c_- y_i, \quad \partial_+ y_i = v_i {}^t c_+ y_i, \quad (10)$$

see, for example, [15]. Using the explicit forms of the matrices c_\pm , we write down the general solutions to (9), (10) as¹

$$u_{i,\beta} = \sum_{\alpha=1}^p \epsilon_p^{\beta\alpha} \exp \left(-\mu_i^{-1} \epsilon_p^{-\alpha} z^- - \mu_i \epsilon_p^\alpha z^+ \right) c_{i,\alpha}, \quad (11)$$

$$y_{i,\beta} = \sum_{\alpha=1}^p \epsilon_p^{\beta\alpha} \exp \left(v_i^{-1} \epsilon_p^\alpha z^- + v_i \epsilon_p^{-\alpha} z^+ \right) d_{i,\alpha}, \quad (12)$$

where $c_{i,\alpha}$ and $d_{i,\alpha}$ are $n_* \times n_*$ complex matrices meaning the initial-value data for the equations (9), (10). With these solutions we immediately obtain for the blocks of the matrix-valued functions \tilde{R}_α the following expression:

$$(\tilde{R}_\alpha)_{ij} = \sum_{\beta,\delta=1}^p e^{Z_\alpha(v_i) - Z_\alpha(\mu_j)} \frac{\epsilon_p^{\alpha(\beta+\delta)}}{1 - \mu_j v_i^{-1} \epsilon_p^{\beta+\delta}} v_i^{-\alpha} ({}^t d_{i,\beta} c_{j,\delta}) \mu_j^\alpha,$$

where we have introduced the notation $Z_\alpha(\mu_i) = \mu_i^{-1} \epsilon_p^{-\alpha} z^- + \mu_i \epsilon_p^\alpha z^+$.

¹It should be instructive to confer these expressions with those obtained in [15] for the case of more general \mathbb{Z} -gradations.

3 Soliton solutions

To construct solutions making sense as r -solitons, that is, by the definition we use here, solutions depending on r linear combinations of independent variables z^+ and z^- , we assume that for each value of the index $i = 1, \dots, r$ the initial-value data of the Toda system under consideration are such that matrix-valued coefficients $c_{i,\alpha}$ are different from zero for only one value of α , which we denote by I_i , and that the matrix-valued coefficients $d_{i,\alpha}$ are different from zero for only two values of α , which we denote by J_i and K_i . We also use for such non-vanishing initial-data $n_* \times n_*$ matrices the notation $d_{J_i} = d_{i,J_i}$, $d_{K_i} = d_{i,K_i}$ and $c_{I_i} = c_{i,I_i}$. For the $n_* \times n_*$ blocks of the matrix-valued functions \tilde{u}_i and \tilde{y}_i this assumption gives

$$\tilde{u}_{i,\alpha} = \mu_i^\alpha \epsilon_p^{\alpha I_i} e^{-Z_{I_i}(\mu_i)} c_{I_i}, \quad (13)$$

$${}^t\tilde{y}_{i,\alpha} = \nu_i^{-\alpha} \epsilon_p^{\alpha J_i} e^{Z_{-J_i}(\nu_i)} {}^t d_{J_i} + \nu_i^{-\alpha} \epsilon_p^{\alpha K_i} e^{Z_{-K_i}(\nu_i)} {}^t d_{K_i}. \quad (14)$$

With these relations, we can write the expression for the mappings Γ_α in the form

$$\Gamma_\alpha = I_{n_\alpha} - \sum_{i,j=1}^r c_{I_i} (\tilde{R}'_\alpha)^{-1}_{ij} ({}^t d_{J_j} + E_{\alpha,j} {}^t d_{K_j}), \quad (15)$$

where we have used the notation

$$(\tilde{R}'_\alpha)_{ij} = \tilde{D}_{ij}(J) + E_{\alpha,i} \tilde{D}_{ij}(K), \quad E_{\alpha,i} = \epsilon_p^{\alpha \rho_i} e^{Z_i(\zeta)}, \quad (16)$$

with the dependence on the variables z^+ and z^- given through the functions

$$Z_i(\zeta) = \kappa_{\rho_i} (\zeta_i^{-1} z^- + \zeta_i z^+),$$

and the convenient parameters $\rho_i = K_i - J_i$, $\zeta_i = -i\nu_i \epsilon_p^{-(K_i+J_i)/2}$, $\kappa_{\rho_i} = 2 \sin(\pi \rho / p)$, and besides

$$\tilde{D}_{ij}(A) = \frac{{}^t d_{A_i} c_{I_j}}{1 - \nu_i^{-1} \mu_j \epsilon_p^{A_i+I_j}}, \quad i, j = 1, \dots, r, \quad (17)$$

for the $n_* \times n_*$ blocks of the $n_* r \times n_* r$ matrices $\tilde{D}(A)$, $A = J, K$ already introduced in our previous paper [15].

We assume that the $n_* \times n_*$ matrices c_{I_i} are non-degenerate. Let us multiply Γ_α in (15) by c_{I_ℓ} from the right hand side and write

$$\Gamma_\alpha c_{I_\ell} = \sum_{i,j=1}^r c_{I_i} (\tilde{R}'_\alpha)^{-1}_{ij} [(\tilde{R}'_\alpha)_{j\ell} - ({}^t d_{J_j} + E_{\alpha,j} {}^t d_{K_j}) c_{I_\ell}]. \quad (18)$$

It is not difficult to see that

$$(\tilde{R}'_\alpha)_{j\ell} - ({}^t d_{J_j} + E_{\alpha,j} {}^t d_{K_j}) c_{I_\ell} = \nu_j^{-1} \epsilon_p^{J_j} (\tilde{R}'_{\alpha+1})_{j\ell} \mu_\ell \epsilon_p^{I_\ell}.$$

Now, multiplying (18) from the right hand side by the inverse matrix $c_{I_\ell}^{-1}$ and summing up over $\ell = 1, \dots, r$, we obtain the following expression:

$$\Gamma_\alpha = \frac{1}{r} \sum_{i,j,k=1}^r c_{I_i} (\tilde{R}'_\alpha)^{-1}_{ij} \nu_j^{-1} \epsilon_p^{J_j} (\tilde{R}'_{\alpha+1})_{jk} \mu_k \epsilon_p^{I_k} c_{I_k}^{-1}.$$

To give the soliton solutions a final form, it is also convenient to make use of simplest symmetries of the Toda equation. It is clear from the relations (3) that the transformations

$$\Gamma_\alpha \rightarrow \xi \Gamma_\alpha, \quad \Gamma_\alpha \rightarrow x^{-1} \Gamma_\alpha x \quad (19)$$

for a nonzero constant ξ and a non-singular constant $n_* \times n_*$ matrix x are symmetry transformations of the Toda system under consideration.

In particular, using the symmetry transformations (19) with $\xi = \mu^{-1} \nu \epsilon_p^{-(I+J)}$ and $x = c_I$, we can write the one-soliton solution as

$$\Gamma_\alpha = \tilde{R}'_\alpha{}^{-1} \tilde{R}'_{\alpha+1}, \quad (20)$$

where the matrices \tilde{R}'_α are explicitly given by the relations (16), (17). It reproduces exactly the one-soliton solution constructed in [13] by means of a different approach.

Also the multi-soliton solutions, $r \geq 2$, can be written in a compact form,

$$\Gamma_\alpha = {}^T c_I (\tilde{R}'_\alpha{}^{-1}) \mathcal{N}_J^{-1} (\tilde{R}'_{\alpha+1}) \mathcal{M}_I c_I^{-1}. \quad (21)$$

Here we use the notation ${}^T c_I$ and c_I^{-1} for $n_* \times n$ and $n \times n_*$ matrices, respectively, defined as ${}^T c_I = (c_{I_1} \dots c_{I_r})$ and $c_I^{-1} = ({}^t c_{I_1}^{-1} \dots {}^t c_{I_r}^{-1})$, and the notation \mathcal{N}_J^{-1} and \mathcal{M}_I for block-diagonal $n_* r \times n_* r$ matrices defined as

$$\mathcal{N}_J^{-1} = \begin{pmatrix} \nu_1^{-1} \epsilon_p^{J_1} I_{n_*} & & & \\ & \ddots & & \\ & & \nu_r^{-1} \epsilon_p^{J_r} I_{n_*} & \\ & & & \ddots \end{pmatrix}, \quad \mathcal{M}_I = \begin{pmatrix} \mu_1 \epsilon_p^{I_1} I_{n_*} & & & \\ & \ddots & & \\ & & \mu_r \epsilon_p^{I_r} I_{n_*} & \\ & & & \ddots \end{pmatrix}.$$

Putting $n_* = 1$ we can recover the abelian case analyzed in [14]. The solutions (20), (21) may be regarded as a novel non-abelian generalization, complementary to that of [15], of the Hirota's soliton constructions [18] successfully used for the abelian loop Toda systems.

To obtain more solutions to these equations, not necessarily soliton-like, one should keep nonzero more initial-value data $c_{i,\alpha}$ and $d_{i,\alpha}$ entering (11), (12), that leads to a more general expressions for $\tilde{u}_{i,\alpha}$ and $\tilde{y}_{i,\alpha}$ than (13) and (14).

Acknowledgments

We express our gratitude to Profs. H. Boos, F. Göhmann and A. Klümper for hospitality at the University of Wuppertal and interesting discussions. This work was supported in part by the by the joint DFG–RFBR grant #08–01–91953.

References

- [1] A. N. Leznov and M. V. Saveliev, *Group-theoretical methods for the integration of non-linear dynamical systems*, (Birkhauser, Basel, 1992).
- [2] A. V. Razumov and M. V. Saveliev, *Lie algebras, geometry and Toda-type systems*, (Cambridge University Press, Cambridge, 1997).
- [3] Kh. S. Nirov and A. V. Razumov, *On \mathbb{Z} -gradations of twisted loop Lie algebras of complex simple Lie algebras*, Commun. Math. Phys. **267** (2006) 587–610, [arXiv:math-ph/0504038].

- [4] Kh. S. Nirov and A. V. Razumov, *Toda equations associated with loop groups of complex classical Lie groups*, Nucl. Phys. **B782** (2007) 241–275, [arXiv:math-ph/0612054].
- [5] Kh. S. Nirov and A. V. Razumov, *\mathbb{Z} -graded loop Lie algebras, loop groups, and Toda equations*, Theor. Math. Phys. **154** (2008) 385–404, [arXiv:0705.2681].
- [6] A. V. Mikhailov, M. A. Olshanetsky and A. M. Perelomov, *Two-dimensional generalized Toda lattice*, Commun. Math. Phys. **79** (1981) 473–488.
- [7] L. A. Ferreira, J. L. Miramontes and J. S. Guillén, *Solitons, τ -functions and hamiltonian reduction for non-Abelian conformal affine Toda theories*, Nucl. Phys. **B449** (1995) 631–679, [arXiv:hep-th/9412127].
- [8] L. A. Ferreira, J. L. Miramontes and J. S. Guillén, *Tau-functions and dressing transformations for zero-curvature affine integrable equations*, J. Math. Phys. **38** (1997) 882–901, [arXiv:hep-th/9606066].
- [9] C. R. Fernández-Pousa, M. V. Gallas, T. J. Hollowood and J. L. Miramontes, *The symmetric space and homogeneous sine-Gordon theories*, Nucl. Phys. **B484** (1997) 609–630, [arXiv:hep-th/9606032].
- [10] L. Miramontes, *Tau-functions generating the conservation laws for generalized integrable hierarchies of KdV and affine-Toda type*, Nucl. Phys. **B547** (1999) 623–663, [arXiv:hep-th/9809052].
- [11] A. V. Mikhailov, *The reduction problem and the inverse scattering method*, Physica **3D** (1981) 73–117.
- [12] Q-Han Park and H. J. Shin, *Classical matrix sine-Gordon theory*, Nucl. Phys. **B458** (1996) 327–354, [arXiv:hep-th/9505017].
- [13] P. Etingof, I. Gelfand, and V. Retakh, *Non-Abelian integrable systems, quasideterminants, and Marchenko Lemma*, Math. Res. Lett. **5** (1998) 1–12, [arXiv:q-alg/9707017].
- [14] Kh. S. Nirov and A. V. Razumov, *Abelian Toda solitons revisited*, [arXiv:0802.0593].
- [15] Kh. S. Nirov and A. V. Razumov, *Solving non-abelian loop Toda equations*, [arXiv:0809.3944].
- [16] A. Kriegl and P. Michor, *Aspects of the theory of infinite dimensional manifolds*, Diff. Geom. Appl. **1** (1991) 159–176, [arXiv:math.DG/9202206].
- [17] A. Kriegl and P. Michor, *The Convenient Setting of Global Analysis*, Mathematical Surveys and Monographs, vol. 53, (American Mathematical Society, Providence, RI, 1997).
- [18] R. Hirota, *The Direct Method in Soliton Theory*, (Cambridge University Press, Cambridge, 2004).