

THE STRATIFIED STRUCTURE OF SPACES OF SMOOTH ORBIFOLD MAPPINGS

JOSEPH E. BORZELLINO AND VICTOR BRUNSDEN

ABSTRACT. We consider four notions of maps between smooth C^r orbifolds \mathcal{O}, \mathcal{P} with \mathcal{O} compact (without boundary). We show that one of these notions is natural and necessary in order to uniquely define the notion of orbifold pullback. For the notion of complete orbifold map, we show that the corresponding set of C^r maps between \mathcal{O} and \mathcal{P} with the C^r topology carries the structure of a smooth C^∞ Banach (r finite)/Fréchet ($r = \infty$) manifold. For the notion of complete reduced orbifold map, the corresponding set of C^r maps between \mathcal{O} and \mathcal{P} with the C^r topology carries the structure of a smooth C^∞ Banach (r finite)/Fréchet ($r = \infty$) orbifold. The remaining two notions carry a stratified structure: The C^r orbifold maps between \mathcal{O} and \mathcal{P} is locally a stratified space with strata modeled on smooth C^∞ Banach (r finite)/Fréchet ($r = \infty$) manifolds while the set of C^r reduced orbifold maps between \mathcal{O} and \mathcal{P} locally has the structure of a stratified space with strata modeled on smooth C^∞ Banach (r finite)/Fréchet ($r = \infty$) orbifolds. Furthermore, we give the explicit relationship between these notions of orbifold map. Applying our results to the special case of orbifold diffeomorphism groups, we show they inherit the structure of C^∞ Banach (r finite)/Fréchet ($r = \infty$) manifolds. In fact, for r finite they are topological groups, and for $r = \infty$ they are convenient Fréchet Lie groups.

CONTENTS

1. Introduction	1
2. Four Notions of Orbifold Map	4
3. Function Space Topologies	10
4. Applications to the Orbifold Diffeomorphism Group	13
5. Why Non-Orbifold Structure Stratifications Arise	14
6. The Tangent Orbifold, Pullbacks and Orbisections	16
7. The exponential map and proof of Theorem 1	22
8. Building Stratified Neighborhoods	25
9. Some Infinite-dimensional Analysis	27
References	30

1. INTRODUCTION

A well-known result in the theory of differentiable dynamical systems states that the set of C^r mappings $C^r(M, N)$ between C^∞ manifolds M and N with M compact

Date: October 26, 2018.

2000 Mathematics Subject Classification. Primary 57S05, 22F50, 54H99; Secondary 22E65.

Key words and phrases. orbifolds, manifolds of mappings, stratified spaces.

has the structure of a C^∞ Banach manifold. If $r = \infty$, $C^\infty(M, N)$ becomes a C^∞ Fréchet manifold. The local model at $f \in C^r(M, N)$ is $\mathcal{D}^r(f^*(TN))$, the space of smooth sections of the pullback tangent bundle $f^*(TN)$ equipped with the C^r topology. $\mathcal{D}^r(f^*(TN))$ is a separable Banach space for $1 \leq r < \infty$ and a separable Fréchet space for $r = \infty$. For reference, see [Eel66], [Pal68], [Nit71], [Fra79], or [Ban97].

We wish to extend these results to the set of C^r maps from a compact smooth orbifold \mathcal{O} (without boundary) to a smooth orbifold \mathcal{P} . Interestingly, there are different useful notions of a C^r map between orbifolds. In [BB02], we defined a notion of (unreduced) C^r orbifold map and the notion of *reduced* orbifold map. In [BB08], we clarified these notions and showed that for a compact orbifold \mathcal{O} (without boundary), both the group $\text{Diff}_{\text{Orb}}^r(\mathcal{O})$ of orbifold diffeomorphisms and the group $\text{Diff}_{\text{red}}^r(\mathcal{O})$ of reduced orbifold diffeomorphisms equipped with the C^r topology, carry the topological structure of a Banach manifold for finite r and Fréchet manifold for $r = \infty$. In fact, we showed that $\text{Diff}_{\text{red}}^r(\mathcal{O})$ is a finite quotient of $\text{Diff}_{\text{Orb}}^r(\mathcal{O})$. While our notion of orbifold map is more general than the one that typically appears in the literature, for example [ALR07], our notion of *reduced* orbifold map agrees with that book's definition 1.3 which is the definition that appears most often.

In order to extend the classical structure result for maps between manifolds to maps between orbifolds, and to generalize our results on the orbifold diffeomorphism group, we will introduce two additional notions of orbifold maps, the *complete orbifold maps* and the *complete reduced orbifold maps*. As simple examples will show, the notion of complete orbifold map is necessary to give a well-defined notion of pullback orbifold bundle. The need to be careful when defining pullback orbifold bundles was already noted in the work of Moerdijk and Pronk [MP97] and Chen and Ruan [CR02].

To reconcile our definitions of orbifold map with the existing literature using Lie groupoid theoretic approach to orbifolds we make the following remarks. Our complete orbifold maps are essentially equivalent to the the groupoid homomorphisms of Moerdijk [Moe02], however, they are independent of any particular groupoid structure one imposes on an orbifold and thus are more natural for the kinds of questions we address here. Moreover, Chen's definition of orbifold map [Che06] agrees with our notion of complete reduced orbifold map up to conjugation.

Lastly, we note that if M and N (as above) are, in addition, Γ -manifolds (Γ , a compact Lie group), then the space $C_\Gamma^r(M, N)$ of C^r equivariant maps from M to N is a closed C^∞ Banach submanifold of $C^r(M, N)$ [Fie70]. In [BB08, Example 3.10], we observed that for a so-called good orbifold $\mathcal{O} = M/\Gamma$, the orbifold diffeomorphism group $\text{Diff}_{\text{Orb}}^r(\mathcal{O})$ is strictly larger than $\text{Diff}_\Gamma^r(M)$, the Γ -equivariant diffeomorphism group of M . The relationship between the space of smooth orbifold maps between good orbifolds $\mathcal{O}_i = M_i/\Gamma$, and the space of equivariant maps $C_\Gamma^r(M_1, M_2)$ will be the focus of a future investigation.

We assume the reader is familiar with the notion of smooth C^r orbifolds, and although there are many nice references for this background material such as the recently published book [ALR07], we will use our previous work [BB08] as our standard reference for notation and needed definitions. We should note, however, that our definition of orbifold is modeled on the definition in Thurston [Thu78] and that the orbifolds that concern us here are referred to as *classical effective orbifolds*

in [ALR07]. More precisely, for our definition of orbifolds, isotropy actions are always effective and we allow for singularities of codimension one. For those notions for which the existing literature is not entirely consistent, we will provide explicit definitions. Our main result is the following

Theorem 1. *Let $r \geq 1$ and let \mathcal{O}, \mathcal{P} be smooth C^r orbifolds (without boundary) with \mathcal{O} compact. Denote by $C_{\star\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ the set of C^r complete orbifold maps between \mathcal{O} and \mathcal{P} equipped with the C^r topology. Let $\star f \in C_{\star\text{Orb}}^r(\mathcal{O}, \mathcal{P})$. Then $C_{\star\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ is a smooth C^∞ manifold modeled locally on the topological vector space $\mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P}))$ of C^r orbisections of the pullback tangent orbibundle of \mathcal{P} equipped with the C^r topology. This separable vector space is a Banach space if $1 \leq r < \infty$ and is a Fréchet space if $r = \infty$.*

As corollaries of theorem 1, we are able to prove the following structure results for our different notions of orbifold map. For the complete reduced orbifold maps we have

Corollary 2. *Let $r \geq 1$ and let \mathcal{O}, \mathcal{P} be as above. Denote by $C_{\blacklozenge\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ the set of complete reduced C^r orbifold maps between \mathcal{O} and \mathcal{P} equipped with the C^r topology inherited from $C_{\star\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ as a quotient space. Then $C_{\blacklozenge\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ carries the structure of a smooth C^∞ Banach (r finite)/Fréchet ($r = \infty$) orbifold.*

This result essentially recovers the result of Chen [Che06] for r finite, where the C^r maps defined there are shown to have the structure of a smooth Banach orbifold. We have the following structure result for orbifold maps.

Corollary 3. *Let $r \geq 1$ and let \mathcal{O}, \mathcal{P} be as above. Denote by $C_{\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ the set of C^r orbifold maps between \mathcal{O} and \mathcal{P} equipped with the C^r topology (as defined in [BB08]). Then $C_{\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ carries the topological structure of a stratified space with strata modeled on smooth C^∞ Banach (r finite)/Fréchet ($r = \infty$) manifolds.*

In section 5, we illustrate this phenomenon with a concrete example. Finally, for the reduced orbifold maps, we conclude

Corollary 4. *Let $r \geq 1$ and let \mathcal{O}, \mathcal{P} be as above. Denote by $C_{\text{red}}^r(\mathcal{O}, \mathcal{P})$ the set of C^r reduced orbifold maps between \mathcal{O} and \mathcal{P} equipped with the C^r topology as a quotient space. Then $C_{\text{red}}^r(\mathcal{O}, \mathcal{P})$ carries the topological structure of a stratified space with strata modeled on smooth C^∞ Banach (r finite)/Fréchet ($r = \infty$) orbifolds.*

We would like to point out in each of the above results we are claiming, in part, the existence of a smooth structure modeled on Banach or Fréchet spaces. While much of the finite dimensional smooth manifold theory carries over to the Banach category, the lack of a general implicit function theorem in Fréchet spaces can cause significant difficulties [Ham82]. In particular, there can be many inequivalent notions of differential calculus [Kel74]. For finite order differentiability, a strong argument can be made that the Lipschitz categories Lip^r are better suited to questions of calculus than the more common C^r category. For our purposes, however, we have chosen to use the C^r category for finite order differentiability and for infinite order differentiability, we use the convenient calculus as detailed in the monographs [FK88, KM97].

The paper is divided into the following sections: Section 2 will define the four notions of orbifold map that we will be considering and how these notions are related. Section 3 defines the C^r topology on $C_{\star\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ with \mathcal{O} compact and

proves corollary 2 assuming theorem 1. Section 4 applies our results to the special case of orbifold diffeomorphisms. Section 5 provides explicit examples to show that non-orbifold structure stratifications naturally arise. Section 6 will construct the pullback orbifold for a smooth complete orbifold map and illustrate the necessity to use complete orbifold maps in order to get a unique notion of pullback. Section 7 recalls some results about the exponential map on orbifolds and contains the proof of theorem 1. Section 8 is devoted to proofs of corollaries 3 and 4. In section 9, we collect the results of infinite-dimensional analysis that we need to substantiate our smoothness claims.

2. FOUR NOTIONS OF ORBIFOLD MAP

We now discuss four related definitions of maps between orbifolds. The first notion we will define is that of a complete orbifold map. It is distinguished from our previous notions of orbifold map and reduced orbifold map [BB08, Section 3] in that we are going to keep track of all defining data. In what follows we use the notation of [BB08, Section 2].

Definition 5. A C^0 complete orbifold map $(f, \{\tilde{f}_x\}, \{\Theta_{f,x}\})$ between locally smooth orbifolds \mathcal{O}_1 and \mathcal{O}_2 consists of the following:

- (1) A continuous map $f : X_{\mathcal{O}_1} \rightarrow X_{\mathcal{O}_2}$ of the underlying topological spaces.
- (2) For each $y \in \mathcal{S}_x$, a group homomorphism $\Theta_{f,y} : \Gamma_{\mathcal{S}_x} \rightarrow \Gamma_{f(y)}$.
- (3) A $\Theta_{f,y}$ -equivariant lift $\tilde{f}_y : \tilde{U}_y \subset \tilde{U}_{\mathcal{S}_x} \rightarrow \tilde{V}_{f(y)}$ where $(\tilde{U}_y, \Gamma_{\mathcal{S}_x}, \rho_y, \phi_y)$ is an orbifold chart at y and $(\tilde{V}_{f(y)}, \Gamma_{f(y)}, \rho_{f(y)}, \phi_{f(y)})$ is an orbifold chart at $f(y)$. That is, the following diagram commutes:

$$\begin{array}{ccc}
 \tilde{U}_y & \xrightarrow{\tilde{f}_y} & \tilde{V}_{f(y)} \\
 \downarrow & & \downarrow \\
 \tilde{U}_y / \Gamma_{\mathcal{S}_x} & \xrightarrow{\tilde{f}_y / \Theta_{f,y}(\Gamma_{\mathcal{S}_x})} & \tilde{V}_{f(y)} / \Theta_{f,y}(\Gamma_{\mathcal{S}_x}) \\
 \downarrow & & \downarrow \\
 U_y \subset U_{\mathcal{S}_x} & \xrightarrow{f} & V_{f(y)}
 \end{array}$$

- (★4) (Equivalence) Two complete orbifold maps $(f, \{\tilde{f}_x\}, \{\Theta_{f,x}\})$ and $(g, \{\tilde{g}_x\}, \{\Theta_{g,x}\})$ are considered equivalent if for each $x \in \mathcal{O}_1$, $f_x = \tilde{g}_x$ as germs and $\Theta_{f,x} = \Theta_{g,x}$. That is, there exists an orbifold chart (\tilde{U}_x, Γ_x) at x such that $\tilde{f}_x|_{\tilde{U}_x} = \tilde{g}_x|_{\tilde{U}_x}$ and $\Theta_{f,x} = \Theta_{g,x}$. Note that this implies that $f = g$.

Definition 6. A complete orbifold map $f : \mathcal{O}_1 \rightarrow \mathcal{O}_2$ of C^r smooth orbifolds is C^r smooth if each of the local lifts \tilde{f}_x may be chosen to be C^r . Given two orbifolds \mathcal{O}_i , $i = 1, 2$, the set of C^r complete orbifold maps from \mathcal{O}_1 to \mathcal{O}_2 will be denoted by $C_{\star\text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2)$.

If we replace (★4) in definition 5 by

- (4) (Equivalence) Two complete orbifold maps $(f, \{\tilde{f}_x\}, \{\Theta_{f,x}\})$ and $(g, \{\tilde{g}_x\}, \{\Theta_{g,x}\})$ are considered equivalent if for each $x \in \mathcal{O}_1$, $\tilde{f}_x = \tilde{g}_x$ as germs. That is, there exists an orbifold chart (\tilde{U}_x, Γ_x) at x such that $\tilde{f}_x|_{\tilde{U}_x} = \tilde{g}_x|_{\tilde{U}_x}$ (which as before implies $f = g$),

where we have dropped the requirement that $\Theta_{f,x} = \Theta_{g,x}$, we recover the notion of *orbifold map* $(f, \{\tilde{f}_x\})$ which appeared in [BB08, Section 3]. Thus, the set of orbifold maps $C_{\text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2)$ can be regarded as the equivalence classes of complete orbifold maps under the less restrictive set-theoretic equivalence (4). The following simple example is illustrative.

Example 7. Let \mathcal{O} be the orbifold \mathbb{R}/\mathbb{Z}_2 where \mathbb{Z}_2 acts on \mathbb{R} via $x \rightarrow -x$ and $f : \mathcal{O} \rightarrow \mathcal{O}$ is the constant map $f \equiv 0$. The underlying topological space $X_{\mathcal{O}}$ of \mathcal{O} is $[0, \infty)$ and the isotropy subgroups are trivial for $x \in (0, \infty)$ and \mathbb{Z}_2 for $x = 0$. The map $\tilde{f}_0 \equiv 0$ is a local equivariant lift of f at $x = 0$ using either of the homomorphisms $\Theta_{f,0} = \text{Id}$ or $\Theta'_{f,0} \equiv e$. Of course, for $x \neq 0$, we set $\tilde{f}_x \equiv 0$ and $\Theta_{f,x} = \Theta'_{f,x} =$ the trivial homomorphism $\Gamma_x = e \mapsto e \in \Gamma_0 = \mathbb{Z}_2$. Thus, as *complete orbifold maps* $(f, \{\tilde{f}_x\}, \{\Theta_{f,x}\}) \neq (f, \{\tilde{f}_x\}, \{\Theta'_{f,x}\})$. However, simply as orbifold maps, they are considered equal.

If we replace (★4) in definition 5 by

- (◆4) (Equivalence) Two complete orbifold maps $(f, \{\tilde{f}_x\}, \{\Theta_{f,x}\})$ and $(g, \{\tilde{g}_x\}, \{\Theta_{g,x}\})$ are considered equivalent if $f = g$ and for each $x \in \mathcal{O}_1$, we have $\Theta_{f,x} = \Theta_{g,x}$,

where we have dropped the requirement that the germs of the lifts \tilde{f}_x and \tilde{g}_x agree, we obtain a new notion of orbifold map $(f, \{\Theta_{f,x}\})$ which we call a *complete reduced orbifold map*. The set of smooth complete reduced orbifold maps will be denoted by $C_{\blacklozenge\text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2)$. As before, it is clear that $C_{\blacklozenge\text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2)$ is a set-theoretic quotient of $C_{\star\text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2)$.

If we replace (4) in the definition of orbifold map, or (◆4) in the definition of complete reduced orbifold map, by

- (●4) (Equivalence) Two orbifold maps $(f, \{\tilde{f}_x\})$ and $(g, \{\tilde{g}_x\})$, (or, complete reduced orbifold maps $(f, \{\Theta_{f,x}\})$ and $(g, \{\Theta_{g,x}\})$) are considered equivalent if $f = g$.

we obtain the notion of *reduced orbifold map* from [BB02]. The set of smooth reduced orbifold maps will be denoted by $C_{\text{red}}^r(\mathcal{O}_1, \mathcal{O}_2)$. Like before, it is clear that $C_{\text{red}}^r(\mathcal{O}_1, \mathcal{O}_2)$ is a set-theoretic quotient of both $C_{\text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2)$ and $C_{\blacklozenge\text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2)$.

Notation. Since we will often need to distinguish between these various notions of orbifold maps, we will denote a complete orbifold map $(f, \{\tilde{f}_x\}, \{\Theta_{f,x}\})$ by $\star f$, and represent an orbifold map $(f, \{\tilde{f}_x\})$ simply by f as in [BB08], a complete reduced orbifold map $(f, \{\Theta_{f,x}\})$ by $\blacklozenge f$, and a reduced orbifold map by $\bullet f$.

Diagrammatically, we have the following:

$$\begin{array}{ccccc}
 & & \star f \in C_{\star\text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2) & & \\
 & \swarrow^{q_\blacklozenge} & \vdots & \searrow^q & \\
 \blacklozenge f \in C_{\blacklozenge\text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2) & & \vdots & & f \in C_{\text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2) \\
 & \searrow^{q_\blacktriangledown} & \vdots & \swarrow^{q_\bullet} & \\
 & & \bullet f \in C_{\text{red}}^r(\mathcal{O}_1, \mathcal{O}_2) & &
 \end{array}$$

where the q 's represent the respective set-theoretic quotient maps. Understanding how these notions are related in the special case of the identity map is crucial in what follows.

Example 8. (Lifts of the Identity Map) Consider the identity map $\text{Id} : \mathcal{O} \rightarrow \mathcal{O}$. Let $x \in \mathcal{O}$ and (\tilde{U}_x, Γ_x) be an orbifold chart at x . From the definition of orbifold map, it follows (since Γ_x is finite) that there exists $\gamma \in \Gamma_x$ such that a lift $\tilde{\text{Id}}_x : \tilde{U}_x \rightarrow \tilde{U}_x$ is given by $\tilde{\text{Id}}_x(\tilde{y}) = \gamma \cdot \tilde{y}$ for all $\tilde{y} \in \tilde{U}_x$. Since $\tilde{\text{Id}}_x$ is $\Theta_{\text{Id},x}$ equivariant we have for $\delta \in \Gamma_x$:

$$\begin{aligned}
 \tilde{\text{Id}}_x(\delta \cdot \tilde{y}) &= \Theta_{\text{Id},x}(\delta) \cdot \tilde{\text{Id}}_x(\tilde{y}) && \text{hence} \\
 \gamma \delta \cdot \tilde{y} &= \Theta_{\text{Id},x}(\delta) \gamma \cdot \tilde{y} && \text{which implies} \\
 & && \text{since } \Gamma_x \text{ acts effectively that} \\
 \gamma \delta &= \Theta_{\text{Id},x}(\delta) \gamma && \text{or, equivalently,} \\
 \Theta_{\text{Id},x}(\delta) &= \gamma \delta \gamma^{-1}
 \end{aligned}$$

Thus, the isomorphism $\Theta_{\text{Id},x}$ is completely determined by the choice of local lift $\tilde{\text{Id}}_x$. This implies that the group $\mathcal{J}\mathcal{D}$ of orbifold maps covering the identity may be regarded as the same as the group $\star\mathcal{J}\mathcal{D}$ of complete orbifold maps covering the identity. That is, we have the bijective correspondence

$$(\text{Id}, \{\tilde{y} \mapsto \gamma_x \cdot \tilde{y}\}, \{\Theta_{\text{Id},x}\}) \longleftrightarrow (\text{Id}, \{\tilde{y} \mapsto \gamma_x \cdot \tilde{y}\}).$$

Suppose now that $\{U_{x_i}\}$ is a countable (possibly finite) cover of \mathcal{O} by charts. Then $\mathcal{J}\mathcal{D}$ can be regarded as a subgroup of the product $\prod \Gamma_{x_i}$ as in the proof of corollary 1.2 in [BB08]. Two inner automorphisms, $\delta \mapsto \gamma_i \delta \gamma_i^{-1}$, give rise to the same automorphism of Γ_x precisely when $\gamma_1 = \zeta \gamma_2$ where $\zeta \in C(\Gamma_x)$, the center of Γ_x . Thus, if we let $C = C(\mathcal{J}\mathcal{D}) \subset \prod C(\Gamma_{x_i})$, then one can see that the complete reduced lifts of the identity $\blacklozenge\mathcal{J}\mathcal{D} \cong \star\mathcal{J}\mathcal{D}/C$, where the free C -action on $\star\mathcal{J}\mathcal{D}$ is defined by

$$(\zeta_i) \cdot (\text{Id}, \{\tilde{y} \mapsto \gamma_{x_i} \cdot \tilde{y}\}, \{\Theta_{\text{Id},x_i}\}) = (\text{Id}, \{\tilde{y} \mapsto (\zeta_i \gamma_{x_i}) \cdot \tilde{y}\}, \{\zeta_i \Theta_{\text{Id},x_i} \zeta_i^{-1} = \Theta_{\text{Id},x_i}\}).$$

Also, note that the correspondence $\star\mathcal{J}\mathcal{D} \leftrightarrow \mathcal{J}\mathcal{D}$ gives an isomorphism $\blacklozenge\mathcal{J}\mathcal{D} \cong \mathcal{J}\mathcal{D}/C$ which in turn is isomorphic to $\text{Inn}(\mathcal{J}\mathcal{D})$, the group of inner automorphisms of $\mathcal{J}\mathcal{D}$. Thus, we have the exact sequence

$$1 \rightarrow C(\mathcal{J}\mathcal{D}) \rightarrow \star\mathcal{J}\mathcal{D} = \mathcal{J}\mathcal{D} \rightarrow \blacklozenge\mathcal{J}\mathcal{D} \rightarrow 1$$

Notation. For a (not necessarily compact) orbifold \mathcal{N} , we will use the notation $\mathcal{J}\mathcal{D}_{\mathcal{N}}$ to denote the group of orbifold lifts of the identity map $\text{Id} : \mathcal{N} \rightarrow \mathcal{N}$. Suppose $f : \mathcal{O}_1 \rightarrow \mathcal{O}_2$ and let the orbifold \mathcal{N} be an open neighborhood of the image $f(\mathcal{O}_1)$.

For an orbifold map $\{\cdot\}f$ (of any type) and $I = (\text{Id}, \{\eta_x \cdot \tilde{y}\}) \in \mathcal{JD}_{\mathcal{N}}$ we can compute $I \circ \{\cdot\}f$. Namely,

$$\begin{aligned} I \circ \star f &= I \circ (f, \{\tilde{f}_x\}, \{\Theta_{f,x}\}) = (f, \{\eta_x \cdot \tilde{f}_x\}, \{\gamma \mapsto \eta_x \Theta_{f,x}(\gamma) \eta_x^{-1}\}) \\ I \circ \blacklozenge f &= I \circ (f, \{\Theta_{f,x}\}) = (f, \{\gamma \mapsto \eta_x \Theta_{f,x}(\gamma) \eta_x^{-1}\}) \\ I \circ f &= I \circ (f, \{\tilde{f}_x\}) = (f, \{\eta_x \cdot \tilde{f}_x\}) \\ I \circ \bullet f &= \bullet f \end{aligned}$$

Suppose $\{\Gamma_x\}$ denotes the family of isotropy groups for an orbifold \mathcal{N} and for subgroups $\Lambda_x \subset \Gamma_x$, let $\{\Lambda_x\}$ denote the corresponding family of subgroups. In what follows, we will use the notation $(\mathcal{JD}_{\mathcal{N}})_{\{\Lambda_x\}}$ for the subgroup of $\mathcal{JD}_{\mathcal{N}}$ defined by

$$\{I \in \mathcal{JD}_{\mathcal{N}} \mid I = (\text{Id}, \{\tilde{y} \mapsto \lambda_x \cdot \tilde{y}\}) \text{ where } \lambda_x \in \Lambda_x \text{ for all } x\}.$$

Lastly, for a fixed orbifold map $\{\cdot\}f$ (of any type), we let $(\mathcal{JD}_{\mathcal{N}}) \cdot \{\cdot\}f$ denote the orbit under the action of $\mathcal{JD}_{\mathcal{N}}$:

$$(\mathcal{JD}_{\mathcal{N}}) \cdot \{\cdot\}f = \{I \circ \{\cdot\}f \mid I \in \mathcal{JD}_{\mathcal{N}}\}$$

and we let $(\mathcal{JD}_{\mathcal{N}})_{\{\cdot\}f}$ denote the corresponding isotropy subgroup of $\{\cdot\}f$ under the action of $\mathcal{JD}_{\mathcal{N}}$:

$$\{I \in \mathcal{JD}_{\mathcal{N}} \mid I \circ \{\cdot\}f = \{\cdot\}f\}.$$

It is also important to note that $\mathcal{JD}_{\mathcal{N}}$ is a finite group in the special case that the source orbifold \mathcal{O}_1 is compact: one may choose the open neighborhood \mathcal{N} of $f(\mathcal{O}_1)$ to be relatively compact and since \mathcal{N} can be covered by finitely many orbifold charts $\{U_{x_i}\}$, the observation that $\mathcal{JD} \subset \prod \Gamma_{x_i}$ from example 8 is enough to show that, in this case, $\mathcal{JD}_{\mathcal{N}}$ is finite.

Implications for the definition of orbifold structure. Recall the following commutative diagram of maps which appears in the definition of a smooth classical effective orbifold [BB08]:

$$\begin{array}{ccc} \tilde{U}_z & \xrightarrow{\tilde{\psi}_{zx}} & \tilde{U}_x \\ \downarrow & & \downarrow \\ U_z \cong \tilde{U}_z/\Gamma_z & \xrightarrow{\psi_{zx}} & \tilde{U}_x/\Gamma_x \cong U_x \end{array}$$

where for a neighborhood $U_z \subset U_x$ with corresponding \tilde{U}_z , and isotropy group Γ_z , there is an open embedding $\tilde{\psi}_{zx} : \tilde{U}_z \rightarrow \tilde{U}_x$ covering the inclusion $\psi_{zx} : U_z \hookrightarrow U_x$ and an injective homomorphism $\theta_{zx} : \Gamma_z \rightarrow \Gamma_x$ so that $\tilde{\psi}_{zx}$ is equivariant with respect to θ_{zx} . For the standard definition of orbifold which appears in the literature, it is understood that $\tilde{\psi}_{zx}$ is defined only up to composition with elements of Γ_x , and θ_{zx} defined only up to conjugation by elements of Γ_x . However, here, we may regard ψ_{zx} as being from any of the notions of orbifold map we have defined, thus giving an orbifold \mathcal{O} , or more precisely, an orbifold atlas for \mathcal{O} , one of four different structures depending on how one keeps track of lifts ψ_{zx} and homomorphisms θ_{zx} . Thus, it makes sense to speak of a *complete orbifold structure* $\star\mathcal{O}$, a *complete reduced orbifold structure* $\blacklozenge\mathcal{O}$, an *orbifold structure* \mathcal{O} , and lastly, a *reduced orbifold structure* $\bullet\mathcal{O}$. Thus, the standard definition of orbifold would correspond to our notion of a reduced orbifold structure. The reader should take care to note that

the term *reduced orbifold* also has been used in the study of so-called noneffective orbifolds [CR02]. Our use of the term *reduced orbifold structure* is unrelated to this. In this paper, the term orbifold will require that the chart maps ψ_{zx} be regarded as orbifold maps in $C_{\text{Orb}}^r(U_z, U_x)$ as defined above. We also point out that there is no fundamental difference between a complete orbifold structure $\star\mathcal{O}$ and an orbifold structure \mathcal{O} and that any reduced orbifold structure $\blacklozenge\mathcal{O}$ is obtained as a quotient of an orbifold structure \mathcal{O} by the action of \mathcal{JD} on orbifold atlases. This follows from example 8 and the fact that any two lifts of ψ_{zx} must differ by a lift of the identity map on U_x . Lastly, we remark that, in general, for an orbifold structure \mathcal{O} , $\tilde{\psi}_{zx} \neq \tilde{\psi}_{yx} \circ \tilde{\psi}_{zy}$ when $U_z \subset U_y \subset U_x$, but there will be an element $\delta \in \Gamma_x$ such that $\delta \cdot \tilde{\psi}_{zx} = \tilde{\psi}_{yx} \circ \tilde{\psi}_{zy}$ and $\delta \cdot \theta_{zx}(\gamma) \cdot \delta^{-1} = \theta_{yx} \circ \theta_{zy}(\gamma)$.

Relationship among the different notions of orbifold map. In this subsection we give a series of lemmas that discuss the relationship among the various notions of orbifold map for a fixed map $f : \mathcal{O}_1 \rightarrow \mathcal{O}_2$. In section 3, we will topologize these sets of mappings and discuss the local structure of these relationships. Our first lemma makes explicit the relationship between the complete reduced orbifold maps and the complete orbifold maps.

Lemma 9. *Let $\star f, \star f' \in C_{\star\text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2)$ be complete orbifold maps which represent the same complete reduced orbifold map. That is, $\blacklozenge f = \blacklozenge f' \in C_{\blacklozenge\text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2)$, so that $\star f = (f, \{\tilde{f}_x\}, \{\Theta_{f,x}\})$ and $\star f' = (f, \{\tilde{f}'_x\}, \{\Theta_{f,x}\})$. Let $C_x = C_{\Gamma_{f(x)}}(\Theta_{f,x}(\Gamma_x))$ denote the centralizer of $\Theta_{f,x}(\Gamma_x)$ in $\Gamma_{f(x)}$. Then there is an orbifold \mathcal{N} which is an open neighborhood of $f(\mathcal{O}_1)$ in \mathcal{O}_2 and an orbifold map $I \in (\mathcal{JD}_{\mathcal{N}})_{\{C_x\}}$ such that $\star f = I \circ \star f'$. Moreover, if the stated condition holds for two complete orbifold maps $\star f$ and $\star f'$, then $\blacklozenge f = \blacklozenge f'$.*

Proof. Since \tilde{f}_x and \tilde{f}'_x are local lifts of the same map f , there exists $\eta_x \in \Gamma_{f(x)}$ such that $\tilde{f}'_x(\tilde{y}) = \eta_x \cdot \tilde{f}_x(\tilde{y})$ for all $\tilde{y} \in \tilde{U}_x$. Thus, for all $\gamma \in \Gamma_x$ we have, on one hand, the equivariance relation $\tilde{f}'_x(\gamma \cdot \tilde{y}) = \Theta_{f,x}(\gamma) \cdot \tilde{f}'_x(\tilde{y})$ while on the other hand, the equivariance relation must be $\tilde{f}'_x(\gamma \cdot \tilde{y}) = \eta_x \cdot \tilde{f}_x(\gamma \cdot \tilde{y}) = \eta_x \Theta_{f,x}(\gamma) \cdot \tilde{f}_x(\tilde{y}) = \eta_x \Theta_{f,x}(\gamma) \eta_x^{-1} \cdot \tilde{f}_x(\tilde{y})$. This implies that $\Theta_{f,x}(\gamma) = \eta_x \Theta_{f,x}(\gamma) \eta_x^{-1}$ and thus $\eta_x \in C_x$. The orbifold \mathcal{N} may be taken to be $\mathcal{N} = \cup_{x \in \mathcal{O}_1} V_{f(x)}$, where $V_{f(x)}$ is an orbifold chart about $f(x) \in \mathcal{O}_2$. We have thus shown the first statement of the lemma, and the last statement is clear from our computation above and the definitions. \square

Example 10. Let \mathcal{O} be as in example 7. Consider the complete orbifold map $\star f = (f, \{\tilde{f}_x\}, \{\Theta_{f,x}\})$ which covers the inclusion map $f : \mathcal{O} \rightarrow \mathcal{O} \times \mathcal{O} \times \mathcal{O}$, $y \mapsto (y, 0, 0)$, where $\tilde{f}_x(\tilde{y}) = (\tilde{y}, 0, 0)$ and $\Theta_{f,0}(\gamma) = (\gamma, e, e) \in \Gamma_{(0,0,0)} = \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ (for $x \neq 0$, $\Theta_{f,x}$ is the trivial homomorphism since $\Gamma_x = \{e\}$). Now, $\eta_0 = (e, \gamma, e) \in C_{\Gamma_{(0,0,0)}}(\Theta_{f,0}(\Gamma_0))$ and $\eta_0 \cdot \tilde{f}_0(\tilde{y}) = \eta_0 \cdot (\tilde{y}, 0, 0) = (\tilde{y}, 0, 0) = \tilde{f}_0(\tilde{y})$. Now let the (finite) group $(\mathcal{JD}_{\mathcal{N}})_{\{C_x\}}$ be as in lemma 9. For fixed $\star f$, let $(\mathcal{JD}_{\mathcal{N}})_{\{C_x\}} \cdot \star f$ denote the orbit of $\star f$. This example shows that the orbit map $(\text{Id}, \{\lambda_x \cdot \tilde{z}\}) \mapsto (\text{Id}, \{\lambda_x \cdot \tilde{z}\}) \circ \star f$ may have nontrivial, (but finite) isotropy. Thus, if we let $(\mathcal{JD}_{\mathcal{N}})_{\star f} \subset (\mathcal{JD}_{\mathcal{N}})_{\{C_x\}}$ denote the isotropy subgroup of $\star f$, then $(\mathcal{JD}_{\mathcal{N}})_{\{C_x\}} / (\mathcal{JD}_{\mathcal{N}})_{\star f} \cong (\mathcal{JD}_{\mathcal{N}})_{\{C_x\}} \cdot \star f$ is a homeomorphism (of discrete sets).

The next lemma describes the relationship between the orbifold maps and the reduced orbifold maps.

Lemma 11. *Let $f, f' \in C_{\text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2)$ be orbifold maps which represent the same reduced orbifold map. That is, $\bullet f = \bullet f' \in C_{\text{red}}^r(\mathcal{O}_1, \mathcal{O}_2)$, so that $f = (f, \{\tilde{f}_x\})$ and $f' = (f, \{\tilde{f}'_x\})$. Then there is an orbifold \mathcal{N} which is an open neighborhood of $f(\mathcal{O}_1)$ in \mathcal{O}_2 and an orbifold map $I = (\text{Id}, \{\eta_x \cdot \tilde{z}\}) \in \mathcal{JD}_{\mathcal{N}}$ with $\eta_x \in \Gamma_{f(x)}$ such that $f = I \circ f'$. Moreover, if the stated condition holds for two orbifold maps f and f' , then $\bullet f = \bullet f'$.*

Proof. \mathcal{N} can be chosen as in lemma 9, and the proof follows from corollary 1.2 in [BB08]. \square

Remark 12. Similar to the situation described in example 10, example 7 shows that the orbit map $\mathcal{JD}_{\mathcal{N}} \cdot f$ may have nontrivial isotropy.

Next, we describe the relationship between the complete orbifold maps and the orbifold maps.

Lemma 13. *Let $\star f, \star f' \in C_{\star\text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2)$ be complete orbifold maps which represent the same orbifold map. That is, $f = f' \in C_{\text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2)$, so that $\star f = (f, \{\tilde{f}_x\}, \{\Theta_{f,x}\})$ and $\star f' = (f, \{\tilde{f}'_x\}, \{\Theta'_{f,x}\})$. Then, for each $x \in \mathcal{O}_1$, $\gamma \in \Gamma_x$, and $\tilde{y} \in \tilde{U}_x$ we have*

$$\left\{ [\Theta'_{f,x}(\gamma)]^{-1} [\Theta_{f,x}(\gamma)] \right\} \cdot \tilde{f}_x(\tilde{y}) = \tilde{f}'_x(\tilde{y}).$$

Moreover, if the stated condition holds for two complete orbifold maps $\star f$ and $\star f'$, then $f = f'$.

Proof. For all $\gamma \in \Gamma_x$ and $\tilde{y} \in \tilde{U}_x$, we have, $\Theta_{f,x}(\gamma) \cdot \tilde{f}_x(\tilde{y}) = \tilde{f}_x(\gamma \cdot \tilde{y}) = \Theta'_{f,x}(\gamma) \cdot \tilde{f}'_x(\tilde{y})$ and the first statement follows. To see the last statement, let $\star f = (f, \{\tilde{f}_x\}, \{\Theta_{f,x}\})$ and $\star f' = (f, \{\tilde{f}'_x\}, \{\Theta'_{f,x}\})$. Note that the condition stated implies that $\tilde{f}_x(\tilde{y}) = \Theta_{f,x}(e) \cdot \tilde{f}_x(\tilde{y}) = \Theta'_{f,x}(e) \cdot \tilde{f}'_x(\tilde{y}) = \tilde{f}'_x(\tilde{y})$. \square

Remark 14. Notice that this relationship is qualitatively different than the relationships described in lemmas 9, 11 and 16, in that it is given as an equality of actions of $\Theta_{f,x}(\Gamma_x)$, $\Theta'_{f,x}(\Gamma_x)$ on the image $\tilde{f}_x(\tilde{U}_x)$ and not as an equality of $\Theta_{f,x}$ and $\Theta'_{f,x}$ as homomorphisms themselves. That is, the representation of $\Theta_{f,x}(\Gamma_x)$ and $\Theta'_{f,x}(\Gamma_x)$ induce actions that when restricted to $\tilde{f}_x(\tilde{U}_x)$ are equal.

Remark 15. Example 7 exhibits the behavior described in lemma 13. A slightly less trivial example is to consider the inclusion map of example 10: $f : \mathcal{O} \rightarrow \mathcal{O} \times \mathcal{O} \times \mathcal{O}$, $y \mapsto (y, 0, 0)$, where $\tilde{f}_x(\tilde{y}) = (\tilde{y}, 0, 0)$. Note that \tilde{f}_0 is equivariant with respect to both $\Theta_{f,0}(\gamma) = (\gamma, e, e)$ and $\Theta'_{f,0}(\gamma) = (\gamma, \gamma, \gamma)$.

The next two lemmas describe the relationship between the complete reduced orbifold maps and the reduced orbifold maps. Given the conclusion of corollary 4, this relationship is necessarily more complicated.

Lemma 16. *Let $\blacklozenge f \in C_{\blacklozenge\text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2)$ be a complete reduced orbifold map and let \mathcal{N} be an open neighborhood of $f(\mathcal{O}_1)$ in \mathcal{O}_2 . Let $I = (\text{Id}, \{\eta_x \cdot \tilde{z}\}) \in \mathcal{JD}_{\mathcal{N}}$ with $\eta_x \in \Gamma_{f(x)}$. If the complete reduced orbifold map $\blacklozenge f' = I \circ \blacklozenge f$, then $\bullet f = \bullet f' \in C_{\text{red}}^r(\mathcal{O}_1, \mathcal{O}_2)$. Furthermore, if $\blacklozenge f'$ is in the orbit $(\mathcal{JD}_{\mathcal{N}}) \cdot \blacklozenge f$, then $\Theta'_{f,x}(\gamma) = \eta_x \Theta_{f,x}(\gamma) \eta_x^{-1}$.*

Proof. Let $\blacklozenge f = (f, \{\Theta_{f,x}\})$ and $\blacklozenge f' = (f', \{\Theta'_{f,x}\})$. Let $\tilde{f}_x, \tilde{f}'_x$ be local lifts equivariant with respect to $\Theta_{f,x}, \Theta'_{f,x}$ respectively. Since $\blacklozenge f' = I \circ \blacklozenge f$, $\tilde{f}'_x(\tilde{y}) = \eta_x \cdot \tilde{f}_x(\tilde{y})$ for all $\tilde{y} \in \tilde{U}_x$, so \tilde{f}_x and \tilde{f}'_x are local lifts of the same map $f = f'$. This implies $\bullet f = \bullet f'$. The last statement follows from the way $\mathcal{J}\mathcal{D}_{\mathcal{N}}$ acts on $\blacklozenge f$. \square

Remark 17. Here, like before, the orbit map $(\text{Id}, \{\eta_x \cdot \tilde{z}\}) \mapsto (\text{Id}, \{\eta_x \cdot \tilde{z}\}) \circ \blacklozenge f$ may have nontrivial, (but finite) isotropy. In fact, $(\mathcal{J}\mathcal{D}_{\mathcal{N}})_{\blacklozenge f} = (\mathcal{J}\mathcal{D}_{\mathcal{N}})_{\{C_x\}}$, the orbifold map lifts of the identity given by elements of $C_{\Gamma_{f(x)}}(\Theta_{f,x}(\Gamma_x))$ described in lemma 9.

In light of lemma 16, we define an equivalence relation the preimage $q_{\blacktriangledown}^{-1}(\bullet f)$:

$$(\dagger) \quad \blacklozenge f = (f, \{\Theta_{f,x}\}) \sim \blacklozenge f' = (f', \{\Theta'_{f,x}\}) \iff \blacklozenge f' = I \circ \blacklozenge f$$

for some $I \in \mathcal{J}\mathcal{D}_{\mathcal{N}}$. That is, for all $x \in \mathcal{O}_1$, there exists $\eta_x \in \Gamma_{f(x)}$ such that $\Theta'_{f,x}(\gamma) = \eta_x \Theta_{f,x}(\gamma) \eta_x^{-1}$ for all $\gamma \in \Gamma_x$. Denote the equivalence class of $\blacklozenge f$ by $[\blacklozenge f]$.

Lemma 18. *Let $[\blacklozenge f = (f, \{\Theta_{f,x}\})] \neq [\blacklozenge f' = (f', \{\Theta'_{f,x}\})]$ be different equivalence classes of complete reduced orbifold maps which represent the same reduced orbifold map. That is, $\bullet f = \bullet f' \in C_{\text{red}}^r(\mathcal{O}_1, \mathcal{O}_2)$. Then there exist local lifts $\{\tilde{f}_x\}$ which are equivariant with respect to both $\{\Theta_{f,x}\}$ and $\{\eta_x \Theta'_{f,x} \eta_x^{-1}\}$ with $\{\Theta_{f,x}\} \neq \{\eta_x \Theta'_{f,x} \eta_x^{-1}\}$ as homomorphisms. However, for each $x \in \mathcal{O}_1$, $\gamma \in \Gamma_x$, and $\tilde{y} \in \tilde{U}_x$ we have as actions on $\tilde{f}_x(\tilde{U}_x)$*

$$(\ddagger) \quad \Theta_{f,x}(\gamma) \cdot \tilde{f}_x(\tilde{y}) = \eta_x \Theta'_{f,x}(\gamma) \eta_x^{-1} \cdot \tilde{f}_x(\tilde{y}).$$

Proof. Since $\bullet f = \bullet f'$, there exists $\eta_x \in \Gamma_{f(x)}$ such that $\tilde{f}_x(\tilde{y}) = \eta_x \cdot \tilde{f}'_x(\tilde{y})$ for all $\tilde{y} \in \tilde{U}_x$. Thus, we conclude that $\tilde{f}_x = \eta_x \cdot \tilde{f}'_x$ is also equivariant with respect to $\eta_x \Theta'_{f,x} \eta_x^{-1}$. Since $[\blacklozenge f] \neq [\blacklozenge f']$, we have $\{\Theta_{f,x}\} \neq \{\eta_x \Theta'_{f,x} \eta_x^{-1}\}$ as homomorphisms. \square

Remark 19. Example 7 illustrates the phenomena dealt with in lemma 18. Lemmas 16 and 18 show that the quotient map $q_{\blacktriangledown} : C_{\blacklozenge \text{Orb}}^r(\mathcal{O}_1, \mathcal{O}_2) \rightarrow C_{\text{red}}^r(\mathcal{O}_1, \mathcal{O}_2)$ factors $q_{\blacktriangledown} = q_{\ddagger} \circ q_{\dagger}$:

$$\begin{array}{ccc} \blacklozenge f & \xrightarrow{q_{\dagger}} & [\blacklozenge f] & \xrightarrow{q_{\ddagger}} & \bullet f \\ & & \searrow & \nearrow & \\ & & & & q_{\blacktriangledown} \end{array}$$

where $q_{\dagger}, q_{\ddagger}$ represent the quotient maps under the equivalences (\dagger) and (\ddagger) , respectively.

3. FUNCTION SPACE TOPOLOGIES

It is easy to define a C^s topology ($1 \leq s \leq r$) on the set of smooth complete orbifold maps $C_{\star \text{Orb}}^r(\mathcal{O}, \mathcal{P})$ with \mathcal{O} compact. Although much of what we do applies to noncompact \mathcal{O} we will assume \mathcal{O} to be compact. As such, implicit in some of the discussion is that \mathcal{O} has been equipped with a finite covering by orbifold charts. The topologies we define have already been shown to be independent of these choices of charts [BB08].

Definition 20. Let $\star f = (f, \{\tilde{f}_x\}, \{\Theta_{f,x}\})$, $\star g = (g, \{\tilde{g}_x\}, \{\Theta_{g,x}\}) \in C_{\star \text{Orb}}^r(\mathcal{O}, \mathcal{P})$. Then a C^s neighborhood of $\star f$ is defined to be

$$\begin{aligned} \mathcal{N}^s(\star f, \varepsilon) &= \{\star g = (g, \{\tilde{g}_x\}, \{\Theta_{g,x}\}) \mid \\ &\quad g \in \mathcal{N}^s(f, \varepsilon) \text{ and } \theta_{f(x)z} \circ \Theta_{f,x} = \theta_{g(x)z} \circ \Theta_{g,x} \text{ for all } x \in \mathcal{O}\} \end{aligned}$$

where $\mathcal{N}^s(f, \varepsilon)$ is the C^s orbifold map neighborhood of f defined in [BB08]. $\theta_{f(x)z} \circ \Theta_{f,x} = \theta_{g(x)z} \circ \Theta_{g,x}$ is to be interpreted as follows: There is a small enough orbifold chart \tilde{U}_x about x , such that the images of both $\tilde{f}_x(\tilde{U}_x)$ and $\tilde{g}_x(\tilde{U}_x)$ are contained in a single orbifold chart \tilde{V}_z and $\theta_{f(x)z} \circ \Theta_{f,x} = \theta_{g(x)z} \circ \Theta_{g,x}$ where $\theta_{f(x)z}, \theta_{g(x)z} : \Gamma_{f(x)}, \Gamma_{g(x)} \hookrightarrow \Gamma_z$ are the injective homomorphisms given in the definition of orbifold. It is important to note that this condition is more than just an isomorphism of groups, but is an equality of their representations as actions on \tilde{V}_z . The collection of sets of this type form a subbasis for the corresponding C^s topology on $C_{\star\text{Orb}}^r(\mathcal{O}, \mathcal{P})$.


Similarly, a C^s neighborhood of $\blacklozenge f = (f, \{\Theta_{f,x}\})$ is defined to be

$$\begin{aligned} \mathcal{N}^s(\blacklozenge f, \varepsilon) = \{ & \blacklozenge g = (g, \{\Theta_{g,x}\}) \in C_{\blacklozenge\text{Orb}}^r(\mathcal{O}, \mathcal{P}) \mid \\ & \bullet g \in \mathcal{N}^s(\bullet f, \varepsilon) \text{ and } \theta_{f(x)z} \circ \Theta_{f,x} = \theta_{g(x)z} \circ \Theta_{g,x} \text{ for all } x \in \mathcal{O} \} \end{aligned}$$

where $\mathcal{N}^s(\bullet f, \varepsilon) = \{\bullet g \in C_{\text{red}}^r(\mathcal{O}, \mathcal{P}) \mid g \in \mathcal{N}^s(I \circ f, \varepsilon) \text{ for some } I \in \mathcal{ID}_{\mathcal{N}}\}$. Here, \mathcal{N} denotes, as usual, an open neighborhood of the image $f(\mathcal{O})$.

Observation. Suppose $\star f = (f, \{\tilde{f}_x\}, \{\Theta_{f,x}\})$ and $\star f' = (f, \{\tilde{f}_x\}, \{\Theta'_{f,x}\})$ are two complete orbifold maps in $C_{\star\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ such that $f = f'$ as orbifold maps. Then $\star f' = (f, \{\tilde{f}_x\}, \{\Theta'_{f,x}\}) \notin \mathcal{N}^s(\star f, \varepsilon)$ for any ε unless $\{\Theta'_{f,x}\} = \{\Theta_{f,x}\}$. Otherwise, it would follow that $\theta_{f(x)z} \circ \Theta_{f,x} = \theta_{f(x)z} \circ \Theta'_{f,x}$, contradicting injectivity of $\theta_{f(x)z}$. Of course the same argument shows that if $g = g'$ as orbifold maps, then $\star g = (g, \{\tilde{g}_x\}, \{\Theta_{g,x}\})$ and $\star g' = (g, \{\tilde{g}_x\}, \{\Theta'_{g,x}\})$ cannot both belong to a neighborhood $\mathcal{N}^s(\star f, \varepsilon)$ unless $\{\Theta_{g,x}\} = \{\Theta'_{g,x}\}$. As a consequence, we see that the preimage $q^{-1}(\mathcal{N}^s(f, \varepsilon)) \subset C_{\star\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ is a (finite) disjoint union of neighborhoods of the form $\mathcal{N}^s(\star f_i, \varepsilon)$ where $\star f_i = (f, \{\tilde{f}_x\}, \{\Theta_{f,x}\}_i)$. Similarly, we see that the preimage $q_{\blacklozenge}^{-1}(\mathcal{N}^s(\bullet f, \varepsilon)) \subset C_{\blacklozenge\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ is a (finite) disjoint union of neighborhoods of the form $\mathcal{N}^s(\blacklozenge f_i, \varepsilon)$ where $\blacklozenge f_i = (f, \{\Theta_{f,x}\}_i)$.

For reference, we have the following diagram of maps:



We now show that the action of identity maps is compatible with the C^s topology on $C_{\star\text{Orb}}^r(\mathcal{O}, \mathcal{P})$. Let $\star f = (f, \{\tilde{f}_x\}, \{\Theta_{f,x}\}) \in C_{\star\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ and let the orbifold \mathcal{N} be an open neighborhood of the image of $f(\mathcal{O})$. Let $\varepsilon > 0$ be chosen so that if $\star g = (g, \{\tilde{g}_x\}, \{\Theta_{g,x}\}) \in \mathcal{N}^s(\star f, \varepsilon)$, then $g(\mathcal{O}) \subset \mathcal{N}$. Let $I \in \mathcal{ID}_{\mathcal{N}}$ be an orbifold map lift of the identity over \mathcal{N} . Then by example 8, I has a representation as $I = (\text{Id}, \{\tilde{w} \mapsto \gamma_z \cdot \tilde{w}\}, \{\delta \mapsto \gamma_z \delta \gamma_z^{-1}\})$.

Lemma 21. *With $\star g \in \mathcal{N}^s(\star f, \varepsilon)$ as above, we have $I \circ \star g \in \mathcal{N}^s(I \circ \star f, \varepsilon)$. Thus, the local action of $\mathcal{J}\mathcal{D}_{\mathcal{N}}$ on a neighborhood of $\star f$ is continuous. In fact, $I : \mathcal{N}^s(\star f, \varepsilon) \rightarrow \mathcal{N}^s(I \circ \star f, \varepsilon)$, $\star g \mapsto I \circ \star g$, is a homeomorphism. In fact, the map I is a C^∞ diffeomorphism.*

Proof. The homeomorphism claim is immediate from the definitions once one realizes that if \tilde{U}_x is chosen as in definition 20, so that $\theta_{f(x)z} \circ \Theta_{f,x} = \theta_{g(x)z} \circ \Theta_{g,x}$, then for $\delta \in \Gamma_x$,

$$\begin{aligned} I \circ \star f &= (f, \{\gamma_z \cdot \tilde{f}_x\}, \{\gamma_z(\theta_{f(x)z} \circ \Theta_{f,x}(\delta))\gamma_z^{-1}\}) \text{ and,} \\ I \circ \star g &= (g, \{\gamma_z \cdot \tilde{g}_x\}, \{\gamma_z(\theta_{g(x)z} \circ \Theta_{g,x}(\delta))\gamma_z^{-1}\}). \end{aligned}$$

The smoothness claims follow from lemma 50 in section 9. \square

Assuming theorem 1, we can now prove corollary 2.

Proof of corollary 2. Let $\blacklozenge f = (f, \{\Theta_{f,x}\}) \in C_{\blacklozenge}^r(\mathcal{O}, \mathcal{P})$. For small enough $\varepsilon > 0$, it is clear that $\mathcal{Q}_\star = q_\blacklozenge^{-1}(\mathcal{N}^r(\blacklozenge f, \varepsilon)) = \bigsqcup \mathcal{N}^r(\star f_i, \varepsilon) \subset C_{\star}^r(\mathcal{O}, \mathcal{P})$, is a finite disjoint union of neighborhoods of the form $\mathcal{N}^r(\star f_i, \varepsilon)$ where $\star f_i = (f, \{\tilde{f}_x\}_i, \{\Theta_{f,x}\})$. If $C_z = C_{\Gamma_z}(\theta_{f(x)z} \circ \Theta_{f,x}(\Gamma_x))$ (which $= C_{\Gamma_z}(\theta_{g(x)z} \circ \Theta_{g,x}(\Gamma_x))$), then lemmas 9 and 21 imply that $(\mathcal{J}\mathcal{D}_{\mathcal{N}})_{\{C_z\}}$ acts smoothly and transitively on fibers of \mathcal{Q}_\star . Example 10 shows that this action is not necessarily free. To understand what happens under these circumstances, suppose that $I \in (\mathcal{J}\mathcal{D}_{\mathcal{N}})_{\{C_z\}}$ fixes $\star f = (f, \{\tilde{f}_x\}, \{\Theta_{f,x}\}) \in q_\blacklozenge^{-1}(\blacklozenge f)$. Let $\star g = (g, \{\tilde{g}_x\}, \{\Theta_{g,x}\}) \in \mathcal{N}^r(\star f, \varepsilon)$. By lemma 21, $I \circ \star g \in \mathcal{N}^r(\star f, \varepsilon)$. This shows that q_\blacklozenge defines an orbifold chart. Since each $\mathcal{N}^r(\star f_i, \varepsilon)$ is an open manifold by theorem 1 and the action of $(\mathcal{J}\mathcal{D}_{\mathcal{N}})_{\{C_z\}}$ is smooth, corollary 2 follows. \square

Proposition 22. *The quotient map $q_\dagger : \mathcal{N}^s(\blacklozenge f, \varepsilon) \rightarrow \mathcal{N}^s([\blacklozenge f], \varepsilon)$, $\blacklozenge f = (f, \{\Theta_{f,x}\}) \mapsto [\blacklozenge f] = (f, \{\Theta_{f,x}\})$ is a local homeomorphism. In fact, it is the quotient map defined by the group action of $\mathcal{J}\mathcal{D}_{\mathcal{N}}$ acting via $\blacklozenge f \mapsto I \circ \blacklozenge f$.*

Proof. It is clear from the definitions that $\mathcal{Q}_\dagger = q_\dagger^{-1}(\mathcal{N}^s([\blacklozenge f], \varepsilon))$ consists of finite disjoint union of neighborhoods of the form $\mathcal{N}^s(\blacklozenge f_i, \varepsilon)$ where $\blacklozenge f_i = (f, \{\eta_{x,i}\Theta_{f,x}\eta_{x,i}^{-1}\})$. The last statement follows by observing that $\mathcal{J}\mathcal{D}_{\mathcal{N}}$ acts transitively on \mathcal{Q}_\dagger and if $\blacklozenge f = I \circ \blacklozenge f$, for $I = (\text{Id}, \{\tilde{y} \rightarrow \eta_x \cdot \tilde{y}\})$, then $I \in (\mathcal{J}\mathcal{D}_{\mathcal{N}})_{\{C_z\}}$ where $C_z = C_{\Gamma_z}(\theta_{f(x)z} \circ \Theta_{f,x}(\Gamma_x))$. Since C_z also is $= C_{\Gamma_z}(\theta_{g(x)z} \circ \Theta_{g,x}(\Gamma_x))$ for any $\blacklozenge g \in \mathcal{N}^s(\blacklozenge f, \varepsilon)$, we see that any such I fixes pointwise the entire neighborhood $\mathcal{N}^s(\blacklozenge f, \varepsilon)$ and the result follows. \square

Later we will have need to refer to the following useful fact about the relation between $\star f$ and maps $\star g \in \mathcal{N}^s(\star f, \varepsilon)$:

Lemma 23. *Let $\star g = (g, \{\tilde{g}_x\}, \{\Theta_{g,x}\}) \in \mathcal{N}^s(\star f, \varepsilon)$. Then for each $x \in \mathcal{O}$, $\Theta_{f,x} = \theta_{g(x)f(x)} \circ \Theta_{g,x} : \Gamma_x \rightarrow \Gamma_{f(x)}$. Moreover, $\tilde{f}_x(\tilde{x})$ and $\tilde{g}_x(\tilde{x})$ both belong to the same connected (closed) stratum $\tilde{V}_{f(x)}^{\Theta_{f,x}(\Gamma_x)} = \{\tilde{y} \in \tilde{V}_{f(x)} \mid \delta \cdot \tilde{y} = \tilde{y} \text{ for all } \delta \in \Theta_{f,x}(\Gamma_x)\}$.*

Proof. In definition 20 we may choose $z = f(x)$. This yields the stated equality of homomorphisms immediately. Recall that $\tilde{\psi}_{g(x)f(x)}$ denotes a lift of the inclusion map $\psi_{g(x)f(x)} : V_{g(x)} \hookrightarrow V_{f(x)}$ given in the definition of orbifold atlas. So, $\tilde{\psi}_{g(x)f(x)} \circ \tilde{g}_x : \tilde{U}_x \rightarrow \tilde{g}_x(\tilde{U}_x) \hookrightarrow \tilde{V}_{f(x)}$ is equivariant relative to $\theta_{g(x)f(x)} \circ \Theta_{g,x}$, which by

hypothesis is the same as $\Theta_{f,x}$. Thus, for each $\gamma \in \Gamma_x$ we have

$$\begin{aligned} \tilde{\psi}_{g(x)f(x)} \circ \tilde{g}_x(\tilde{x}) &= \tilde{\psi}_{g(x)f(x)} \circ \tilde{g}_x(\gamma \cdot \tilde{x}) = (\theta_{g(x)f(x)} \circ \Theta_{g,x}(\gamma)) \cdot \tilde{g}_x(\tilde{x}) \\ &= \Theta_{f,x}(\gamma) \cdot \tilde{g}_x(\tilde{x}) \end{aligned}$$

from which it follows that $\tilde{g}_x(\tilde{x}) \in \tilde{V}_{f(x)}^{\Theta_{f,x}(\Gamma_x)}$. \square

4. APPLICATIONS TO THE ORBIFOLD DIFFEOMORPHISM GROUP

In this section, we show how the discussion of the previous sections applies to orbifold diffeomorphisms. For simplicity, we will continue to assume that the orbifold \mathcal{O} is compact. In [BB08], we studied the group of orbifold diffeomorphisms $\text{Diff}_{\text{Orb}}^r(\mathcal{O})$ and the reduced orbifold diffeomorphisms $\text{Diff}_{\text{red}}^r(\mathcal{O})$ showing that each carried the structure of a (topological) Banach/Fréchet manifold. In fact we expressed $\text{Diff}_{\text{red}}^r(\mathcal{O})$ as the quotient $\text{Diff}_{\text{Orb}}^r(\mathcal{O})/\mathcal{J}\mathcal{D}$, where, of course, $\mathcal{J}\mathcal{D} \subset \text{Diff}_{\text{Orb}}^r(\mathcal{O})$ represents the (finite) group of orbifold map lifts of the identity on \mathcal{O} .

For diffeomorphism groups, it is not hard to see that the group of complete orbifold diffeomorphisms $\text{Diff}_{\star\text{Orb}}^r(\mathcal{O})$ may be regarded as the same as $\text{Diff}_{\text{Orb}}^r(\mathcal{O})$ in much the same way that example 8 illustrated the correspondence $\star\mathcal{J}\mathcal{D} \leftrightarrow \mathcal{J}\mathcal{D}$. This follows from the proof of corollary 1.2 in [BB08], where it is shown that if $f_1, f_2 \in \text{Diff}_{\text{Orb}}^r(\mathcal{O})$ represent the same reduced diffeomorphism $\bullet f \in \text{Diff}_{\text{red}}^r(\mathcal{O})$, then $f_1 \circ f_2^{-1} \in \mathcal{J}\mathcal{D}$. In the diffeomorphism case, one should note that since all homomorphisms $\Theta_{f,x}$ are actually isomorphisms and we assume isotropy groups act effectively, the behavior exhibited in lemmas 13 and 18 cannot occur. There can never be multiple $\Theta_{f,x}$'s corresponding to a particular local lift \tilde{f}_x . Collecting the results of example 8 and lemmas 9, 16 and 21, and exploiting the fact that, in the case of diffeomorphism groups, we have a *global* (C^∞ -) smooth action of $\mathcal{J}\mathcal{D}$, we get the following algebraic and topological structure result.

Theorem 24. *Let \mathcal{O} be a compact smooth C^r orbifold. Then the following sequences are exact:*

$$\begin{aligned} 1 &\longrightarrow \mathcal{J}\mathcal{D} \longrightarrow \text{Diff}_{\text{Orb}}^r(\mathcal{O}) \longrightarrow \text{Diff}_{\text{red}}^r(\mathcal{O}) \longrightarrow 1 \\ 1 &\longrightarrow C(\mathcal{J}\mathcal{D}) \longrightarrow \text{Diff}_{\text{Orb}}^r(\mathcal{O}) \longrightarrow \text{Diff}_{\blacklozenge\text{Orb}}^r(\mathcal{O}) \longrightarrow 1 \\ 1 &\longrightarrow \blacklozenge\mathcal{J}\mathcal{D} = \mathcal{J}\mathcal{D}/C(\mathcal{J}\mathcal{D}) \longrightarrow \text{Diff}_{\blacklozenge\text{Orb}}^r(\mathcal{O}) \longrightarrow \text{Diff}_{\text{red}}^r(\mathcal{O}) \longrightarrow 1 \end{aligned}$$

where $C(\mathcal{J}\mathcal{D})$ denotes the center of $\mathcal{J}\mathcal{D}$. Moreover, each of the diffeomorphism groups $\text{Diff}_{\text{Orb}}^r(\mathcal{O})$, $\text{Diff}_{\blacklozenge\text{Orb}}^r(\mathcal{O})$ and $\text{Diff}_{\text{red}}^r(\mathcal{O})$ carries the structure of a smooth C^∞ Banach ($r < \infty$)/Fréchet ($r = \infty$) manifold.

Theorem 25. *Each of the diffeomorphism groups $\text{Diff}_{\text{Orb}}^r(\mathcal{O})$, $\text{Diff}_{\blacklozenge\text{Orb}}^r(\mathcal{O})$ and $\text{Diff}_{\text{red}}^r(\mathcal{O})$ is a topological group. That is, composition and inversion are continuous. Furthermore, when $r = \infty$, $\text{Diff}_{\text{Orb}}^\infty(\mathcal{O})$, $\text{Diff}_{\blacklozenge\text{Orb}}^\infty(\mathcal{O})$ and $\text{Diff}_{\text{red}}^\infty(\mathcal{O})$ are convenient Fréchet Lie groups.*

Proof. For $0 < r < \infty$, the group multiplication $\mu(f, g) = f \circ g$ and inversion $\text{inv}(f) = f^{-1}$ in the diffeomorphism group $\text{Diff}_{\text{Orb}}^r(\mathcal{O})$, corresponds to composition and inversion of the C^r local equivariant lifts. These operations are known only to be C^0 . This follows by the so-called Ω -lemma of Palais [Pal68], a suitable version of which is stated as lemma 52 for completeness. Thus, $\text{Diff}_{\text{Orb}}^r(\mathcal{O})$ is a

topological group. The structure result of theorem 24 then yields the topological group structure for $\text{Diff}_{\blacklozenge\text{Orb}}^r(\mathcal{O})$ and $\text{Diff}_{\text{red}}^r(\mathcal{O})$.

For $r = \infty$, by lemma 51, group multiplication $\mu(f, g) = f \circ g$ is smooth since $\text{Diff}_{\text{Orb}}^\infty(\mathcal{O})$ is an open submanifold of $C_{\text{Orb}}^\infty(\mathcal{O}, \mathcal{O})$ by [BB08, section 7]. To show that inversion is smooth, we use the argument given in [KM97, Theorem 43.1]. Let $c = (c^t, \{\tilde{c}_x^t\}) : \mathbb{R} \rightarrow \text{Diff}_{\text{Orb}}^\infty(\mathcal{O}) \subset C_{\text{Orb}}^\infty(\mathcal{O}, \mathcal{O})$ be a smooth curve. Then, in a local orbifold chart by corollary 49, the mapping $\tilde{c}_x^\wedge : (0, 1) \times \tilde{U}_x \rightarrow \tilde{V}_z$ is smooth and $(\text{inv} \circ \tilde{c}_x)^\wedge$ satisfies the finite dimensional implicit equation $\tilde{c}_x^\wedge(t, (\text{inv} \circ \tilde{c}_x)^\wedge(t, \tilde{y})) = \tilde{y}$ for all $t \in \mathbb{R}$ and $\tilde{y} \in \tilde{U}_x$. By the finite dimensional implicit function theorem, $(\text{inv} \circ \tilde{c}_x)^\wedge$ is smooth in (t, \tilde{y}) . Hence, by corollary 49, inv maps smooth curves to smooth curves and is thus smooth. This shows that $\text{Diff}_{\text{Orb}}^\infty(\mathcal{O})$ is a convenient Fréchet Lie group and thus, by the structure results of theorem 24, so are $\text{Diff}_{\blacklozenge\text{Orb}}^\infty(\mathcal{O})$ and $\text{Diff}_{\text{red}}^\infty(\mathcal{O})$. \square

5. WHY NON-ORBIFOLD STRUCTURE STRATIFICATIONS ARISE

In this section, we wish to give an example on why non-orbifold structure stratifications arise in the topological structure of our orbifold maps. We first recall a definition of stratification in the infinite-dimensional setting. We will use the definition found in [Fis70] or [Bou75] for infinite-dimensional stratifications although we do not need the full generality presented in these references. In our case, each point with a stratified neighborhood has only a finite number of strata coming together.

Definition 26 ([Fis70, Bou75]). Let X be a topological space and \mathcal{A} a countable set with partial order \prec . A *partition* of X is a collection of non-empty pairwise disjoint subspaces $\{X_\alpha\}$ indexed by \mathcal{A} such that $X = \cup_{\alpha \in \mathcal{A}} X_\alpha$. A partition $\{X_\alpha\}_{\alpha \in \mathcal{A}}$ is a *stratification* of X if

- (1) each X_α is a submanifold when given the topology induced by X and,
- (2) $X_\alpha \cap \overline{X_\beta} \neq \emptyset$, $\alpha \neq \beta$, then $\beta \prec \alpha$ and $X_\alpha \subset \overline{X_\beta}$.

The X_α are called the *strata* of the stratification and may have many connected components. Moreover, condition (2) implies that $\overline{X_\beta} - X_\beta \subset \cup_{\alpha \succ \beta} X_\alpha$.

Before we show how these stratifications arise, we first present a simple example to help motivate the discussion.

Example 27. Consider the situation described in Example 7: \mathcal{O} is the orbifold \mathbb{R}/\mathbb{Z}_2 where \mathbb{Z}_2 acts on \mathbb{R} via $x \rightarrow -x$ and $f : \mathcal{O} \rightarrow \mathcal{O}$ is the constant map $f \equiv 0$. The map $\tilde{f}_0 \equiv 0$ is a local equivariant lift of f at $x = 0$ using either of the homomorphisms $\Theta_{f,0} = \text{Id}$ or $\Theta'_{f,0} \equiv e$. Of course, for $x \neq 0$, we set $\tilde{f}_x \equiv 0$ and $\Theta_{f,x} = \Theta'_{f,x} =$ the trivial homomorphism $\Gamma_x = e \mapsto e \in \Gamma_0 = \mathbb{Z}_2$. Thus, we have two complete orbifold maps $\star f = (f, \{\tilde{f}_x\}, \{\Theta_{f,x}\})$ and $\star f' = (f, \{\tilde{f}_x\}, \{\Theta'_{f,x}\})$ which cover the same orbifold map $f = (f, \{\tilde{f}_x\})$.

We need to first compute $\mathcal{N}^r(\star f, \varepsilon)$. We will do this in detail since this is the first time we have done an explicit computation of this type. Using definition 20 and the notation there, let $\star g \in \mathcal{N}^r(\star f, \varepsilon)$. For all $x \in \mathcal{O}$ we may choose $z = 0$ and thus $\tilde{V}_z = \tilde{V}_0$ may be chosen to be the interval $(-\varepsilon, \varepsilon)$ as a chart about 0 in the target. There are two cases to consider: $x = 0$ and $x \neq 0$. For $x = 0$, let \tilde{U}_0 be any orbifold chart about 0. It follows that the local lift \tilde{g}_0 over $x = 0$ must take $0 \in \tilde{U}_0$ to $0 \in \tilde{V}_0$. To see this, suppose to the contrary that $\tilde{g}_0(0) = \tilde{y} \neq 0$. By

definition 20, we must have the following equality of homomorphisms from $\mathbb{Z}_2 = \Gamma_0$ to Γ_0 :

$$\begin{aligned} \theta_{f(0)0} \circ \Theta_{f,0} &= \theta_{g(0)0} \circ \Theta_{g,0} \iff \\ \text{Id} &= \theta_{y_0} \circ \Theta_{g,0} \end{aligned}$$

However, $\Theta_{g,0} : \Gamma_0 \rightarrow \Gamma_y = \{e\}$ has nontrivial kernel which contradicts the last line above. We thus may conclude that for $x = 0$, $\tilde{g}_0(0) = 0$ and $\Theta_{g,0} = \Theta_{f,0} = \text{Id}$. From $\Theta_{g,0} = \text{Id}$, it follows that the local lift \tilde{g}_0 must be an odd function. For $x \neq 0$, there is no restriction on \tilde{g}_x arising from equivariance since $\Gamma_x = \{e\}$ and $\theta_{f(x)0} \circ \Theta_{f,x} = \theta_{g(x)0} \circ \Theta_{g,x} : \Gamma_x \rightarrow \Gamma_0$ will always be the trivial homomorphism $e \mapsto e$. Putting this all together we have shown that

$$q(\mathcal{N}^r(\star f, \varepsilon)) = \{g \in C_{\text{Orb}}^r(\mathcal{O}) \mid \|\tilde{g}_x\| < \varepsilon \text{ and } \tilde{g}_0 \text{ is an odd function}\}.$$

We now use a similar argument to compute $\mathcal{N}^r(\star f', \varepsilon)$. Let $\star g' \in \mathcal{N}^r(\star f', \varepsilon)$. For $x \neq 0$, $\Theta'_{f',x} = \Theta_{f,x}$, so we conclude as above that there is no restriction on \tilde{g}'_x arising from equivariance. On the other hand, for $x = 0$ we must have the equality of homomorphisms $\theta_{f(0)0} \circ \Theta'_{f',0} = \theta_{g'(0)0} \circ \Theta_{g',0} : \Gamma_0 \rightarrow \Gamma_0$. Since $\Theta'_{f',0} \equiv e$, injectivity of $\theta_{g'(0)0}$ implies that $\Theta_{g',0} \equiv e$. Thus, there is no restriction on \tilde{g}'_0 arising from equivariance either and we can conclude that

$$q(\mathcal{N}^r(\star f', \varepsilon)) = \{g' \in C_{\text{Orb}}^r(\mathcal{O}) \mid \|\tilde{g}'_x\| < \varepsilon\}.$$

Here, it is clear that $q(\mathcal{N}^r(\star f, \varepsilon))$ is a proper subset (later, a submanifold) of $q(\mathcal{N}^r(\star f', \varepsilon))$, and that any orbifold map $g \in \mathcal{N}^r(f, \varepsilon)$ must be in $q(\mathcal{N}^r(\star f', \varepsilon))$ so that the topological structure of a neighborhood of f is completely determined from only an understanding of the topological structure of $q(\mathcal{N}^r(\star f', \varepsilon))$ which in turn is determined by the structure of $\mathcal{N}^r(\star f', \varepsilon)$, which will be shown to be a manifold.

Unfortunately, in general, the topological structure of a neighborhood of an orbifold map f is rarely determined completely by the topological structure of a *single* neighborhood of one of its complete orbifold lifts $q^{-1}(f)$. This is illustrated in the next example.

Example 28. Let $\mathcal{O} = \mathbb{R}/\mathbb{Z}_2$ with \mathbb{Z}_2 acting with generator α , where $\alpha \cdot x = -x$ as above. Let $\mathcal{P} = \mathbb{R}^3/(\mathbb{Z}_2 \times \mathbb{Z}_2)$ where $\mathbb{Z}_2 \times \mathbb{Z}_2 = \langle j, k \mid j^2 = k^2 = 1 = [j, k] \rangle$ with the action defined by $j \cdot (x, y, z) = (-x, y, -z)$ and $k \cdot (x, y, z) = (-x, -y, z)$. Consider $\bullet f \in C_{\text{red}}^r(\mathcal{O}, \mathcal{P})$ defined by $\bullet f(y_1) = (y_1, 0, 0)$ and choose the orbifold map $f \in q_{\bullet}^{-1}(\bullet f)$ given by $f = (f, \{\tilde{y}_1 \mapsto (\tilde{y}_1, 0, 0)\})$. That is, for each $x \in \mathcal{O}$, the local lift $\tilde{f}_x(\tilde{y}_1) = (\tilde{y}_1, 0, 0)$ on \tilde{U}_x . Since Γ_x is trivial when $x \neq 0$ and $\Gamma_0 = \mathbb{Z}_2$, there are precisely two complete maps in $q^{-1}(f)$:

$$\begin{aligned} \star f &= (f, \{\tilde{y}_1 \mapsto (\tilde{y}_1, 0, 0)\}, \{\Theta_{f,0} : \alpha \mapsto j\}) \\ \star f' &= (f, \{\tilde{y}_1 \mapsto (\tilde{y}_1, 0, 0)\}, \{\Theta'_{f,0} : \alpha \mapsto k\}). \end{aligned}$$

Note that since $\Theta_{f,x}$ is the trivial homomorphism $e \mapsto e$ for all $x \neq 0$, we have only indicated the two possible homomorphisms at $x = 0$, namely, $\Theta_{f,0}, \Theta'_{f,0} : \Gamma_0 = \mathbb{Z}_2 \rightarrow \Gamma_{(0,0,0)} = \mathbb{Z}_2 \times \mathbb{Z}_2$.

We will proceed as in example 27 and first compute $\mathcal{N}^r(\star f, \varepsilon)$. Let $\star g \in \mathcal{N}^r(\star f, \varepsilon)$. Then $\star g$ has a representation

$$\star g = (g, \{\tilde{g}_x = (\tilde{y}_1 + (\tilde{g}_1)_x(\tilde{y}_1), (\tilde{g}_2)_x(\tilde{y}_1), (\tilde{g}_3)_x(\tilde{y}_1))\}, \{\Theta_{g,x}\}).$$

For $x \neq 0$ we have $\Gamma_x = \{e\}$ so, like before, there is no restriction on $(\tilde{g}_i)_x, i = 1, 2, 3$ arising from equivariance. Thus, we focus on lifts $(\tilde{g}_i)_0$ over a chart \tilde{U}_0 about $x = 0$. We may assume that $\tilde{V}_z = \tilde{V}_0$ where we have shortened the subscript $(0, 0, 0)$ to $\vec{0} \in \mathbb{R}^3$. We will continue to do this for the remainder of this example.

By lemma 23, $\tilde{g}_0(0) \in \tilde{V}_0^{\{e, j\}} = y$ -axis and $\Theta_{g,0} : \Gamma_0 \rightarrow \Gamma_{g(0)}$ is $\alpha \mapsto j$. We now compute

$$\tilde{g}_0(\alpha \cdot \tilde{y}_1) = \tilde{g}_0(-y_1) = (-\tilde{y}_1 + (\tilde{g}_1)_0(-\tilde{y}_1), (\tilde{g}_2)_0(-\tilde{y}_1), (\tilde{g}_3)_0(-\tilde{y}_1)).$$

On the other hand,

$$\begin{aligned} \tilde{g}_0(\alpha \cdot \tilde{y}_1) &= \Theta_{g,0}(\alpha) \cdot \tilde{g}_0(y_1) = j \cdot \tilde{g}_0(y_1) \\ &= (-\tilde{y}_1 - (\tilde{g}_1)_0(\tilde{y}_1), (\tilde{g}_2)_0(\tilde{y}_1), -(\tilde{g}_3)_0(\tilde{y}_1)). \end{aligned}$$

Thus,

$$\begin{aligned} q(\mathcal{N}^r(\star f, \varepsilon)) &= \{g \in C_{\text{Orb}}^r(\mathcal{O}, \mathcal{P}) \mid \\ &\|\tilde{g}_x - \tilde{f}_x\| < \varepsilon \text{ with } (\tilde{g}_1)_0, (\tilde{g}_3)_0 \text{ odd functions and } (\tilde{g}_2)_0 \text{ an even function}\}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} q(\mathcal{N}^r(\star f', \varepsilon)) &= \{g' \in C_{\text{Orb}}^r(\mathcal{O}, \mathcal{P}) \mid \\ &\|\tilde{g}'_x - \tilde{f}_x\| < \varepsilon \text{ with } (\tilde{g}'_1)_0, (\tilde{g}'_2)_0 \text{ odd functions and } (\tilde{g}'_3)_0 \text{ an even function}\}. \end{aligned}$$

Thus, the corresponding neighborhood of the orbifold map f is the union of two sets $\mathcal{N}^r(f, \varepsilon) = q(\mathcal{N}^r(\star f, \varepsilon)) \cup q(\mathcal{N}^r(\star f', \varepsilon))$ each of which will later be shown to carry a Banach/Fréchet manifold structure. Their intersection is along the submanifold

$$\begin{aligned} \mathcal{H} &= q(\mathcal{N}^r(\star f, \varepsilon)) \cap q(\mathcal{N}^r(\star f', \varepsilon)) = \\ &\{h \in C_{\text{Orb}}^r(\mathcal{O}, \mathcal{P}) \mid \|\tilde{h}_x - \tilde{f}_x\| < \varepsilon \text{ with } \tilde{h}_0(\tilde{y}_1) = (\tilde{y}_1 + (\tilde{h}_1)_0(\tilde{y}_1), 0, 0) \\ &\text{where } (\tilde{h}_1)_0 \text{ is an odd function}\}. \end{aligned}$$

Thus, the neighborhood $\mathcal{N}^r(f, \varepsilon)$ has a stratified structure (see figure 1): Just let $\mathcal{A} = \{\alpha, \beta, \gamma\}$ with partial order $\beta \prec \alpha, \gamma \prec \alpha$ and define $X = \mathcal{N}^r(f, \varepsilon)$, $X_\alpha = \mathcal{H}$, $X_\beta = q(\mathcal{N}^r(\star f, \varepsilon)) - \mathcal{H}$, and $X_\gamma = q(\mathcal{N}^r(\star f', \varepsilon)) - \mathcal{H}$. Moreover, since $\mathcal{N}^r(f, \varepsilon) - \mathcal{H}$ is not connected we see that this stratified structure is not that of an orbifold structure as removal of the singular set of an orbifold never disconnects a connected component of the orbifold [Bor92, Bor93]. Furthermore, if we let \mathcal{N} denote an open neighborhood of the image $f(\mathcal{O})$, then from [BB08] a neighborhood of the reduced orbifold map $\bullet f$ is given by $\mathcal{N}^r(\bullet f, \varepsilon) = \mathcal{N}^r(f, \varepsilon)/\mathcal{J}\mathcal{D}_{\mathcal{N}}$ where $\mathcal{J}\mathcal{D}_{\mathcal{N}}$ acts in such a way that the quotient map restricts on each stratum to give a smooth orbifold chart (see proof of corollary 4 which appears at the end of section 8). Thus, $\mathcal{N}^r(\bullet f, \varepsilon)$ has a non-orbifold structure stratification also.

6. THE TANGENT ORBIBUNDLE, PULLBACKS AND ORBISECTIONS

The tangent orbibundle. We recall the definition of the tangent orbibundle of a smooth C^{r+1} orbifold.

Definition 29. Let \mathcal{O} be an n -dimensional C^{r+1} orbifold. The *tangent orbibundle* of \mathcal{O} , $p : T\mathcal{O} \rightarrow \mathcal{O}$, is the C^r orbibundle defined as follows. If (\tilde{U}_x, Γ_x) is an orbifold chart around $x \in \mathcal{O}$ then $p^{-1}(U_x) \cong (\tilde{U}_x \times \mathbb{R}^n)/\Gamma_x$ where Γ_x acts on $\tilde{U}_x \times \mathbb{R}^n$ via




FIGURE 1. A stratified neighborhood

$\gamma \cdot (\tilde{y}, \tilde{v}) = (\gamma \cdot \tilde{y}, d\gamma_{\tilde{y}}(\tilde{v}))$. In keeping with tradition, we denote the fiber $p^{-1}(x)$ over $x \in U_x$ by $T_x\mathcal{O} \cong \mathbb{R}^n/\Gamma_x$. Note that, in general, if Γ_x is non-trivial then $T_x\mathcal{O}$ will be a convex cone rather than a vector space. Locally we have the diagram:

$$\begin{array}{ccc} T\tilde{U}_x \cong \tilde{U}_x \times \mathbb{R}^n & \xrightarrow{\Pi_x} & (\tilde{U}_x \times \mathbb{R}^n)/\Gamma_x \\ \text{pr}_1 \downarrow & & \downarrow p \\ \tilde{U}_x & \xrightarrow{\pi_x} & U_x \end{array}$$

where $\text{pr}_1 : \tilde{U}_x \times \mathbb{R}^n \rightarrow \tilde{U}_x$ denotes the projection onto the first factor $(\tilde{y}, \tilde{v}) \mapsto \tilde{y}$ (which is a specific choice of lift of p).

Pulling back an orbundle. The definition of the pullback of an orbundle depends crucially on the notion of orbifold map. In simple examples, we will see that a unique notion of pullback exists only when using complete orbifold maps. On the other hand, we will see that once one has a pullback bundle defined via a complete orbifold map $*f$, there is no difference between the notion of an orbisection and a complete orbisection. Not surprisingly, if one tries to define a useful notion of reduced or complete reduced orbisection one loses the vector space structure on the space of such sections. As in the case of the tangent orbifold, the pullback bundle will be an example of the more general notion of a linear orbundle given in [BB02].

Definition 30. Let \mathcal{O}, \mathcal{P} be C^{r+1} orbifolds of dimension n and m , respectively. Given $*f \in C_{*Orb}^{r+1}(\mathcal{O}, \mathcal{P})$ we define the pullback of the tangent orbifold to \mathcal{P} by $*f$, $*f^*(T\mathcal{P})$ as follows: Let $*f = (f, \{\tilde{f}_x\}, \{\Theta_{f,x}\})$ and let \tilde{U}_x and $\tilde{V}_{f(x)}$ be orbifold charts about $x \in \mathcal{O}$ and $f(x) \in \mathcal{P}$ respectively. Define the pullback $*f^*(T\mathcal{P})$ to be the orbifold with charts of the form (a fibered product)

$$\tilde{U}_x \times_{\tilde{V}_{f(x)}} T\tilde{V}_{f(x)} = \{(\tilde{y}, \tilde{\xi}) \in \tilde{U}_x \times T\tilde{V}_{f(x)} \mid \tilde{f}_x(\tilde{y}) = \text{pr}_1(\tilde{\xi})\}$$

where $\tilde{f}_x : \tilde{U}_x \rightarrow \tilde{V}_{f(x)}$ and $\text{pr}_1 : T\tilde{V}_{f(x)} \rightarrow \tilde{V}_{f(x)}$ is the tangent bundle projection. If we write $\tilde{\xi} = [\tilde{f}_x(\tilde{y}), \tilde{v}] \in T\tilde{V}_{f(x)} = \tilde{V}_{f(x)} \times \mathbb{R}^m$, the action of Γ_x is specified in

local coordinates by:

$$\begin{aligned} \gamma \cdot (\tilde{y}, \tilde{\xi}) &= (\gamma \cdot \tilde{y}, \Theta_{f,x}(\gamma) \cdot \tilde{\xi}) = (\gamma \cdot \tilde{y}, [\Theta_{f,x}(\gamma) \cdot \tilde{f}_x(\tilde{y}), d(\Theta_{f,x}(\gamma))_{\tilde{f}_x(\tilde{y})} \cdot \tilde{v}]) \\ &= (\gamma \cdot \tilde{y}, [\tilde{f}_x(\gamma \cdot \tilde{y}), d(\Theta_{f,x}(\gamma))_{\tilde{f}_x(\tilde{y})} \cdot \tilde{v}]) \in \tilde{U}_x \times_{\tilde{V}_{f(x)}} T\tilde{V}_{f(x)} \end{aligned}$$

where $\tilde{y} \in \tilde{U}_x$ and $\tilde{v} \in \text{pr}_1^{-1}(\tilde{f}_x(\tilde{y}))$. Also, we let $\text{pr}_2 : T\tilde{V}_{f(x)} \cong \tilde{V}_{f(x)} \times \mathbb{R}^m \rightarrow \mathbb{R}^m$, $\tilde{\xi} \mapsto \tilde{v}$ be the fiber projection. This gives $\star f^*(T\mathcal{P})$ the structure of a smooth C^r m -dimensional linear orbifold over \mathcal{O} . In an abuse of notation, $p : \star f^*(T\mathcal{P}) \rightarrow \mathcal{O}$ will denote the orbifold projection. Denote the fiber over x , by $p^{-1}(x) = \star f^*(T\mathcal{P})_x$. In local coordinates, we have the diagram (all vertical arrows are quotient maps by respective group actions):

$$\begin{array}{ccc} & \tilde{U}_x \times_{\tilde{V}_{f(x)}} T\tilde{V}_{f(x)} & \xrightarrow{\tilde{F}_x} & T\tilde{V}_{f(x)} \cong (\tilde{V}_{f(x)} \times \mathbb{R}^m) \\ & \swarrow \text{pr}_1 & & \swarrow \text{pr}_1 \\ \tilde{U}_x & \xrightarrow{\tilde{f}_x} & \tilde{V}_{f(x)} & \xrightarrow{\Pi_{\Theta_{f,x}}} & T\tilde{V}_{f(x)}/\Theta_{f,x}(\Gamma_x) \\ & \downarrow \Pi_x & & \downarrow \Pi_{f(x)} & \\ & \star f^*(TV_{f(x)}) \cong (\tilde{U}_x \times_{\tilde{V}_{f(x)}} T\tilde{V}_{f(x)})/\Gamma_x & \xrightarrow{F} & TV_{f(x)}/\Gamma_{f(x)} & \\ & \downarrow \pi_x & & \downarrow \pi_{f(x)} & \\ U_x & \xrightarrow{p} & V_{f(x)} & \xrightarrow{p} & \\ & \downarrow p & & \downarrow p & \\ & U_x & \xrightarrow{f} & V_{f(x)} & \end{array}$$

The map $\tilde{F}_x : \tilde{U}_x \times_{\tilde{V}_{f(x)}} T\tilde{V}_{f(x)} \rightarrow T\tilde{V}_{f(x)}$ given by $\tilde{F}_x(\tilde{y}, \tilde{\xi}) = \tilde{\xi}$ induces a map $F : \star f^*(TV_{f(x)}) \rightarrow TV_{f(x)}/\Theta_{f,x}(\Gamma_x)$ defined by $F(y, \xi) = \Pi_{\Theta_{f,x}} \circ \tilde{F}_x \circ \Pi_x^{-1}(y, \xi)$. This is well defined since, for any $\gamma \in \Gamma_x$, $\tilde{F}_x(\gamma \cdot (\tilde{y}, \tilde{\xi})) = \tilde{F}_x(\gamma \cdot \tilde{y}, \Theta_{f,x}(\gamma) \cdot \tilde{\xi}) = \Theta_{f,x}(\gamma) \cdot \tilde{\xi}$ and $\Pi_{\Theta_{f,x}}(\Theta_{f,x}(\gamma) \cdot \tilde{\xi}) = \Pi_{\Theta_{f,x}}(\tilde{\xi})$.

Note that the pullback is defined only if we have all the information contained in both the choices of local lifts $\{\tilde{f}_x\}$ and the choices of the homomorphisms $\Theta_{f,x} \in \text{Hom}(\Gamma_x, \Gamma_{f(x)})$. That is, all of the information of a complete orbifold map is used. As an illustration of the necessity for needing to use complete orbifold maps to define pullbacks we give two examples. The first example shows that, unless complete orbifold maps are used, the pullback orbifold is not well-defined even up to a reasonable notion of equivalence.

Example 31. Consider the situation from example 27: $\mathcal{O} = \mathcal{P}$ is the orbifold \mathbb{R}/\mathbb{Z}_2 where \mathbb{Z}_2 acts on \mathbb{R} via $x \rightarrow -x$ and $f : \mathcal{O} \rightarrow \mathcal{P}$ is the constant map $f \equiv 0$. Note that $T\mathcal{O} = T\mathcal{P} = \mathbb{R}^2/\mathbb{Z}_2$ where the generator α of \mathbb{Z}_2 acts via $(x, y) \mapsto (-x, -y)$ and the bundle projection p is just projection onto the first factor. We note that for $x \neq 0$, $p^{-1}(x) = \mathbb{R}$ and that $p^{-1}(0) = \mathbb{R}/\mathbb{Z}_2$. Let $\star f$ and $\star f'$ be the two complete

orbifold maps from example 27 which cover the orbifold map $f = (f, \{\tilde{f}_x\})$. Then we claim that

$${}_*\mathbf{f}^*(T\mathcal{P}) \cong T\mathcal{O}$$

while

$${}_*\mathbf{f}'^*(T\mathcal{P}) \cong \mathcal{O} \times \mathbb{R}$$

To see this we work in local coordinates: Since $\tilde{f}_x \equiv 0$, we may take \tilde{V}_0 as a chart about $f(x)$ for all x . Thus for each x ,

$$\begin{aligned} \tilde{U}_x \times_{\tilde{V}_0} T\tilde{V}_0 &= \{(\tilde{y}, \tilde{\xi}) \in \tilde{U}_x \times T\tilde{V}_0 \mid 0 = \text{pr}_1(\tilde{\xi})\} \\ &= \tilde{U}_x \times T_0\tilde{V}_0 \cong \tilde{U}_x \times \mathbb{R} \end{aligned}$$

Now for $x \neq 0$, $\Gamma_x = \{e\}$ and so the action of Γ_x on $\tilde{U}_x \times_{\tilde{V}_0} T\tilde{V}_0$ is necessarily trivial. If we denote the orbundle projections $p : {}_*\mathbf{f}^*(T\mathcal{P}) \rightarrow \mathcal{O}$ and $p' : {}_*\mathbf{f}'^*(T\mathcal{P}) \rightarrow \mathcal{O}$, then $p^{-1}(U_x) = p'^{-1}(U_x) \cong \tilde{U}_x \times \mathbb{R}$. On the other hand, for $x = 0$, since $\Theta_{f,0}(\alpha) = \alpha$ and $\Theta'_{f,0}(\alpha) = e$ we see that

$$\begin{aligned} p^{-1}(U_0) &\cong (\tilde{U}_0 \times \mathbb{R})/\Gamma_0 \text{ where the action of } \Gamma_0 \text{ is} \\ \alpha \cdot (\tilde{y}, \tilde{v}_0) &= (\alpha \cdot \tilde{y}, d(\Theta_{f,0}(\alpha))_{\tilde{f}_0(\tilde{y})} \cdot \tilde{v}_0) = (\alpha \cdot \tilde{y}, d(-\text{Id})_0 \cdot \tilde{v}_0) = (-\tilde{y}, -\tilde{v}_0) \end{aligned}$$

while

$$\begin{aligned} p'^{-1}(U_0) &\cong (\tilde{U}_0 \times \mathbb{R})/\Gamma_0 \text{ where the action of } \Gamma_0 \text{ is} \\ \alpha \cdot (\tilde{y}, \tilde{v}_0) &= (\alpha \cdot \tilde{y}, d(\Theta'_{f,0}(\alpha))_{\tilde{f}_0(\tilde{y})} \cdot \tilde{v}_0) = (\alpha \cdot \tilde{y}, d(\text{Id})_0 \cdot \tilde{v}_0) = (-\tilde{y}, \tilde{v}_0) \end{aligned}$$

which is enough to substantiate our claim. Note that these orbundles are not equivalent in any reasonable sense.

To further illustrate the complexity involved in pulling back the tangent bundle by an orbifold map, the following is instructive.

Example 32. Consider the situation from example 28: $\mathcal{O} = \mathbb{R}/\mathbb{Z}_2$ with \mathbb{Z}_2 acting with generator α , where $\alpha \cdot x = -x$ as above and $\mathcal{P} = \mathbb{R}^3/(\mathbb{Z}_2 \times \mathbb{Z}_2)$ where $\mathbb{Z}_2 \times \mathbb{Z}_2 = \langle j, k \mid j^2 = k^2 = 1 = [j, k] \rangle$ with the action defined by $j \cdot (x, y, z) = (-x, y, -z)$ and $k \cdot (x, y, z) = (-x, -y, z)$. We consider the two complete orbifold maps ${}_*\mathbf{f}$ and ${}_*\mathbf{f}'$ from example 28 which cover the orbifold map $f = (f, \{\tilde{y}_1 \mapsto (\tilde{y}_1, 0, 0)\})$ where $\bullet f(y_1) = (y_1, 0, 0)$. We have for all x , $\tilde{U}_x \times_{\tilde{V}_{f(x)}} T\tilde{V}_{f(x)} \cong \tilde{U}_x \times \mathbb{R}^3$. Like in example 31, for $x \neq 0$, $\Gamma_x = \{e\}$, and so the action of Γ_x on $\tilde{U}_x \times_{\tilde{V}_{f(x)}} T\tilde{V}_{f(x)}$ is necessarily trivial. If we denote, as before, the orbundle projections $p : {}_*\mathbf{f}^*(T\mathcal{P}) \rightarrow \mathcal{O}$ and $p' : {}_*\mathbf{f}'^*(T\mathcal{P}) \rightarrow \mathcal{O}$, then $p^{-1}(U_x) = p'^{-1}(U_x) \cong \tilde{U}_x \times \mathbb{R}^3$. On the other hand, for $x = 0$, since $\Theta_{f,0}(\alpha) = j$ and $\Theta'_{f,0}(\alpha) = k$ we see that

$$\begin{aligned} p^{-1}(U_0) &\cong (\tilde{U}_0 \times \mathbb{R}^3)/\Gamma_0 \text{ where the action of } \Gamma_0 \text{ is} \\ \alpha \cdot (\tilde{y}, \tilde{v}_0) &= (\alpha \cdot \tilde{y}, dj_{\tilde{f}_0(\tilde{y})} \cdot \tilde{v}_0) = (-\tilde{y}, -(\tilde{v}_1)_0, (\tilde{v}_2)_0, -(\tilde{v}_3)_0) \end{aligned}$$

where $\tilde{v}_0 = ((\tilde{v}_1)_0, (\tilde{v}_2)_0, (\tilde{v}_3)_0) \in \mathbb{R}^3$. Similarly,

$$\begin{aligned} p'^{-1}(U_0) &\cong (\tilde{U}_0 \times \mathbb{R}^3)/\Gamma_0 \text{ where the action of } \Gamma_0 \text{ is} \\ \alpha \cdot (\tilde{y}, \tilde{v}_0) &= (\alpha \cdot \tilde{y}, dk_{\tilde{f}_0(\tilde{y})} \cdot \tilde{v}_0) = (-\tilde{y}, -(\tilde{v}_1)_0, -(\tilde{v}_2)_0, (\tilde{v}_3)_0). \end{aligned}$$

Although the pullback orbundles $\star f^*(T\mathcal{P})$ and $\star f'^*(T\mathcal{P})$ are naturally isomorphic, we will later see that neighborhoods of the zero section are taken by the Riemannian exponential map to the neighborhoods $\mathcal{N}^r(\star f, \varepsilon)$ and $\mathcal{N}^r(\star f', \varepsilon)$ of example 28, respectively. This illustrates why it is necessary to use *complete* orbifold maps in order to fully understand the topological structure of a neighborhood $\mathcal{N}^r(f, \varepsilon)$ of an orbifold map.

Orbisections. We now define a natural notion of section of a linear orbundle. For a definition, see for example [BB02].

Definition 33. A C^r orbisection of a m -dimensional linear orbundle $p : \mathcal{E} \rightarrow \mathcal{O}$ is a C^r orbifold map $\sigma : \mathcal{O} \rightarrow \mathcal{E}$ such that $p \circ \sigma = \text{Id}_{\mathcal{O}}$ and for any $x \in \mathcal{O}$ and chart U_x about x , we have $\text{pr}_1 \circ \tilde{\sigma}_x = \text{Id}_{\tilde{U}_x}$. That is, we take the *identity* lift of the identity $\text{Id}_{\mathcal{O}}$ in \tilde{U}_x . Locally we have the diagram:

$$\begin{array}{ccc}
 & & \tilde{U}_x \times \mathbb{R}^m \\
 & \nearrow \tilde{\sigma}_x & \downarrow \text{pr}_1 \\
 \tilde{U}_x & \xrightarrow{\text{Id}_{\tilde{U}_x}} & \tilde{U}_x \\
 \downarrow \pi_x & & \downarrow \pi_x \\
 U_x & \xrightarrow{\text{Id}} & U_x \\
 & \nearrow \sigma & \downarrow p \\
 & & p^{-1}(U_x) \cong (\tilde{U}_x \times \mathbb{R}^m) / \Gamma_x \\
 & & \downarrow \Pi_x
 \end{array}$$

Note that the action of Γ_x on $\tilde{U}_x \times \mathbb{R}^m$ is given as part of the data defining \mathcal{E} . Although, in general, the class of complete orbifold maps is different from the class of orbifold maps, as in the case for diffeomorphisms (section 4), in the case of orbisections of the pullback of a tangent orbundle, these notions coincide.

Proposition 34. *Let $\star f : \mathcal{O} \rightarrow \mathcal{P}$ be a C^{r+1} complete orbifold map between C^{r+1} orbifolds and let $\star f^*(T\mathcal{P})$ denote the pullback of the tangent orbundle. Let $\sigma = (\sigma, \{\tilde{\sigma}_x\})$ be a C^r orbisection of $\star f^*(T\mathcal{P})$. Then there is a unique homomorphism $\Theta_{\sigma, x}$ for which $\tilde{\sigma}_x$ is Γ_x equivariant. In other words, the set of orbisections can be identified with the set of complete orbisections $\sigma \leftrightarrow \star \sigma$.*

Proof. Given an orbifold chart \tilde{U}_x around x and an orbundle chart for $\star f^*(T\mathcal{P})$ with local product coordinates $(\tilde{y}, \tilde{\xi}) = (\tilde{y}, [\tilde{f}_x(\tilde{y}), \tilde{v}]) \in \tilde{U}_x \times_{\tilde{V}_{f(x)}} T\tilde{V}_{f(x)}$, the local action of Γ_x in these coordinates is given by $\gamma \cdot (\tilde{y}, \tilde{\xi}) = (\gamma \cdot \tilde{y}, \Theta_{f, x}(\gamma) \cdot \tilde{\xi}) = (\gamma \cdot \tilde{y}, [\Theta_{f, x}(\gamma) \cdot \tilde{f}_x(\tilde{y}), d(\Theta_{f, x}(\gamma))_{\tilde{f}_x(\tilde{y})} \cdot \tilde{v}])$. With respect to these local coordinates, $\tilde{\sigma}_x$ has the form $\tilde{\sigma}_x(\tilde{y}) = (\tilde{y}, [\tilde{f}_x(\tilde{y}), \tilde{s}_x(\tilde{y})])$ and if $\Theta_{\sigma, x} : \Gamma_x \rightarrow \Gamma_{\sigma(x)} = \Gamma_x$ is some homomorphism for which $\tilde{\sigma}_x$ is equivariant with respect to, then

$$\begin{aligned}
 \tilde{\sigma}_x(\gamma \cdot \tilde{y}) &= (\gamma \cdot \tilde{y}, [\tilde{f}_x(\gamma \cdot \tilde{y}), \tilde{s}_x(\gamma \cdot \tilde{y})]) \\
 &= \Theta_{\sigma, x}(\gamma) \cdot (\tilde{y}, [\tilde{f}_x(\tilde{y}), \tilde{s}_x(\tilde{y})]) \\
 &= (\Theta_{\sigma, x}(\gamma) \cdot \tilde{y}, [(\Theta_{f, x}(\Theta_{\sigma, x}(\gamma))) \cdot \tilde{f}_x(\tilde{y}), d(\Theta_{f, x}(\Theta_{\sigma, x}(\gamma)))_{\tilde{f}_x(\tilde{y})} \cdot \tilde{s}_x(\tilde{y})])
 \end{aligned}$$

Therefore, since Γ_x acts effectively on \tilde{U}_x , $\Theta_{\sigma,x}(\gamma) = \gamma$ and $\Theta_{\sigma,x} = \text{Id} : \Gamma_x \rightarrow \Gamma_x$ for all $\gamma \in \Gamma_x$ and $x \in \mathcal{O}$. Furthermore, we get the equivariance relation $\tilde{s}_x(\gamma \cdot \tilde{y}) = d(\Theta_{f,x}(\gamma))_{\tilde{f}_x(\tilde{y})}(\tilde{s}_x(\tilde{y}))$. \square

Just as in the case of orbisections of the tangent orbifold, the set of orbisections of the pullback tangent orbifold carry a vector space structure.

Proposition 35. *Let $\star f \in C_{\star\text{Orb}}^{r+1}(\mathcal{O}, \mathcal{P})$. The set $\mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P}))$ of C^r orbisections of the pullback tangent orbifold $\star f^*(T\mathcal{P})$ is naturally a real vector space with the vector space operations being defined pointwise.*

Proof. The argument here is basically the same as the corresponding argument for orbisections of the tangent orbifold [BB08]. Let $\sigma \in \mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P}))$. Let $\tilde{\sigma}_x$ be the lift of σ . Then $\tilde{\sigma}_x(\tilde{y}) = (\tilde{y}, [\tilde{f}_x(\tilde{y}), \tilde{s}_x(\tilde{y})])$. By proposition 34, $\tilde{s}_x : \tilde{U}_x \rightarrow \mathbb{R}^m$ is such that $\tilde{s}_x(\gamma \cdot \tilde{y}) = d(\Theta_{f,x}(\gamma))_{\tilde{f}_x(\tilde{y})}(\tilde{s}_x(\tilde{y}))$. In particular, since \tilde{x} is a fixed point of the Γ_x action on \tilde{U}_x , we have $\tilde{s}_x(\tilde{x}) = \tilde{s}_x(\gamma \cdot \tilde{x}) = d(\Theta_{f,x}(\gamma))_{\tilde{f}_x(\tilde{x})}(\tilde{s}_x(\tilde{x}))$. Thus $\tilde{s}_x(\tilde{x})$ is a fixed point of the (linear) action of Γ_x on \mathbb{R}^m as defined in the pullback orbifold. Note that the set of such fixed points forms a *vector subspace* of \mathbb{R}^m . As a result we may define a real vector space structure on $\mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P}))$ as follows: For $\sigma_i \in \mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P}))$, let $\tilde{\sigma}_{i,x}$ be local lifts at x as above. Define

$$\begin{aligned} (\sigma_1 + \sigma_2)(y) &= \Pi_x((\tilde{\sigma}_{1,x} + \tilde{\sigma}_{2,x})(\tilde{y})) = \Pi_x((\tilde{y}, \tilde{s}_1(\tilde{y}) + \tilde{s}_2(\tilde{y}))) = \sigma_1(y) + \sigma_2(y) \\ (\lambda\sigma)(y) &= \Pi_x((\lambda\tilde{\sigma}_x)(\tilde{y})) = \Pi_x((\tilde{y}, \lambda\tilde{s}(\tilde{y}))) = \lambda(\sigma(y)) \end{aligned}$$

\square

Proposition 36. *Let $\star f \in C_{\star\text{Orb}}^{r+1}(\mathcal{O}, \mathcal{P})$ with \mathcal{O} compact (without boundary). The inclusion map $\mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P})) \hookrightarrow C_{\text{Orb}}^r(\mathcal{O}, \star f^*(T\mathcal{P}))$ induces a separable Banach space structure on $\mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P}))$ for $1 \leq r < \infty$ and a separable Fréchet space structure if $r = \infty$.*

Proof. The argument here is also similar to the corresponding argument for orbisections of the tangent orbifold [BB08]. Let $\mathcal{D} = \{D_i\}_{i=1}^N$ be a cover of $f(\mathcal{O})$ by a finite number of compact orbifold charts over each of which the tangent orbifold $T\mathcal{P}$ is trivialized. Then the collection $\mathcal{C} = \{C_i = f^{-1}(D_i)\}$ is a finite cover of \mathcal{O} by compact subsets. By reindexing and shrinking D_i if necessary, we may assume each C_i is connected and is contained in a orbifold chart of \mathcal{O} and so, $C_i \cong \tilde{C}_i/\Gamma_i$. Let $\tilde{C}_i \times_{\tilde{D}_i} T\tilde{D}_i \cong \tilde{C}_i \times \mathbb{R}^m$ be the corresponding orbifold charts for $\star f^*(T\mathcal{P})$ with action of $\Gamma_i : \gamma \cdot (\tilde{y}, \tilde{\xi}) = (\gamma \cdot \tilde{y}, [\Theta_{f,i}(\gamma) \cdot \tilde{f}_x(\tilde{y}), d(\Theta_{f,i}(\gamma))_{\tilde{f}_x(\tilde{y})} \cdot \tilde{v}])$. Let $V_{i,r} = C^r(\tilde{C}_i, \mathbb{R}^m)$ for $i = 1, \dots, N$ and $0 \leq r \leq \infty$ with topology of uniform convergence of derivatives of order $\leq r$. This is a Banach space for finite r and a Fréchet space for $r = \infty$. For finite r , let $\|\cdot\|_{i,r}$ be a C^r norm on $V_{i,r}$. Define a linear map $L : \mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P})) \rightarrow \bigoplus_{i=1}^N V_{i,r}$ by

$$L(\sigma) = (\text{pr}_2(\tilde{\chi}_1\tilde{\sigma}), \dots, \text{pr}_2(\tilde{\chi}_N\tilde{\sigma}))$$

where $\chi_i \in C_{\text{Orb}}^r(\mathcal{O}, [0, 1])$, $i = 1, \dots, N$, is a partition of unity subordinate to the cover \mathcal{C} ([BB08, proposition 6.1]) and $\text{pr}_2 : \tilde{C}_i \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ is bundle projection onto the second factor. Continuity of L is immediate from the definitions of the C^r topology on $\mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P}))$ and the topology on $\bigoplus_{i=1}^N V_{i,r}$. Moreover, given a neighborhood of the zero section $\mathbf{0} \in \mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P}))$ of the

form $\mathcal{N}^r(\mathbf{0}, \varepsilon_i; \mathcal{C})$, it is apparent that there is a neighborhood of the zero section $\mathbf{0}$ in $\bigoplus_{i=1}^N V_{i,r}$ of the form $\max\{\|s_1\|_{1,r}, \dots, \|s_N\|_{N,r}\} < \delta$ where $\delta \leq \min\{\varepsilon_1, \dots, \varepsilon_N\}$ contained in $L(\mathcal{N}^r(\mathbf{0}, \varepsilon_i; \mathcal{C}))$. Thus, with the subspace topology on $L(\mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P})))$, $L : \mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P})) \rightarrow L(\mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P})))$ is a linear homeomorphism. Since $\mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P})) \subset C_{\text{Orb}}^r(\mathcal{O}, \star f^*(T\mathcal{P}))$ is a closed subset, we see that $L(\mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P})))$ is a closed subspace of the direct sum and thus $\mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P}))$ inherits a Banach space structure if $r < \infty$ and a Fréchet space structure if $r = \infty$. \square

The following is the analogue of the notion of admissible tangent vector as defined in [BB08].

Definition 37. Let \mathcal{O}, \mathcal{P} and f be as in proposition 36. Let $x \in \mathcal{O}$. Denote by $A_x(\star f^*(T\mathcal{P}))$ the set of *admissible vectors at x*

$$A_x(\star f^*(T\mathcal{P})) = \{v \in \star f^*(T\mathcal{P})_x \mid (x, v) = \sigma(x) \text{ for some } \sigma \in \mathcal{D}_{\text{Orb}}^0(\star f^*(T\mathcal{P}))\}$$

By proposition 35, $A_x(\star f^*(T\mathcal{P}))$ is a vector space for each x , and a suborbifold of $\star f^*(T\mathcal{P})_x$. The *admissible pullback bundle of $T\mathcal{P}$* is the subset $A(\star f^*(T\mathcal{P})) = \bigcup_{x \in \mathcal{O}} A_x(\star f^*(T\mathcal{P})) \subset \star f^*(T\mathcal{P})$ with the subspace topology. In general, $A(\star f^*(T\mathcal{P}))$ will not be an orbifold. Recall that the set of admissible *tangent* vectors at z , $A_z(\mathcal{P})$ as defined in [BB08] are obtained from definition 37 by replacing $\star f^*(T\mathcal{P})$ by $T\mathcal{P}$.

7. THE EXPONENTIAL MAP AND PROOF OF THEOREM 1

In this section we will need several facts about Riemannian orbifolds and the exponential map. Our reference for this material will be [BB08]. Throughout this section, we assume that \mathcal{O}, \mathcal{P} are smooth orbifolds and that \mathcal{O} is compact (without boundary). Without loss of generality, we may assume, by [BB08, propositions 3.11, 6.4], that both \mathcal{O} and \mathcal{P} are C^∞ orbifolds with C^∞ Riemannian metrics.

The exponential map. Recall the construction of the exponential map for a smooth Riemannian orbifold \mathcal{P} [BB08, section 6]. Assume that the collection $\{V_\alpha\}$ is a locally finite open covering of \mathcal{P} by orbifold charts that are relatively compact. Let $TV_\alpha \cong (\tilde{V}_\alpha \times \mathbb{R}^m)/\Gamma_\alpha$ be a local trivialization of the tangent bundle over V_α . Denote the Riemannian exponential map on $T\tilde{V}_\alpha$ by $\widetilde{\text{exp}}_{\tilde{V}_\alpha} : T\tilde{V}_\alpha \rightarrow \tilde{V}_\alpha$. Thus, for $\tilde{z} \in \tilde{V}_\alpha$ and $\tilde{v} \in T_{\tilde{z}}\tilde{V}_\alpha$ we have $\widetilde{\text{exp}}_{\tilde{V}_\alpha}(\tilde{z}, t\tilde{v}) = \tilde{c}_{\tilde{z}, \tilde{v}}(t)$ where $\tilde{c}_{\tilde{z}, \tilde{v}}$ is the unit speed geodesic in \tilde{V}_α which starts at \tilde{z} and has initial velocity \tilde{v} . Recall that there is an open neighborhood $\tilde{\Omega}_{\tilde{V}_\alpha} \subset T\tilde{V}_\alpha$ of the 0-section of $T\tilde{V}_\alpha$ such that $\tilde{c}_{\tilde{z}, \tilde{v}}(1)$ is defined for $\tilde{v} \in T_{\tilde{z}}\tilde{V}_\alpha \cap \tilde{\Omega}_{\tilde{V}_\alpha}$. Furthermore, by shrinking $\tilde{\Omega}_{\tilde{V}_\alpha}$ if necessary, we may assume that on $T_{\tilde{z}}\tilde{V}_\alpha \cap \tilde{\Omega}_{\tilde{V}_\alpha}$, $\widetilde{\text{exp}}_{\tilde{V}_\alpha}(\tilde{z}, \cdot)$ is a local diffeomorphism onto a neighborhood of $\tilde{z} \in \tilde{V}_\alpha$ for each $\tilde{z} \in \tilde{V}_\alpha$. Let $\Omega_\alpha = \Pi_\alpha(\tilde{\Omega}_{\tilde{V}_\alpha})$, an open subset of $T\mathcal{P}$, and define $\Omega = \bigcup_\alpha \Omega_\alpha$. Ω is an open neighborhood of the 0-orbisection of $T\mathcal{P}$.

Definition 38. Let $z \in V_\alpha$, and $(z, v) \in \Omega_\alpha$. Choose $(\tilde{z}, \tilde{v}) \in \Pi_\alpha^{-1}(z, v)$. Then the *Riemannian exponential map* $\text{exp} : \Omega \subset T\mathcal{P} \rightarrow \mathcal{P}$ is defined by $\text{exp}(z, v) = \pi_\alpha \circ \widetilde{\text{exp}}_{\tilde{V}_\alpha}(\tilde{z}, \tilde{v})$.

By [BB08, proposition 6.7], this exponential map is well-defined and $\widetilde{\text{exp}}_{\tilde{V}_\alpha}$ satisfies, for all $\delta \in \Gamma_\alpha$, the equivariance relation:

$$\widetilde{\text{exp}}_{\tilde{V}_\alpha}[\delta \cdot (\tilde{z}, \tilde{v})] = \delta \cdot \widetilde{\text{exp}}_{\tilde{V}_\alpha}(\tilde{z}, \tilde{v}).$$

As usual we denote by \exp_z the restriction of \exp to a single tangent cone $T_z\mathcal{P}$. We let $B(x, r)$ denote the metric r -ball centered at z and use tildes to denote corresponding points in local coverings.

The relation between orbisections, the exponential map and complete orbifold maps. The composition of the exponential map with an orbisection of the pullback tangent orbibundle via a complete orbifold map $\star f$ turns out to be a smooth complete orbifold map with the same equivariance relation as $\star f$.

Proposition 39. *Let \mathcal{O}, \mathcal{P} be smooth Riemannian orbifolds and let $\star f \in C_{\star\text{Orb}}^{r+1}(\mathcal{O}, \mathcal{P})$. Let σ be a C^r orbisection of the pullback tangent orbibundle $\star f^*(T\mathcal{P})$ with $F \circ \sigma(x) \in \Omega$. Then the map $E^\sigma(x) = (\exp \circ F \circ \sigma)(x) : \mathcal{O} \rightarrow \mathcal{P}$ is a complete C^r orbifold map with representation $\star E^\sigma = (E^\sigma, \{\tilde{E}_x^\sigma\}, \{\Theta_{E^\sigma, x}\})$ where $\tilde{E}_x^\sigma = \widetilde{\exp}_{V_{f(x)}} \circ \tilde{F}_x \circ \tilde{\sigma}_x$ and $\Theta_{E^\sigma, x} = \Theta_{f, x}$ for all $x \in \mathcal{O}$. In particular, $\star E^\sigma \in C_{\star\text{Orb}}^r(\mathcal{O}, \mathcal{P})$.*

Proof. Let (\tilde{U}_x, Γ_x) be an orbifold chart at $x \in \mathcal{O}$. For $y \in U_x$, $\sigma(y) = (y, \xi(y)) \in \star f^*(TV_{f(x)})$. Let $\tilde{\sigma}_x = (\tilde{y}, \tilde{\xi}(\tilde{y}))$ be a lift of σ and \tilde{F}_x the map defined after the diagram of definition 30. Then the map $\tilde{E}_x^\sigma = \widetilde{\exp}_{V_{f(x)}} \circ \tilde{F}_x \circ \tilde{\sigma}_x$ is a C^r lift of E^σ and using the equivariance relations for σ_x , \tilde{F}_x and $\widetilde{\exp}_{V_{f(x)}}$ we have for all $\gamma \in \Gamma_x$:

$$\begin{aligned} \tilde{E}_x^\sigma(\gamma \cdot \tilde{y}) &= \widetilde{\exp}_{V_{f(x)}} \circ \tilde{F}_x \circ \tilde{\sigma}_x(\gamma \cdot \tilde{y}) \\ &= \widetilde{\exp}_{V_{f(x)}} \circ \tilde{F}_x(\gamma \cdot \tilde{\sigma}_x(\tilde{y})) \\ &= \widetilde{\exp}_{V_{f(x)}}(\Theta_{f, x}(\gamma) \cdot \tilde{\xi}(\tilde{y})) \\ &= \Theta_{f, x}(\gamma) \cdot \widetilde{\exp}_{V_{f(x)}}(\tilde{\xi}(\tilde{y})) \\ &= \Theta_{f, x}(\gamma) \cdot \widetilde{\exp}_{V_{f(x)}} \circ \tilde{F}_x \circ \tilde{\sigma}_x(\tilde{y}) \\ &= \Theta_{f, x}(\gamma) \cdot \tilde{E}_x^\sigma(\tilde{y}). \end{aligned}$$

□

The local manifold structure. Let \mathcal{O}, \mathcal{P} and $\star f$ be as in proposition 39. Denote by $\mathbf{0} : \mathcal{O} \rightarrow \star f^*(T\mathcal{P})$, $\mathbf{0}(x) = 0_x \in \star f^*(T\mathcal{P})_x$, the 0-orbisection of $\star f^*(T\mathcal{P})$. Then $E^{\mathbf{0}}(x) = (\exp \circ F \circ \mathbf{0})(x) = f(x)$. We let $\mathcal{B}_{\star f}^r(\sigma, \varepsilon) = \mathcal{N}^r(\sigma, \varepsilon) \cap \mathcal{D}_{\text{Orb}}^r(\star f^*(T\mathcal{P}))$. That is, $\mathcal{B}_{\star f}^r(\sigma, \varepsilon)$ is the set of C^r orbisections ε -close to σ in the C^r topology on $C_{\text{Orb}}^r(\mathcal{O}, \star f^*(T\mathcal{P}))$. Proposition 39 and definition 20 immediately yield the following

Proposition 40. *Let \mathcal{O}, \mathcal{P} and $\star f$ be as in proposition 39 with \mathcal{O} compact. There exists $\varepsilon > 0$ and continuous map $E : \mathcal{B}_{\star f}^r(\mathbf{0}, \varepsilon) \rightarrow \mathcal{N}^r(\star f, \varepsilon)$ defined by $E(\sigma) = E^\sigma$.*

Since $\mathcal{B}_{\star f}^r(\mathbf{0}, \varepsilon)$ is an open subset of a Banach/Fréchet space by proposition 36, the proof theorem 1 will be complete if E is shown to be a homeomorphism. We first show that E is injective.

Proposition 41. *The map $E : \mathcal{B}_{\star f}^r(\mathbf{0}, \varepsilon) \rightarrow \mathcal{N}^r(\star f, \varepsilon)$ is injective.*

Proof. Suppose $E(\sigma) = E(\tau)$ for $\sigma, \tau \in \mathcal{B}_{\star f}^r(\mathbf{0}, \varepsilon)$. Since these are to be considered equal as complete orbifold maps, in each orbifold chart (\tilde{U}_x, Γ_x) , we must have equal local lifts: $\widetilde{\exp}_{V_{f(x)}} \circ \tilde{F}_x \circ \tilde{\sigma}_x(\tilde{y}) = \widetilde{\exp}_{V_{f(x)}} \circ \tilde{F}_x \circ \tilde{\tau}_x(\tilde{y})$. If we write in local coordinates $\tilde{\sigma}_x(\tilde{y}) = (\tilde{y}, \tilde{\xi}(\tilde{y}))$ and $\tilde{\tau}_x(\tilde{y}) = (\tilde{y}, \tilde{\eta}(\tilde{y}))$ where $\tilde{\xi}(\tilde{y}) = [\tilde{f}_x(\tilde{y}), v_{\tilde{f}_x(\tilde{y})}]$ and

$\eta(\tilde{y}) = [\tilde{f}_x(\tilde{y}), w_{\tilde{f}_x(\tilde{y})}]$ then

$$\begin{aligned} \widetilde{\exp}_{\tilde{V}_{f(x)}} \circ \tilde{F}_x \circ \tilde{\sigma}_x(\tilde{y}) &= \widetilde{\exp}_{\tilde{V}_{f(x)}} \circ \tilde{F}_x \circ \tilde{\tau}_x(\tilde{y}) \iff \\ \widetilde{\exp}_{\tilde{V}_{f(x)}} \circ \tilde{F}_x(\tilde{y}, \tilde{\xi}(\tilde{y})) &= \widetilde{\exp}_{\tilde{V}_{f(x)}} \circ \tilde{F}_x(\tilde{y}, \tilde{\eta}(\tilde{y})) \iff \\ \widetilde{\exp}_{\tilde{V}_{f(x)}}[\tilde{f}_x(\tilde{y}), v_{\tilde{f}_x(\tilde{y})}] &= \widetilde{\exp}_{\tilde{V}_{f(x)}}[\tilde{f}_x(\tilde{y}), w_{\tilde{f}_x(\tilde{y})}] \end{aligned}$$

Since $\widetilde{\exp}_{\tilde{V}_{f(x)}}(\tilde{f}_x(\tilde{y}), \cdot)$ is a local C^r diffeomorphism we must have $v_{\tilde{f}_x(\tilde{y})} = w_{\tilde{f}_x(\tilde{y})}$. Hence $\sigma = \tau$ (as orbifold maps) and E is injective. \square

The proof of the following proposition is a slightly modified version of [BB08, proposition 7.3].

Proposition 42. *The map $E : \mathcal{B}_{\star f}^r(\mathbf{0}, \varepsilon) \rightarrow \mathcal{N}^r(\star f, \varepsilon)$ is surjective.*

Proof. Let $\star g = (g, \{\tilde{g}_x\}, \{\Theta_{g,x}\}) \in \mathcal{N}^r(\star f, \varepsilon)$. Let $\{C_i\}$ be a finite covering of \mathcal{O} by compact sets such that C_i is an orbifold chart and $g(C_i) \subset V_i$ where V_i is a relatively compact orbifold chart of \mathcal{P} . Let $x \in C_i$, and $\tilde{U}_x \subset \text{int } \tilde{C}_i$ an orbifold chart at x where the local lift \tilde{g}_x to \tilde{U}_x is C^0 ε -close to the local lift \tilde{f}_x . By lemma 23 and its proof we have $\Theta_{f,x} = \theta_{g(x)f(x)} \circ \Theta_{g,x} : \Gamma_x \rightarrow \Gamma_{f(x)}$. In particular, the action of $\Theta_{f,x}$ is the same as action $\Theta_{g,x}$ on the image $\tilde{g}_x(\tilde{U}_x) \subset \tilde{V}_{f(x)}$.

We wish to define a C^r orbisection σ so that $E(\sigma) = \star g$. We do this by defining appropriate local lifts $\tilde{\sigma}_x$. In particular, let

$$\tilde{\sigma}_x(\tilde{y}) = (\tilde{y}, \tilde{\xi}(\tilde{y})) = \left(\tilde{y}, \left[\tilde{f}_x(\tilde{y}), \widetilde{\exp}_{\tilde{V}_{f(x), \tilde{f}_x(\tilde{y})}}^{-1}(\tilde{g}_x(\tilde{y})) \right] \right) \in \star f^*(T\tilde{V}_{f(x)}).$$

With this definition, we see that

$$\begin{aligned} \tilde{E}_x^\sigma(\tilde{y}) &= \widetilde{\exp}_{\tilde{V}_{f(x)}} \circ \tilde{F}_x \circ \tilde{\sigma}_x(\tilde{y}) \\ &= \widetilde{\exp}_{\tilde{V}_{f(x)}} \circ \widetilde{\exp}_{\tilde{V}_{f(x), \tilde{f}_x(\tilde{y})}}^{-1}(\tilde{g}_x(\tilde{y})) = \tilde{g}_x(\tilde{y}). \end{aligned}$$

This shows that $E(\sigma) = \star g$. All that remains to show is that $\tilde{\sigma}_x$ satisfies the correct equivariance relation for an orbisection. Before we do that, observe that, in general, for $\delta \in \Gamma_{f(x)}$ we have (essentially for any exponential map)

$$\begin{aligned} \widetilde{\exp}_{\tilde{V}_{f(x), \tilde{f}_x(\tilde{y})}}^{-1}(\delta \cdot \tilde{z}) &= (d\delta)_{\delta^{-1}\tilde{f}_x(\tilde{y})} \circ \widetilde{\exp}_{\tilde{V}_{f(x), \delta^{-1}\tilde{f}_x(\tilde{y})}}^{-1}(\tilde{z}) \\ &= \delta \cdot \widetilde{\exp}_{\tilde{V}_{f(x), \tilde{f}_x(\tilde{y})}}^{-1}(\tilde{z}). \end{aligned}$$

Thus,

$$\begin{aligned} \tilde{\sigma}_x(\gamma \cdot \tilde{y}) &= (\gamma \cdot \tilde{y}, \tilde{\xi}(\gamma \cdot \tilde{y})) = \left(\gamma \cdot \tilde{y}, \left[\tilde{f}_x(\gamma \cdot \tilde{y}), \widetilde{\exp}_{\tilde{V}_{f(x), \tilde{f}_x(\gamma \cdot \tilde{y})}}^{-1}(\tilde{g}_x(\gamma \cdot \tilde{y})) \right] \right) \\ &= \left(\gamma \cdot \tilde{y}, \left[\Theta_{f,x}(\gamma) \cdot \tilde{f}_x(\tilde{y}), \widetilde{\exp}_{\tilde{V}_{f(x), \Theta_{f,x}(\gamma)\tilde{f}_x(\tilde{y})}}^{-1}(\Theta_{g,x}(\gamma) \cdot \tilde{g}_x(\tilde{y})) \right] \right) \\ &= \left(\gamma \cdot \tilde{y}, \left[\Theta_{f,x}(\gamma) \cdot \tilde{f}_x(\tilde{y}), \Theta_{f,x}(\gamma) \cdot \widetilde{\exp}_{\tilde{V}_{f(x), (\Theta_{f,x}(\gamma))^{-1}\Theta_{f,x}(\gamma)\tilde{f}_x(\tilde{y})}}^{-1}(\tilde{g}_x(\tilde{y})) \right] \right) \\ &= \left(\gamma \cdot \tilde{y}, \left[\Theta_{f,x}(\gamma) \cdot \tilde{f}_x(\tilde{y}), \Theta_{f,x}(\gamma) \cdot \widetilde{\exp}_{\tilde{V}_{f(x), \tilde{f}_x(\tilde{y})}}^{-1}(\tilde{g}_x(\tilde{y})) \right] \right) \\ &= \left(\gamma \cdot \tilde{y}, \Theta_{f,x}(\gamma) \cdot \tilde{\xi}(\tilde{y}) \right) \\ &= \gamma \cdot \tilde{\sigma}_x(\tilde{y}) \end{aligned}$$

which is the correct equivariance relation for an orbisection. As a result we see that the map $\sigma(x) = \Pi_x \circ \tilde{\sigma}_x(\tilde{x})$ defines a C^r orbisection of $\star f^*(T\mathcal{P})$ and that $E(\sigma) = \star g$. \square

The following proposition is the last step to complete the proof of the theorem 1. It gives a C^0 manifold structure to $C^r_{\star\text{Orb}}(\mathcal{O}, \mathcal{P})$ where the model space for a neighborhood of $\star f$ is the topological vector space of C^r orbisections of a pullback tangent orbundle of \mathcal{P} via $\star f$ with the C^r topology.

Proposition 43. *The map $E : \mathcal{B}_{\star f}^r(\mathbf{0}, \varepsilon) \rightarrow \mathcal{N}^r(\star f, \varepsilon)$ is a homeomorphism.*

Proof. Propositions 41 and 42 show that E is bijective. Continuity of E follows from the formula for a local lift of E given in proposition 39 and continuity of E^{-1} follows from the formula for $\tilde{\sigma}_x$ given in the proof of proposition 42. \square

8. BUILDING STRATIFIED NEIGHBORHOODS

Our first task of this section will be to prove corollary 3. Let $f \in C^r_{\text{Orb}}(\mathcal{O}, \mathcal{P})$. From the observation following definition 20, we have that

$$q^{-1}(\mathcal{N}^r(f, \varepsilon)) = \mathcal{N}^r(\star f_1, \varepsilon) \cup \cdots \cup \mathcal{N}^r(\star f_k, \varepsilon)$$

is a disjoint union of neighborhoods where each complete map $\star f_i = (f, \{\tilde{f}_x\}, \{\Theta_{f,x}\}_i)$. We first partition the neighborhood $\mathcal{N}^r(f, \varepsilon)$. For each $g \in \mathcal{N}^r(f, \varepsilon)$, define $J_g \subset \{1, \dots, k\}$ to be the set of indices j such that $q^{-1}(g) \cap \mathcal{N}^r(\star f_j, \varepsilon) \neq \emptyset$. For $J \subset \{1, \dots, k\}$, define

$$\mathcal{X}_J = \{g \in \mathcal{N}^r(f, \varepsilon) \mid J = J_g\}.$$

This is a partition of $\mathcal{N}^r(f, \varepsilon)$. Of course, the partial ordering is from set inclusion: $J' \prec J \Leftrightarrow J' \subset J$. We now verify conditions (1) and (2) of definition 26 in the next two lemmas.

Lemma 44. *Each \mathcal{X}_J is a submanifold of $\mathcal{N}^r(f, \varepsilon)$.*

Proof. Let $J = \{j_1, \dots, j_l\}$. For any $g \in \mathcal{X}_J$, we have

$$q^{-1}(g) \cap \mathcal{N}^r(\star f_{j_1}, \varepsilon) \cap \cdots \cap \mathcal{N}^r(\star f_{j_l}, \varepsilon) \neq \emptyset.$$

By proposition 43, there exists unique orbisections $\sigma_{j_1}, \dots, \sigma_{j_l}$ (of the respective pullback tangent orbundles $\star f_{j_i}^*(T\mathcal{P})$) such that $q(E(\sigma_{j_i})) = g$ for $i = 1, \dots, l$. Let (\tilde{U}_x, Γ_x) be a local chart about $x \in \mathcal{O}$ and let $\tilde{\sigma}_{j_i, x}$ denote the local lifts of the orbisection σ_{j_i} and let $\tilde{F}_{j_i, x} : \tilde{U}_x \times_{\tilde{V}_{f(x)}} T\tilde{V}_{f(x)} \rightarrow T\tilde{V}_{f(x)}$ denote the linear isomorphism $(\tilde{y}, \tilde{\xi}) \mapsto \tilde{\xi}$ given in definition 30 of the pullback tangent orbundle. Since $\widetilde{\text{exp}}_{\tilde{V}_{f(x)}} \circ \tilde{F}_{j_i, x} \circ \tilde{\sigma}_{j_i, x}(\tilde{y}) = \tilde{g}_x(\tilde{y})$ for all $i = 1, \dots, l$ and since $\widetilde{\text{exp}}_{\tilde{V}_{f(x)}}(\tilde{f}_x(\tilde{y}), \cdot)$ is a local diffeomorphism, we must have

$$\tilde{F}_{j_1, x} \circ \tilde{\sigma}_{j_1, x}(\tilde{y}) = \cdots = \tilde{F}_{j_l, x} \circ \tilde{\sigma}_{j_l, x}(\tilde{y})$$

for all $\tilde{y} \in \tilde{U}_x$. Because $\tilde{F}_{j_i, x}$ is a linear isomorphism, this relation is preserved under addition and scalar multiplication of local lifts of orbisections $\sigma_{j_i, x}$. From the proof of proposition 35, this relation descends to

$$(*) \quad F_{j_1} \circ \sigma_{j_1}(y) = \cdots = F_{j_l} \circ \sigma_{j_l}(y)$$

for $y \in U_x$. Since F_{j_i} is a linear isomorphism when restricted to the vector space of admissible vectors $A_x(\star f_{j_i}^*(T\mathcal{P}))$, the set of orbisections satisfying these relations is a linear submanifold of each $\mathcal{B}_{\star f_{j_i}}(\mathbf{0}, \varepsilon)$. From this we may conclude that each

$g \in \mathcal{X}_J$ has a neighborhood modeled on a linear submanifold of $\mathcal{B}_{\star f_{j_i}}(\mathbf{0}, \varepsilon)$, which is enough to prove that \mathcal{X}_J is a submanifold of $\mathcal{N}^r(f, \varepsilon)$. \square

Lemma 45. *If $\mathcal{X}_J \cap \overline{\mathcal{X}_{J'}} \neq \emptyset$, $J \neq J'$, then $J' \prec J$ and $\mathcal{X}_J \subset \overline{\mathcal{X}_{J'}}$.*

Proof. Let $J' = \{j_1, \dots, j_l\}$. For $i = 1, \dots, l$, suppose $\{\sigma_{j_i}^{(k)}\}_{k=1}^\infty$ is a sequence of orbisections which converges to $\overline{\sigma_{j_i}} \in \mathcal{D}_{\text{Orb}}^r(\star f_{j_i}^*(T\mathcal{P}))$. Further suppose each $\sigma_{j_i}^{(k)}$ satisfies condition (*) of lemma 44. Then, by continuity, each $\overline{\sigma_{j_i}}$ satisfies (*) also. If we let $q(E(\overline{\sigma_{j_i}})) = \overline{g}$, and $J = J_{\overline{g}}$ we have shown that if $\mathcal{X}_J \cap \overline{\mathcal{X}_{J'}} \neq \emptyset$, $J \neq J'$, then $J' \prec J$ and $\mathcal{X}_J \subset \overline{\mathcal{X}_{J'}}$. \square

Theorem 1 with lemmas 44 and 45 together prove corollary 3. Finally the proof of corollary 4 follows from corollary 3 and lemmas 11 and 21. That is, $\mathcal{N}^r(\bullet f, \varepsilon)$ is the quotient of the finite group $\mathcal{J}\mathcal{D}_{\mathcal{N}}$ acting on $\mathcal{N}^r(f, \varepsilon)$. That the corresponding quotient map q_\bullet restricts on each stratum to give a smooth orbifold chart follows from an argument almost identical to the argument in the proof of corollary 2 from section 3 that q_\bullet defined a smooth orbifold chart.

An alternative view of the stratification. Up to this point, the notion of pull-back bundle (definition 30) required the use of a *complete* orbifold map. Although not necessary for our results, we present a more global view of the stratification obtained above by defining directly an appropriate notion of pullback for an orbifold map f . We will use the setup of this section and the notation of definition 30. However, for convenience we will write a complete orbifold map $\star f_i = (f, \{\tilde{f}_x\}, \{\Theta_{f,x,i}\})$. To begin, we let $f^*(T\mathcal{P})$ be the space defined by:

$$f^*(T\mathcal{P}) = \left(\bigsqcup_{i=1}^k \star f_i^*(T\mathcal{P}) \right) / \sim$$

where the equivalence relation \sim is defined as follows: Let

$$\bigsqcup_{i=1}^k \left(\tilde{U}_x \times_{\tilde{V}_{f(x)}} T\tilde{V}_{f(x)} \right)_i$$

denote the disjoint union of k copies of $\tilde{U}_x \times_{\tilde{V}_{f(x)}} T\tilde{V}_{f(x)}$. Then in local bundle charts, for $(\tilde{y}_i, \tilde{\xi}_i) \in \left(\tilde{U}_x \times_{\tilde{V}_{f(x)}} T\tilde{V}_{f(x)} \right)_i$,

$$\begin{aligned} (\tilde{y}_i, \tilde{\xi}_i) \sim (\tilde{y}_j, \tilde{\xi}_j) &\iff \\ \tilde{y}_i = \tilde{y}_j \text{ and,} & \end{aligned}$$

$$\Theta_{f,x,i}(\gamma) \cdot \tilde{\xi}_i = \Theta_{f,x,j}(\gamma) \cdot \tilde{\xi}_j \text{ for all } \gamma \in \Gamma_x.$$

There is an obvious projection map onto \mathcal{O} and the total space of $f^*(T\mathcal{P})$ is a bundle over \mathcal{O} . Note that there are standard continuous injections $\iota_i : \star f_i^*(T\mathcal{P}) \rightarrow f^*(T\mathcal{P})$ and that the bundle maps $F_i : \star f_i^*(T\mathcal{P}) \rightarrow T\mathcal{P}$ glue together to give a continuous bundle map $F : f^*(T\mathcal{P}) \rightarrow T\mathcal{P}$ satisfying $F \circ \iota_i = F_i$.

We also define for $J \subset \{1, \dots, k\}$ the suborbifold $f^*(T\mathcal{P})_J$ of $f^*(T\mathcal{P})$ by

$$f^*(T\mathcal{P})_J = \bigsqcup_{i \in J} \star f_i^*(T\mathcal{P}_J) / \sim$$

where $T\mathcal{P}_J$ is the subspace of $T\mathcal{P}$ covered in bundle charts by

$$(T\tilde{V}_{f(x)})_J = \left\{ (\tilde{f}_x(\tilde{y}), \tilde{\xi}) \in T\tilde{V}_{f(x)} \mid \Theta_{f,x,i}(\gamma) \cdot \tilde{\xi} = \Theta_{f,x,j}(\gamma) \cdot \tilde{\xi} \text{ for all } i, j \in J \right\}.$$

Finally, let

$$\begin{aligned} \mathcal{D}_{\text{Orb}}^r(\star f_i^*(T\mathcal{P}))_J = \\ \{ \sigma \in \mathcal{D}_{\text{Orb}}^r(\star f_i^*(T\mathcal{P})) \mid \tilde{F}_{x,i} \circ \tilde{\sigma}_x(\tilde{y}) \in (T\tilde{V}_{f(x)})_J \text{ for all } \tilde{y} \in \tilde{U}_x \text{ and } x \in \mathcal{O} \}. \end{aligned}$$

Note that $\tilde{F}_{x,i} \circ \tilde{F}_{x,j}^{-1} : \mathcal{D}_{\text{Orb}}^r(\star f_j^*(T\mathcal{P}))_J \rightarrow \mathcal{D}_{\text{Orb}}^r(\star f_i^*(T\mathcal{P}))_J$ is a linear isomorphism of Banach (r finite)/Fréchet ($r = \infty$) subspaces for all $i, j \in J$ and $0 \leq r \leq \infty$. By abuse of notation, we write $\mathcal{D}_{\text{Orb}}^r(f^*(T\mathcal{P}))$ for the space of C^r orbisections $\sigma : \mathcal{O} \rightarrow f^*(T\mathcal{P})$ equipped with the C^r topology. From the construction of $f^*(T\mathcal{P})$ it is clear that the Riemannian exponential map on $T\mathcal{P}$ induces a map $E : \mathcal{D}_{\text{Orb}}^r(f^*T\mathcal{P}) \rightarrow C_{\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ as in proposition 40. For $f \in C_{\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ and $\star f_i \in C_{\star\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ mapping to f we let $\Theta(f)_x = \{\Theta_{f,x,i}\}$ where $\star f_i = (f, \{\tilde{f}_x\}, \{\Theta_{f,x,i}\})$.

Lemma 46. *There are neighborhoods $\mathcal{B}_f^r(\mathbf{0}, \varepsilon)$ of $\mathbf{0} \in \mathcal{D}_{\text{Orb}}^r(f^*T\mathcal{P})$ and $\mathcal{N}^r(f, \varepsilon)$ of f in $C_{\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ so that $E : \mathcal{B}_f^r(\mathbf{0}, \varepsilon) \rightarrow \mathcal{N}^r(f, \varepsilon)$ is a homeomorphism.*

Proof. The proof follows from observing that $\{g \in C_{\text{Orb}}^r(\mathcal{O}, \mathcal{P}) \mid \Theta(g)_x \subset \Theta(f)_x \text{ for all } x \in \mathcal{O}\}$ is an open subset (since the homomorphisms $\Theta_{f,x}$ are locally constant). By theorem 1, there is a neighborhood of each $\star f_i$ for which the map E of proposition 40 is a homeomorphism. By taking $g \in C_{\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ as above and sufficiently C^r close to f , all of its preimages $\star g_j = (g, \{\tilde{g}_x\}, \{\Theta_{g,x,j}\})$ will lie in such neighborhoods. \square

9. SOME INFINITE-DIMENSIONAL ANALYSIS

In this section we recall the results of global analysis that we need in order to substantiate our various claims of smoothness. For finite order differentiability, a strong argument can be made that the Lipschitz categories Lip^r are better suited to questions of calculus than the more common C^r category [FK88]. For our purposes, however, we have chosen to use the C^r category for questions of finite order differentiability, and for questions of infinite order differentiability, we use the convenient calculus as detailed in the monographs [FK88, KM97].

Review of the convenient calculus. For any topological vector space E the notion of a smooth curve $c \in C^\infty(\mathbb{R}, E)$ makes sense using the usual difference quotient and iterating. A mapping $f : E \rightarrow F$ between locally convex vector spaces is called *smooth* if it maps smooth curves to smooth curves. That is, if $f \circ c \in C^\infty(\mathbb{R}, F)$ for all $c \in C^\infty(\mathbb{R}, E)$. For E, F finite dimensional this yields the usual notion of (C^∞ -) smoothness. Unfortunately, such a characterization fails for finite order (C^r -) differentiability [Bom67]. Generalizing the fact that a map f between finite dimensional vector spaces is smooth if and only if its component functions are smooth, Frölicher, Kriegl and Michor [FK88, KM97] introduce the notion of a *convenient* vector space: A locally convex vector space is *convenient* if every scalarwise smooth curve $c : \mathbb{R} \rightarrow E$ is smooth. c is a scalarwise smooth curve if $\ell \circ c : \mathbb{R} \rightarrow \mathbb{R}$ is smooth for all continuous linear functionals ℓ on E . For our purposes, we remark that if E is a Fréchet space then E is convenient and the locally convex topology agrees with the c^∞ -topology or final topology with respect to the set of mappings $C^\infty(\mathbb{R}, E)$. It then follows from these definitions that smooth

mappings between Fréchet (or, convenient vector) spaces are continuous. At this point, one can work on open subsets of convenient vector spaces and introduce the notions of smooth manifold, smooth tangent bundle and smooth Lie group modeled on convenient vector spaces in a straightforward way.

An important feature of the convenient setting is the following theorem on Cartesian closedness:

Theorem 47 ([KM97, Theorem 3.12]). *Let $A_i \subset E_i$ be c^∞ -open subsets in locally convex spaces, which need not be c^∞ -complete. Then a mapping $f : A_1 \times A_2 \rightarrow F$ is smooth if and only if the canonically associated mapping $f^\vee : A_1 \rightarrow C^\infty(A_2, F)$ exists and is smooth.*

Our first use of the convenient calculus will be to substantiate the smoothness claims of Theorem 1 completing the proof. Throughout the remainder, we assume, as in section 7, that all orbifolds are C^∞ with C^∞ Riemannian metric. Further, the orbifold \mathcal{O} will be compact (without boundary).

Lemma 48. $C_{*\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ has the structure of a smooth (C^∞) Banach/Fréchet manifold.

Proof. We have already shown in section 7 that $C_{*\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ has the required structure as a topological Banach/Fréchet manifold. Let $*f \in C_{*\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ and let $E_{*f} : \mathcal{B}_{*f}^r(\mathbf{0}, \varepsilon') \rightarrow \mathcal{N}^r(*f, \varepsilon')$ be a manifold chart about $*f$ given by proposition 43. Let $*g \in \mathcal{N}^r(*f, \varepsilon')$ and choose $0 < \varepsilon < \varepsilon'$ so that the manifold chart $E_{*g} : \mathcal{B}_{*g}^r(\mathbf{0}, \varepsilon) \rightarrow \mathcal{N}^r(*g, \varepsilon)$ is contained entirely within $\mathcal{N}^r(*f, \varepsilon')$. Then the chart transition map

$$E_{*f}^{-1} \circ E_{*g} : \mathcal{B}_{*g}^r(\mathbf{0}, \varepsilon) \rightarrow \left(E_{*f}^{-1} \circ E_{*g} \right) \left(\mathcal{B}_{*g}^r(\mathbf{0}, \varepsilon) \right) \subset \mathcal{B}_{*f}^r(\mathbf{0}, \varepsilon')$$

is a homeomorphism between open subsets of Banach/Fréchet spaces. Thus to show smoothness, we need to show that $E_{*f}^{-1} \circ E_{*g}$ takes smooth curves to smooth curves. So, let $\sigma^t : \mathbb{R} \rightarrow \mathcal{B}_{*g}^r(\mathbf{0}, \varepsilon) \subset C_{*\text{Orb}}^r(\mathcal{O}, *g^*(T\mathcal{P}))$ be a smooth curve and let $\tilde{\sigma}_x^t : (0, 1) \rightarrow \mathcal{D}^r(*g^*(T\tilde{V}_{g(x)})) \subset C^r(\tilde{U}_x, \tilde{U}_x \times_{\tilde{V}_{g(x)}} T\tilde{V}_{g(x)})$ be a smooth local equivariant lift over an orbifold chart U_x . The interval $(0, 1)$ is being chosen for convenience to make clear that we want the image of the lift $\tilde{\sigma}_x^t$ to lie in a single trivializing bundle chart. The key observation is that the computations of difference quotients for σ^t are identical in the local lift $\tilde{\sigma}_x^t$ since an orbisection must take values in the admissible bundle $A(*g^*(T\mathcal{P}))$ whose fibers are the vector spaces fixed by the action of the local isotropy subgroups Γ_x (section 6). In particular, it follows that σ^t is smooth if and only if each local lift $\tilde{\sigma}_x^t$ is smooth. Using lemma 47, it follows that $\tilde{\sigma}_x^\wedge(t, \tilde{y}) : (0, 1) \times \tilde{U}_x \rightarrow \tilde{U}_x \times_{\tilde{V}_{g(x)}} T\tilde{V}_{g(x)}$ is smooth. From the formulas in section 7 for E and its local lifts and using the (C^∞ -) smoothness of the Riemannian exponential map, it follows that the map $\tilde{\eta}_x^\wedge(t, \tilde{y}) = \tilde{E}_{*f,x}^{-1} \circ \tilde{E}_{*g,x} \circ \tilde{\sigma}_x^\wedge(t, \tilde{y}) : (0, 1) \times \tilde{U}_x \rightarrow \tilde{U}_x \times_{\tilde{V}_{f(x)}} T\tilde{V}_{f(x)}$ is smooth. Another application of lemma 47 implies that $\tilde{\eta}_x^t = \tilde{E}_{*f,x}^{-1} \circ \tilde{E}_{*g,x} \circ \tilde{\sigma}_x^t : (0, 1) \rightarrow \mathcal{D}^r(*f^*(T\tilde{V}_{f(x)})) \subset C^r(\tilde{U}_x, \tilde{U}_x \times_{\tilde{V}_{f(x)}} T\tilde{V}_{f(x)})$ is smooth. Thus, by our earlier observation, $\eta^t = \left(E_{*f}^{-1} \circ E_{*g} \right) (\sigma^t) : \mathbb{R} \rightarrow \mathcal{B}_{*f}^r(\mathbf{0}, \varepsilon') \subset C_{*\text{Orb}}^r(\mathcal{O}, *f^*(T\mathcal{P}))$ is a smooth curve. \square

A useful consequence of the observation made in lemma 48 is the following (compare [KM97, Lemma 42.5]):

Corollary 49. *The following conditions on a curve $c = (f^t, \{\tilde{f}_x^t\}, \{\Theta_{f^t, x}\}) : \mathbb{R} \rightarrow C_{\star\text{Orb}}^r(\mathcal{O}, \mathcal{P})$ are equivalent:*

- (1) c is smooth
- (2) each local equivariant lift $\tilde{f}_x^t : (0, 1) \rightarrow C^r(\tilde{U}_x, \tilde{V}_z)$ is smooth
- (3) each local equivariant lift $\tilde{f}^\wedge(t, x) : (0, 1) \times \tilde{U}_x \rightarrow \tilde{V}_z$ is smooth

Proof. Note that the interval $(0, 1)$ is being chosen for convenience to make clear that we want the image of the lift \tilde{f}_x^t to lie in a single orbifold chart. (1) \iff (2) follows from the observation that smoothness of c is equivalent in local charts to smoothness of a curve into a space of orbisections. This in turn is equivalent to smoothness of the local equivariant lifts of the orbisections as in lemma 48. This, of course, is equivalent to smoothness of the local equivariant lifts \tilde{f}_x^t . (2) \iff (3) follows from theorem 47. \square

Lemma 50. *Let $I : \mathcal{N}^s(\star f, \varepsilon) \rightarrow \mathcal{N}^s(I \circ \star f, \varepsilon)$, $\star g \mapsto I \circ \star g$, be the homeomorphism from lemma 21. Then I is a C^∞ diffeomorphism.*

Proof. By lemma 21 and theorem 1 via lemma 48, we know that I is a map between open subsets of a smooth C^∞ Banach/Fréchet manifold. By [KM97, section 27], it suffices to show that

$$\sigma^t = E_{I \circ \star f}^{-1} \circ I \circ \star g^t : \mathbb{R} \rightarrow \mathcal{D}_{\text{Orb}}^s((I \circ \star f)^*(T\mathcal{P}))$$

is smooth for any smooth curve $\star g^t = (g^t, \{\tilde{g}_x^t\}, \{\Theta_{g^t, x}\})$. From an argument similar to that given in lemma 48, it follows that each local equivariant lift $\tilde{g}_x^t : (0, 1) \rightarrow C^s(\tilde{U}_x, \tilde{V}_{f(x)})$ is smooth. Using the discussion after example 8, for $I = (\text{Id}, \{\eta_x \cdot \tilde{y}\})$ we have

$$I \circ \star g^t = (g^t, \{\eta_x \cdot \tilde{g}_x^t\}, \{\gamma \mapsto \eta_x \Theta_{g^t, x}(\gamma) \eta_x^{-1}\}).$$

Then by the formulas in proposition 42, σ^t has local equivariant lifts

$$\begin{aligned} \tilde{\sigma}_x^t(\tilde{y}) &= (\tilde{y}, \tilde{\xi}^t(\tilde{y})) = \\ &= \left(\tilde{y}, \left[\eta_x \cdot \tilde{f}_x(\tilde{y}), \widetilde{\exp}_{\tilde{V}_{f(x), \eta_x \cdot \tilde{f}_x(\tilde{y})}^{-1}}(\eta_x \cdot \tilde{g}_x^t(\tilde{y})) \right] \right) \in (I \circ \star f)^*(T\tilde{V}_{f(x)}). \end{aligned}$$

Since the exponential map and action of local isotropy subgroups are (C^∞ -) smooth in charts and \tilde{g}_x^t is smooth, we see that $\tilde{\sigma}_x^t$ is smooth. It then follows that σ^t is smooth by corollary 49 or the observation in the proof of lemma 48, and this completes the proof. \square

The next result we will need is that composition in our spaces of smooth orbifold maps is smooth.

Lemma 51. *Let \mathcal{O} , \mathcal{P} and \mathcal{R} be smooth C^∞ compact orbifolds without boundary. Then the composition mappings*

$$\begin{aligned} \text{comp} : C_{\star\text{Orb}}^\infty(\mathcal{P}, \mathcal{R}) \times C_{\star\text{Orb}}^\infty(\mathcal{O}, \mathcal{P}) &\rightarrow C_{\star\text{Orb}}^\infty(\mathcal{O}, \mathcal{R}), & (\star f, \star g) &\mapsto \star f \circ \star g, \\ \text{comp} : C_{\blacklozenge\text{Orb}}^\infty(\mathcal{P}, \mathcal{R}) \times C_{\blacklozenge\text{Orb}}^\infty(\mathcal{O}, \mathcal{P}) &\rightarrow C_{\blacklozenge\text{Orb}}^\infty(\mathcal{O}, \mathcal{R}), & (\blacklozenge f, \blacklozenge g) &\mapsto \blacklozenge f \circ \blacklozenge g, \\ \text{comp} : C_{\text{Orb}}^\infty(\mathcal{P}, \mathcal{R}) \times C_{\text{Orb}}^\infty(\mathcal{O}, \mathcal{P}) &\rightarrow C_{\text{Orb}}^\infty(\mathcal{O}, \mathcal{R}), & (f, g) &\mapsto f \circ g, \\ \text{comp} : C_{\text{red}}^\infty(\mathcal{P}, \mathcal{R}) \times C_{\text{red}}^\infty(\mathcal{O}, \mathcal{P}) &\rightarrow C_{\text{red}}^\infty(\mathcal{O}, \mathcal{R}), & (\bullet f, \bullet g) &\mapsto \bullet f \circ \bullet g, \end{aligned}$$

are smooth.

Proof. By lemmas 9, 11, 21, and the observation following definition 20, it suffices to prove the result for the complete orbifold maps. We use an argument analogous to [KM97, Theorem 42.13]. Namely, let $(c_1, c_2) : \mathbb{R} \rightarrow C_{\star\text{Orb}}^{\infty}(\mathcal{P}, \mathcal{R}) \times C_{\star\text{Orb}}^{\infty}(\mathcal{O}, \mathcal{P})$ be a smooth curve. Then $(\text{comp} \circ (c_1, c_2))(t)(x) = c_1^{\wedge}(t, c_2^{\wedge}(t, x))$ is smooth by corollary 49. Hence comp is smooth. \square

Finally, for finite order differentiability, we will need to refer to the Ω -lemma of Palais [Pal68] as stated in [Sch04]:

Lemma 52 (Ω -lemma). *Let M be a C^{∞} compact manifold and let $\tau : E \rightarrow M$ and $\tau' : E' \rightarrow M$ be C^r vector bundles over M . Let $U \subset E$ be open and let $\omega : U \subset E \rightarrow E'$ be a C^{∞} vector bundle map. Then the induced map*

$$\Omega_{\omega} : \mathcal{D}^r(U) \subset \mathcal{D}^r(\tau) \rightarrow \mathcal{D}^r(\tau'), \quad \Omega_{\omega}(\xi) = \omega \circ \xi$$

is a C^{∞} map. If ω is only C^{r+k} , then Ω_{ω} is C^k .

REFERENCES

- [ALR07] Alejandro Adem, Johann Leida, and Yongbin Ruan, *Orbifolds and stringy topology*, Cambridge Tracts in Mathematics, vol. 171, Cambridge University Press, Cambridge, 2007. MR 2359514
- [Ban97] Augustin Banyaga, *The structure of classical diffeomorphism groups*, Mathematics and its Applications, vol. 400, Kluwer Academic Publishers Group, Dordrecht, 1997. MR 1445290 (98h:22024)
- [Bom67] Jan Boman, *Differentiability of a function and of its compositions with functions of one variable*, Math. Scand. **20** (1967), 249–268. MR 0237728 (38 #6009)
- [Bor92] Joseph E. Borzellino, *Riemannian Geometry of Orbifolds*, University of California, Los Angeles, June 1992.
- [Bor93] ———, *Orbifolds of maximal diameter*, Indiana Univ. Math. J. **42** (1993), no. 1, 37–53. MR 1218706 (94d:53053)
- [BB02] Joseph E. Borzellino and Victor Brunsten, *Orbifold homeomorphism and diffeomorphism groups*, (Washington, DC, 2000), World Sci. Publ., River Edge, NJ, 2002, pp. 116–137. MR 1926425 (2003g:58013)
- [BB08] ———, *A manifold structure for the group of orbifold diffeomorphisms of a smooth orbifold*, J. Lie Theory **18** (2008), no. 4, 979–1007, available at [arXiv:math.DG/0608526](https://arxiv.org/abs/math/0608526).
- [Bou75] Jean-Pierre Bourguignon, *Une stratification de l'espace des structures riemanniennes*, Compositio Math. **30** (1975), 1–41 (French). MR 0418147 (54 #6189)
- [CR02] Weimin Chen and Yongbin Ruan, *Orbifold Gromov-Witten theory*, Orbifolds in mathematics and physics (Madison, WI, 2001), Contemp. Math., vol. 310, Amer. Math. Soc., Providence, RI, 2002, pp. 25–85. MR 1950941 (2004k:53145)
- [Che06] Weimin Chen, *On a notion of maps between orbifolds. I. Function spaces*, Commun. Contemp. Math. **8** (2006), no. 5, 569–620. MR 2263948 (2007f:57055)
- [Eel66] James Eells Jr., *A setting for global analysis*, Bull. Amer. Math. Soc. **72** (1966), 751–807. MR 0203742 (34 #3590)
- [Fie70] Mike Field, *Equivariant dynamical systems*, Bull. Amer. Math. Soc. **76** (1970), 1314–1318. MR 0277850 (43 #3583)
- [Fis70] Arthur Elliot Fischer, *The theory of superspace*, Relativity (Proc. Conf. Midwest, Cincinnati, Ohio, 1969), Plenum, New York, 1970, pp. 303–357. MR 0347323 (49 #12043)
- [Fra79] John Franks, *Manifolds of C^r mappings and applications to differentiable dynamical systems*, Studies in analysis, Adv. in Math. Suppl. Stud., vol. 4, Academic Press, New York, 1979, pp. 271–290. MR 546811 (81a:58018)
- [FK88] Alfred Frölicher and Andreas Kriegl, *Linear spaces and differentiation theory*, Pure and Applied Mathematics (New York), John Wiley & Sons Ltd., Chichester, 1988. A Wiley-Interscience Publication. MR 961256 (90h:46076)
- [Ham82] Richard S. Hamilton, *The inverse function theorem of Nash and Moser*, Bull. Amer. Math. Soc. (N.S.) **7** (1982), no. 1, 65–222. MR 656198 (83j:58014)

- [Kel74] Hans Heinrich Keller, *Differential calculus in locally convex spaces*, Lecture Notes in Mathematics, Vol. 417, Springer-Verlag, Berlin, 1974. MR 0440592 (55 #13466)
- [KM97] Andreas Kriegl and Peter W. Michor, *The convenient setting of global analysis*, Mathematical Surveys and Monographs, vol. 53, American Mathematical Society, Providence, RI, 1997. MR **1471480** (**98i**:58015)
- [Moe02] Ieke Moerdijk, *Orbifolds as groupoids: an introduction*, Orbifolds in mathematics and physics (Madison, WI, 2001), Contemp. Math., vol. 310, Amer. Math. Soc., Providence, RI, 2002, pp. 205–222. MR **1950948** (**2004c**:22003)
- [MP97] I. Moerdijk and D. A. Pronk, *Orbifolds, sheaves and groupoids*, *K-Theory* **12** (1997), no. 1, 3–21. MR **1466622** (**98i**:22004)
- [Nit71] Zbigniew Nitecki, *Differentiable dynamics. An introduction to the orbit structure of diffeomorphisms*, The M.I.T. Press, Cambridge, Mass.-London, 1971. MR 0649788 (58 #31210)
- [Pal68] Richard S. Palais, *Foundations of global non-linear analysis*, W. A. Benjamin, Inc., New York-Amsterdam, 1968. MR 0248880 (40 #2130)
- [Sch04] Rudolf Schmid, *Infinite dimensional Lie groups with applications to mathematical physics*, *J. Geom. Symmetry Phys.* **1** (2004), 54–120. MR **2096566** (**2006a**:22019)
- [Thu78] William Thurston, *The geometry and topology of 3-manifolds*, Princeton University Mathematics Department, 1978. Lecture Notes.

DEPARTMENT OF MATHEMATICS, CALIFORNIA POLYTECHNIC STATE UNIVERSITY, 1 GRAND AVENUE, SAN LUIS OBISPO, CALIFORNIA 93407
E-mail address: jborzell@calpoly.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, PENN STATE ALTOONA, 3000 IVYSIDE PARK, ALTOONA, PENNSYLVANIA 16601
E-mail address: vw2@psu.edu