# Towards a three dimensional solution for 3N bound states with 3N forces

M. R. Hadizadeh<sup>\*</sup> and S. Bayegan<sup>†</sup>

Department of Physics, University of Tehran, P.O.Box 14395-547, Tehran, Iran

(Dated: November 25, 2018)

### Abstract

After a brief discussion about the necessity of using the 3D approach, we present the non PW formalism for 3N bound state with the inclusion of 3N force (3NF). As an example the evaluation of 3NF matrix elements, which appear in the obtained coupled three dimensional integral equations, for  $2\pi$ -exchange Tucson-Melbourne 3NF show how would be this formalism efficient and less cumbersome in comparison with the PW formalism.

PACS numbers: 21.45.-v, 21.30.-x, 21.10.Dr, 27.10.+h, 21.10.Hw

Keywords: three dimensional approach; 3N bound state; Tucson-Melbourne 3N force.

<sup>\*</sup>Electronic address: hadizade@khayam.ut.ac.ir

<sup>&</sup>lt;sup>†</sup>Electronic address: bayegan@khayam.ut.ac.ir

### I. WHY DO WE USE 3D APPROACH INSTEAD OF PW APPROACH?

The answer to this main question indicates the motivation for using this approach. Fewbody calculations are traditionally carried out by solving the relevant equations in a PW basis which after truncation they lead to coupled equations on angular momentum quantum numbers. A few PWs often provide qualitative insight, but modern calculations need many different spin, isospin and angular momentum combinations. It is clear that in PW approach one should sum all PWs to infinite order, but in practice one truncates the sum to a finite angular momentum number which is dependent to the energy that one is working. It means that in higher energies one will need more PWs to obtain a convergence. In contrast to the traditional PW representation, the novel 3D approach replaces the discrete angular momentum quantum numbers with continuous angle variables and consequently it considers automatically all PW components to infinite order [1]-[13]. So the number of equations in 3D formalism is energy independent, also this formalism avoids the very involved angular momentum algebra occurring for the permutations, transformations and especially for the 3NFs.

### **II. 3D REPRESENTATION OF FADDEEV EQUATIONS WITH 3NF**

We have recently applied the 3D approach to the 3N bound state, where the Faddeev equations with NN interactions are successfully solved with Bonn-B potential [10]. In this article we extend this formalism by considering the 3NF, which helps us to reach to the full solution of the 3N bound state in a straightforward manner. The 3N bound state in the presence of the 3NF is described by the Faddeev equation:

$$|\psi\rangle = G_0 t P |\psi\rangle + (1 + G_0 t) G_0 V_{123}^{(3)} |\Psi\rangle, \qquad (1)$$

where the quantity  $V_{123}^{(3)}$  defines a part of the 3NF which is symmetric under the exchange of the particles 1 and 2. In order to solve the Eq. (1) in momentum space we introduce the 3N basis states in a 3D formalism as:

$$|\mathbf{p}\mathbf{q}\,\alpha\rangle \equiv |\mathbf{p}\mathbf{q}\,\alpha_S\,\alpha_T\rangle \equiv |\mathbf{p}\mathbf{q}\,(s_{12}\,\frac{1}{2})S\,M_S\,(t_{12}\,\frac{1}{2})T\,M_T\rangle.$$
(2)

Evaluation of the transition and permutation operators need to the free 3N basis states  $|\mathbf{p}\mathbf{q}\gamma\rangle$ , where the spin-isospin parts  $\gamma$  are given as:  $|\gamma\rangle \equiv |\gamma_S\gamma_T\rangle \equiv$   $|m_{s_1} m_{s_2} m_{s_3} m_{t_1} m_{t_2} m_{t_3} \rangle$ . To this aim by changing the 3N basis states  $|\alpha\rangle$  to the free 3N basis states  $|\gamma\rangle$  one needs to calculate the usual Clebsch-Gordan coefficients  $\langle \gamma | \alpha \rangle = g_{\gamma\alpha} \equiv g_{\gamma\alpha}^S g_{\gamma\alpha}^T$ , see appendix (A) of Ref. [10]. The evaluation of Faddeev equation with the inclusion of 3NF will be exactly the same as Eq. (19) of Ref. [10] except that an extra term with  $V_{123}^{(3)}$ . This is

$$\langle \mathbf{p} \mathbf{q} \alpha | (1 + G_0 t) G_0 V_{123}^{(3)} | \Psi \rangle = \frac{1}{E - \frac{p^2}{m} - \frac{3q^2}{4m}} \bigg\{ \langle \mathbf{p} \mathbf{q} \alpha | V_{123}^{(3)} | \Psi \rangle + \langle \mathbf{p} \mathbf{q} \alpha | t G_0 V_{123}^{(3)} | \Psi \rangle \bigg\}.$$
(3)

The matrix elements of the second term can be evaluated as:

$$\langle \mathbf{p} \mathbf{q} \alpha | t G_0 V_{123}^{(3)} | \Psi \rangle = \sum_{\gamma', \gamma'', \alpha'''} \int d^3 p''' \int d^3 q''' \frac{g_{\gamma'\alpha} g_{\gamma''\alpha'''}}{E - \frac{p'''^2}{m} - \frac{3q'''^2}{4m}} \\ \times \langle \mathbf{p} \mathbf{q} \gamma' | t | \mathbf{p}''' \mathbf{q}''' \gamma'' \rangle \langle \mathbf{p}''' \mathbf{q}''' \alpha''' | V_{123}^{(3)} | \Psi \rangle,$$

$$(4)$$

after evaluating the matrix elements of the NN t-matrix and integrating over  $\mathbf{q}'''$  vector, one obtains:

$$\langle \mathbf{p} \mathbf{q} \alpha | t G_0 V_{123}^{(3)} | \Psi \rangle = \sum_{\gamma', \gamma'', \alpha'''} \int d^3 p''' \frac{g_{\gamma'\alpha} g_{\gamma''\alpha'''}}{E - \frac{p'''^2}{m} - \frac{3q^2}{4m}} \delta_{m'_{s_3} m''_{s_3}} \delta_{m'_{t_3} m''_{t_3}} \\ \times \langle \mathbf{p} m'_{s_1} m'_{s_2} m'_{t_1} m'_{t_2} | t(\epsilon) | \mathbf{p}''' m''_{s_1} m''_{s_2} m''_{t_1} m''_{t_2} \rangle \langle \mathbf{p}''' \mathbf{q} \alpha''' | V_{123}^{(3)} | \Psi \rangle.$$
 (5)

By using the symmetry property of 3NF and the anti-symmetry property of the total wave function under exchange of nucleons 1 and 2, one can rewrite Eq. (5) as:

$$\langle \mathbf{p} \mathbf{q} \alpha | t G_0 V_{123}^{(3)} | \Psi \rangle = \sum_{\gamma', \gamma'', \alpha'''} \int d^3 p''' \frac{g_{\gamma'\alpha} g_{\gamma''\alpha'''} \,\delta_{m'_{s_3}m''_{s_3}} \,\delta_{m'_{t_3}m''_{t_3}}}{E - \frac{p'''^2}{m} - \frac{3q^2}{4m}} \left( -(-)^{s'''_{12} + t''_{12}} \right) \\ \times \langle \mathbf{p} m'_{s_1} m'_{s_2} m'_{t_1} m'_{t_2} | t(\epsilon) P_{12} | - \mathbf{p}''' m''_{s_2} m''_{s_1} m''_{t_2} m''_{t_1} \rangle \langle -\mathbf{p}''' \mathbf{q} \alpha''' | V_{123}^{(3)} | \Psi \rangle,$$
 (6)

under the exchange of the labels  $m''_{s_1}, m''_{t_1}$  to  $m''_{s_2}, m''_{t_2}$ , reverse of it and changing  $\mathbf{p}'''$  to  $-\mathbf{p}'''$  one finds:

$$\langle \mathbf{p} \mathbf{q} \alpha | t G_0 V_{123}^{(3)} | \Psi \rangle = \sum_{\gamma', \gamma'', \alpha'''} \int d^3 p''' \frac{g_{\gamma'\alpha} g_{\gamma''\alpha''}}{E - \frac{p'''^2}{m} - \frac{3q^2}{4m}} \, \delta_{m'_{s_3} m''_{s_3}} \, \delta_{m'_{t_3} m''_{t_3}} \\ \times \langle \mathbf{p} m'_{s_1} m'_{s_2} m'_{t_1} m'_{t_2} | - t(\epsilon) P_{12} | \mathbf{p}''' m''_{s_1} m''_{s_2} m''_{t_1} m''_{t_2} \rangle \, \langle \mathbf{p}''' \mathbf{q} \, \alpha''' \, | V_{123}^{(3)} | \Psi \rangle.$$
 (7)

Now one can consider Eqs. (5) and (7) together to achieve:

by applying the introduction of the anti-symmetrized NN *t*-matrix one can obtain the coupled three-dimensional Faddeev integral equations as:

$$\langle \mathbf{p} \mathbf{q} \alpha | \psi \rangle = \frac{1}{E - \frac{p^2}{m} - \frac{3q^2}{4m}} \\ \times \left[ \int d^3 q' \sum_{\gamma', \gamma'', \alpha''} g_{\alpha \gamma'''} g_{\gamma' \alpha''} \delta_{m_{s_3}''' m_{s_1}'} \delta_{m_{t_3}''' m_{t_1}'''} \\ \times {}_a \langle \mathbf{p} \, m_{s_1}''' m_{s_2}''' m_{t_1}''' m_{t_2}''' | t(\epsilon) | \frac{-1}{2} \mathbf{q} - \mathbf{q}' \, m_{s_2}' m_{s_3}' m_{t_2}' m_{t_3}' \rangle_a \, \langle \mathbf{q} + \frac{1}{2} \mathbf{q}' \, \mathbf{q}' \, \alpha'' | \psi \rangle \\ + \left\{ \left. \langle \mathbf{p} \, \mathbf{q} \, \alpha \, | V_{123}^{(3)} | \Psi \right\rangle \right. \\ \left. + \frac{1}{2} \sum_{\gamma', \gamma'', \alpha'''} g_{\alpha \gamma'} g_{\gamma'' \alpha'''} \int d^3 p' \, \frac{\delta_{m_{s_3}'' m_{s_3}'} \delta_{m_{t_3}' m_{t_3}''}}{E - \frac{p'^2}{m} - \frac{3q^2}{4m}} \\ \left. \times {}_a \langle \mathbf{p} \, m_{s_1}' m_{s_2}' \, m_{t_1}' m_{t_2}' | t(\epsilon) | \mathbf{p}' \, m_{s_1}'' m_{s_2}'' \, m_{t_1}'' m_{t_1}' \rangle_a \, \langle \mathbf{p}' \, \mathbf{q} \, \alpha''' | V_{123}^{(3)} | \Psi \rangle \right\} \right].$$

To represent the generality of our 3D formalism we can simplify the Eq. (9) to the bosonic case by switching off the spin-isospin quantum numbers, see Ref. [3]. In order to show the efficiency of the presented formalism the realistic  $2\pi$ -exchange TM 3NF [15] has been used to evaluate the matrix elements of  $\langle \mathbf{p} \mathbf{q} \alpha | V_{123}^{(3)} | \Psi \rangle$ .

## III. THE EVALUATION OF $\langle \, {f p} \, {f q} \, lpha \, | V^{(3)}_{123} | \Psi angle$ for the TM 3NF

For evaluation of the coupled three dimensional Faddeev equations, the matrix elements  $\langle \mathbf{p} \mathbf{q} \alpha | V_{123}^{(3)} | \Psi \rangle$  need to be calculated. In this section these matrix elements have been evaluated for the TM 2 $\pi$ -exchange 3NF. To this aim one should first prepare this force in the form that can be evaluated easily in 3D representation as:

$$V_{123}^{(3)} = V_0 Q \ Q' \ \sigma_1 \cdot \widehat{\mathbf{Q}} \ \sigma_2 \cdot \widehat{\mathbf{Q}}' \frac{F(Q^2)}{Q^2 + m_\pi^2} \frac{F(Q'^2)}{Q'^2 + m_\pi^2}$$

$$\times \left[ \tau_1 \cdot \tau_2 \left( A + B \ Q Q' \gamma + C \ (Q^2 + Q'^2) \right) + D \ \tau_3 \cdot \tau_1 \times \tau_2 \left\{ \frac{3ia}{4\gamma} - \frac{i}{2\gamma a} \ (\sigma_3 \cdot \widehat{\mathbf{Q}}')^2 + \frac{i}{2a} \ \sigma_3 \cdot \widehat{\mathbf{Q}}' \ \sigma_3 \cdot \widehat{\mathbf{Q}} + \frac{i}{8\gamma a} \ (\sigma_3 \cdot \widehat{\mathbf{Q}}')^2 \ (\sigma_3 \cdot \widehat{\mathbf{Q}})^2 - \frac{i}{2\gamma a} \ (\sigma_3 \cdot \widehat{\mathbf{Q}})^2 \right\} \right], \quad (10)$$

where  $\gamma = \widehat{\mathbf{Q}} \cdot \widehat{\mathbf{Q}}'$  and  $a = \sqrt{1 - \gamma^2}$ . The  $\mathbf{Q}$  and  $\mathbf{Q}'$  are momentum transfers,  $\sigma_i$ 's are Pauli spin matrices and  $F(Q^2)$  are form factors. One distinguishes four A-, B-, C- and D-terms in the TM force. The scalar product of the spin-momentum vectors can be evaluated as:

$$\langle \hat{z} \, m'_s | \sigma \cdot \widehat{\mathbf{Q}} | \hat{z} \, m''_s \rangle = \sum_{m_s} m_s \, D^{\frac{1}{2}}_{m'_s m_s}(\widehat{\mathbf{Q}}) \quad D^{\frac{1}{2}*}_{m''_s m_s}(\widehat{\mathbf{Q}}) = \mathcal{O}^{\widehat{\mathbf{Q}}}_{m'_s m''_s}, \tag{11}$$

where  $D_{m'_s m_s}^{\frac{1}{2}}$  is Wigner D-function which is defined generally as  $D_{m'_s m_s}^s(\hat{q}) = \langle \hat{z}sm'_s | \hat{q}sm_s \rangle$ . The application of the TM 3NF to the total wave function  $|\Psi\rangle$  can be considered as sum of the the four independent terms:

$$V_{123}^{(3)}|\Psi\rangle = \sum_{i=A}^{D} V_{31}^{(i)} I^{(i)} V_{23}^{(i)}|\Psi\rangle = \sum_{i=A}^{D} |\psi^{i}\rangle, \quad I^{(i)} = \begin{cases} \tau_{1} \cdot \tau_{2} & \text{i=A, B, C} \\ \tau_{3} \cdot \tau_{1} \times \tau_{2} & \text{i=D.} \end{cases}$$
(12)

In the following the matrix elements of  $\langle \mathbf{p} \mathbf{q} \alpha | \psi^i \rangle$  terms have been evaluated. It is clear that  $V_{123}^{(3)}$  can be splitted into two parts and each part contains a meson exchange, the mesons are exchanged in the subsystems (31) and (23), which are called for convenience subsystem 2 and 1 correspondingly. So it is convenient to insert a complete set of states of type 2 between  $V_{123}^{(3)}$  and  $|\Psi\rangle$  and another complete set of states of type 1 between the two meson exchanges. Then the matrix elements of  $\psi^i$  can be written as:

$${}_{3}\langle \mathbf{p} \mathbf{q} \alpha | \psi^{i} \rangle = \sum_{\alpha'} \int d^{3}p' \int d^{3}q' {}_{3}\langle \mathbf{p} \mathbf{q} \alpha | \mathbf{p}' \mathbf{q}' \alpha' \rangle_{1} \\ \times \sum_{\alpha''} \int d^{3}p'' \int d^{3}q'' {}_{1}\langle \mathbf{p}' \mathbf{q}' \alpha' | V_{31}^{(i)} | \mathbf{p}'' \mathbf{q}'' \alpha'' \rangle_{1} \\ \times \sum_{\alpha'''} \int d^{3}p''' \int d^{3}q''' {}_{1}\langle \mathbf{p}'' \mathbf{q}'' \alpha'' | I^{(i)} | \mathbf{p}''' \mathbf{q}''' \alpha''' \rangle_{2} \\ \times \sum_{\alpha''''} \int d^{3}p'''' \int d^{3}q'''' {}_{2}\langle \mathbf{p}''' \mathbf{q}''' \alpha''' | V_{23}^{(i)} | \mathbf{p}'''' \alpha'''' \rangle_{2} \\ \times {}_{2}\langle \mathbf{p}'''' \mathbf{q}'''' \alpha''' | \Psi \rangle,$$
(13)

Here the subscripts 1, 2, 3 of the bra and ket vectors stand for different 3N chains. The coordinate transformation from the system of type 1 to one of type 3 can be evaluated explicitly as:

$${}_{3}\langle \mathbf{p}\mathbf{q}\alpha | \mathbf{p'}\mathbf{q'}\alpha' \rangle_{1} = g_{\alpha_{3}\alpha_{1}'}\delta^{3}(\mathbf{p'} + \frac{1}{2}\mathbf{p} + \frac{3}{4}\mathbf{q})\delta^{3}(\mathbf{q'} - \mathbf{p} + \frac{1}{2}\mathbf{q}).$$
(14)

The matrix elements of the isospin coordinate transformations can be evaluated as:

$${}_{1}\langle \mathbf{p}'' \mathbf{q}'' \alpha'' | I^{(i)} | \mathbf{p}''' \mathbf{q}''' \alpha''' \rangle_{2} = {}_{1}\langle \mathbf{p}'' \mathbf{q}'' \alpha''_{S} | \mathbf{p}''' \mathbf{q}''' \alpha'''_{S} \rangle_{2} {}_{1}\langle \alpha''_{T} | I^{(i)} | \alpha'''_{T} \rangle_{2},$$
(15)

$${}_{1}\langle \mathbf{p}'' \mathbf{q}'' \alpha_{S}'' | \mathbf{p}''' \mathbf{q}''' \alpha_{S}''' \rangle_{2} = g_{\alpha_{1}'' \alpha_{2}''}^{S} \delta^{3}(\mathbf{p}''' + \frac{1}{2}\mathbf{p}'' + \frac{3}{4}\mathbf{q}'')\delta^{3}(\mathbf{q}''' - \mathbf{p}'' + \frac{1}{2}\mathbf{q}'').$$
(16)

The matrix elements of  $_1\langle \alpha_T'' | I^{(i)} | \alpha_T''' \rangle_2$  have been derived in Ref. [14].

In the following the matrix elements for the different V's are evaluated. In the first step the matrix elements of  $V_{31}^i$  are evaluated. Since both pion-exchange propagators in the 3NF term depend only on the momentum transfer in a two-nucleon subsystem and also because of the separation of the isospin parts of 3NF, the matrix elements of  $V_{31}^i$  can be evaluated as:

$${}_{1}\langle \mathbf{p}' \mathbf{q}' \alpha' | V_{31}^{(i)} | \mathbf{p}'' \mathbf{q}'' \alpha'' \rangle_{1} = \delta^{3} (\mathbf{p}' - \mathbf{p}'') \, \delta_{\alpha'_{T} \alpha''_{T} \ 1} \langle \mathbf{q}' \alpha'_{S} | V_{31}^{(i)} | \mathbf{q}'' \alpha''_{S} \rangle_{1}, \tag{17}$$

and the spin-space parts can be more simplified for the A-,B-,C- and D-terms separately as:

$${}_{1} \langle \mathbf{q}' \, \alpha_{S}' | V_{31}^{(A,B,C)} | \mathbf{q}'' \, \alpha_{S}'' \rangle_{1} = \sum_{\gamma_{S}', \gamma_{S}''} g_{\alpha_{S}' \gamma_{S}'}^{1} g_{\alpha_{S}' \gamma_{S}''}^{1} \delta_{m_{s_{2}}'' m_{s_{2}}''} \delta_{m_{s_{3}}'' m_{s_{3}}''} \\ \times {}_{1} \langle \mathbf{q}' \, \alpha_{S}' | V_{31}^{(D)} | \mathbf{q}'' \, \alpha_{S}'' \rangle_{1} = \sum_{\gamma_{S}', \gamma_{S}''} g_{\alpha_{S}' \gamma_{S}'}^{1} g_{\alpha_{S}' \gamma_{S}''}^{1} \delta_{m_{s_{2}}'' m_{s_{1}}''} \\ \times {}_{1} \langle \mathbf{q}' \, m_{s_{1}}' \, m_{s_{1}}' \, N_{31}^{(D)} | \mathbf{q}'' \, m_{s_{1}}'' \, n_{s_{3}}'' \rangle_{1}.$$
(18)

The A-, B- and C-terms can be evaluated as:

$${}_{1}\langle \mathbf{q}' \, m_{s_{1}}' | V_{31}^{(A,B,C)} | \mathbf{q}'' \, m_{s_{1}}'' \rangle_{1} = \frac{F((\mathbf{q}' - \mathbf{q}'')^{2})}{(\mathbf{q}' - \mathbf{q}'')^{2} + m_{\pi}^{2}} | \mathbf{q}' - \mathbf{q}'' |^{N} \, \mathscr{O}_{m_{s_{1}}' m_{s_{1}}''}^{(\widehat{\mathbf{q}' - \mathbf{q}''})}, \tag{19}$$

where the value of N is 1, 2 and 3 corresponding to A-, B- and C-terms. For the different parts, five parts, of the D-term one obtains:

$${}_{1}\langle \mathbf{q}' \, m_{s_{1}}' \, m_{s_{3}}' | V_{31}^{(D)} | \mathbf{q}'' \, m_{s_{1}}'' \, m_{s_{3}}'' \rangle_{1} = {}_{1}\langle \mathbf{q}' \, m_{s_{1}}' | V_{31}^{(B)} | \mathbf{q}'' \, m_{s_{1}}'' \rangle_{1} \, \mathcal{O}_{m_{s_{3}}' m_{s_{3}}''}^{(\widehat{\mathbf{q}'-\mathbf{q}''}) \, M}, \tag{20}$$

where the value of M is 0 for the first and second parts of this term, it is 1 for the third part and it is 2 for the fourth and fifth parts. The matrix elements of  $V_{23}^i$  can be evaluated by following the same algorithm as above. So one can obtain:

$${}_{2}\langle \mathbf{p}^{\prime\prime\prime\prime} \mathbf{q}^{\prime\prime\prime\prime} \alpha^{\prime\prime\prime\prime} | V_{23}^{(i)} | \mathbf{p}^{\prime\prime\prime\prime\prime} \mathbf{q}^{\prime\prime\prime\prime\prime} \alpha^{\prime\prime\prime\prime\prime} \rangle_{2} = \delta^{3}(\mathbf{p}^{\prime\prime\prime\prime} - \mathbf{p}^{\prime\prime\prime\prime\prime}) \, \delta_{\alpha_{T}^{\prime\prime\prime\prime} \alpha_{T}^{\prime\prime\prime\prime} \, 2} \langle \mathbf{q}^{\prime\prime\prime\prime} \alpha_{S}^{\prime\prime\prime\prime} | V_{23}^{(i)} | \mathbf{q}^{\prime\prime\prime\prime\prime} \alpha_{S}^{\prime\prime\prime\prime} \rangle_{2},$$

$$(21)$$

$${}_{2}\langle \mathbf{q}^{\prime\prime\prime\prime} \alpha_{S}^{\prime\prime\prime\prime} | V_{23}^{(A,B,C)} | \mathbf{q}^{\prime\prime\prime\prime\prime} \alpha_{S}^{\prime\prime\prime\prime\prime} \rangle_{2} = \sum_{\gamma_{S}^{\prime\prime\prime\prime}, \gamma_{S}^{\prime\prime\prime\prime\prime}} g_{\alpha_{S}^{\prime\prime\prime\prime} \gamma_{S}^{\prime\prime\prime\prime}}^{2} g_{\alpha_{S}^{\prime\prime\prime\prime} \gamma_{S}^{\prime\prime\prime\prime}}^{2} \delta_{m_{s_{1}}^{\prime\prime\prime} m_{s_{1}}^{\prime\prime\prime\prime}} \delta_{m_{s_{3}}^{\prime\prime\prime\prime} m_{s_{3}}^{\prime\prime\prime\prime\prime}} \\ \times {}_{2}\langle \mathbf{q}^{\prime\prime\prime\prime} m_{s_{2}}^{\prime\prime\prime\prime} | V_{23}^{(A,B,C)} | \mathbf{q}^{\prime\prime\prime\prime\prime} m_{s_{2}}^{\prime\prime\prime\prime} \rangle_{2}, \qquad (22)$$

$${}_{2}\langle \mathbf{q}^{\prime\prime\prime\prime} \, \alpha_{S}^{\prime\prime\prime\prime} | V_{23}^{(D)} | \mathbf{q}^{\prime\prime\prime\prime\prime} \, \alpha_{S}^{\prime\prime\prime\prime\prime} \rangle_{2} = \sum_{\gamma_{S}^{\prime\prime\prime\prime}, \gamma_{S}^{\prime\prime\prime\prime}} g_{\alpha_{S}^{\prime\prime\prime\prime} \gamma_{S}^{\prime\prime\prime\prime}}^{2} g_{\alpha_{S}^{\prime\prime\prime\prime} \gamma_{S}^{\prime\prime\prime\prime}}^{2} \delta_{m_{s_{1}}^{\prime\prime\prime} m_{s_{1}}^{\prime\prime\prime\prime}} \\ \times {}_{2}\langle \mathbf{q}^{\prime\prime\prime\prime} \, m_{s_{2}}^{\prime\prime\prime\prime} \, m_{s_{3}}^{\prime\prime\prime\prime} | V_{23}^{(D)} | \mathbf{q}^{\prime\prime\prime\prime\prime} \, m_{s_{2}}^{\prime\prime\prime\prime} \, m_{s_{3}}^{\prime\prime\prime\prime} \rangle_{2}.$$
(23)

The A-, B- and C-terms which have been considered in the right side of Eq. (22) can be evaluated as:

$${}_{2}\langle \mathbf{q}^{\prime\prime\prime\prime} m_{s_{2}}^{\prime\prime\prime\prime} | V_{23}^{(A,B,C)} | \mathbf{q}^{\prime\prime\prime\prime\prime} m_{s_{2}}^{\prime\prime\prime\prime\prime} \rangle_{2} = \frac{F((\mathbf{q}^{\prime\prime\prime\prime\prime} - \mathbf{q}^{\prime\prime\prime\prime})^{2})}{(\mathbf{q}^{\prime\prime\prime\prime\prime} - \mathbf{q}^{\prime\prime\prime\prime})^{2} + m_{\pi}^{2}} | \mathbf{q}^{\prime\prime\prime\prime} - \mathbf{q}^{\prime\prime\prime\prime} |^{N} \mathcal{O}_{m_{s_{2}}^{\prime\prime\prime} m_{s_{2}}^{\prime\prime\prime\prime\prime}}^{(\mathbf{q}^{\prime\prime\prime\prime\prime} - \mathbf{q}^{\prime\prime\prime\prime})^{2}},$$
(24)

where the value of N is 1, 2 and 3 corresponding to A-, B- and C-terms. Also the different parts of the D-term which have been considered in the right side of Eq. (23) can be evaluated in the following:

$${}_{2}\langle \mathbf{q}^{\prime\prime\prime\prime} m_{s_{2}}^{\prime\prime\prime\prime} m_{s_{3}}^{\prime\prime\prime\prime} | V_{23}^{(D)} | \mathbf{q}^{\prime\prime\prime\prime\prime} m_{s_{1}}^{\prime\prime\prime\prime\prime} m_{s_{3}}^{\prime\prime\prime\prime\prime} \rangle_{2} = {}_{2}\langle \mathbf{q}^{\prime\prime\prime\prime} m_{s_{2}}^{\prime\prime\prime\prime} | V_{23}^{(B)} | \mathbf{q}^{\prime\prime\prime\prime\prime} m_{s_{1}}^{\prime\prime\prime\prime} \rangle_{2} \mathcal{O}_{m_{s_{2}}^{\prime\prime\prime\prime} m_{s_{2}}^{\prime\prime\prime\prime\prime}}^{(\mathbf{q}^{\prime\prime\prime\prime} - \mathbf{q}^{\prime\prime\prime\prime}) M},$$
(25)

where the value of M is 0 for the first and fifth parts of this term, it is 1 for the third part and it is 2 for the second and fourth parts.

#### IV. SUMMARY

A new formalism of three dimensional Faddeev integral equations for the 3N bound state including the 3NF is proposed. This formulation leads to a coupled set of a strictly finite number of equations in two vector variables for the amplitudes. The comparison of 3D and PW formalisms shows that this non PW formalism avoids the very involved angular momentum algebra occurring for the permutations and transformations and it is more efficient and less cumbersome for considering the 3NF. This formalism enables us to handle the realistic 2N and 3N forces with all their complexity in 3N bound state calculations.

#### Acknowledgments

M. R. H. would like to thank the organizers of the APFB08 conference for their kind invitation, warm hospitality and support, and the very pleasant and stimulating atmosphere at the conference. We would like to express our acknowledgements to our colleagues M. Harzchi and M. A. Shalchi for the enriching discussions. This work was supported by the research council of the University of Tehran.

- [1] I. Fachruddin, Ch. Elster, and W. Glöckle, *Phys. Rev.* C 62, 044002 (2000).
- [2] I. Fachruddin, Ch. Elster, and W. Glöckle, *Phys. Rev.* C 63, 054003 (2001).
- [3] H. Liu, Ch. Elster, and W. Glöckle, Few Body Systems 33, 241 (2003).
- [4] I. Fachruddin, Ch. Elster, and W. Glöckle, *Phys. Rev.* C 68, 054003 (2003).
- [5] I. Fachruddin, W. Glöckle, Ch. Elster, and A. Nogga, Phys. Rev. C 69, 064002 (2004).
- [6] H. Liu, Ch. Elster, and W. Glöckle, *Phys. Rev.* C 72, 054003 (2005).
- [7] T. Lin, Ch. Elster, W. N. Polyzou, and W. Glöckle, Phys. Rev. C 76, 014010 (2007).
- [8] M. R. Hadizadeh and S. Bayegan, Few Body Systems 40, 171 (2007).
- [9] M. R. Hadizadeh and S. Bayegan, Eur. Phys. J. A 36, 201 (2008).
- [10] S. Bayegan, M. R. Hadizadeh, and M. Harzchi, Phys. Rev. C 77, 064005 (2008).
- [11] Ch. Elster, T. Lin, W. Glöckle, and S. Jeschonnek, Phys. Rev. C 78, 034002 (2008).
- [12] S. Bayegan, M. Harzchi, and M. R. Hadizadeh, accepted in Nuclear Physics A. arXiv:0809.0200
- S. Bayegan, M. R. Hadizadeh, and W. Glöckle, submitted to Progress of Theoretical Physics. arXiv:0806.1520
- [14] D. Hüber, H. Witala, A. Nogga, W. Glöckle, and H. Kamada, Few-Body Systems 22, 107 (1997).
- [15] S. A. Coon and W. Glöckle, *Phys. Rev.* C 23 (1981) 1790.