

How to De-Rotate the Cosmic Microwave Background Polarization

Marc Kamionkowski

California Institute of Technology, Mail Code 130-33, Pasadena, CA 91125

(Dated: April 21, 2019)

The linear polarization of the cosmic microwave background (CMB) might be rotated as it propagates from the surface of last scatter. While Faraday rotation can be subtracted with multifrequency observations, more exotic mechanisms may give rise to a frequency-independent rotation. This rotation introduces a B-mode polarization pattern, even if there was none at the surface of last scatter. Here I consider the effects of a rotation angle that varies across the sky. I show that measurement of higher-order TE, EE, EB, and TB correlations induced by this rotation can be used to reconstruct the rotation angle as a function of position on the sky. This technique can be used to distinguish primordial B modes from those induced by rotation, and it may be useful as a diagnostic for systematic artifacts in the data. The rotation may also contaminate cosmic-shear maps, but can be distinguished by TT, which is unaltered by rotation.

PACS numbers: 98.80.-k

The polarization of the cosmic microwave background (CMB) can be decomposed into gradient (E mode) and curl (B mode) components [1]. Primordial density perturbations produce a polarization pattern that is purely E mode at the surface of last scatter, while primordial gravitational waves, such as those from inflation, produce a B mode [2]. There is now an active experimental effort to detect B modes, as these would constitute a “smoking gun” for inflationary gravitational waves [3].

There may, however, be other mechanisms—apart from gravitational waves—for producing B modes. The most widely considered is cosmic shear: the deflection of CMB photons due to weak gravitational lensing by density perturbations along the line of sight will convert some of the E mode at the surface of last scatter to a B mode [4]. Cosmic shear of the CMB is no longer the realm of futuristic theorists; it has recently been detected [14].

Another possibility is a rotation of the linear polarization of the CMB as it travels from the surface of last scatter [5]. This could occur, for example, if there is a quintessence field that couples to the pseudoscalar of electromagnetism [6]. In this case, the polarization would be rotated by an angle α that is uniform across the sky. A fraction $\sin \alpha$ of the E mode would thus be converted into a B mode [5]. This B mode could be distinguished from that due to gravitational waves by the parity breaking EB and TB cross-correlations that it produces, as well as by the shape of the TB, EB, and BB power spectra. Searches with current data already constrain such a uniform rotation to be no more than a few degrees [7].

But what if the rotation angle α varies across the sky? Refs. [9] have recently proposed scalar-field models in which this might occur. (Faraday rotation could also rotate the polarization [8], but this can be identified with multifrequency maps.) If the distribution of rotation angles is symmetric about zero, then there will be a B mode induced with no parity-breaking TB or EB correlations [9]. Can this type of B mode be distinguished from that

due to primordial gravitational waves?

In this *Letter*, I show that a position-dependent rotation of the polarization induces higher-order correlations in the temperature-polarization of the CMB. I then show how these new correlations can be used to measure the rotation angle $\alpha(\hat{n})$ as a function of position \hat{n} on the sky. The observed polarization pattern can then be corrected to construct the primordial polarization pattern. It will thus be possible to distinguish whether a B-mode pattern, if detected, is primordial or due to a post-recombination rotation. The techniques I will discuss can also be used to test an experiment for systematic artifacts.

The reconstruction algorithm will be similar to that developed to reconstruct the cosmic-shear deflection angle [10] and also to that developed to test for statistical isotropy (SI) [11]. In fact, the effects of rotation may be viewed as a possible contaminant for cosmic-shear maps. However, the effects of cosmic shear (or SI violations) and rotation can be distinguished, as rotation acts only on the polarization, while cosmic shear affects the temperature as well.

The linear polarization at each point \hat{n} on the sky is quantified by Stokes parameters $Q(\hat{n})$ and $U(\hat{n})$, measured with respect to the $\hat{\theta}$ - $\hat{\phi}$ axes. These Stokes parameters are components of a symmetric trace-free 2×2 polarization tensor field $\mathcal{P}_{ab}(\hat{n})$ which can be expanded in terms of tensor spherical harmonics as,

$$\mathcal{P}_{ab}(\hat{n}) = \sum_{l=2}^{\infty} \sum_{m=-l}^l \left[E_{lm} Y_{(lm)ab}^E(\hat{n}) + B_{lm} Y_{(lm)ab}^B(\hat{n}) \right]. \quad (1)$$

Here, $Y_{(lm)ab}^E(\hat{n})$ and $Y_{(lm)ab}^B(\hat{n})$ are complete sets of basis functions for the gradient (E mode) and curl (B mode) components of the polarization. Orthonormality of the basis functions allows us to write the expansion coeffi-

cients as

$$E_{lm} = \int d\hat{\mathbf{n}} \mathcal{P}_{ab}(\hat{\mathbf{n}}) Y_{(lm)}^{Eab*}(\hat{\mathbf{n}}), \quad (2)$$

$$B_{lm} = \int d\hat{\mathbf{n}} \mathcal{P}_{ab}(\hat{\mathbf{n}}) Y_{(lm)}^{Bab*}(\hat{\mathbf{n}}). \quad (3)$$

Suppose now that the polarization pattern at the surface of last scatter has no B mode and that the polarization at each point $\hat{\mathbf{n}}$ is rotated by an angle $\alpha(\hat{\mathbf{n}})$. In that case, the observed Stokes parameters will be

$$\begin{pmatrix} Q_{\text{obs}}(\hat{\mathbf{n}}) \\ U_{\text{obs}}(\hat{\mathbf{n}}) \end{pmatrix} = \begin{pmatrix} \cos 2\alpha(\hat{\mathbf{n}}) & \sin 2\alpha(\hat{\mathbf{n}}) \\ \sin 2\alpha(\hat{\mathbf{n}}) & \cos 2\alpha(\hat{\mathbf{n}}) \end{pmatrix} \begin{pmatrix} Q(\hat{\mathbf{n}}) \\ U(\hat{\mathbf{n}}) \end{pmatrix} \\ = \cos 2\alpha \begin{pmatrix} Q \\ U \end{pmatrix} + \sin 2\alpha \begin{pmatrix} U \\ -Q \end{pmatrix}. \quad (4)$$

The concordance of measured TT, TE, and EE power spectra suggest that α is relatively small, and that the uniform component of α is small [7]. We therefore work to linear order in α . The change to the polarization tensor induced by the rotation is then $\delta\mathcal{P}_{ab}(\hat{\mathbf{n}}) = 2\alpha(\hat{\mathbf{n}})\mathcal{P}_{ab}^r(\hat{\mathbf{n}})$, where $\mathcal{P}_{ab}^r(\hat{\mathbf{n}})$ is rotated from the original polarization \mathcal{P}_{ab} by 45° . Thus, if \mathcal{P}_{ab} is pure E mode, then \mathcal{P}_{ab}^r is pure B mode and can be written as

$$\mathcal{P}_{ab}^r(\hat{\mathbf{n}}) = \sum_{l=2}^{\infty} \sum_{m=-l}^l E_{lm} Y_{(lm)ab}^B. \quad (5)$$

To calculate the curl component induced by rotation, we insert $\delta\mathcal{P}_{ab}$ into the expression, Eq. (3), for B_{lm} . We then expand the rotation angle in terms of (scalar) spherical harmonics $Y_{(lm)}$,

$$\alpha(\hat{\mathbf{n}}) = \sum_{LM} \alpha_{LM} Y_{(LM)}(\hat{\mathbf{n}}), \quad (6)$$

to obtain

$$\begin{aligned} B_{lm} &= \sum_{LM} \sum_{l_2 m_2} \alpha_{LM} E_{l_2 m_2} \int d\hat{\mathbf{n}} Y_{(lm)}^{B,ab*} Y_{(LM)} Y_{(l_2 m_2)ab}^B \\ &= 2(-1)^m \sum_{LM} \sum_{l_2 m_2} \alpha_{LM} E_{l_2 m_2} \xi_{lm l_2 m_2}^{LM} H_{ll_2}^L, \end{aligned} \quad (7)$$

where the sum is taken only over l_2 values that satisfy $L + l + l_2 = \text{even}$,

$$H_{ll'}^L \equiv \begin{pmatrix} l & l' & L \\ -2 & 2 & 0 \end{pmatrix} \begin{pmatrix} l & l' & L \\ 0 & 0 & 0 \end{pmatrix}^{-1}, \quad (8)$$

in terms of Wigner-3j symbols, and

$$\xi_{lm l' m'}^{LM} = \int d\hat{\mathbf{n}} Y_{(lm)}^*(\hat{\mathbf{n}}) Y_{(l' m')}(\hat{\mathbf{n}}) Y_{(LM)}(\hat{\mathbf{n}}). \quad (9)$$

We thus see that rotation induces a B mode [5, 9]. If there is a power spectrum for α_{LM} , then the CMB power spectrum C_l^{BB} can be calculated from Eq. (7) [9].

There is also an $O(\alpha)$ change in the E mode induced by rotation with precisely the same form as Eq. (7), but with contributions only from $L + l + l_2 = \text{odd}$, rather than even. In the discussion below, we focus for brevity and clarity on the induced EB and TB correlations. However, the entire discussion applies (with differences to be pointed out below) to induced EE and TE correlations.

The next step is to consider the correlation of the induced B mode with the original E mode, as well as with the temperature T. Recalling that the original E modes have (assuming Gaussian initial conditions) expectation values $\langle E_{lm}^* E_{l' m'} \rangle = C_l^{\text{EE}} \delta_{ll'} \delta_{mm'}$ (where C_l^{EE} is the EE power spectrum), the correlation between the observed E and B modes is [12],

$$\begin{aligned} \langle B_{lm} E_{l' m'}^* \rangle &= 2 \frac{\alpha_{00}}{\sqrt{4\pi}} C_l^{\text{EE}} \delta_{ll'} \delta_{mm'} \\ &+ \sum_{L \geq 1} \sum_{M=-L}^L \alpha_{LM} C_l^{\text{EE}} \xi_{lm l' m'}^{LM} H_{ll'}^L. \end{aligned} \quad (10)$$

I have split off the $L = 0$ term in Eq. (10) to show that the result for a uniform rotation angle α is $\langle B_{lm} E_{l' m'}^* \rangle = 2\alpha C_l^{\text{EE}} \delta_{ll'} \delta_{mm'}$, as it should be [5]. The sum in Eq. (10) is taken only over $L + l + l' = \text{even}$. The expression for the induced correlations $\langle E_{lm} E_{l' m'}^* \rangle$ is the same as that in Eq. (10), but with $L + l + l' = \text{odd}$, and no $L = 0$ contribution.

Likewise, given that the temperature T and E-mode polarization are correlated at the surface of last scatter, there will also be a nonzero TB correlation induced by rotation. The expression for $\langle T_{lm} B_{l' m'}^* \rangle$ is identical to Eq. (10) with the replacement $C_l^{\text{EE}} \rightarrow C_l^{\text{TE}}$, again for $L + l + l' = \text{even}$. There will also be TE correlations of the same form, but with $L + l + l' = \text{odd}$ and no $L = 0$ term.

If $\alpha(\hat{\mathbf{n}})$ varies with $\hat{\mathbf{n}}$, then $\alpha_{LM} \neq 0$ for $L \geq 1$, and if so, there will be correlations between E_{lm} and $B_{l' m'}$ (and T_{lm} and $B_{l' m'}$) of different lm and $l' m'$. The existence of these off-diagonal correlations can be used to measure each of the rotation multipole moments α_{lm} , and thus $\alpha(\hat{\mathbf{n}})$. The relevant formalism is similar to that for measuring the cosmic-shear deflection angle [10], or for searching for SI violations [11], so we can adopt results from prior work. To do so, we note that Eq. (10) is identical to Eq. (A1) in Ref. [11] with the identification $X = B$ (or T in place of B), $X' = E$, and $D_{ll'}^{LM, XX'} = \alpha_{LM} C_l^{\text{EE}} H_{ll'}^L$ (or C_l^{TE} in place of C_l^{EE} for TB).

Our goal is to obtain the minimum-variance estimator $\hat{\alpha}_{LM}$ that can be obtained from a full-sky polarization map or a temperature-polarization map. We suppose that the maps are provided as a measured temperature $T^{\text{map}}(\hat{\mathbf{n}})$ and Stokes parameters $Q^{\text{map}}(\hat{\mathbf{n}})$ and $U^{\text{map}}(\hat{\mathbf{n}})$ in N_{pix} pixels on the sky. The temperature (polarization) in each pixel receives contributions from the signal, which is the temperature (polarization) on the sky

smoothed with a Gaussian beam of full-width half maximum (FWHM) θ_{fwhm} , and a Gaussian noise with variance σ_T^2 (σ_P^2). The power spectra for the map are then $C_l^{\text{A,map}} = |W_l^2| C_l^{\text{A}} + C_l^{\text{A,n}}$, where $C_l^{\text{A,n}}$ is the noise power spectrum for A (e.g., A = {TT, EE, BB, TE, TB, EB}). These are $C_l^{\text{TT,n}} = (4\pi/N_{\text{pix}})\sigma_T^2$, $C_l^{\text{EE,n}} = C_l^{\text{BB,n}} = (4\pi/N_{\text{pix}})\sigma_P^2$, and $C_l^{\text{TE,n}} = C_l^{\text{TB,n}} = C_l^{\text{EB,n}} = 0$. Beam smearing is taken into account with the window function $W_l = \exp(-l^2\sigma_b^2/2)$, with $\sigma_b = \theta_{\text{fwhm}}/\sqrt{8\ln 2} = 0.0742$ ($\theta_{\text{fwhm}}/1^\circ$).

We now derive the minimum-variance estimator $\hat{\alpha}_{LM}$ that can be obtained from EB correlations; the results for the estimator that can be obtained from TB correlations will be identical with the replacement E \rightarrow B; TE and EE estimators are similarly derived.

Following Ref. [11], the minimum-variance estimator for each $D_{ll'}^{LM} = \alpha_{LM} C_l^{\text{EE}} H_{ll'}^L$ that can be obtained from the polarization map is (see also Refs. [13])

$$\hat{D}_{ll'}^{LM,\text{map}} = (G_{ll'}^L)^{-1} \sum_{mm'} B_{lm}^{\text{map}} (E_{l'm'}^{\text{map}})^* \xi_{lm'l'm'}^{LM}, \quad (11)$$

with

$$G_{ll'}^L = \frac{(2l+1)(2l'+1)}{4\pi(2L+1)} (C_{l0l'0}^{L0})^2, \quad (12)$$

where $C_{lm'l'm'}^{LM}$ is a Clebsch-Gordan coefficient.

The coefficient α_{LM} can be estimated from measurement of $D_{ll'}^{LM}$ from each ll' pair through $\alpha_{LM} = D_{ll'}^{LM} / (C_l^{\text{EE}} H_{ll'}^L)$. One can then average the estimates of $\alpha_{ll'}$ from all of the ll' pairs. The trick, though, is to weight these all in a manner that minimizes the variance to α_{LM} . If each estimator $\hat{D}_{ll'}^{LM,\text{map}}$ were statistically independent, then we could simply weight by the inverse variance. However, things are a bit (though not much) more complicated.

Each of the estimators $\hat{D}_{ll'}^{LM,\text{map}}$ are statistically independent for different LM . They are also statistically independent for different ll' , *except* that $\hat{D}_{ll'}^{LM,\text{map}}$ is correlated with $\hat{D}_{l'l}^{LM,\text{map}}$. To take this into account, we take $l' \geq l$ and then consider for $l \neq l'$ EB modes as well as BE modes. For $l = l'$, there is only a single variance; for $l' > l$, there is a 2×2 covariance matrix in the EB-BE space.

Write the covariances between the different $\hat{D}_{ll'}^{LM,\text{map}}$ as

$$C_{AA'}^{ll'} \equiv \frac{G_{ll'}^L}{(1 + \delta_{ll'})} \left\langle \hat{D}_{ll'}^{LM,\text{map}} \hat{D}_{l'l}^{LM,\text{map}} \right\rangle, \quad (13)$$

for A = {EB, BE}. For $l = l'$, there is no distinction between EB and BE; the variance in this case is then

$$C_{\text{BE,BE}}^{ll} = \frac{1}{2} \left[C_l^{\text{BB,map}} C_l^{\text{EE,map}} + \left(C_l^{\text{BE,map}} \right)^2 \right]. \quad (14)$$

For $l' > l$, the covariances are $C_{\text{BE,BE}}^{ll'} = C_l^{\text{BB,map}} C_{l'}^{\text{EE,map}}$, $C_{\text{EB,EB}}^{ll'} = C_{l'}^{\text{BB,map}} C_l^{\text{EE,map}}$, and $C_{\text{BE,EB}}^{ll'} = C_l^{\text{BE,map}} C_{l'}^{\text{BE,map}}$.

We now write two estimators, $\hat{\alpha}_{LM}^{l=l'}$ and $\hat{\alpha}_{LM}^{l<l'}$, the first coming from EB correlations with $l = l'$ and the second from those with $l < l'$. We will then average them, with inverse-variance weighting, to obtain the final estimator. The first is

$$\hat{\alpha}_{LM}^{l=l'} = \frac{\sum_l F_{ll}^L (W_l)^2 \hat{D}_{ll}^{LM,\text{map}} G_{ll}^L / C_{\text{BE,BE}}^{ll}}{\sum_l [F_{ll}^L (W_l)^2]^2 G_{ll}^L / C_{\text{BE,BE}}^{ll}}, \quad (15)$$

where $F_{ll'}^L \equiv C_l^{\text{EE}} H_{ll'}^L$. The second is

$$\hat{\alpha}_{LM}^{l<l'} = \frac{\sum_{l'>l} W_l W_{l'} G_{ll'}^L \sum_{AA'} F_{ll'}^{L,A} \hat{D}_{ll'}^{LM,A,\text{map}} \left[(C^{ll'})^{-1} \right]_{AA'}}{\sum_{l'>l} (W_l W_{l'})^2 G_{ll'}^L \sum_{AA'} F_{ll'}^{L,A} F_{ll'}^{L,A'} \left[(C^{ll'})^{-1} \right]_{AA'}}, \quad (16)$$

where the matrix inversion is the in the 2×2 EB-BE basis. Note that the superscripts A on F and \hat{D} are necessary in these expressions, as these quantities differ for EB and BE. The variance to the first estimator is

$$(\sigma_{\alpha_{LM}^{l=l'}})^2 = \left(\sum_l \frac{[F_{ll}^L (W_l)^2]^2 G_{ll}^L}{2C_{\text{BE,BE}}^{ll}} \right)^{-1}. \quad (17)$$

The variance $(\sigma_{\alpha_{LM}^{l<l'}})^2$ to the second estimator is obtained from

$$(\sigma_{\alpha_{LM}^{l<l'}})^{-2} = \sum_{l'>l} G_{ll'}^L \sum_{AA'} F_{ll'}^{L,A} F_{ll'}^{L,A'} (W_l W_{l'})^2 \left[(C^{ll'})^{-1} \right]_{AA'}. \quad (18)$$

The final minimum-variance estimator $\hat{\alpha}_{LM}$ is then obtained by averaging, with inverse-variance weighting, the two estimators above:

$$\hat{\alpha}_{LM} = \frac{\hat{\alpha}_{LM}^{l=l'} (\sigma_{\alpha_{LM}^{l=l'}})^{-2} + \hat{\alpha}_{LM}^{l<l'} (\sigma_{\alpha_{LM}^{l<l'}})^{-2}}{(\sigma_{\alpha_{LM}^{l=l'}})^{-2} + (\sigma_{\alpha_{LM}^{l<l'}})^{-2}}. \quad (19)$$

The variance $(\sigma_{\alpha_{LM}})^2$ for this estimator is given by $(\sigma_{\alpha_{LM}})^{-2} = (\sigma_{\alpha_{LM}^{l=l'}})^{-2} + (\sigma_{\alpha_{LM}^{l<l'}})^{-2}$. For small L (e.g., a rotation dipole), the two terms will contribute comparably to the statistical weight. For larger L , the $l' > l$ estimator should carry most of the statistical weight.

The estimators described so far have been for EB correlations. However, estimators from TB correlations are identical with the replacement E \rightarrow B. Likewise, there will be EE and TE correlations similarly induced. (Things simplify for EE, as the 2×2 AA' covariance matrix becomes a single variance.) The estimators for α_{LM} for EE and TE and their variances can be constructed analogously. There will also be BB correlations, but they will be higher order in α . There will be no TT correlations

induced, as the rotation does not act on the temperature. Since T and E are correlated in the primordial polarization field, there will be cross correlations between the estimators $\hat{\alpha}_{LM}$ from EB, TB, EE, and TE. This may be an order-unity effect if the statistical weights of the various estimators are comparable. The expressions for the complete covariances are long, and so I leave them for future work.

Here I have shown that CMB temperature-polarization statistics can be developed to measure the angle $\alpha(\hat{n})$ by which CMB photons were rotated, en route from the surface of last scatter, as a function of position \hat{n} on the sky. Explicit formulas to obtain the coefficients α_{LM} in a spherical-harmonic expansion of $\alpha(\hat{n})$ from a full-sky CMB map were provided. This technique can then determine whether B modes, if detected, occur at the surface of last scatter or are due to a post-recombination rotation of the polarization. It is interesting to know that the rotation angle can be determined from the data, rather than by assumption. And if the rotation angle is assumed to be zero, then the techniques developed here can provide a test for systematic artifacts in the data.

The rotation and cosmic-shear formalisms share some similarities, and so rotation, if it exists, could show up as an artifact in a cosmic-shear analysis. However, the detailed effects are different and can be distinguished in the data. Most significantly, unlike cosmic shear, which acts the same on temperature and polarization, rotation will leave the temperature map unaltered.

Since the CMB is never really measured on the entire sky, the precise formulas for the estimators $\hat{\alpha}_{LM}$ will need to be modified to take into account partial sky coverage. However, it will be straightforward to adapt the techniques that have been developed to measure cosmic shear of the CMB on a partial sky to measure the rotation angle. Likewise, it is straightforward to simplify the full-sky analysis performed here to the flat-sky limit, which may be appropriate for sub-orbital experiments that map the CMB on a small patch of sky. Since the formalism to reconstruct the rotation angle resembles that to determine the cosmic-shear deflection angle, there may be other cosmic-shear techniques that can be adapted for rotation. For example, maximum-likelihood techniques [15] may be developed to provide even more sensitive probes of rotation than the quadratic estimators discussed here.

Finally, what I have done here is to simply show that the rotation can be measured. It will be interesting to apply the formalism developed here to quantitative estimates of the detectability of the effect for specific models of the rotation, such as those in Refs. [9], or perhaps other models yet to be developed.

I thank Asantha Cooray and Chris Hirata for useful comments. This work was supported by DoE DE-FG03-92-ER40701 and the Gordon and Betty Moore Foundation.

-
- [1] M. Kamionkowski, A. Kosowsky and A. Stebbins, Phys. Rev. D **55**, 7368 (1997) [arXiv:astro-ph/9611125]; M. Zaldarriaga and U. Seljak, Phys. Rev. D **55**, 1830 (1997) [arXiv:astro-ph/9609170].
 - [2] M. Kamionkowski, A. Kosowsky and A. Stebbins, Phys. Rev. Lett. **78**, 2058 (1997) [arXiv:astro-ph/9609132]; U. Seljak and M. Zaldarriaga, Phys. Rev. Lett. **78**, 2054 (1997) [arXiv:astro-ph/9609169].
 - [3] J. Bock *et al.*, arXiv:astro-ph/0604101.
 - [4] M. Zaldarriaga and U. Seljak, Phys. Rev. D **58**, 023003 (1998) [arXiv:astro-ph/9803150].
 - [5] A. Lue, L. M. Wang and M. Kamionkowski, Phys. Rev. Lett. **83**, 1506 (1999) [arXiv:astro-ph/9812088]; N. F. Lepora, arXiv:gr-qc/9812077.
 - [6] S. M. Carroll, Phys. Rev. Lett. **81**, 3067 (1998) [arXiv:astro-ph/9806099].
 - [7] B. Feng *et al.*, Phys. Rev. Lett. **96**, 221302 (2006) [arXiv:astro-ph/0601095]; T. Kahniashvili, R. Durrer and Y. Maravin, arXiv:0807.2593 [astro-ph]; J. Q. Xia *et al.*, arXiv:0803.2350 [astro-ph]; E. Komatsu *et al.* [WMAP Collaboration], arXiv:0803.0547 [astro-ph]; J. Q. Xia *et al.*, arXiv:0710.3325 [hep-ph]; E. Y. .S. Wu *et al.* [QUaD Collaboration], in preparation.
 - [8] A. Kosowsky and A. Loeb, Astrophys. J. **469**, 1 (1996) [arXiv:astro-ph/9601055].
 - [9] M. Pospelov, A. Ritz and C. Skordis, arXiv:0808.0673 [astro-ph]; M. Li and X. Zhang, arXiv:0810.0403 [astro-ph].
 - [10] U. Seljak and M. Zaldarriaga, Phys. Rev. Lett. **82**, 2636 (1999) [arXiv:astro-ph/9810092]; M. Zaldarriaga and U. Seljak, Phys. Rev. D **59**, 123507 (1999) [arXiv:astro-ph/9810257]; W. Hu, Phys. Rev. D **62**, 043007 (2000) [arXiv:astro-ph/0001303]; T. Okamoto and W. Hu, Phys. Rev. D **67**, 083002 (2003) [arXiv:astro-ph/0301031]; W. Hu and T. Okamoto, Astrophys. J. **574**, 566 (2002) [arXiv:astro-ph/0111606]; M. Kesden, A. Cooray and M. Kamionkowski, Phys. Rev. Lett. **89**, 011304 (2002) [arXiv:astro-ph/0202434]; L. Knox and Y. S. Song, Phys. Rev. Lett. **89**, 011303 (2002) [arXiv:astro-ph/0202286]; M. H. Kesden, A. Cooray and M. Kamionkowski, Phys. Rev. D **67**, 123507 (2003) [arXiv:astro-ph/0302536]; P. Cabella and M. Kamionkowski, arXiv:astro-ph/0403392; A. Lewis and A. Challinor, Phys. Rept. **429**, 1 (2006) [arXiv:astro-ph/0601594].
 - [11] A. R. Pullen and M. Kamionkowski, Phys. Rev. D **76**, 103529 (2007) [arXiv:0709.1144 [astro-ph]].
 - [12] C. Scoccola, D. Harari and S. Mollerach, Phys. Rev. D **70**, 063003 (2004) [arXiv:astro-ph/0405396].
 - [13] A. Hajian and T. Souradeep, arXiv:astro-ph/0501001.
 - [14] K. M. Smith, O. Zahn and O. Dore, Phys. Rev. D **76**, 043510 (2007) [arXiv:0705.3980 [astro-ph]]; C. M. Hirata *et al.*, Phys. Rev. D **78**, 043520 (2008) [arXiv:0801.0644 [astro-ph]].
 - [15] C. M. Hirata and U. Seljak, Phys. Rev. D **67**, 043001 (2003) [arXiv:astro-ph/0209489]; C. M. Hirata and U. Seljak, Phys. Rev. D **68**, 083002 (2003) [arXiv:astro-ph/0306354].