

Approximation theorems for Banach-valued almost periodic and semi-almost periodic holomorphic functions

Alexander Brudnyi*

Department of Mathematics and Statistics
University of Calgary, Calgary
Canada

Damir Kinzebulatov

Department of Mathematics
University of Toronto, Toronto
Canada

Abstract

The paper studies semi-almost periodic holomorphic functions on a poly-disk which have, in a sense, the weakest possible discontinuities on the boundary torus. The basic result used in the proofs is an extension of the classical Bohr approximation theorem for almost periodic holomorphic functions on a strip to the case of Banach-valued almost periodic holomorphic functions.

1 Introduction and Formulation of Main Results

1.1. Let B be a complex Banach space and $C_b^B(U)$ be the space of bounded continuous B -valued functions f on a closed subset $U \subset \mathbb{C}$ endowed with sup-norm $\|f\| := \sup_{t \in U} \|f(t)\|_B$.

*Research supported in part by NSERC.

2000 *Mathematics Subject Classification.* Primary 30H05, Secondary 46J20.

Key words and phrases. Banach-valued holomorphic functions, almost periodic functions, approximation property

Definition 1.1. A function $f \in C_b^B(\mathbb{R})$ is said to be *almost periodic* if the family of shifts $\{S_\tau f\}_{\tau \in \mathbb{R}}$, $S_\tau f(t) := f(t + \tau)$, $t \in \mathbb{R}$, is relatively compact in $C_b^B(\mathbb{R})$.

By $AP_B(\mathbb{R}) \subset C_b^B(\mathbb{R})$ we denote the Banach space of B -valued almost periodic functions on \mathbb{R} endowed with sup-norm.

The following result due to H. Bohr, commonly called the *approximation theorem*, is a cornerstone of the theory of Banach-valued almost periodic functions.

Theorem 1.2 (H. Bohr). *The vector subspace of $C_b^B(\mathbb{R})$ spanned over B by functions $t \mapsto e^{i\lambda t}$, $\lambda \in \mathbb{R}$, is dense in $AP_B(\mathbb{R})$.*

Similarly one defines B -valued holomorphic almost periodic functions on the strip $\Sigma := \{z \in \mathbb{C} : 0 \leq \text{Im}(z) \leq \pi\}$.

Definition 1.3. A function $f \in C_b^B(\Sigma)$ is called *holomorphic almost periodic* on Σ if it is holomorphic in the interior of Σ and the family of shifts $\{S_x f\}_{x \in \mathbb{R}}$ is relatively compact in $C_b^B(\Sigma)$.

By $APH_B(\Sigma) \subset C_b^B(\Sigma)$ we denote the Banach space of B -valued almost periodic holomorphic functions on Σ endowed with sup-norm.

Our first result is an analog of Theorem 1.2 for functions from $APH_B(\Sigma)$.

Theorem 1.4. *The complex vector subspace of $C_b^B(\Sigma)$ spanned over B by functions $z \mapsto e^{i\lambda z}$, $\lambda \in \mathbb{R}$, is dense in $APH_B(\Sigma)$.*

This result is equivalent to the fact that the Banach space $APH(\Sigma) := APH_{\mathbb{C}}(\Sigma)$ of holomorphic almost periodic functions on Σ has the *approximation property* along with the scalar version of Theorem 1.4, see Grothendieck [5, Section 5.1]. (Similarly Theorem 1.2 can be derived from the fact that $AP(\mathbb{R}) := AP_{\mathbb{R}}(\mathbb{R})$ has the approximation property.) Let us recall

Definition 1.5. A Banach space B is said to have the approximation property, if, for every compact set $K \subset B$ and every $\varepsilon > 0$, there is an operator $T : B \rightarrow B$ of finite rank so that $\|Tx - x\|_B \leq \varepsilon$ for every $x \in K$. If such T has norm at most 1, then B is said to have metric approximation property.

It is easy to see that every space B with a Schauder basis has the approximation property. However, there are Banach spaces without this property, the first such example was constructed by Enflo [3]. In Proposition 3.1 we will show that $APH(\Sigma)$ has metric approximation property.

1.2. We apply Theorem 1.4 to study the algebra of bounded holomorphic functions on the unit polydisk $\mathbb{D}^n \subset \mathbb{C}^n$ with semi-almost periodic boundary values. Originally

the algebra of semi-almost periodic functions on the real line \mathbb{R} was introduced and studied by Sarason [6] in connection with some important problems of operator theory. By the definition it is a closed subalgebra of $C_b(\mathbb{R})$ generated by algebras $AP(\mathbb{R})$ and PC_∞ (the algebra of continuous functions on \mathbb{R} that have finite limits at $+\infty$ and at $-\infty$). In [2] we introduced the algebra $SAP(\partial\mathbb{D})$ of semi-almost periodic functions on the unit circle $\partial\mathbb{D}$ which generalizes the Sarason algebra. Then we studied the algebra $H^\infty(\mathbb{D}) \cap SAP(\partial\mathbb{D})$ of bounded holomorphic functions on the unit disk \mathbb{D} having boundary values in $SAP(\partial\mathbb{D})$ which we called *the algebra of semi-almost periodic holomorphic functions*. (Recall that for a function $f \in H^\infty(\mathbb{D})$, one defines its boundary values a.e. on $\partial\mathbb{D}$ by taking limits in non-tangential directions, so that $f|_{\partial\mathbb{D}} \in L^\infty(\partial\mathbb{D})$.)

Our interest in algebra $H^\infty(\mathbb{D}) \cap SAP(\partial\mathbb{D})$ is motivated by the following problem:

Find a closed subalgebra $A \subset H^\infty(\mathbb{D})$ whose elements have, in a sense, the weakest possible discontinuities at the boundary $\partial\mathbb{D}$.

An obvious candidate would be the subalgebra of bounded holomorphic functions having discontinuities of at most first kind at the boundary. Unfortunately, as it follows from the Lindelöf Theorem, see [4], any such function must be continuous on $\bar{\mathbb{D}}$. Moreover, the same result holds if we consider bounded holomorphic functions with first kind discontinuous on $\partial\mathbb{D}$ of their real or imaginary parts.

Another way to measure singularities of a function from $H^\infty(\mathbb{D})$ is to consider discontinuities of its modulus on $\partial\mathbb{D}$. Assuming that $A \subset H^\infty(\mathbb{D})$ consists of functions f such that $|f|_{\partial\mathbb{D}} \in L^\infty(\partial\mathbb{D})$ has first-kind discontinuous only, we obtain that A contains all inner functions. Then by the Marshall theorem, see [4], $A = H^\infty(\mathbb{D})$. Thus to get a nontrivial answer we correct the above problem restricting ourselves to the case of algebras $A \subset H^\infty(\mathbb{D})$ generating by subgroups G_A of invertible elements of $H^\infty(\mathbb{D})$ such that for each $f \in G_A$ the function $|f|$ (or, equivalently, the harmonic function $\ln|f|$) has finitely many discontinuities of at most first kind on $\partial\mathbb{D}$. The main result in [2] (Theorem 1.8) establishes a connection between such algebras A and certain subalgebras of $H^\infty(\mathbb{D}) \cap SAP(\partial\mathbb{D})$.

The purpose of the present paper is to define semi-almost periodic holomorphic functions on \mathbb{D}^n and to extend the results of [2] to this class of functions.

First we recall some definitions from [2].

In what follows, we consider $\partial\mathbb{D}$ with the counterclockwise orientation. For $t_0 \in \mathbb{R}$ let

$$\gamma_{t_0}^k(s) := \{e^{i(t_0+kt)} : 0 \leq t < s < 2\pi\}, \quad k \in \{-1, 1\},$$

be two open arcs having e^{it_0} as the right and the left endpoints, respectively.

Definition 1.6 ([2]). A function $f \in L^\infty(\partial\mathbb{D})$ is called semi-almost periodic on $\partial\mathbb{D}$ if for any $t_0 \in [0, 2\pi)$ and any $\varepsilon > 0$ there exist a number $s = s(t_0, \varepsilon) \in (0, \pi)$ and functions $f_k : \gamma_{t_0}^k(s) \rightarrow \mathbb{C}$, $k \in \{-1, 1\}$, such that functions

$$t \mapsto f_k(e^{i(t_0 + kse^t)}), \quad -\infty < t < 0, \quad k \in \{-1, 1\},$$

are restrictions of some almost periodic functions from $AP(\mathbb{R})$, and

$$\sup_{z \in \gamma_{t_0}^k(s)} |f(z) - f_k(z)| < \varepsilon, \quad k \in \{-1, 1\}.$$

We denote by $SAP(\partial\mathbb{D})$ the Banach algebra of semi-almost periodic functions on $\partial\mathbb{D}$ endowed with sup-norm. For S a closed subset of $\partial\mathbb{D}$ we denote by $SAP(S)$ the Banach algebra of semi-almost periodic functions on $\partial\mathbb{D}$ that are continuous on $\partial\mathbb{D} \setminus S$. (Note the the Sarason algebra is isomorphic to $SAP(\{z_0\})$ for any $z_0 \in \partial\mathbb{D}$.)

Next, let $A(\mathbb{D})$ be the algebra consisting of holomorphic functions in $H^\infty(\mathbb{D})$ that are continuous on $\bar{\mathbb{D}}$. Suppose that S contains at least two points. By A_S we denote the closure in $L^\infty(\partial\mathbb{D})$ of the algebra generated by $A(\mathbb{D})$ and holomorphic functions of the form e^f , where $\text{Re}(f)|_{\partial\mathbb{D}}$ is a finite linear combination (over \mathbb{R}) of characteristic functions of closed arcs in $\partial\mathbb{D}$ whose endpoints belong to S . If S consists of a single point, then we define A_S to be the closure of the algebra generated by $A(\mathbb{D})$ and functions ge^f , where $\text{Re}(f)|_{\partial\mathbb{D}}$ is the characteristic function of a closed arc with an endpoint in S and $g \in A(\mathbb{D})$ is a function such that ge^f has discontinuity on S only.

Now, Theorem 1.8 of [2] describes the structure of the algebra of semi-almost periodic holomorphic functions:

Theorem 1.7 ([2]). $H^\infty(\mathbb{D}) \cap SAP(S) = A_S$.

(Here and below we identify the elements of algebra $H^\infty(\mathbb{D}^n)$ with their boundary values defined on $(\partial\mathbb{D})^n$.)

Remark 1.8. Suppose that $S \subset \partial\mathbb{D}$ contains at least 2 points. Let $e^{\lambda f} \in A_S$, $\lambda \in \mathbb{R}$, where $\text{Re}(f)$ is the characteristic function of an arc $[x, y]$ with $x, y \in S$. Let $\phi_{x,y} : \mathbb{D} \rightarrow \mathbb{H}_+$ be the bilinear map onto the upper half-plane that maps x to 0, the midpoint of the arc $[x, y]$ to 1 and y to ∞ . Then there is a constant C such that

$$e^{\lambda f(z)} = e^{-\frac{i\lambda}{\pi} \text{Log } \phi_{x,y}(z) + \lambda C}, \quad z \in \mathbb{D},$$

where Log is the principal branch of the logarithmic function. Thus Theorem 1.7 implies that the algebra $H^\infty(\mathbb{D}) \cap SAP(S)$ is the uniform closure of the algebra generated by $A(\mathbb{D})$ and the family of functions $e^{i\lambda(\text{Log} \circ \phi_{x,y})}$, $\lambda \in \mathbb{R}$, $x, y \in S$.

Let us define now multi-dimensional analogues of algebras $SAP(S)$ and A_S . Namely, if $S_k \subset \partial\mathbb{D}$ are closed sets, $1 \leq k \leq n$, and $S = \prod_{k=1}^n S_k \subset (\partial\mathbb{D})^n$, then we define

$$SAP^n(S) := \bigotimes_{k=1}^n SAP(S_k), \quad A_S^n := \bigotimes_{k=1}^n A_{S_k}.$$

Here \bigotimes stands for completion of symmetric tensor product of the corresponding algebras. In particular, $SAP^n(S)$ and A_S^n are uniform closures of the algebras of complex polynomials in variables $f_1(z_1), \dots, f_n(z_n)$ with $f_k(z_k) \in SAP(S_k)$ and $f_k(z_k) \in A_{S_k}$, $1 \leq k \leq n$, respectively, where $z = (z_1, \dots, z_n) \in (\partial\mathbb{D})^n$.

Now the extension of Theorem 1.7 is read as follows.

Theorem 1.9. $H^\infty(\mathbb{D}^n) \cap SAP^n(S) = A_S^n$.

Remark 1.10. We apply Theorem 1.4 and some arguments from [2] to show that the algebra $H^\infty(\mathbb{D}) \cap SAP(S)$ has the approximation property. This will imply Theorem 1.9.

Let M_S be the maximal ideal space of the algebra $H^\infty(\mathbb{D}^n) \cap SAP^n(S)$ and M_{S_k} be the maximal ideal space of the algebra $H^\infty(\mathbb{D}) \cap SAP(S_k)$, $1 \leq k \leq n$. As a corollary of Theorem 1.9 we obtain

Theorem 1.11. M_S is homeomorphic to $M_{S_1} \times \dots \times M_{S_n}$. Moreover, the Corona Theorem is valid for $H^\infty(\mathbb{D}^n) \cap SAP^n(S)$, i.e., \mathbb{D}^n is dense in M_S in the Gelfand topology.

Remark 1.12. The structure of M_{S_k} is described in [2], Theorems 1.7 and 1.14. Let us recall this result. In what follows $M(A)$ stands for the maximal ideal space of the Banach algebra A .

Since $A(\mathbb{D}) \hookrightarrow A_{S_k}$, there is a continuous surjection of the maximal ideal spaces

$$a_{S_k} : M_{S_k} \rightarrow M(A(\mathbb{D})) \cong \bar{\mathbb{D}}.$$

Recall that the *Šilov boundary* of A_{S_k} is the smallest compact subset $K \subset M_{S_k}$ such that for each $f \in A_{S_k}$

$$\sup_{z \in M_{S_k}} |f(z)| = \sup_{\xi \in K} |f(\xi)|.$$

Here we assume that every $f \in A_{S_k}$ is also defined on M_{S_k} where its extension to $M_{S_k} \setminus \mathbb{D}$ is given by the *Gelfand transform*: $f(\xi) := \xi(f)$, $\xi \in M_{S_k}$.

Theorem 1.13 ([2]). (1) $a_{S_k} : M_{S_k} \setminus a_{S_k}^{-1}(S_k) \rightarrow \bar{\mathbb{D}} \setminus S_k$ is a homeomorphism.

(2) The Šilov boundary K_{S_k} of A_{S_k} is naturally homeomorphic to $M(SAP(S_k))$. Moreover, $K_{S_k} \setminus a_{S_k}^{-1}(S_k) = \partial\mathbb{D} \setminus S_k$ and $K_{S_k} \cap a_{S_k}^{-1}(z)$, $z \in S_k$, is homeomorphic to the disjoint union $b\mathbb{R} \sqcup b\mathbb{R}$ of the Bohr compactifications $b\mathbb{R}$ of \mathbb{R} .

(3) For each $z \in S_k$ preimage $a_{S_k}^{-1}(z)$ is homeomorphic to the maximal ideal space of the algebra $APH(\Sigma)$ of holomorphic almost periodic functions on the strip $\Sigma := \{z \in \mathbb{C} : \text{Im}(z) \in [0, \pi]\}$.

Acknowledgment. We are grateful to S. Favorov for useful discussions and valuable comments improving the presentation.

2 Auxiliary Results

2.1. In our proofs we use some results on Bohr's compactifications.

First recall that the algebra of almost periodic functions $AP(\mathbb{R}) := AP_{\mathbb{C}}(\mathbb{R})$ is naturally isomorphic to the algebra $C(b\mathbb{R})$ of complex continuous functions on the Bohr compactification $b\mathbb{R}$ of \mathbb{R} (the maximal ideal space of $AP(\mathbb{R})$). That is, a complex continuous function on \mathbb{R} is almost periodic if and only if it admits a continuous extension to $b\mathbb{R}$ by means of the Gelfand transform. (Recall that the maximal ideal space $b\mathbb{R}$ is defined as the space of continuous non-zero characters $AP(\mathbb{R}) \mapsto \mathbb{C}$ endowed with the Gelfand (i.e. weak*) topology. Then $b\mathbb{R}$ is a compact abelian group, \mathbb{R} is naturally embedded into $b\mathbb{R}$ as a dense subset, so that the action of \mathbb{R} on itself by translations extends uniquely to the continuous action of \mathbb{R} on $b\mathbb{R}$.)

Let us describe now the maximal ideal space of the algebra $APH(\Sigma)$.

Let $R = \{z \in \mathbb{C} : e^{-2\pi^2} \leq |z| \leq 1\}$. Then $\pi(z) := e^{2\pi iz}$, $z \in \mathbb{C}$, determines a projection of the strip Σ onto the annulus R , so that the triple (Σ, R, π) forms a principal bundle on R with fibre \mathbb{Z} . Suppose that U_1 and U_2 are compact simply connected subsets of R which cover R . Then we can represent Σ as a quotient space of $(U_1 \times \mathbb{Z}) \sqcup (U_2 \times \mathbb{Z})$ by the equivalence relation \sim determined by a locally constant function $c_{12} : U_1 \cap U_2 \mapsto \mathbb{Z}$ such that $U_1 \times \mathbb{Z} \ni (z, n) \sim (z, n + c_{12}(z)) \in U_2 \times \mathbb{Z}$ for all $z \in U_1 \cap U_2$, $n \in \mathbb{Z}$.

Let $b\mathbb{Z}$ be the Bohr compactification of \mathbb{Z} . Then $b\mathbb{Z}$ is a compact abelian group with $\mathbb{Z} \subset b\mathbb{Z}$ acting continuously on $b\mathbb{Z}$. In particular, given $z \in R$ we have a continuous mapping $b\mathbb{Z} \mapsto b\mathbb{Z}$ determined by the formula $\xi \mapsto \xi + c_{12}(z)$, $\xi \in b\mathbb{Z}$. So, we can define

$$b\Sigma := (U_1 \times b\mathbb{Z}) \sqcup (U_2 \times b\mathbb{Z}) / \sim$$

where, by definition, $U_1 \times b\mathbb{Z} \ni (z, \xi) \sim (z, \xi + c_{12}(z)) \in U_2 \times b\mathbb{Z}$ for all $z \in U_1 \cap U_2$, $\xi \in b\mathbb{Z}$. The local embeddings $U_k \times \mathbb{Z} \hookrightarrow U_k \times b\mathbb{Z}$ ($k = 1, 2$) determine an embedding

$\iota_0 : \Sigma \hookrightarrow b\Sigma$. Similarly, an embedding $\iota_\xi : \Sigma \hookrightarrow b\Sigma$, where $\xi \in b\mathbb{Z}$, is determined by the local embedding $(z, n) \mapsto (z, n + \xi)$. Since \mathbb{Z} is dense in $b\mathbb{Z}$, $\iota_\xi(\Sigma)$ is dense in $b\Sigma$ for each $\xi \in b\mathbb{Z}$. Furthermore, the sets $\iota_\xi(\Sigma)$ are mutually disjoint and cover $b\Sigma$.

We say that a function $f : b\Sigma \mapsto \mathbb{C}$ is *holomorphic* if its pullback $\iota_\xi^* f$ is holomorphic on Σ for each $\xi \in b\mathbb{Z}$. The space of functions holomorphic on $b\Sigma$ is denoted by $\mathcal{O}(b\Sigma)$. Now, we have

Proposition 2.1 ([2]). *$APH(\Sigma)$ is naturally isomorphic to $\mathcal{O}(b\Sigma)$.*

The result states that $\mathcal{O}(b\Sigma)|_\Sigma = APH(\Sigma)$. Moreover, as it is shown in [2], if $\iota_\xi^* f$ is holomorphic on Σ for a certain $\xi \in b\mathbb{Z}$, then f is holomorphic on $b\Sigma$. This gives another definition of a holomorphic almost periodic function (cf. Definition 1.3).

Proposition 2.2 ([2]). *$b\Sigma$ is homeomorphic to the maximal ideal space of the algebra $APH(\Sigma)$.*

2.2. In the proof of Theorem 1.4 we use the following equivalent definition of B -valued holomorphic functions on Σ (cf. Definition 1.3).

Proposition 2.3. *A continuous function $f : \Sigma \mapsto B$ is holomorphic almost periodic on Σ if and only if it is holomorphic in the interior of Σ and admits a continuous extension to $b\Sigma$.*

Proof. Assume that f is a B -valued continuous almost periodic function on Σ . Since $g \circ f \in APH(\Sigma)$ for every continuous functional $g \in B^*$, the function f admits a continuous extension $\hat{f} : b\Sigma \rightarrow B^{**}$; here B^{**} is equipped with weak* topology. Next, due to Definition 1.3, the closure $\overline{f(\Sigma)}$ of $f(\Sigma)$ in the strong topology of B is compact. We naturally identify B with a subspace of B^{**} . Since any compact subset of B is also compact in the weak* topology of B^{**} , $\hat{f}(b\Sigma) = \overline{f(\Sigma)} \subset B$. Hence \hat{f} maps $b\Sigma$ into B and is continuous.

Conversely, assume that $f : b\Sigma \rightarrow B$ is continuous and $f|_\Sigma$ is holomorphic. Since f is uniformly continuous on $b\Sigma$ and the natural action of group \mathbb{R} on Σ by shifts is extended to a continuous action of $b\mathbb{R}$ on $b\Sigma$, the family of shifts $\{S_x f\}_{x \in \mathbb{R}}$ is relatively compact in $C_b^B(b\Sigma) \subset C_b^B(\Sigma)$. \square

Now, suppose that $f \in APH_B(\Sigma)$. According to Proposition 2.3, there exists a continuous extension \hat{f} of f to $b\Sigma$. For each continuous functional $g \in B^*$ the function $g \circ f$ belongs to $APH(\Sigma)$. By Proposition 2.1, $g \circ f$ admits a continuous extension $\widehat{g \circ f} \in \mathcal{O}(b\Sigma)$. Since Σ is dense in $b\Sigma$, the identity $g \circ \hat{f} = \widehat{g \circ f}$ is valid for all $g \in B^*$. This implies that for each $\xi \in b\mathbb{Z}$, the function $g \circ \hat{f} \circ \iota_\xi \in APH(\Sigma)$ for all $g \in B^*$. Therefore the continuous B -valued function $\hat{f} \circ \iota_\xi$ is holomorphic in the interior of Σ , i.e., it belongs to $APH_B(\Sigma)$.

3 Proof of Theorem 1.4

Proposition 3.1. *$APH(\Sigma)$ has metric approximation property.*

Proof. We refer to the book of Besicovich [1] for the corresponding definitions and facts from the theory of almost periodic functions.

Let $K \subset APH(\Sigma)$ be compact. Given $\varepsilon > 0$ consider an $\frac{\varepsilon}{3}$ -net $\{f_1, \dots, f_l\} \subset K$. Let

$$\mathcal{K}(t) := \sum_{|\nu_1| \leq n_1, \dots, |\nu_r| \leq n_r} \left(1 - \frac{\nu_1}{n_1}\right) \dots \left(1 - \frac{\nu_r}{n_r}\right) e^{-i\left(\frac{\nu_1}{n_1}\beta_1 + \dots + \frac{\nu_r}{n_r}\beta_r\right)t}$$

be a Bochner-Fejer kernel such that for all $1 \leq k \leq l$

$$\sup_{z \in \Sigma} |f_k(z) - M_t\{f_k(z+t)\mathcal{K}(t)\}| \leq \frac{\varepsilon}{3}. \quad (3.1)$$

Here β_1, \dots, β_r are elements of a basis over \mathbb{Q} of the union of spectra of functions f_1, \dots, f_l , $\nu_1, \dots, \nu_r \in \mathbb{Z}$, $n_1, \dots, n_r \in \mathbb{N}$ and

$$M_t\{f_k(z+t)\mathcal{K}(t)\} := \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T f_k(z+t)\mathcal{K}(t) dt$$

are the corresponding holomorphic Bochner-Fejer polynomials.

We define an operator $T : APH(\Sigma) \rightarrow APH(\Sigma)$ from Definition 1.5 by the formula

$$(Tf)(z) := M_t\{f(z+t)\mathcal{K}(t)\}, \quad f \in APH(\Sigma). \quad (3.2)$$

Then T is a linear projection onto a finite-dimensional space generated by functions $e^{i\left(\frac{\nu_1}{n_1}\beta_1 + \dots + \frac{\nu_r}{n_r}\beta_r\right)z}$, $|\nu_1| \leq n_1, \dots, |\nu_r| \leq n_r$. Moreover, since $\mathcal{K}(t) \geq 0$ for all $t \in \mathbb{R}$ and $M_t\{\mathcal{K}(t)\} = 1$, the norm of T is 1. Finally, given $f \in K$ choose k such that $\|f - f_k\|_{APH(\Sigma)} \leq \frac{\varepsilon}{3}$. Then we have by (3.1)

$$\|Tf - f\|_{APH(\Sigma)} \leq \|T(f - f_k)\|_{APH(\Sigma)} + \|Tf_k - f_k\|_{APH(\Sigma)} + \|f_k - f\|_{APH(\Sigma)} < \varepsilon.$$

This completes the proof of the proposition. \square

Let us prove now Theorem 1.4. According to Section 2.2 each B -valued almost periodic function f on Σ admits a continuous extension $\hat{f} \in C_b^B(b\Sigma)$. Also, for each $g \in B^*$ the function $g \circ \hat{f} \in \mathcal{O}(b\Sigma)$. Since $\mathcal{O}(b\Sigma) \cong APH(\Sigma)$ has the approximation property by Proposition 3.1 and since $b\Sigma$ is compact, the results of Section 5.1 of [5] imply that each \hat{f} belongs to the closure in $C_b^B(b\Sigma)$ of the symmetric tensor product of $APH(\Sigma)$ and B . Finally, the classical Bohr theorem, see, e.g., [1], asserts that each element of $APH(\Sigma)$ is the uniform limit of finite linear combinations of functions $e^{i\lambda z}$, $\lambda \in \mathbb{R}$. These facts complete the proof of the theorem. \square

4 Proofs of Theorems 1.9 and 1.11

4.1 Semi-almost periodic holomorphic functions on \mathbb{D} with values in a Banach space

To prove Theorem 1.9 we generalize results of [2] to the Banach space of semi-almost periodic holomorphic functions on the unit disk \mathbb{D} with values in a Banach space B .

4.1.1. Let $L_B^\infty(\partial\mathbb{D})$ be the Banach space of B -valued bounded measurable functions on $\partial\mathbb{D}$ equipped with sup-norm.

Definition 4.1. A function $f \in L_B^\infty(\partial\mathbb{D})$ is called semi-almost periodic on $\partial\mathbb{D}$ if for any $t_0 \in [0, 2\pi)$ and any $\varepsilon > 0$ there exist a number $s = s(t_0, \varepsilon) \in (0, \pi)$ and functions $f_k : \gamma_{t_0}^k(s) \rightarrow B$, $\gamma_{t_0}^k(s) := \{e^{i(t_0+kt)} : 0 \leq t < s < 2\pi\}$, $k \in \{-1, 1\}$, such that functions

$$t \mapsto f_k(e^{i(t_0+kse^t)}), \quad -\infty < t < 0, \quad k \in \{-1, 1\},$$

are restrictions of some B -valued almost periodic functions from $AP_B(\mathbb{R})$ and

$$\sup_{z \in \gamma_{t_0}^k(s)} \|f(z) - f_k(z)\|_B < \varepsilon, \quad k \in \{-1, 1\}.$$

Analogously, we denote by $SAP_B(S) \subset L_B^\infty(S)$ the Banach space of semi-almost periodic functions that are continuous on $\partial\mathbb{D} \setminus S$.

Using the Poisson integral formula we can extend each function from $SAP_B(S)$ to a bounded B -valued harmonic function on \mathbb{D} having the same sup-norm. Below we identify $SAP_B(S)$ with its harmonic extension.

Suppose that S contains at least two points. Recall that A_S is the closure in $H^\infty(\mathbb{D})$ of the algebra generated by $A(\mathbb{D})$ and holomorphic functions of the form e^f , where $\operatorname{Re}(f)|_{\partial\mathbb{D}}$ is a finite linear combination (over \mathbb{R}) of characteristic functions of closed arcs in $\partial\mathbb{D}$ whose endpoints belong to S . If S consists of a single point, then A_S is the closure of the algebra generated by $A(\mathbb{D})$ and functions $ge^{\lambda f}$, where $\operatorname{Re}(f)|_{\partial\mathbb{D}}$ is the characteristic function of a closed arc with an endpoint in S and $g \in A(\mathbb{D})$ is a function such that ge^f has discontinuity on S only. (Here $A(\mathbb{D}) \subset H^\infty(\mathbb{D})$ is the disk-algebra, see Section 1.)

We define the Banach space A_S^B as

$$A_S^B = A_S \otimes B.$$

Here \otimes stands for completion of symmetric tensor product with respect to norm

$$\left\| \sum_{k=1}^m f_k b_k \right\| := \sup_{z \in \partial\mathbb{D}} \left\| \sum_{k=1}^m f_k(z) b_k \right\|_B \quad \text{with } f_k \in A_S, b_k \in B. \quad (4.1)$$

Let $H_B^\infty(\mathbb{D})$ be the Banach space of bounded B -valued holomorphic functions on \mathbb{D} equipped with sup-norm. The Banach-valued analogue of Theorems 1.9 is now formulated as follows.

Theorem 4.2. $SAP_B(S) \cap H_B^\infty(\mathbb{D}) = A_S^B$.

4.1.2. For the proof we require some auxiliary results.

Let $APC(\Sigma)$ be the Banach algebra of functions $f : \Sigma \mapsto \mathbb{C}$ uniformly continuous on $\Sigma := \{z \in \mathbb{C} : \text{Im}(z) \in [0, \pi]\}$ and almost periodic on each horizontal line.

Proposition 4.3 ([2]). $M(APH(\Sigma)) = M(APC(\Sigma))$.

We define $APC_B(\Sigma) = APC(\Sigma) \otimes B$.

Lemma 4.4. *Suppose that $f_1 \in AP_B(\mathbb{R})$, $f_2 \in AP_B(\mathbb{R} + i\pi)$. Then there exists a function $F \in APC_B(\Sigma)$ which is harmonic in the interior of Σ whose boundary values are f_1 and f_2 . Moreover, F admits a continuous extension to the maximal ideal space $M(APH(\Sigma))$.*

Proof. By definition f_1 and f_2 can be approximated on \mathbb{R} and $\mathbb{R} + i\pi$ by functions of the form

$$q_1 = \sum_{l=0}^{k_1} b_l e^{i\lambda_l t}, \quad q_2 = \sum_{l=0}^{k_2} c_l e^{i\mu_l t},$$

respectively, where $b_l, c_l \in B$, $\lambda_l, \mu_l \in \mathbb{R}$. Using the Poisson integral formula we extend q_1 and q_2 from the boundary to a function $H \in APC_B(\Sigma)$ harmonic in the interior of Σ , see, e.g., [2], Lemma 4.3 for similar arguments. Also, by the definition of the algebra $APC_B(\Sigma)$ and by Proposition 4.3, H admits a continuous extension to $M(APH(\Sigma))$. Now, by the maximum principle for harmonic functions the sequence of functions H converges in $APC_B(\Sigma)$ to a certain function F harmonic in the interior of Σ which satisfies the required properties of the lemma. \square

Let $\phi_{z_0} : \mathbb{D} \rightarrow \mathbb{H}_+$,

$$\phi_{z_0}(z) := \frac{2i(z_0 - z)}{z_0 + z}, \quad z \in \mathbb{D}, \quad (4.2)$$

be a conformal map of \mathbb{D} onto the upper half-plane \mathbb{H}_+ . Then ϕ_{z_0} is also continuous on $\partial\mathbb{D} \setminus \{-z_0\}$ and maps it diffeomorphically onto \mathbb{R} (the boundary of \mathbb{H}_+) so that $\phi_{z_0}(z_0) = 0$. Let Σ_0 be the interior of the strip Σ . Consider the conformal map $\text{Log} : \mathbb{H}_+ \rightarrow \Sigma_0$, $z \mapsto \text{Log}(z) := \ln|z| + i\text{Arg}(z)$, where $\text{Arg} : \mathbb{C} \setminus \mathbb{R}_- \rightarrow (-\pi, \pi)$ is the principal branch of the multi-function \arg , the argument of a complex number. The function Log is extended to a homeomorphism of $\overline{\mathbb{H}_+} \setminus \{0\}$ onto Σ ; here $\overline{\mathbb{H}_+}$ stands for the closure of \mathbb{H}_+ .

The proof of the next statement uses Lemma 4.4 and is very similar to the proof of Lemma 4.2 (for $B = \mathbb{C}$) in [2], so we omit it.

Suppose that $z_0 = e^{it_0}$. For $s \in (0, \pi)$ we set $\gamma_1(z_0, s) := \text{Log}(\phi_{z_0}(\gamma_{t_0^1}(s))) \subset \mathbb{R}$ and $\gamma_{-1}(z_0, s) := \text{Log}(\phi_{z_0}(\gamma_{t_0^{-1}}(s))) \subset \mathbb{R} + i\pi$.

Lemma 4.5. *Let $z_0 \in S$, suppose that $f \in \text{SAP}_B(\{-z_0, z_0\})$. We put $f_k = f|_{\gamma_{t_0^k}(\pi)}$, and define on arc $\gamma_k(z_0, s)$*

$$h_k = f_k \circ \varphi_{z_0}^{-1} \circ \text{Log}^{-1}, \quad k \in \{-1, 1\}.$$

Then for any $\varepsilon > 0$ there exist a number $s_\varepsilon \in (0, s)$ and a function $H \in \text{APC}_B(\Sigma)$ harmonic on Σ_0 such that

$$\sup_{z \in \gamma_k(z_0, s_\varepsilon)} \|h_k(z) - H(z)\|_B < \varepsilon, \quad k \in \{-1, 1\}.$$

Let $z_0 \in \partial\mathbb{D}$ and U_{z_0} be the intersection of an open disk of radius ≤ 1 centered at z_0 with $\bar{\mathbb{D}} \setminus z_0$. We call such U_{z_0} a *circular neighbourhood* of z_0 .

We say that a bounded continuous function $f : \mathbb{D} \mapsto B$ is *almost-periodic near* z_0 if there exist a circular neighbourhood U_{z_0} , and a function $\hat{f} \in \text{APC}_B(\Sigma)$ such that

$$f(z) = \hat{f}(\text{Log}(\varphi_{z_0}(z))), \quad z \in U_{z_0}.$$

Let M_S be the maximal ideal space of $H^\infty(\mathbb{D}) \cap \text{SAP}(S)$ and $a_S : M_S \rightarrow \bar{\mathbb{D}}$ be the natural continuous surjection, see Remark 1.12. Recall that for $z_0 \in S$, $a_S^{-1}(z_0)$ can be naturally identified with $b\Sigma := M(\text{APH}(\Sigma))$. Therefore the algebra $\mathcal{O}(a_S^{-1}(z_0))$ of holomorphic functions on $a_S^{-1}(z_0)$ can be defined similarly to $\mathcal{O}(b\Sigma)$ from Section 2 (by means of this identification), see [2] for details.

In the proof of Theorem 1.8 of [2] (see Lemmas 4.4, 4.6 there) we established

- (1) Any scalar harmonic function f on \mathbb{D} almost periodic near z_0 admits a continuous extension to $a_S^{-1}(\bar{U}_{z_0}) \subset M_S$ for some circular neighbourhood U_{z_0} .
- (2) For any holomorphic function $f \in \mathcal{O}(a_S^{-1}(z_0))$, $z_0 \in S$, there is a bounded holomorphic function \hat{f} on \mathbb{D} of the same sup-norm almost periodic near z_0 such that its extension to $a_S^{-1}(z_0)$ coincides with f .

Similarly to $\mathcal{O}(b\Sigma)$ we define the Banach space $\mathcal{O}_B(b\Sigma)$ of B -valued holomorphic functions on $b\Sigma$, cf. Section 2. Then by Theorem 1.4, $\mathcal{O}_B(b\Sigma) := \mathcal{O}(b\Sigma) \otimes B$. This together with the B -valued Bohr approximation theorem for the subalgebra of harmonic functions in $\text{APC}_B(\Sigma)$ (see Lemma 4.4) and statements (1), (2) imply

- (3) Any B -valued harmonic function f on \mathbb{D} almost periodic near z_0 admits a continuous extension to $a_S^{-1}(\bar{U}_{z_0}) \subset M_S$ for some circular neighbourhood U_{z_0} .

- (4) For any holomorphic function $f \in \mathcal{O}_B(a_S^{-1}(z_0))$, $z_0 \in S$, there is a bounded B -valued holomorphic function \hat{f} on \mathbb{D} of the same sup-norm almost periodic near z_0 such that its extension to $a_S^{-1}(z_0)$ coincides with f .

From (3) and (4) we obtain

Lemma 4.6. *Let $f \in SAP_B(S) \cap H_B^\infty(\mathbb{D})$ and $z_0 \in \partial\mathbb{D}$. There is a bounded B -valued holomorphic function \hat{f} on \mathbb{D} almost periodic near z_0 such that for any $\varepsilon > 0$ there is a circular neighbourhood $U_{z_0;\varepsilon}$ of z_0 so that*

$$\sup_{z \in U_{z_0;\varepsilon}} \|f(z) - \hat{f}(z)\|_B < \varepsilon.$$

Proof. Assume, first, that $z_0 \in S$. By Lemma 4.5, for any $n \in \mathbb{N}$ there exist a number $s_n \in (0, s)$ and a function $H_n \in APC_B(\Sigma)$ harmonic on Σ_0 such that

$$\sup_{z \in \gamma_k(z_0, s_n)} \|f_k(z) - H_n(z)\|_B < \frac{1}{n}, \quad k \in \{-1, 1\}. \quad (4.3)$$

Using the Poisson integral formula for the bounded B -valued harmonic function $f - H_n$ on \mathbb{D} we easily obtain from (4.3) that there is a circular neighbourhood $V_{z_0;n}$ of z_0 such that

$$\sup_{z \in V_{z_0;n}} \|f(z) - H_n(z)\|_B < \frac{2}{n}. \quad (4.4)$$

According to (3) each H_n admits a continuous extension \hat{H}_n to $a_S^{-1}(z_0) \cong b\Sigma$. Moreover, (4.4) implies that the restriction of the sequence $\{\hat{H}_n\}_{n \in \mathbb{N}}$ to $a_S^{-1}(z_0)$ forms a Cauchy sequence in $C_B(a_S^{-1}(z_0))$. Let $\hat{H} \in C_B(a_S^{-1}(z_0))$ be the limit of this sequence.

Further, for any functional $\phi \in B^*$ the function $\phi \circ f \in SAP(S) \cap H^\infty(\mathbb{D})$ and therefore admits a continuous extension f_ϕ to $a_S^{-1}(z_0)$ such that on $a_S^{-1}(z_0)$ the extended function belongs to $\mathcal{O}(a_S^{-1}(z_0))$. Now, (4.4) implies directly that $f_\phi = \phi \circ \hat{H}$ for any $\phi \in B^*$. Then from the definition of $\mathcal{O}_B(a_S^{-1}(z_0))$, see Section 2, follows that $\hat{H} \in \mathcal{O}_B(a_S^{-1}(z_0))$. Thus by (4) we find a bounded B -valued holomorphic function \hat{f} on \mathbb{D} of the same sup-norm as \hat{H} almost periodic near z_0 such that its extension to $a_S^{-1}(z_0)$ coincides with \hat{H} . Now by the definition of the topology of M_S , see [2], Lemma 4.4 (a), we obtain that for any $\varepsilon > 0$ there is a number $N \in \mathbb{N}$ such that for all $n \geq N$,

$$\sup_{z \in V_{z_0;n}} \|\hat{f}(z) - H_n(z)\|_B < \frac{\varepsilon}{2}.$$

Finally, choose $n \geq N$ in (4.4) such that the right-hand there $< \frac{\varepsilon}{2}$. For this n we set $U_{z_0;\varepsilon} := V_{z_0;n}$. Then the previous inequality and (4.4) imply the required

$$\sup_{z \in U_{z_0;\varepsilon}} \|f(z) - \hat{f}(z)\|_B < \varepsilon.$$

If, now, $z_0 \notin S$, then, by definition, $f|_{\partial\mathbb{D}}$ is continuous at z_0 . In this case as the function \hat{f} we can choose the constant B -valued function equal to $f(z_0)$ on \mathbb{D} . Then the required result follows from the Poisson integral formula for $f - \hat{f}$. We leave the details to the reader. \square

4.1.3. We are now ready to prove Theorem 4.2.

Proof. As it was shown in [2], $A_S \subset SAP(S) \cap H^\infty(\mathbb{D})$. So $A_S^B \subset SAP_B(S) \cap H_B^\infty(\mathbb{D})$. Let us prove the opposite inclusion.

(A) Consider first the case $S = F$, where $F = \{z_i\}_{i=1}^m$ is a finite subset of $\partial\mathbb{D}$.

Let $f \in SAP_B(F) \cap H_B^\infty(\mathbb{D})$. Then according to Lemma 4.6 there exists a function $f_{z_1} \in APH_B(\Sigma)$ such that the bounded B -valued holomorphic function $g_{z_1} - f$, $g_{z_1} := f_{z_1} \circ \text{Log} \circ \varphi_{z_0}$, on \mathbb{D} is continuous and equals 0 at z_1 .

Let us show that $g_{z_1} \in A_{\{z_1, -z_1\}}^B$. Indeed, since $f_{z_1} \in APH_B(\Sigma)$, by Theorem 1.4 it can be approximated in $APH_B(\Sigma)$ by finite sums of functions $be^{i\lambda z}$, $b \in B$, $\lambda \in \mathbb{R}$, $z \in \Sigma$. In turn, g_{z_1} can be approximated by finite sums of functions $be^{i\lambda \text{Log} \circ \varphi_{z_1}}$. As it was shown in [2], $e^{i\lambda \text{Log} \circ \varphi_{z_1}} \in A_{\{z_1, -z_1\}}$. Hence, $g_{z_1} \in A_{\{z_1, -z_1\}}^B$.

We define

$$\hat{g}_{z_1} = \frac{g_{z_1}(z)(z + z_1)}{2z_1}.$$

Then, since the function $z \mapsto (z + z_1)/(2z_1) \in A(\mathbb{D})$ and equals 0 at $-z_1$, and $g_{z_1} \in A_{\{z_1, -z_1\}}^B$, the function $\hat{g}_{z_1} \in A_{\{z_1\}}^B$. Moreover, by the definitions of g_{z_1} and \hat{g}_{z_1} , the function $\hat{g}_{z_1} - f$ is continuous and equal to zero at z_1 . Thus,

$$\hat{g}_{z_1} - f \in SAP(F \setminus \{z_1\}) \cap H_B^\infty(\mathbb{D}).$$

We proceed in this way to get functions $\hat{g}_{z_k} \in A_{\{z_k\}}^B$ such that

$$f - \sum_{k=1}^m \hat{g}_{z_k} \in A_B(\mathbb{D}).$$

Here $A_B(\mathbb{D})$ is the Banach space of B -valued bounded holomorphic functions on \mathbb{D} continuous up to the boundary. As in the scalar case using the Taylor expansion at 0 of functions from $A_B(\mathbb{D})$ one can easily show that $A_B(\mathbb{D}) = A(\mathbb{D}) \otimes B$. This and the above implication imply that $f \in A_F^B := A_F \otimes B$, as required.

(B) Let us consider the general case of $S \subset \partial\mathbb{D}$ an arbitrary closed set.

Let $f \in SAP_B(S) \cap H_B^\infty(\mathbb{D})$. As it follows from Lemma 4.6 and the arguments presented in part (A), given an $\varepsilon > 0$ there exist points $z_k \in \partial\mathbb{D}$, functions $f_k \in A_{\{z_k\}}^B$ and circular neighbourhoods U_{z_k} ($1 \leq k \leq m$) such that $\{U_k\}_{k=1}^m$ forms an open cover of set $\partial\mathbb{D} \setminus \{z_k\}_{k=1}^m$ and

$$\|f(z) - f_k(z)\|_B < \varepsilon \quad \text{on } U_{z_k}, \quad 1 \leq k \leq m. \quad (4.5)$$

Since S is closed, for $z_k \notin S$ we may assume that f_k is continuous in \bar{U}_{z_k} .

Let us define a B -valued 1-cocycle $\{c_{kj}\}_{k,j=1}^m$ on intersections of the sets in $\{U_{z_k}\}_{k=1}^m$ by the formula

$$c_{kj}(z) := f_k(z) - f_j(z), \quad z \in U_{z_k} \cap U_{z_j}. \quad (4.6)$$

Then (4.5) implies

$$\sup_{k,j,z} \|c_{kj}(z)\|_B < 2\varepsilon. \quad (4.7)$$

Since $m < \infty$, we may assume, without loss of generality, that none of the intersections $U_{z_k} \cap U_{z_j}$, $k \neq j$, contains points z_k . Further, we may choose the above functions and sets such that c_{kj} is holomorphic in $U_{z_k} \cap U_{z_j}$ and continuous in the closure of $U_{z_k} \cap U_{z_j}$. Let $\{\rho_k\}_{k=1}^m$ be a smooth partition of unity subordinate to the open cover $\{U_{z_k}\}_{k=1}^m$ of an open annulus $A \subset\subset \cup_{k=1}^m U_{z_k}$ with outer boundary $\partial\mathbb{D}$ such that each ρ_k is the restriction of a smooth function defined on a neighbourhood of $\partial\mathbb{D}$ and $\rho_k(z_k) = 1$, $1 \leq k \leq m$. We resolve the cocycle $\{c_{kj}\}_{k,j=0}^m$ using this partition of unity by formulas

$$\tilde{f}_j(z) = \sum_{k=1}^m \rho_k(z) c_{kj}(z), \quad z \in U_{z_j} \cap A,$$

so, by definition,

$$c_{kj}(z) = \tilde{f}_k(z) - \tilde{f}_j(z), \quad z \in U_{z_k} \cap U_{z_j} \cap A. \quad (4.8)$$

Now since c_{kj} are B -valued holomorphic functions in $U_{z_i} \cap U_{z_j}$ continuous up to the boundary, ρ_{kj} are smooth functions on a neighbourhood of $\partial\mathbb{D}$ and $A \subset\subset \cup_{k=1}^m U_{z_k}$,

$$h(z) := \frac{\partial \tilde{f}_j(z)}{\partial \bar{z}}, \quad z \in U_{z_j} \cap A, \quad 1 \leq j \leq m,$$

is a smooth global B -valued function on A continuous up to the boundary.

We define

$$H(z) := \frac{1}{2\pi i} \int_{\zeta \in \mathbb{D}} \frac{h(\zeta)}{\zeta - z} d\zeta \wedge d\bar{\zeta}, \quad z \in \bar{\mathbb{D}}. \quad (4.9)$$

Similarly to the scalar case we use polar coordinates in (4.9) to obtain

$$\sup_{z \in A} \|H(z)\|_B \leq Cw(A) \sup_{z \in A} \|h(z)\|_B,$$

where $w(A)$ is width of A and $C > 0$ is a numerical constant. Furthermore, H is continuous on \bar{A} and $\frac{\partial H}{\partial \bar{z}} = h$ on A . Without loss of generality (see (4.7) and the definition of functions ρ_j) we may assume that $w(A)$ is sufficiently small, so that

$$\sup_{z \in A} \|H(z)\|_B < \varepsilon.$$

Let us define

$$c_i(z) := \tilde{f}_i(z) - H(z), \quad z \in \bar{U}_{z_i} \cap \bar{A}. \quad (4.10)$$

Then c_i is a B -valued continuous function holomorphic in $U_{z_i} \cap A$ satisfying (see (4.7))

$$\sup_{z \in U_{z_i} \cap A} \|c_i(z)\|_B \leq 3\varepsilon. \quad (4.11)$$

By (4.8),

$$c_i(z) - c_j(z) = c_{ij}(z), \quad z \in U_{z_i} \cap U_{z_j} \cap A. \quad (4.12)$$

Let us determine a function f_ε defined on $\bar{A} \setminus \{z_i\}_{i=1}^m$ by formulas

$$f_\varepsilon(z) := f_i(z) - c_i(z), \quad z \in U_{z_i} \cap \bar{A}.$$

According to (4.6) and (4.12), f_ε is a bounded continuous B -valued function on $\bar{A} \setminus \{z_i\}_{i=1}^m$ holomorphic in A . Furthermore, since c_i is continuous on $\bar{U}_{z_i} \cap \bar{A}$, and $f_i \in A_{\{z_i\}}^B$, for $z_i \in S$ and $f_i \in A_B(\mathbb{D})$ otherwise, $f_\varepsilon|_{\partial\mathbb{D}} \in SAP_B(F)$ where $F = \{z_i\}_{i=1}^m \cap S$. Also, from inequalities (4.5) and (4.11) we have

$$\sup_{z \in A} \|f(z) - f_\varepsilon(z)\|_B < 4\varepsilon. \quad (4.13)$$

Now let D' be an open disk centered at 0 whose intersection with A is an annulus of width $\frac{w(A)}{2}$. Let A' be the open annulus with the interior boundary coinciding with the interior boundary of A and with the outer boundary $\{z \in \mathbb{C} : |z| = 2\}$. Then $\{A', D'\}$ forms an open cover of \mathbb{D} . Let us define a 1-cocycle on $A' \cap D' = A \cap D'$ by the formula

$$c(z) = f(z) - f_\varepsilon(z), \quad z \in A \cap D'.$$

As it follows from (4.13), $\|c(z)\|_B < 4\varepsilon$ for all $z \in D' \cap A'$. Consider a smooth partition of unity subordinate to the cover $\{A', D'\}$ of \mathbb{D} which consists of smooth radial functions $\rho_{A'}, \rho_{D'}$ defined on \mathbb{C} such that

$$\max\{\|\nabla \rho_{A'}\|_{L^\infty(\mathbb{C})}, \|\nabla \rho_{D'}\|_{L^\infty(\mathbb{C})}\} \leq \frac{C}{w(D' \cap A')} = \frac{2C}{w(A)} \quad (4.14)$$

where $C > 0$ is a numerical constant. We resolve the cocycle c as follows:

$$\begin{aligned} f_{A'}(z) &= -\rho_{D'}(z)c(z), \quad z \in A', \\ f_{D'}(z) &= \rho_{A'}(z)c(z), \quad z \in D'. \end{aligned}$$

Then $c(z) = f_{D'}(z) - f_{A'}(z)$, $z \in A' \cap D'$. Since c is holomorphic in $A' \cap D'$,

$$g(z) = \begin{cases} \frac{\partial f_{A'}(z)}{\partial \bar{z}}, & z \in A', \\ \frac{\partial f_{D'}(z)}{\partial \bar{z}}, & z \in D' \end{cases} \quad (4.15)$$

is a bounded smooth B -valued function on \mathbb{D} with support in A .

Next,

$$G(z) = \frac{1}{2\pi i} \int_{\zeta \in D} \frac{g(\zeta)}{\zeta - z} d\zeta \wedge d\bar{\zeta} \quad (4.16)$$

is a smooth B -valued function on \mathbb{D} , continuous up to the boundary such that $\frac{\partial G}{\partial \bar{z}} = g$ on \mathbb{D} . Moreover, by (4.13), (4.14) and the fact that $\text{supp}(g) \subset A$ we obtain from (4.16)

$$\sup_{z \in \mathbb{D}} \|G(z)\|_B \leq C' w(A) \frac{\varepsilon}{w(A)} = C' \varepsilon \quad (4.17)$$

for a numerical constant $C' > 0$.

Now we define

$$\begin{aligned} c_{A'}(z) &= f_{A'}(z) - G(z), \quad z \in A, \\ c_{D'}(z) &= f_{D'}(z) - G(z), \quad z \in D'. \end{aligned}$$

Then $c_{A'}$ and $c_{D'}$ are B -valued holomorphic functions in A and D' , respectively. Clearly, we have $c_{D'}(z) - c_{A'}(z) = c(z)$ for all $z \in D' \cap A'$. Furthermore, as it follows from (4.13), (4.17) there exists a numerical constant $\bar{C} > 0$ such that

$$\|c_{A'}(z)\|_B < \bar{C}\varepsilon, \quad z \in A, \quad \|c_{D'}(z)\|_B < \bar{C}\varepsilon, \quad z \in D'.$$

Finally, let us define

$$F_\varepsilon(z) := \begin{cases} f(z) - c_{D'}(z), & z \in D', \\ f_\varepsilon(z) - c_{A'}(z), & z \in A. \end{cases}$$

Clearly, F_ε is a B -valued holomorphic function on \mathbb{D} (since for every $z \in D' \cap A$ we have $f(z) - f_\varepsilon(z) - c_{D'}(z) + c_{A'}(z) = f(z) - f_\varepsilon(z) - c(z) = 0$). Also,

$$\sup_{z \in \mathbb{D}} \|f(z) - F_\varepsilon(z)\|_B < \hat{C}\varepsilon$$

for a numerical constant $\hat{C} > 0$, and by definition $F_\varepsilon \in SAP_B(F) \cap H_B^\infty(\mathbb{D})$, where $F = \{z_1, \dots, z_m\} \cap S$.

The last inequality and part (A) of the proof show that the complex vector space generated by spaces A_F^B for all possible finite subsets $F \subset S$ is dense in $SAP_B(S) \cap H_B^\infty(\mathbb{D})$. Since the closure of all such A_F^B is A_S^B , we obtain the required: $SAP_B(S) \cap H_B^\infty(\mathbb{D}) = A_S^B$. \square

Remark 4.7. From the proof of Theorem 4.2 one obtains also that each function from $SAP_B(S) \cap H_B^\infty(\mathbb{D})$ admits a continuous extension to the maximal ideal space of the algebra $SAP(S) \cap H^\infty(\mathbb{D})$. Then this theorem and the results of Section 5.1 of [5] imply that $SAP(S) \cap H^\infty(\mathbb{D})$ has the approximation property.

4.2 Proofs

Proof of Theorem 1.9. Let $S = \prod_{k=1}^n S_k \subset (\partial\mathbb{D})^n$ where $S_k \subset \partial\mathbb{D}$ is a closed set. We must prove that $H^\infty(\mathbb{D}^n) \cap SAP^n(S) = A_S^n$.

We prove this statement by induction over n . For $n = 1$ the identity was already proved. Suppose that it is true for $n - 1$, that is, $H^\infty(\mathbb{D}^{n-1}) \cap SAP^{n-1}(S') = A_{S'}^{n-1}$ where $S' := \prod_{k=1}^{n-1} S_k$, and prove it for n . Let $S = S' \times S_n$.

Lemma 4.8.

$$SAP^n(S) = SAP_B(S_n)$$

for $B = SAP^{n-1}(S')$ and

$$A_S^n = A_{S_n}^C$$

for $C = A_{S'}^{n-1}$.

Proof. The proof follows directly from the definitions of $SAP^n(S)$ and A_S^n . \square

Let $f \in H^\infty(\mathbb{D}^n) \cap SAP^n(S)$. According to Lemma 4.8, $f \in SAP_B(S_n)$ for $B = SAP^{n-1}(S')$. Also, by the Poisson integral formula for bounded polyharmonic functions on \mathbb{D}^n , and the fact that $f \in H^\infty(\mathbb{D}^n)$ we obtain that $f \in H_B^\infty(\mathbb{D})$. Thus, in virtue of Theorem 4.2,

$$f \in SAP_B(S_n) \cap H_B^\infty(\mathbb{D}) = A_{S_n}^B. \quad (4.18)$$

Let $\xi = (\xi_1, \dots, \xi_n)$ be coordinates on $(\partial\mathbb{D})^n$. Applying again the Poisson integral formula for bounded polyharmonic functions on \mathbb{D}^n we obtain that the functions $f(\cdot, \dots, \cdot, \xi_n) \in H^\infty(\mathbb{D}^{n-1})$ for almost all $\xi_n \in \partial\mathbb{D}$. Thus, by the induction hypothesis, for almost all $\xi_n \in \mathbb{D}$ we have $f(\cdot, \dots, \cdot, \xi_n) \in H^\infty(\mathbb{D}^{n-1}) \cap B = A_{S'}^{n-1}$. From here and (4.18) we get $f \in A_{S_n}^C$, $C := A_{S'}^{n-1}$. Now, Lemma 4.8 implies that $f \in A_S^n$. This shows that $H^\infty(\mathbb{D}^n) \cap SAP^n(S) \subset A_S^n$. The converse embedding $A_S^n \subset H^\infty(\mathbb{D}^n) \cap SAP^n(S)$ follows immediately from the definition of A_S^n . \square

Proof of Theorem 1.11. According to Theorem 1.9 and the scalar version of Theorem 4.2, the algebra $H^\infty(\mathbb{D}^n) \cap SAP^n(S)$ is symmetric tensor product of algebras $H^\infty(\mathbb{D}) \cap SAP(S_k)$, $1 \leq k \leq n$. Therefore by a standard result of the theory of Banach algebras we obtain that M_S is homeomorphic to $M_{S_1} \times \dots \times M_{S_n}$ for the corresponding maximal ideal spaces of these algebras. As it was proved in [2], \mathbb{D} is dense in each M_{S_k} in the corresponding Gelfand topology. Therefore \mathbb{D}^n is dense in M_S in the Gelfand topology. \square

References

- [1] A. S. Besicovich, *Almost periodic functions*. Dover Publications, 1958.
- [2] A. Brudnyi and D. Kinzebulatov, On uniform subalgebras of L^∞ on the unit circle generated by almost periodic functions. *Algebra and Analysis* **19** (2007), 1–33.
- [3] P. Enflo, A counterexample to the approximation property in Banach spaces. *Acta Math.* **130** (1973), 309–317.
- [4] J. Garnett, *Bounded analytic functions*. Academic Press, 1981.
- [5] A. Grothendieck, Produits tensoriels topologiques et espaces nucléaires. *Memoirs Amer. Math. Society* **16**, 1955.
- [6] D. Sarason, Toeplitz operators with semi-almost periodic kernels. *Duke Math J.* **44** (1977), 357–364.