

Reflection matrices for the  $U_q[spo(2n|2m)]$  vertex model

A. Lima-Santos

*Universidade Federal de São Carlos**Departamento de Física**C.P. 676, 13565-905, São Carlos-SP, Brasil***Abstract**

We propose a classification of the solutions of the graded reflection equations to the  $U_q[spo(2n|2m)]$  vertex model. We find twelve distinct classes of reflection matrices such that four of them are diagonal. In the non-diagonal matrices the number of free parameters depending on the number of bosonic ( $2n$ ) and fermionic ( $2m$ ) degrees of freedom while in the diagonal ones we find solutions with at most one free parameter.

PACS numbers: 05.50+q, 02.30.IK, 75.10.Jm

Keywords: Reflection Equations, K-matrices, Superalgebras

October 31, 2018

# 1 Introduction

The present work is the third of three papers devoted to the classification of the integrable reflection  $K$ -matrices for the vertex models associated with the  $U_q[sl(r|2m)^{(2)}]$ ,  $U_q[osp(r|2m)^{(1)}]$  and  $U_q[spo(2n|2m)]$  Lie superalgebras. In the first paper [1] we have considered the  $U_q[sl(r|2m)^{(2)}]$  vertex model and in the second paper [2] the  $U_q[osp(r|2m)^{(1)}]$  vertex model ones. In this paper we have presented the general set of regular solutions of the graded reflection equation for the  $U_q[spo(2n|2m)]$  vertex model.

It is well-known that the Lie superalgebra  $osp(2m|2n)$  and  $spo(2n|2m)$  are naturally isomorphic: From the classical Lie superalgebras[3], we have the notation  $D(m|n) = osp(2m|2n) = spo(2n|2m)$ . Thus, the transition from  $osp(2m|2n)^{(1)}$  to  $spo(2n|2m)$  amounts to shift of the gradation of the vector representation[4]. But, we will work with the natural gradations in order to avoid any shift of gradations. Moreover, for trivial values of  $m$  and  $n$ , a Lie superalgebra coincides with a Lie algebra:  $D(m|0) = D_m^{(1)}$ ,  $D(0|n) = C_n^{(1)}$ . Therefore, by considering the  $U_q[spo(2n|2m)]$  vertex model we are completing the study of the integrable  $K$ -matrices for the graded version of the vertex models associated with the  $B_n^{(1)}$ ,  $C_n^{(1)}$ ,  $D_n^{(1)}$ ,  $A_{2n}^{(2)}$  and  $A_{2n-1}^{(2)}$  Lie algebras.

Our findings can be summarized into four classes of diagonal solutions and eight classes of non-diagonal ones. This paper is organized as follows. In the next section we present the  $R$ -matrix of the  $U_q[spo(2n|2m)]$  vertex model in terms of standard Weyl matrices. In the section 3 we present four classes of diagonal solutions. In the section 4 we present eight classes of non-diagonal solutions what we hope to be the most general set of  $K$ -matrices for the vertex model here considered. Concluding remarks are discussed in the section 5, and in the appendix A we present solutions associated with the  $U_q[spo(2|2)]$  case.

## 2 The $U_q[spo(2n|2m)]$ vertex model

The  $U_q[spo(2n|2m)]$  invariant  $R$ -matrices are given by[5]

$$\begin{aligned}
 R(x) = & \sum_{\substack{i=1 \\ i \neq i'}}^N (-1)^{p_i} a_i(x) \hat{e}_{ii} \otimes \hat{e}_{ii} + b(x) \sum_{\substack{i,j=1 \\ i \neq j, i \neq j'}}^N \hat{e}_{ii} \otimes \hat{e}_{jj} \\
 & + \bar{c}(x) \sum_{\substack{i,j=1 \\ i < j, i \neq j'}} (-1)^{p_i p_j} \hat{e}_{ji} \otimes \hat{e}_{ij} + c(x) \sum_{\substack{i,j=1 \\ i > j, i \neq j'}}^N (-1)^{p_i p_j} \hat{e}_{ji} \otimes \hat{e}_{ij} \\
 & + \sum_{i,j=1}^N (-1)^{p_i} d_{i,j}(x) \hat{e}_{ij} \otimes \hat{e}_{i'j'} \tag{1}
 \end{aligned}$$

where  $N = 2n + 2m$  is the dimension of the graded space with  $2n$  bosonic and  $2m$  fermionic degrees of freedom. Here  $i' = N + 1 - i$  corresponds to the conjugated index of  $i$  and  $\hat{e}_{ij}$  refers to a usual  $N \times N$  Weyl matrix with only one non-null entry with value 1 at the row  $i$  and column  $j$ .

In what follows we shall adopt the grading structure

$$p_i = \begin{cases} 0 & \text{for } i = 1, \dots, m \text{ and } i = 2n + m + 1, \dots, N \\ 1 & \text{for } i = m + 1, \dots, 2n + m \end{cases}, \quad (2)$$

and the corresponding Boltzmann weights  $a_i(x)$ ,  $b(x)$ ,  $c(x)$ ,  $\bar{c}(x)$  and  $d_{ij}(x)$  are then given by

$$\begin{aligned} a_i(x) &= (x - \zeta)(x^{1-p_i} - q^2 x^{p_i}), & b(x) &= q(x - 1)(x - \zeta), \\ c(x) &= (1 - q^2)(x - \zeta), & \bar{c}(x) &= x(1 - q^2)(x - \zeta) \end{aligned} \quad (3)$$

and

$$d_{i,j}(x) = \begin{cases} q(x - 1)(x - \zeta) + x(q^2 - 1)(\zeta - 1), & (i = j = j') \\ (x - 1)[(x - \zeta)(-1)^{p_i} q^{2p_i} + x(q^2 - 1)], & (i = j \neq j') \\ (q^2 - 1)[\zeta(x - 1) \frac{\theta_i q^{t_i}}{\theta_j q^{t_j}} - \delta_{i,j'}(x - \zeta)], & (i < j) \\ (q^2 - 1)x[(x - 1) \frac{\theta_i q^{t_i}}{\theta_j q^{t_j}} - \delta_{i,j'}(x - \zeta)], & (i > j) \end{cases} \quad (4)$$

where  $\zeta = q^{2n-2m+2}$ . The remaining variables  $\theta_i$  and  $t_i$  depend strongly on the grading structure considered and they are determined by the relations

$$\theta_i = \begin{cases} (-1)^{-\frac{p_i}{2}}, & 1 \leq i \leq \frac{N}{2} \\ -(-1)^{\frac{p_i}{2}}, & \frac{N}{2} + 1 \leq i \leq N \end{cases} \quad (5)$$

$$t_i = \begin{cases} i - [\frac{1}{2} + p_i - 2 \sum_{j=i}^{\frac{N}{2}} p_j], & 1 \leq i \leq \frac{N}{2} \\ i + [\frac{1}{2} + p_i - 2 \sum_{j=\frac{N}{2}+1}^i p_j], & \frac{N}{2} + 1 \leq i \leq N \end{cases} \quad (6)$$

The  $R$ -matrix (1) satisfies important symmetry relations, besides the standard properties of regularity and unitarity, namely

$$\begin{aligned} \text{PT - Symmetry :} & \quad R_{21}(x) = R_{12}^{st_1 st_2}(x) \\ \text{Crossing Symmetry :} & \quad R_{12}(x) = \frac{\rho(x)}{\rho(x^{-1}\eta^{-1})} V_1 R_{12}^{st_2}(x^{-1}\eta^{-1}) V_1^{-1}, \end{aligned} \quad (7)$$

where the symbol  $st_k$  stands for the supertransposition operation in the space with index  $k$ . In its turn  $\rho(x)$  is an appropriate normalization function given by  $\rho(x) = q(x - 1)(x - \zeta)$  and the crossing

parameter is  $\eta = \zeta^{-1}$ . The corresponding crossing matrix  $V$  is an anti-diagonal matrix with the following non-null entries  $V_{ii'}$

$$V_{ii'} = \begin{cases} (-1)^{\frac{1-p_i}{2}}, & i = 1, \\ (-1)^{\frac{1-p_i}{2}} q^{F_1^{(i)}}, & 1 < i < \frac{N+1}{2}, \\ (-1)^{\frac{1+p_i}{2}} q^{F_2^{(i)}}, & \frac{N+1}{2} < i \leq N. \end{cases} \quad (8)$$

where

$$F_1^{(i)} = i - 1 - p_1 - p_i - 2 \sum_{j=1}^{i-1} p_j, \quad F_2^{(i)} = i - p_1 - p_i - 2 \sum_{j=2 \neq \frac{N}{2} + 1}^{i-1} p_j. \quad (9)$$

The construction of integrable models with open boundaries was largely impulsed by Sklyanin's pioneer work [6]. In Sklyanin's approach the construction of such models are based on solutions of the so-called reflection equations [7] for a given integrable bulk system. The reflection equations determine the boundary conditions compatible with the bulk integrability and it reads

$$R_{21}(x/y)K_2^-(x)R_{12}(xy)K_1^-(y) = K_1^-(y)R_{21}(xy)K_2^-(x)R_{12}(x/y), \quad (10)$$

where the tensor products appearing in (10) should be understood in the graded sense. The matrix  $K^-(x)$  describes the reflection at one of the ends of an open chain while a similar equation should also hold for a matrix  $K^+(x)$  describing the reflection at the opposite boundary. As discussed above, the  $U_q[spo(2n|2m)]$   $R$ -matrix satisfies important symmetry relations such as the PT-symmetry and crossing symmetry. When these properties are fulfilled one can follow the scheme devised in [8, 9] and the matrix  $K^-(x)$  is obtained by solving the Eq. (10) while the matrix  $K^+(x)$  can be obtained from the isomorphism  $K^-(x) \mapsto K^+(x)^{st} = K^-(x^{-1}\eta^{-1})V^{st}V$ .

The purpose of this work is to investigate the general families of regular solutions of the graded reflection equation (10). Regular solutions mean that the  $K$ -matrices have the general form

$$K^-(x) = \sum_{i,j=1}^N k_{i,j}(x) \hat{e}_{ij}, \quad (11)$$

such that the condition  $k_{i,j}(1) = \delta_{ij}$  holds for all matrix elements.

The direct substitution of (11) and the  $U_q[spo(2n|2m)]$   $R$ -matrix (1)-(9) in the graded reflection equation (10), leave us with a system of  $N^4$  functional equations for the entries  $k_{i,j}(x)$ . In order to solve these equations we shall make use of the derivative method. Thus, by differentiating the equation (10) with respect to  $y$  and setting  $y = 1$ , we obtain a set of algebraic equations for the matrix elements  $k_{i,j}(x)$ . Although we obtain a large number of equations only a few of them are actually independent and a direct inspection of those equations, in the lines described in [10], allows us to find the branches of regular solutions. In what follows we shall present our findings for the regular solutions of the reflection equation associated with the  $U_q[spo(2n|2m)]$  vertex model.

### 3 Diagonal K-matrix solutions

The diagonal solutions of the graded reflection equation (10) is characterized by a  $K$ -matrix of the form

$$K^-(x) = \sum_{i=1}^N k_{i,i}(x) \hat{e}_{ii}. \quad (12)$$

with the entries  $k_{i,j}(x)$  related with  $k_{1,1}(x)$  in a general form given by

$$k_{i,i}(x) = \frac{(\beta_{i,i} - \beta_{1,1})(x-1) + 2}{(\beta_{i,i} - \beta_{1,1})(x^{-1}-1) + 2} k_{1,1}(x) \quad (13)$$

for  $i = 2, \dots, N-1$  and

$$k_{N,N}(x) = \frac{(\beta_{N,N} - \beta_{1,1})(x-1) + 2}{(\beta_{N,N} - \beta_{1,1})(x^{-1}-1) + 2} k_{N-1,N-1}(x) \quad (14)$$

The parameters  $\beta_{i,i} = \frac{d}{dx}[k_{i,i}(x)]_{x=1}$  are fixed in order to give us four families of diagonal  $K$ -matrices such that half of them with one free parameter  $\beta$ .

$$\begin{aligned} k_{1,1}(x) &= \frac{\beta(x^{-1}-1) + 2}{\beta(x-1) + 2}, \\ k_{22}(x) &= \dots = k_{N-1,N-1}(x) = 1, \\ k_{N,N}(x) &= x \frac{\beta(1-xq^2\zeta) - 2}{\beta(x-q^2\zeta) - 2x}. \end{aligned} \quad (15)$$

and

$$\begin{aligned} k_{1,1}(x) &= \dots = k_{n+m,n+m}(x) = 1, \\ k_{n+m+1,m+m+1}(x) &= \dots = k_{N,N}(x) = \frac{\beta(x-1) + 2}{\beta(x^{-1}-1) + 2}. \end{aligned} \quad (16)$$

while the other half without free parameters:

$$\begin{aligned} k_{1,1}(x) &= \dots = k_{p-1,p-1}(x) = 1, \\ k_{p,p}(x) &= \dots = k_{N+1-p,N+1-p}(x) = \frac{x + \epsilon q^{2p-3} \sqrt{\zeta}}{x^{-1} + \epsilon q^{2p-3} \sqrt{\zeta}}, \\ k_{N+2-p,N+2-p}(x) &= \dots = k_{N,N}(x) = x^2. \end{aligned} \quad (17)$$

where the discrete label  $p$  assumes values in the interval  $2 \leq p \leq m+1$  and

$$\begin{aligned} k_{1,1}(x) &= \dots = k_{p-1,p-1}(x) = 1, \\ k_{p,p}(x) &= \dots = k_{N+1-p,N+1-p}(x) = \frac{x + \epsilon q^{4m+1-2p} \sqrt{\zeta}}{x^{-1} + \epsilon q^{4m+1-2p} \sqrt{\zeta}}, \\ k_{N+2-p,N+2-p}(x) &= \dots = k_{N,N}(x) = x^2. \end{aligned} \quad (18)$$

for the discrete label  $p$  with values in the interval  $m+2 \leq p \leq n+m$ .

Here and in what follows,  $\epsilon$  is a discrete parameter assuming the values  $\pm 1$ . We also notice that the case  $p = 2$  is limit of the solution of the solution (15) and we have  $(n+m)$  diagonal solutions.

## 4 Non-Diagonal K-matrix Solutions

Analyzing the reflection matrix equation (10) we can see that the non-diagonal matrix elements  $k_{i,j}(x)$  have the form

$$k_{i,j}(x) = \begin{cases} \beta_{i,j}xG(x), & i > j' \\ \beta_{i,j}xG(x), & i > j' \\ \beta_{i,j}G(x)H_f(x), & i = j' \text{ (fermionic)} \\ \beta_{i,j}G(x)H_b(x), & i = j' \text{ (bosonic)} \end{cases} \quad (19)$$

Here and in what follows  $G(x)$  is an arbitrary function satisfying the regular condition  $k_{i,j}(1) = \delta_{ij}$  and

$$\beta_{i,j} = \frac{d}{dx}[k_{i,j}(x)]_{x=1}, \quad H_f(x) = \frac{x - \epsilon q \sqrt{\zeta}}{1 - \epsilon q \sqrt{\zeta}}, \quad H_b(x) = \frac{qx + \epsilon \sqrt{\zeta}}{q + \epsilon \sqrt{\zeta}}. \quad (20)$$

Moreover, the diagonal entries  $k_{i,i}(x)$  satisfy defined recurrence relations which depend on the bosonic and fermionic degree of freedom. However, for the  $U_q[s\mathfrak{po}(2n|2m)]$  model we have found  $K$ -matrix solutions with both degree of freedom only for the cases  $(n \neq 1, m = 1)$  and  $(n = 1, m \neq 1)$ .

### 4.1 K-matrices with fermionic degree of freedom

Here we shall focus on the non-diagonal solutions of the graded reflection equation (10) with  $k_{i,j}(x) = 0$  for the bosonic degree of freedom *i.e.*, for  $i \neq j = m + 1, \dots, m + 2n$ . We have found three classes of non-diagonal solutions that we refer in what follows as solutions of type  $\mathcal{P}_1$  to type  $\mathcal{P}_3$ :

#### 4.1.1 Solution $\mathcal{P}_1$

The solution of type  $\mathcal{P}_1$  is valid only for the  $U_q[s\mathfrak{op}(2n|2)]$  models with  $n \geq 1$  and the  $K$ -matrix has the following block structure

$$K^- = \begin{pmatrix} k_{1,1} & \mathbb{O}_{1 \times 2n} & k_{1,N} \\ \mathbb{O}_{2n \times 1} & \mathbb{K}_1 & \mathbb{O}_{2n \times 1} \\ k_{N,1} & \mathbb{O}_{1 \times 2n} & k_{N,N} \end{pmatrix}, \quad (21)$$

where  $\mathbb{O}_{a \times b}$  is a  $a \times b$  null matrix and

$$\mathbb{K}_1(x) = \frac{x^2 - \zeta}{1 - \zeta} \mathbb{I}_{2n \times 2n}. \quad (22)$$

Here and in what follows  $\mathbb{I}_{2n \times 2n}$  denotes a  $2n \times 2n$  identity matrix and the remaining non-null entries are given by

$$\begin{aligned}
k_{1,1}(x) &= 1, \\
k_{1,N}(x) &= \frac{1}{2}\beta_{1,N}(x^2 - 1), \quad k_{N,1}(x) = \frac{2\zeta}{(1-\zeta)^2} \frac{(x^2 - 1)}{\beta_{1,N}}, \\
k_{N,N}(x) &= x^2.
\end{aligned} \tag{23}$$

where  $\beta_{1,N}$  is the free parameter. We remark here that this solution for  $n = 1$  consist of a particular case of the three parameter solution given in the appendix for the  $U_q[spo(2|2)]$  vertex model.

#### 4.1.2 Solution $\mathcal{P}_2$

The  $U_q[spo(2n|4)]$  vertex models admit the solution  $\mathcal{P}_2$  whose corresponding  $K$ -matrix has the following structure

$$K^- = \begin{pmatrix} k_{1,1} & k_{1,2} & & k_{1,N-1} & k_{1,N} \\ k_{2,1} & k_{2,2} & \mathbb{O}_{2 \times 2n} & k_{2,N-1} & k_{2,N} \\ & \mathbb{O}_{2n \times 2} & \mathbb{K}_2 & \mathbb{O}_{2n \times 2} & \\ k_{N-1,1} & k_{N-1,2} & \mathbb{O}_{2 \times 2n} & k_{N-1,N-1} & k_{N-1,N} \\ k_{N,1} & k_{N,2} & & k_{N,N-1} & k_{N,N} \end{pmatrix}, \tag{24}$$

where  $\mathbb{K}_2 = k_{3,3}(x)\mathbb{I}_{2n \times 2n}$ . The non-diagonal entries can be written as

$$\begin{aligned}
k_{1,2}(x) &= \beta_{1,2}G_1(x), & k_{2,1}(x) &= \beta_{2,1}G_1(x) \\
k_{1,N-1}(x) &= \beta_{1,N-1}G_1(x) & k_{N-1,1}(x) &= q^2\zeta \frac{\beta_{1,2}\beta_{2,1}}{\beta_{1,N-1}}G_1(x) \\
k_{2,N-1}(x) &= -\frac{\beta_{2,1}\beta_{1,N}}{\beta_{1,2}}G_1(x)H(x) & k_{N-1,2}(x) &= -q^2\zeta \frac{\beta_{1,2}\beta_{2,1}\beta_{1,N}}{\beta_{1,N-1}^2}G_1(x)H(x) \\
k_{2,N}(x) &= -\Gamma_n \frac{\beta_{1,N}}{\beta_{1,2}}xG_2(x) & k_{N,2}(x) &= -q^2\zeta\Gamma_n \frac{\beta_{2,1}\beta_{1,N}}{\beta_{1,N-1}^2}xG_2(x) \\
k_{N-1,N}(x) &= -q^2\zeta\Gamma_n \frac{\beta_{1,N}}{\beta_{1,N-1}}xG_2(x) & k_{N,N-1}(x) &= -q^2\zeta\Gamma_n \frac{\beta_{2,1}\beta_{1,N}}{\beta_{1,2}\beta_{1,N-1}}xG_2(x) \\
k_{N,1}(x) &= q^2\zeta \frac{\beta_{2,1}^2\beta_{1,N}}{\beta_{1,N-1}^2}G_1(x)H(x) & k_{1,N}(x) &= \beta_{1,N}G_1(x)H(x),
\end{aligned} \tag{25}$$

and the auxiliary functions are given by

$$\begin{aligned}
G_1(x) &= \left[ -q^2\zeta\Gamma_n \frac{\beta_{1,N}}{\beta_{1,2}\beta_{1,N-1}} + \frac{x - q^2\zeta}{x - 1} \right] \left( \frac{x - 1}{x^2 - q^2\zeta} \right) \frac{k_{1,N}(x)}{\beta_{1,N}}, \\
G_2(x) &= \left[ -\frac{1}{\Gamma_n} \frac{\beta_{1,2}\beta_{1,N-1}}{\beta_{1,N}} + \frac{x - q^2\zeta}{x - 1} \right] \left( \frac{x - 1}{x^2 - q^2\zeta} \right) \frac{k_{1,N}(x)}{\beta_{1,N}}, \\
H(x) &= \frac{\beta_{1,2}\beta_{1,N-1}(x^2 - q^2\zeta)}{\beta_{1,2}\beta_{1,N-1}(x - q^2\zeta) - q^2\zeta\Gamma_n\beta_{1,N}(x - 1)}
\end{aligned} \tag{26}$$

and

$$\Gamma_n = \frac{\beta_{2,1}\beta_{1,N}}{\beta_{1,N-1}} - \frac{2}{1 - q^2\zeta} \quad (27)$$

With respect to the diagonal matrix elements, we have the following expressions

$$\begin{aligned} k_{11}(x) = & \left\{ (x - q^2\zeta) \left[ (1 + q^2\zeta) \frac{\beta_{2,1}\beta_{1,N}}{\beta_{1,N-1}} - \frac{\beta_{1,2}\beta_{1,N-1}}{\beta_{1,N}} \right] + q^2\zeta\Gamma_n^2(1 - xq^2\zeta) \frac{\beta_{1,N}}{\beta_{1,2}\beta_{1,N-1}} \right. \\ & \left. + \frac{2(1 + q^2\zeta)^2x - 4q^2\zeta(x^2 + 1)}{(1 - q^2\zeta)(x - 1)} \right\} \frac{1}{(x^2 - q^2\zeta)(x + 1)} \frac{k_{1,N}(x)}{\beta_{1,N}} \end{aligned} \quad (28)$$

for the recurrence relation

$$\begin{aligned} k_{2,2}(x) &= k_{1,1}(x) + (\beta_{2,2} - \beta_{1,1})G_1(x), \\ k_{3,3}(x) &= k_{1,1}(x) + (\beta_{3,3} - \beta_{1,1})G(x) + \Delta_1(x), \\ k_{N-1,N-1}(x) &= k_{3,3}(x) + (\beta_{N-1,N-1} - \beta_{3,3})xG_2(x) + \Delta_2(x), \\ k_{N,N}(x) &= k_{N-1,N-1}(x) + (\beta_{N,N} - \beta_{N-1,N-1})xG_2(x). \end{aligned} \quad (29)$$

where

$$\begin{aligned} \Delta_1(x) &= \frac{\beta_{2,1}}{\beta_{1,N-1}} \left( x + q^2\zeta\Gamma_n \frac{\beta_{1,N}}{\beta_{1,2}\beta_{1,N-1}} \right) \left( \frac{x - 1}{x^2 - q^2\zeta} \right) k_{1,N}(x) \\ \Delta_2(x) &= -\frac{q^2\zeta}{\Gamma_n} \beta_{1,2} \frac{\beta_{1,N-1}}{\beta_{1,N}} \Delta_1(x). \end{aligned} \quad (30)$$

The diagonal entries (29) depend on the variables  $\beta_{\alpha,\alpha}$  which are related to the four free parameters  $\beta_{1,2}, \beta_{2,1}, \beta_{1,N-1}$  and  $\beta_{1,N}$  through the expressions

$$\begin{aligned} \beta_{2,2} &= \beta_{1,1} + \frac{\beta_{1,2}\beta_{1,N-1}}{\beta_{1,N}} - \Gamma_n, \\ \beta_{3,3} &= \beta_{1,1} + \frac{\beta_{1,2}\beta_{1,N-1}}{\beta_{1,N}} + \frac{2}{1 - q^2\zeta}, \\ \beta_{N-1,N-1} &= \beta_{1,1} + 2 + \frac{\beta_{1,2}\beta_{1,N-1}}{\beta_{1,N}} - q^2\zeta\Gamma_n, \\ \beta_{N,N} &= \beta_{1,1} + 2 + \frac{\beta_{1,2}\beta_{1,N-1}}{\beta_{1,N}} - q^2\zeta\Gamma_n^2 \frac{\beta_{1,N}}{\beta_{1,2}\beta_{1,N-1}}. \end{aligned} \quad (31)$$

### 4.1.3 Solution $\mathcal{P}_3$

This class of solution is valid for all  $U_q[\mathfrak{spo}(2n|2m)]$  vertex models with  $m \geq 3$  and the corresponding  $K$ -matrix possess the following general form

$$K^- = \begin{pmatrix} k_{1,1} & \cdots & k_{1,m} & \cdots & k_{1,2n+m+1} & \cdots & k_{1,N} \\ \vdots & \ddots & \vdots & \mathbb{O}_{m \times 2n} & \vdots & \ddots & \vdots \\ k_{m,1} & \cdots & k_{m,m} & \cdots & k_{m,2n+m+1} & \cdots & k_{m,N} \\ & \mathbb{O}_{2n \times m} & & \mathbb{K}_3 & & \mathbb{O}_{2n \times m} & \\ k_{2n+m+1,1} & \cdots & k_{2n+m+1,m} & \cdots & k_{2n+m+1,2n+m+1} & \cdots & k_{2n+m+1,N} \\ \vdots & \ddots & \vdots & \mathbb{O}_{m \times 2n} & \vdots & \ddots & \vdots \\ k_{N,1} & \cdots & k_{N,m} & \cdots & k_{N,2n+m+1} & \cdots & k_{N,N} \end{pmatrix}, \quad (32)$$

where  $\mathbb{K}_3$  is a diagonal matrix given by

$$\mathbb{K}_3(x) = k_{m+1,m+1}(x) \mathbb{I}_{2n \times 2n}. \quad (33)$$

With respect to the elements of the last column, we have the following expression

$$\begin{aligned} k_{i,N}(x) &= -\frac{\epsilon}{\sqrt{\zeta}} q^{t_i-t_1} \beta_{1,i'} x G(x) \\ i &= 2, \dots, m \text{ and } i = 2n+m+1, \dots, N-1, \end{aligned} \quad (34)$$

In their turn the entries of the first column are mainly given by

$$\begin{aligned} k_{i,1}(x) &= q^{t_i-t_2} \frac{\beta_{2,1} \beta_{1,i'}}{\beta_{1,N-1}} G(x), \\ i &= 3, \dots, m \text{ and } i = 2n+m+1, \dots, N-1. \end{aligned} \quad (35)$$

In the last row we have

$$\begin{aligned} k_{N,j}(x) &= -\frac{\epsilon}{\sqrt{\zeta}} q^{t_N-t_2} \frac{\beta_{2,1} \beta_{1,j}}{\beta_{1,N-1}} x G(x) \\ j &= 2, \dots, m \text{ and } j = 2n+m+1, \dots, N-1, \end{aligned} \quad (36)$$

while the elements of the first row are  $k_{1,j}(x) = \beta_{1,j} G(x)$  for  $j = 2, \dots, m$  and  $j = 2n+m+1, \dots, N-1$ .

Concerning the elements of the secondary diagonal, they are given by

$$\begin{aligned} k_{i,i'}(x) &= q^{t_1-t_{i'}} \left( \frac{1-\epsilon q \sqrt{\zeta}}{q+1} \right)^2 \frac{\beta_{1,i'}^2}{\beta_{1,N}} G(x) H_f(x) \\ i &= 2, \dots, m, \quad i \neq i' \text{ and } i = 2n+m+1, \dots, N-1, \end{aligned} \quad (37)$$

while the remaining entries  $k_{1,N}(x)$  and  $k_{N,1}(x)$  are determined by the following expressions

$$\begin{aligned} k_{1,N}(x) &= \beta_{1,N}G(x)H_f(x) \\ k_{N,1}(x) &= q^{t_{N-1}-t_2} \frac{\beta_{1,N}\beta_{2,1}^2}{\beta_{1,N-1}^2} G(x)H_f(x) \end{aligned} \quad (38)$$

where  $H_f(x)$  is given by (20).

The remaining matrix elements  $k_{i,j}(x)$  with  $i \neq j$  are then

$$k_{i,j}(x) = \begin{cases} -\frac{\epsilon}{\sqrt{\zeta}} q^{t_i-t_1} \left( \frac{1-\epsilon q\sqrt{\zeta}}{q+1} \right) \frac{\beta_{1,i'}\beta_{1,j}}{\beta_{1,N}} G(x), & i < j' \quad 2 \leq i, j \leq N-1 \\ \frac{1}{\zeta} q^{t_i-t_1} \left( \frac{1-\epsilon q\sqrt{\zeta}}{q+1} \right) \frac{\beta_{1,i'}\beta_{1,j}}{\beta_{1,N}} xG(x), & i > j' \quad 2 \leq i, j \leq N-1 \end{cases} \quad (39)$$

and

$$\begin{aligned} k_{2,1}(x) &= \frac{2(-1)^m(1-\epsilon q\sqrt{\zeta})q^{m-2}}{(1+\epsilon q\sqrt{\zeta})(\epsilon q^{m-1}\sqrt{\zeta}+(-1)^m)(\epsilon q^m\sqrt{\zeta}+(-1)^m)} \frac{\beta_{1,N-1}}{\beta_{1,N}} G(x), \\ k_{1,m}(x) &= -\frac{2\zeta q^{m-1}(q+1)^2}{(1-\epsilon q\sqrt{\zeta})(1+\epsilon q\sqrt{\zeta})(\epsilon q^{m-1}\sqrt{\zeta}+(-1)^m)(\epsilon q^m\sqrt{\zeta}+(-1)^m)} \frac{\beta_{1,N}}{\beta_{1,m'}} G(x), \end{aligned} \quad (40)$$

and the parameters  $\beta_{1,j}$  are constrained by the relation

$$\beta_{1,j} = -\beta_{1,j+1} \frac{\beta_{1,N-j}}{\beta_{1,N+1-j}} \quad j = 2, \dots, m-1. \quad (41)$$

With regard to the diagonal matrix elements, they are given by

$$k_{i,i}(x) = \begin{cases} k_{1,1}(x) + (\beta_{i,i} - \beta_{1,1})G(x), & 2 \leq i \leq m \\ k_{1,1}(x) + (\beta_{m+1,m+1} - \beta_{1,1})G(x) + \Delta(x), & i = m+1 \\ k_{m+1,m+1}(x) + (\beta_{2n+m+1,2n+m+1} - \beta_{m+1,m+1})xG(x) - \epsilon q^{2n+1}\sqrt{\zeta}\Delta(x), & i = 2n+m+1 \\ k_{i-1,i-1}(x) + (\beta_{i,i} - \beta_{i-1,i-1})xG(x), & 2n+m+2 \leq i \leq N \end{cases} \quad (42)$$

The last term of the recurrence relation (42) is identified with

$$k_{N,N}(x) = x^2 k_{1,1}(x) \quad (43)$$

to find

$$k_{1,1}(x) = \left[ \frac{2x}{x^2-1} - \frac{\beta_{m+1,m+1} - \beta_{1,1}}{x+1} \right] G(x) + [1 - \epsilon q^{2n+1}\sqrt{\zeta}] \frac{\Delta(x)}{x^2-1} \quad (44)$$

where

$$\Delta(x) = \frac{2(x-1)G(x)}{(1+\epsilon q\sqrt{\zeta})(\epsilon q^{m-1}\sqrt{\zeta}+(-1)^m)(\epsilon q^m\sqrt{\zeta}+(-1)^m)}. \quad (45)$$

In their turn the diagonal parameters  $\beta_{i,i}$  are fixed by the relations

$$\beta_{i,i} = \begin{cases} \beta_{1,1} + \Lambda_m \sum_{k=0}^{i-2} \left(-\frac{1}{q}\right)^k, & 2 \leq i \leq m \\ \beta_{1,1} + 2 - \epsilon q\sqrt{\zeta}\Lambda_m \sum_{k=0}^{N-1-i} (-q)^k, & 2n+m+1 \leq i \leq N-1 \\ \beta_{1,1} + 2, & i = N \end{cases} \quad (46)$$

and

$$\beta_{m+1,m+1} = \beta_{1,1} + \Lambda_m \left( \frac{q}{q+1} + \frac{(-1)^m}{q^{m-2}(q+1)^2} \frac{1 + \epsilon\sqrt{\zeta}}{\epsilon\sqrt{\zeta}} \right), \quad (47)$$

with

$$\Lambda_m = \frac{2(-1)^m q^{m-2} (q+1)^2 \epsilon \sqrt{\zeta}}{(1 + \epsilon\sqrt{\zeta})(\epsilon q^{m-1} \sqrt{\zeta} + (-1)^m)(\epsilon q^m \sqrt{\zeta} + (-1)^m)}. \quad (48)$$

The class of solution  $\mathcal{P}_3$  has a total amount of  $m$  free parameters namely  $\beta_{1,2n+m+1}, \dots, \beta_{1,N}$ .

Here we note that we can take the limit  $m = 2$  in the class  $\mathcal{P}_3$  to get a two parameter solution which is a particular case of the three parameter solution of the class  $\mathcal{P}_2$ . This complete the set of solutions with the fermionic degree of freedom.

## 4.2 K-matrices with bosonic degree of freedom

Here we consider  $k_{i,j}(x) = 0$ , for  $i \neq j = \{1, \dots, m\}$  and  $i \neq j = \{2n + m + 1, \dots, N\}$  in order to get  $K$ -matrix solutions with only bosonic degree of freedom. We have found three classes of non-diagonal solutions that we refer in what follows as solutions of type  $\mathcal{P}_4$  to type  $\mathcal{P}_6$ :

### 4.2.1 Solution $\mathcal{P}_4$ :

This family of solutions is valid only for the  $U_q[spo(2|2m)]$  vertex model with  $m \geq 1$  and the corresponding  $K$ -matrix has the following block diagonal structure

$$K^- = \begin{pmatrix} k_{1,1} \mathbb{I}_{m \times m} & \mathbb{O}_{m \times 2} & \mathbb{O}_{m \times m} \\ \mathbb{O}_{2 \times m} & k_{m+1,m+1} & k_{m+1,m+2} & \mathbb{O}_{2 \times m} \\ & k_{m+2,m+1} & k_{m+2,m+2} & \\ \mathbb{O}_{m \times m} & \mathbb{O}_{m \times 2} & k_{N,N} \mathbb{I}_{m \times m} \end{pmatrix} \quad (49)$$

The non-null entries are given by

$$\begin{aligned} k_{1,1}(x) &= 1, \\ k_{m+1,m+1}(x) &= x \frac{\alpha[q^2 \zeta^{-1} + 1](x-1) + 2}{\alpha[xq^2 \zeta^{-1} - 1](x-1) + 2x}, \\ k_{m+1,m+2}(x) &= \beta \frac{x(x^2 - 1)}{\alpha[xq^2 \zeta^{-1} - 1](x-1) + 2x}, \\ k_{m+2,m+1}(x) &= -\frac{\alpha^2}{\beta} \frac{xq^2 \zeta^{-1} (x^2 - 1)}{\alpha[xq^2 \zeta^{-1} - 1](x-1) + 2x}, \\ k_{m+2,m+2}(x) &= -x^2 \frac{\alpha[q^2 \zeta^{-1} + 1](x-1) - 2x}{\alpha[xq^2 \zeta^{-1} - 1](x-1) + 2x}, \\ k_{N,N}(x) &= x^2. \end{aligned} \quad (50)$$

where  $\alpha = \beta_{m+1,m+1}$  and  $\beta = \beta_{m+1,m+2}$  are the two free parameters.

### 4.2.2 Solution $\mathcal{P}_5$ :

The solution  $\mathcal{P}_5$  is admitted for the  $U_q[\mathfrak{spo}(4|2m)]$  models with the following  $K$ -matrix

$$K^- = \begin{pmatrix} k_{1,1}\mathbb{I}_{m \times m} & & \mathbb{O}_{m \times 4} & & \mathbb{O}_{m \times m} \\ & k_{m+1,m+1} & k_{m+1,m+2} & k_{m+1,m+3} & k_{m+1,m+4} \\ \mathbb{O}_{4 \times m} & k_{m+2,m+1} & k_{m+2,m+2} & k_{m+2,m+3} & k_{m+2,m+4} \\ & k_{m+3,m+1} & k_{m+3,m+2} & k_{m+3,m+3} & k_{m+3,m+4} \\ & k_{m+4,m+1} & k_{m+4,m+2} & k_{m+4,m+3} & k_{m+4,m+4} \\ \mathbb{O}_{m \times m} & & \mathbb{O}_{m \times 4} & & k_{N,N}\mathbb{I}_{m \times m} \end{pmatrix}. \quad (51)$$

The non-diagonal elements are all grouped in the  $4 \times 4$  central block matrix. With respect to this central block, the entries of the secondary diagonal are given by

$$\begin{aligned} k_{m+1,m+4}(x) &= \beta_{m+1,m+4} G(x)H_b(x) \\ k_{m+2,m+3}(x) &= \epsilon q^{m-2} \frac{\beta_{m+1,m+3}}{\beta_{m+1,m+2}} \Gamma_m G(x)H_b(x), \\ k_{m+3,m+2}(x) &= -\epsilon q^m \frac{\beta_{m+1,m+2}}{\beta_{m+1,m+3}} \Gamma_m G(x)H_b(x), \\ k_{m+4,m+1}(x) &= -q^{2m-2} \frac{1}{\beta_{m+1,m+4}} \Gamma_m^2 G(x)H_b(x), \end{aligned} \quad (52)$$

where we recall  $H_b(x) = \frac{qx + \epsilon\sqrt{\zeta}}{q + \epsilon\sqrt{\zeta}}$  and  $G(x)$  arbitrary. The remaining non-diagonal elements can be written as

$$\begin{aligned} k_{m+1,m+2}(x) &= \beta_{m+1,m+2} G(x) & k_{m+2,m+1}(x) &= -\epsilon q^{m-2} \frac{\beta_{m+1,m+3}}{\beta_{m+1,m+4}} \Gamma_m G(x), \\ k_{m+1,m+3}(x) &= \beta_{m+1,m+3} G(x) & k_{m+3,m+1}(x) &= \epsilon q^m \frac{\beta_{m+1,m+2}}{\beta_{m+1,m+4}} \Gamma_m G(x) \\ k_{m+2,m+4}(x) &= \epsilon q^{m-2} \beta_{m+1,m+3} xG(x) & k_{m+4,m+2}(x) &= q^{2m-2} \frac{\beta_{m+1,m+2}}{\beta_{m+1,m+4}} \Gamma_m xG(x) \\ k_{m+3,m+4}(x) &= -\epsilon q^m \beta_{m+1,m+2} xG(x) & k_{m+4,m+3}(x) &= q^{2m-2} \frac{\beta_{m+1,m+3}}{\beta_{m+1,m+4}} \Gamma_m xG(x), \end{aligned} \quad (53)$$

where

$$\Gamma_m = \frac{\beta_{m+1,m+2}\beta_{m+1,m+3}}{\beta_{m+1,m+4}} - \frac{2}{(q^{m-2} + \epsilon)(q^m + \epsilon)}. \quad (54)$$

In their turn the diagonal entries are given by the following expressions

$$k_{1,1}(x) = \left\{ \frac{2(xq^{m-2} + \epsilon)^2}{(q^{m-2} + \epsilon)(q^m + \epsilon)(xq^{2m-4} - 1)(x - 1)} - \frac{\beta_{m+1,m+2}\beta_{m+1,m+3}}{\beta_{m+1,m+4}} \right\} \frac{(xq^m + \epsilon)(xq^{2m-4} - 1)}{(q^{m-2} + \epsilon)x(x + 1)} G(x) \quad (55)$$

with the recurrence relation

$$\begin{aligned}
k_{m+1,m+1}(x) &= k_{1,1}(x) + (\beta_{m+1,m+1} - \beta_{1,1})G(x) + \Delta_1(x), \\
k_{m+2,m+2}(x) &= k_{m+1,m+1}(x) + (\beta_{m+2,m+2} - \beta_{m+1,m+1})G(x), \\
k_{m+3,m+3}(x) &= k_{m+2,m+2}(x) + \Delta_2(x), \\
k_{m+4,m+4}(x) &= k_{m+3,m+3}(x) + (\beta_{m+4,m+4} - \beta_{m+3,m+3})xG(x), \\
k_{N,N}(x) &= x^2k_{1,1}(x),
\end{aligned} \tag{56}$$

where

$$\begin{aligned}
\Delta_1(x) &= \frac{\epsilon\Gamma_m}{q^{m-2} + \epsilon} \frac{(x-1)G(x)}{x} \\
\Delta_2(x) &= -\epsilon q^{m-2}(q^2 + 1)\Gamma_m \left( \frac{q^{m-2}x + \epsilon}{q^{m-2} + \epsilon} \right) G(x).
\end{aligned} \tag{57}$$

The variables  $\beta_{i,j}$  are given in terms of the free parameters  $\beta_{m+1,m+2}$ ,  $\beta_{m+1,m+3}$  and  $\beta_{m+1,m+4}$  through the relations here

$$\begin{aligned}
\beta_{m+1,m+1} &= \beta_{1,1} - \Gamma_m \\
\beta_{m+2,m+2} &= \beta_{1,1} + \frac{2}{(q^{m-2} + \epsilon)(q^m + \epsilon)} + \epsilon q^{m-2} \frac{\beta_{m+1,m+2}\beta_{m+1,m+3}}{\beta_{m+1,m+4}} \\
\beta_{m+3,m+3} &= \beta_{1,1} + 2 - \frac{2q^{2m-2}}{(q^{m-2} + \epsilon)(q^m + \epsilon)} - \epsilon q^m \frac{\beta_{m+1,m+2}\beta_{m+1,m+3}}{\beta_{m+1,m+4}}
\end{aligned} \tag{58}$$

$$\beta_{m+4,m+4} = \beta_{1,1} + 2 - \frac{2q^{2m-2}}{(q^{m-2} + \epsilon)(q^m + \epsilon)} + q^{2m-2} \frac{\beta_{m+1,m+2}\beta_{m+1,m+3}}{\beta_{m+1,m+4}}. \tag{59}$$

We remark here that the form of this class of solution differs from general form (19) by the number of free parameters.

### 4.2.3 Solution $\mathcal{P}_6$ :

The vertex model  $U_q[spo(2n|2m)]$  admits the solution  $\mathcal{P}_6$  for  $n \geq 3$ , whose  $K$ -matrix has the following block structure

$$K^- = \begin{pmatrix} k_{1,1}\mathbb{I}_{m \times m} & & \mathbb{O}_{m \times 2n} & & \mathbb{O}_{m \times m} \\ & k_{m+1,m+1} & \cdots & k_{m+1,2n+m} & \\ \mathbb{O}_{2n \times m} & \vdots & \ddots & \vdots & \mathbb{O}_{2n \times m} \\ & k_{2n+m,m+1} & \cdots & k_{2n+m,2n+m} & \\ \mathbb{O}_{m \times m} & & \mathbb{O}_{m \times 2n} & & k_{N,N}\mathbb{I}_{m \times m} \end{pmatrix} \tag{60}$$

The central block matrix cluster all non-diagonal elements different from zero. Concerning

that central block, we have the following expressions determining entries of the borders,

$$\begin{aligned}
k_{i,2n+m}(x) &= \frac{\epsilon}{\sqrt{\zeta}} \frac{\theta_i q^{t_i}}{\theta_{m+1} q^{t_{m+1}}} \beta_{m+1,i'} x G(x), & i = m+2, \dots, 2n+m-1 \\
k_{2n+m,j}(x) &= \frac{\epsilon}{\sqrt{\zeta}} \frac{\theta_{2n+m} q^{t_{2n+m}}}{\theta_{m+2} q^{t_{m+2}}} \frac{\beta_{m+2,m+1} \beta_{m+1,j}}{\beta_{m+1,2n+m-1}} x G(x), & j = m+2, \dots, 2n+m-1 \\
k_{i,m+1}(x) &= \frac{\theta_i q^{t_i}}{\theta_{m+2} q^{t_{m+2}}} \frac{\beta_{m+2,m+1} \beta_{m+1,i'}}{\beta_{m+1,2n+m-1}} G(x), & i = m+3, \dots, 2n+m-1 \\
k_{m+1,j}(x) &= \beta_{m+1,j} G(x). & j = m+2, \dots, 2n+m-1
\end{aligned} \tag{61}$$

The entries of the secondary diagonal are given by

$$k_{i,i'}(x) = \begin{cases} \beta_{m+1,2n+m} G(x) H_b(x), & i = m+1 \\ -q^{2(m-1)} \frac{\theta_{m+1} q^{t_{m+1}}}{\theta_{i'} q^{t_{i'}}} \left( \frac{q + \epsilon \sqrt{\zeta}}{q+1} \right)^2 \frac{\beta_{m+1,i'}}{\beta_{m+1,2n+m}} G(x) H_b(x), & i = m+2, \dots, 2n+m-1 \\ \frac{\theta_{2n+m-1} q^{t_{2n+m-1}}}{\theta_{m+2} q^{t_{m+2}}} \frac{\beta_{m+2,m+1} \beta_{m+1,2n+m}}{\beta_{m+1,2n+m-1}^2} G(x) H_b(x), & i = 2n+m \end{cases} \tag{62}$$

and the remaining non-diagonal elements are determined by the expression

$$k_{i,j}(x) = \begin{cases} \frac{\epsilon}{\sqrt{\zeta}} \frac{\theta_i q^{t_i}}{\theta_{m+1} q^{t_{m+1}}} \left( \frac{q + \epsilon \sqrt{\zeta}}{q+1} \right) \frac{\beta_{m+1,i'} \beta_{m+1,j}}{\beta_{m+1,2n+m}} G(x), & i < j', \quad m+1 < i, j < 2n+m \\ \frac{1}{\zeta} \frac{\theta_i q^{t_i}}{\theta_{m+1} q^{t_{m+1}}} \left( \frac{q + \epsilon \sqrt{\zeta}}{q+1} \right) \frac{\beta_{m+1,i'} \beta_{m+1,j}}{\beta_{m+1,2n+m}} x G(x), & i > j', \quad m+1 < i, j < 2n+m \end{cases} \tag{63}$$

and

$$\begin{aligned}
k_{m+1,m+n}(x) &= -\frac{2\epsilon(-1)^n \sqrt{\zeta} (1+q)^2}{(1-\epsilon\sqrt{\zeta})(q+\epsilon\sqrt{\zeta})} \frac{1}{(q^m + (-1)^n \epsilon)(q^{m-1} - (-1)^n \epsilon)} \frac{\beta_{m+1,2n+m}}{\beta_{m+1,n+m+1}} G(x), \\
k_{m+2,m+1}(x) &= \frac{2\epsilon q (q + \epsilon \sqrt{\zeta})}{\sqrt{\zeta} (1 - \epsilon \sqrt{\zeta})} \frac{1}{(q^m + (-1)^n \epsilon)(q^{m-1} - (-1)^n \epsilon)} \frac{\beta_{m+1,2n+m-1}}{\beta_{m+1,2n+m}} G(x).
\end{aligned} \tag{64}$$

In their turn the diagonal entries  $k_{i,i}(x)$  are given by

$$k_{1,1}(x) = \left( \frac{x - \epsilon \sqrt{\zeta}}{1 - \epsilon \sqrt{\zeta}} \right) \left( \frac{x q^m + (-1)^n \epsilon}{q^m + (-1)^n \epsilon} \right) \left( \frac{x q^{m-1} - (-1)^n \epsilon}{q^{m-1} - (-1)^n \epsilon} \right) \frac{2G(x)}{x(x^2 - 1)} \tag{65}$$

$$k_{i,i}(x) = \begin{cases} k_{1,1}(x) + \Gamma_{n,m}(x), & i = m+1 \\ k_{m+1,m+1}(x) + (\beta_{i,i} - \beta_{m+1,m+1}) G(x), & i = m+2, \dots, m+n \\ k_{n+m,n+m}(x) + (\beta_{n+m+1,n+m+1} - \beta_{n+m,n+m}) x G(x) + \Delta_{n,m}(x), & i = n+m+1 \\ k_{n+m+1,n+m+1}(x) + (\beta_{i,i} - \beta_{n+m+1,n+m+1}) x G(x), & i = n+m+2, \dots, 2n+m \\ x^2 k_{1,1}(x), & i = N \end{cases} \tag{66}$$

where

$$\Gamma_{n,m}(x) = \frac{2(qx + \epsilon \sqrt{\zeta})}{(1 - \epsilon \sqrt{\zeta})(q^m + (-1)^n \epsilon)(q^{m-1} - (-1)^n \epsilon)} \frac{G(x)}{x}. \tag{67}$$

and

$$\Delta_{n,m}(x) = -\frac{2(-1)^n q^{n-1}(q^2+1)}{(1-\epsilon\sqrt{\zeta})(q^m+(-1)^n\epsilon)(q^{m-1}-(-1)^n\epsilon)}(x-1)G(x)$$

The diagonal parameters are then fixed by the relations

$$\beta_{i,i} = \begin{cases} \beta_{1,1} - \frac{q+\epsilon\sqrt{\zeta}}{(1+q)^2}Q_{n,m}, & i = m+1 \\ \beta_{m+1,m+1} + Q_{n,m} \sum_{k=0}^{i-m-2} (-q)^k, & i = m+2, \dots, n+m \\ \beta_{n+m,n+m} - \epsilon(-1)^n(1+q^2)q^{m-2} \frac{q+\epsilon\sqrt{\zeta}}{(1+q)^2}Q_{n,m}, & i = n+m+1 \\ \beta_{n+m+1,n+m+1} + \epsilon(-1)^n q^m Q_{n,m} \sum_{k=0}^{i-n-m-2} (-q)^k, & i = m+2, \dots, n+m \end{cases} \quad (68)$$

and the auxiliary parameter  $Q_{n,m}$  is given by

$$Q_{n,m} = -\frac{2(1+q)^2}{(1-\epsilon\sqrt{\zeta})(q^m+(-1)^n\epsilon)(q^{m-1}-(-1)^n\epsilon)}. \quad (69)$$

Besides the above relations the following constraints should also holds

$$\beta_{m+1,m+j} = -\beta_{m+1,j+m+1} \frac{\beta_{m+1,2n+m-j}}{\beta_{m+1,2n+m+1-j}} \quad j = 2, \dots, n-1, \quad (70)$$

and  $\beta_{m+1,m+n+1}, \dots, \beta_{m+1,2n+m}$  are regarded as the  $n$  free parameters. The case  $n = 2$  is a particular solution of the three parameter class  $\mathcal{P}_6$ .

### 4.3 Complete K-matrices

The complete  $K$ -matrices are solutions with all entries different from zero. This kind of solution will be present only in two class: the models with two bosonic degree of freedom,  $U_q[spo(2|2m)]$  and those models with only two fermionic degree of freedom  $U_q[spo(2n|2)]$ .

#### 4.3.1 Solution $\mathcal{P}_7$ :

The series of solutions  $\mathcal{P}_7$  is valid for the  $U_q[spo(2|2m)]$  model and the corresponding  $K$ -matrix also possess all entries different from zero. In the first and last columns, the matrix elements are mainly given by

$$\begin{aligned} k_{i,1}(x) &= \frac{\theta_i q^{t_i}}{\theta_2 q^{t_2}} \frac{\beta_{2,1} \beta_{1,i'}}{\beta_{1,N-1}} G(x) & i = 3, \dots, N-1 \\ k_{i,N}(x) &= -\frac{\epsilon_m}{\sqrt{\zeta}} \frac{\theta_i q^{t_i}}{\theta_1 q^{t_1}} \beta_{1,i'} x G(x) & i = 2, \dots, N-1 \end{aligned} \quad (71)$$

while the ones in the first and last rows are respectively

$$\begin{aligned} k_{1,j}(x) &= \beta_{1,j} G(x) & j = 2, \dots, N-1 \\ k_{N,j}(x) &= -\frac{\epsilon_m}{\sqrt{\zeta}} \frac{\theta_N q^{t_N}}{\theta_2 q^{t_2}} \frac{\beta_{2,1} \beta_{1,j}}{\beta_{1,N-1}} x G(x) & j = 2, \dots, N-1 \end{aligned} \quad (72)$$

In the secondary diagonal we have the following expression determining the matrix elements

$$k_{i,i'}(x) = \begin{cases} \beta_{1,N}G(x)H_f(x), & i = 1 \\ \frac{\theta_1 q^{t_1}}{\theta_{i'} q^{t_{i'}}} \left( \frac{1 - \epsilon_m \epsilon q \sqrt{\zeta}}{q+1} \right)^2 \frac{\beta_{1,i'}^2}{\beta_{1,N}} G(x)H_f(x), & i \neq \{1, m+1, m+2, N\} \\ \frac{\theta_1 q^{t_1}}{\theta_{i'} q^{t_{i'}}} \left( \frac{q \epsilon_m \epsilon \sqrt{\zeta}}{q-1} \right) \left( \frac{1 - \epsilon_m q \sqrt{\zeta}}{q+1} \right) \frac{\beta_{1,i'}^2}{\beta_{1,N}} G(x)H_b(x), & i = \{m+1, m+2\} \\ \frac{\theta_{N-1} q^{t_{N-1}}}{\theta_2 q^{t_2}} \frac{\beta_{1,N} \beta_{2,1}^2}{\beta_{1,N-1}^2} G(x)H_f(x), & i = N. \end{cases} \quad (73)$$

recalling that

$$H_b(x) = \frac{qx + \epsilon_m \sqrt{\zeta}}{q + \epsilon_m \sqrt{\zeta}} \quad \text{and} \quad H_f(x) = \frac{x - \epsilon_m q \sqrt{\zeta}}{1 - \epsilon_m q \sqrt{\zeta}}. \quad (74)$$

In their turn the other non-diagonal entries satisfy the relation

$$k_{i,j}(x) = \begin{cases} -\frac{\epsilon_m}{\sqrt{\zeta}} \frac{\theta_i q^{t_i}}{\theta_1 q^{t_1}} \left( \frac{1 - \epsilon_m q \sqrt{\zeta}}{q+1} \right) \frac{\beta_{1,i'} \beta_{1,j}}{\beta_{1,N}} G(x), & i < j', \quad 2 < i, j < N-1 \\ \frac{1}{\zeta} \frac{\theta_i q^{t_i}}{\theta_1 q^{t_1}} \left( \frac{1 - \epsilon_m q \sqrt{\zeta}}{q+1} \right) \frac{\beta_{1,i'} \beta_{1,j}}{\beta_{1,N}} x G(x), & i > j', \quad 2 < i, j < N-1 \end{cases}, \quad (75)$$

and

$$\begin{aligned} k_{1,m}(x) &= \frac{i(q+1)}{q-1} \frac{\beta_{1,m+1} \beta_{1,m+2}}{\beta_{1,m+3}} G(x), \\ k_{1,m+1}(x) &= (-1)^{\delta_{m,2}} \frac{i \sqrt{\zeta} (q + \epsilon_m)}{(1 + \epsilon_m \sqrt{\zeta})(1 + q \sqrt{\zeta})} \frac{\beta_{1,N}}{\beta_{1,m+2}} G(x), \\ k_{2,1}(x) &= (-1)^{\delta_{m,2}} \frac{\epsilon_m}{q \sqrt{\zeta}} \frac{(1 - \epsilon_m q \sqrt{\zeta})^2}{(q - \epsilon_m)(1 + \epsilon_m \sqrt{\zeta})(1 + q \sqrt{\zeta})} \frac{\beta_{1,N-1}}{\beta_{1,N}} G(x), \end{aligned} \quad (76)$$

and the parameters  $\beta_{1,j}$  are required to satisfy the recurrence relation

$$\beta_{1,j} = -\frac{\beta_{1,j+1} \beta_{1,N-j}}{\beta_{1,N-j+1}} \quad j = 2, \dots, m-1. \quad (77)$$

Considering now the diagonal entries, they are given by

$$k_{i,i}(x) = \begin{cases} k_{1,1} + (\beta_{i,i} - \beta_{1,1})G(x), & i = 2, \dots, m+1 \\ k_{1,1}(x) + (\beta_{m+2,m+2} - \beta_{1,1})xG(x) + \Delta(x), & i = m+2 \\ k_{m+2,m+2}(x) + (\beta_{i,i} - \beta_{m+2,m+2})xG(x), & i = m+3, \dots, N-1 \\ x^2 k_{1,1}(x). & i = N \end{cases}, \quad (78)$$

where

$$k_{1,1}(x) = \frac{G(x)}{x-1} \quad \text{and} \quad \Delta(x) = (1-x)G(x).$$

The diagonal parameters are determined by the expressions

$$\beta_{i,i} = \begin{cases} \beta_{1,1} + \frac{(q+1)}{q(1+\epsilon_m\sqrt{\zeta})} \sum_{k=0}^{i-2} \left(-\frac{1}{q}\right)^k, & i = 2, \dots, m \\ \beta_{m,m} - \frac{2\epsilon_m\sqrt{\zeta}}{q(q+1)(1+\epsilon_m\sqrt{\zeta})}, & i = m+1 \\ \beta_{m+1,m+1} + \frac{(q^2+1)(q+\epsilon_m\sqrt{\zeta})}{q(q+1)(1+\epsilon_m\sqrt{\zeta})}, & i = m+2 \\ \beta_{m+2,m+2} - \frac{2q^2}{(q+1)(1+\epsilon_m\sqrt{\zeta})}, & i = m+3 \\ \beta_{m+3,m+3} + \left(\frac{q+1}{1+\epsilon_m\sqrt{\zeta}}\right)^{i-m-4} \sum_{k=0}^{i-m-4} \left(-\frac{1}{q}\right)^k. & i = m+4, \dots, N \end{cases} \quad (79)$$

This solution has altogether  $m+1$  free parameters, namely  $\beta_{1,m+2}, \dots, \beta_{1,N}$ . This complete solution depend on the parity of  $m$  through the relation  $\epsilon_m = (-1)^m$ .

Here we notice that the case  $m = 2$  is also special because  $\beta_{1,m} = \beta_{1,2}$  and we have an exchange of sign for  $k_{2,1}(x)$  and  $k_{1,3}(x)$  as indicate in equation (76).

#### 4.3.2 Solution $\mathcal{P}_8$ :

The class of solutions  $\mathcal{P}_8$  is valid for the vertex model  $U_q[spo(2n|2)]$  and the corresponding  $K$ -matrix contains only non-null entries. The border elements are mainly given by the following expressions

$$\begin{aligned} k_{i,N}(x) &= -\frac{\epsilon_n}{\sqrt{\zeta}} \frac{\theta_i q^{t_i}}{\theta_1 q^{t_1}} \beta_{1,i'} x G(x) & i = 2, \dots, N-1 \\ k_{i,1}(x) &= \frac{\theta_i q^{t_i}}{\theta_2 q^{t_2}} \frac{\beta_{2,1} \beta_{1,i'}}{\beta_{1,N-1}} G(x) & i = 3, \dots, N-1 \\ k_{N,j}(x) &= -\frac{\epsilon_n}{\sqrt{\zeta}} \frac{\theta_N q^{t_N}}{\theta_2 q^{t_2}} \frac{\beta_{2,1} \beta_{1,j}}{\beta_{1,N-1}} x G(x) & j = 2, \dots, N-1 \\ k_{1,j}(x) &= \beta_{1,j} G(x) & j = 2, \dots, N-1. \end{aligned} \quad (80)$$

The secondary diagonal is constituted by elements  $k_{i,i'}(x)$  given by

$$k_{i,i'}(x) = \begin{cases} \beta_{1,N} G(x) H_f(x), & i = 1 \\ -\frac{\theta_1 q^{t_1}}{\theta_{i'} q^{t_{i'}}} \left(\frac{1-\epsilon_n q \sqrt{\zeta}}{q-1}\right) \left(\frac{q+\epsilon_n \sqrt{\zeta}}{q+1}\right) \frac{\beta_{1,i'}^2}{\beta_{1,N}} G(x) H_b(x), & i \neq \{1, N\} \\ \frac{\theta_{N-1} q^{t_{N-1}}}{\theta_2 q^{t_2}} \frac{\beta_{1,N} \beta_{2,1}^2}{\beta_{1,N-1}^2} G(x) H_f(x), & i = N. \end{cases} \quad (81)$$

where the functions  $H_b(x)$  and  $H_f(x)$  were already given in (74). The remaining non-diagonal entries are determined by the expression

$$k_{i,j}(x) = \begin{cases} \frac{\epsilon_n}{\sqrt{\zeta}} \frac{\theta_i q^{t_i}}{\theta_1 q^{t_1}} \left(\frac{1-\epsilon_n q \sqrt{\zeta}}{q-1}\right) \frac{\beta_{1,i'} \beta_{1,j}}{\beta_{1,N}} G(x), & i < j', 2 < i, j < N-1 \\ \frac{1}{\zeta} \frac{\theta_i q^{t_i}}{\theta_1 q^{t_1}} \left(\frac{1-\epsilon_n q \sqrt{\zeta}}{q-1}\right) \frac{\beta_{1,i'} \beta_{1,j}}{\beta_{1,N}} x G(x), & i > j', 2 < i, j < N-1 \end{cases}, \quad (82)$$

and

$$k_{1,n+1}(x) = \frac{-i\sqrt{\zeta}(q+1)}{(q\sqrt{\zeta} - \epsilon_n)(\sqrt{\zeta} - \epsilon_n)} \frac{\beta_{1,N}}{\beta_{1,n+2}} G(x)$$

$$k_{2,1}(x) = \frac{-i\epsilon_n(q\sqrt{\zeta} - \epsilon_n)}{(\sqrt{\zeta} - \epsilon_n)(q-1)} \frac{\beta_{1,N-1}}{\beta_{1,N}} G(x).$$

and the parameters  $\beta_{1,j}$  are required to satisfy

$$\beta_{1,j} = -\frac{\beta_{1,j+1}\beta_{1,N-j}}{\beta_{1,N+1-j}} \quad j = 2, \dots, n. \quad (83)$$

With respect to the diagonal entries, they are given by

$$k_{i,i}(x) = \begin{cases} k_{1,1}(x) + (\beta_{i,i} - \beta_{1,1})G(x), & i = 2, \dots, n+1 \\ k_{n+1,n+1}(x) + (\beta_{n+2,n+2} - \beta_{n+1,n+1})xG(x) + \Delta(x), & i = n+2 \\ k_{n+2,n+2}(x) + (\beta_{i,i} - \beta_{n+2,n+2})xG(x), & i = n+3, \dots, N-1 \\ x^2k_{1,1}(x). & i = N \end{cases}, \quad (84)$$

where

$$k_{1,1}(x) = \frac{G(x)}{x-1} \quad \text{and} \quad \Delta(x) = \epsilon_n \left( \frac{\sqrt{\zeta} + q}{\sqrt{\zeta} + \epsilon_n q} \right) \left( \frac{q^2 + \epsilon_n}{\sqrt{\zeta} - \epsilon_n} \right) \left( \frac{x-1}{q-1} \right) G(x).$$

The parameters  $\beta_{i,i}$  are determined by the expressions

$$\beta_{i,i} = \begin{cases} \beta_{1,1} + \frac{(-1)^{n+i}[q^{i-1} + q^{i-2} - (-1)^i(q-1)]}{(q-1)(\sqrt{\zeta} - \epsilon_n)}, & i = 2, \dots, n+1 \\ \beta_{n+1,n+1} + \frac{(\sqrt{\zeta} + q)(q^2 + \epsilon_n)}{q(q-1)(\sqrt{\zeta} - \epsilon_n)}, & i = n+2 \\ \beta_{n+2,n+2} - \frac{\epsilon_n q(q+1)^2}{(q-1)(\sqrt{\zeta} - \epsilon_n)} \sum_{k=0}^{i-n-3} (-q)^k, & i = n+3, \dots, N-1 \\ \beta_{1,1} + 2. & i = N \end{cases} \quad (85)$$

The variables  $\beta_{1,n+2}, \dots, \beta_{1,N}$  give us a total amount of  $n+1$  free parameters. This complete solution depend on the parity of  $n$  through the relation  $\epsilon_n = (-1)^n$ .

## 5 Concluding Remarks

In this work we have presented the general set of regular solutions of the graded reflection equation for the  $U_q[\mathfrak{osp}(2n|2m)]$  vertex model. Our findings can be summarized into four classes of diagonal solutions and eight classes of non-diagonal ones. Although the  $R$  matrix of the  $U_q[\mathfrak{osp}(2m|2n)^{(1)}]$  vertex model is isomorphic to the  $R$  matrix of the  $U_q[\mathfrak{spo}(2n|2m)]$  vertex model, we have find additional different  $K$ -matrices from those obtained by the exchange of the degree of freedom in these models (see, for example, the diagonal solutions of the  $U_q[\mathfrak{spo}(2|2)]$  vertex model).

We expect the results presented here to motivate further developments on the subject of integrable open boundaries for vertex models based on  $q$ -deformed Lie superalgebras. In particular, the classification of the solutions of the graded reflection equation for others  $q$ -deformed Lie superalgebras, for instance, the  $U_q[osp(2n|2m)^{(2)}]$  Lie superalgebra, which we hope to report on a future work.

## 6 Acknowledgments

We thank to W. Galleas and M.J. Martins by the discussions. This work is partially supported by the Brazilian research councils CNPq and FAPESP.

## References

- [1] A. Lima-Santos and W.Galleas, *Reflection matrices for the  $U_q[sl(r|2m)^{(2)}]$  vertex model*, [ArXiv: nlin.SI/08063659].
- [2] A. Lima-Santos, *Reflection matrices for the  $U_q[osp(r|2m)^{(1)}]$  vertex model*, [ArXiv: nlin.SI/08090421].
- [3] V. G. Kac, *Lie superalgebras*, Adv. Math. **26** (1977) 8.
- [4] M. Scheunert, *The theory of Lie superalgebras*, Lecture Notes in Mathematics 716, Springer-Verlag, Berlin, Heidelberg, New York, 1979.
- [5] W. Galleas and M.J. Martins, *Nucl. Phys. B* 732, (2006) 444.
- [6] E.K. Sklyanin, *J. Phys. A: Math. Gen.* 21, (1988) 2375.
- [7] I.V. Cherednik, *Theor. Math. Phys.* 61, (1984) 977.
- [8] A.J. Bracken, X.Y. Ge, Y.Z. Zhang and H.Q. Zhou, *Nucl. Phys. B* 516, (1998) 588.
- [9] L. Mezincescu and R.I. Nepomechie, *J. Phys. A: Math. Gen.* 24, (1991) 17; L. Mezincescu and R.I. Nepomechie, *Int. J. Mod. Phys. A*6, (1991) 5231
- [10] R. Malara and A. Lima-Santos, *J. Stat. Mech.: Theor. Exp.*, (2006) P09013.s. B 424, (1994) 468.

## Appendix A: The $U_q[spo(2|2)]$ case

The set of  $K$ -matrices associated with the  $U_q[spo(2|2)]$  vertex model includes both diagonal and non-diagonal solutions. The four solutions intrinsically diagonal three contain only one free parameter  $\beta$  and they are given by

$$\begin{aligned}
K^-(x) &= \text{Diag}\left(\frac{2x - \beta(x-1)}{2x + \beta x(x-1)}, 1, 1, \frac{2x + \beta x(xq^4 - 1)}{2x + \beta(q^4 - x)}\right), \\
K^-(x) &= \text{Diag}\left(1, 1, \frac{2x + \beta x(x-1)}{2x - \beta(x-1)}, \frac{2x + \beta x(x-1)}{2x - \beta(x-1)}\right) \\
K^-(x) &= \text{Diag}\left(1, \frac{2x + \beta x(x-1)}{2x - \beta(x-1)}, x^2, \frac{2x + \beta x(x-1)}{2x - \beta(x-1)}x^2\right)
\end{aligned} \tag{B.1}$$

and one without free parameters

$$K^-(x) = \text{Diag}\left(1, x \frac{q^2 + \epsilon x}{xq^2 + \epsilon}, x \frac{q^2 + \epsilon x}{xq^2 + \epsilon}, x^2\right). \tag{B.2}$$

We have also found the following non-diagonal solutions

$$K^-(x) = \begin{pmatrix} 1 - \frac{(x^{-1}\beta^2 + \alpha)(x-1)}{2\beta} & 0 & 0 & \frac{1}{2}\beta_{23}(x^2 - 1) \\ 0 & 1 + \frac{(\beta^2 - \alpha)(x-1)}{2\beta} & 0 & 0 \\ 0 & 0 & x^2 - \frac{(\beta^2 - \alpha)x(x-1)}{2\beta} & 0 \\ \frac{\alpha}{2\beta_{23}}(x^2 - 1) & 0 & 0 & x^2 + \frac{(x\beta^2 + \alpha)x(x-1)}{2\beta} \end{pmatrix} \tag{B.3}$$

containing three free parameters  $\alpha, \beta$  and  $\beta_{23}$ , and another solution with one free parameter

$$K^-(x) = \begin{pmatrix} 1 & 0 & 0 & \frac{1}{2}\beta_{1,4}(x^2 - 1) \\ 0 & \frac{q^2 - x^2}{q^2 - 1} & 0 & 0 \\ 0 & 0 & \frac{q^2 - x^2}{q^2 - 1} & 0 \\ 0 & \frac{2}{\beta_{1,4}} \frac{q^2}{(q^2 - 1)^2} (x^2 - 1) & 0 & x^2 \end{pmatrix}. \tag{B.4}$$

We don't find any complete solution for this model.