

Topological stability of anyonic quantum spin chains and formation of new topological liquids

Charlotte Gils,¹ Eddy Ardonne,² Simon Trebst,³ Andreas W. W. Ludwig,⁴ Matthias Troyer,¹ and Zhenghan Wang³

¹*Theoretische Physik, Eidgenössische Technische Hochschule Zürich, 8093 Zürich, Switzerland*

²*Nordita, Roslagstullsbacken 23, SE-106 91 Stockholm, Sweden*

³*Microsoft Research, Station Q, University of California, Santa Barbara, CA 93106*

⁴*Physics Department, University of California, Santa Barbara, CA 93106*

(Dated: October 29, 2022)

Quantum mechanical systems, whose degrees of freedom are so-called $\text{su}(2)_k$ anyons, form a bridge between ordinary $\text{SU}(2)$ spin systems and systems of interacting non-Abelian anyons. Such a connection can be made for arbitrary spin- S systems, and we explicitly discuss spin- $1/2$ and spin- 1 systems. Anyonic spin- $1/2$ chains exhibit a topological protection mechanism that stabilizes their gapless ground states and which vanishes only in the limit ($k \rightarrow \infty$) where the system turns into the ordinary spin- $1/2$ Heisenberg chain. For anyonic spin- 1 chains we show that their phase diagrams closely mirror the one of the biquadratic $\text{SU}(2)$ spin- 1 chain. This includes generalizations of the Haldane phase, of the AKLT point, and the appearance of several stable critical phases described by (super)conformal field theories. Our results describe at the same time nucleation of different 2D topological quantum fluids within a ‘parent’ non-Abelian quantum Hall state, arising from a macroscopic occupation with interacting anyons. The edge states between the ‘nucleated’ and the ‘parent’ liquids are neutral, and correspond precisely to the gapless modes of the anyonic chains.

PACS numbers: 05.30.Pr, 73.43.?f, 03.65.Vf

$\text{SU}(2)$ spin degrees of freedom are ubiquitous in condensed matter physics describing the elementary quantum mechanical properties of many magnetic materials. For ordinary $\text{SU}(2)$ spins there is – mathematically spoken – an infinite number of representations, or in other words arbitrarily large spins. Here we consider a ‘quantum deformation’ of $\text{SU}(2)$ [1], where we limit the number of representations to $k+1$ ‘angular momenta’ that take the values $j = 0, \frac{1}{2}, 1, \dots, \frac{k}{2}$. The degrees of freedom in these so-called $\text{su}(2)_k$ theories are examples of non-Abelian anyons with the Ising ($k = 2$) and Fibonacci ($k = 3$) anyons being studied in a variety of contexts such as unconventional $p_x + ip_y$ superconductors [2], fractional quantum Hall states [3], and proposals for topological quantum computation [4]. We will therefore refer to these generalized angular momenta also as anyon types. The analog of combining two ordinary spins, and reducing the tensor product, corresponds to the ‘fusion’ of two anyons which obey the fusion rules

$$j_1 \times j_2 = |j_1 - j_2| + (|j_1 - j_2| + 1) + \dots + \min(j_1 + j_2, k - j_1 - j_2). \quad (1)$$

For example, fusing two anyons with generalized angular momenta $j_{1,2} = \frac{1}{2}$, these rules imply $\frac{1}{2} \times \frac{1}{2} = 0 + 1$ for $k \geq 2$, which in the limit $k \rightarrow \infty$ describes the coupling of two ordinary spin- $1/2$'s into a singlet or triplet. For finite ‘level’ k the occurrence of several possible fusion outcomes is the hallmark of what is known as non-Abelian particle statistics.

Like the Heisenberg Hamiltonian for ordinary spins we can model a pairwise interaction between non-Abelian anyons that energetically splits the possible fusion outcomes. A first step towards understanding the collective ground states formed by a set of interacting anyons has been taken by studying chains of interacting Fibonacci anyons [5, 6, 7]: Uniform chains with pairwise interactions that favor either the ‘singlet’ or ‘triplet’-channel are gapless and can be mapped exactly onto the tricritical Ising and critical 3-state Potts models, respectively [5]. In

conventional systems these gapless theories would be completely unstable to the formation of a gap, but a more subtle mechanism is at play in the Fibonacci chain where an additional topological symmetry stabilizes the gaplessness of the system against local perturbations [5]: This topological symmetry is sensitive to a topological flux in the center of a periodic chain. For Fibonacci anyons there are two distinct sectors, a flux-free ($j = 0$) sector and a sector with topological flux $j = 1$. While the ground state is flux-free, all uniform, relevant perturbations carry non-trivial flux and are thereby prohibited from driving the system into a gapped phase.

However, if we consider the more general anyonic systems described by $\text{su}(2)_k$ theories and take the ‘undeformed’ limit $k \rightarrow \infty$, we recover a system of ordinary $\text{SU}(2)$ spins for which there is no such notion of a topological symmetry. This observation naturally raises the question whether the observed topological protection is unique to the Fibonacci theory $\text{su}(2)_3$. In this Letter, we show that the existence of a topological symmetry is a common feature in all $\text{su}(2)_k$ anyonic theories and that it protects the gaplessness of generalized spin- $1/2$ chains in these theories for *all finite levels* k . In fact, it is the ordinary $\text{SU}(2)$ spin- $1/2$ chain that stands out in this series as it loses this special symmetry and protection mechanism.

We also discuss $\text{su}(2)_k$ generalizations of the spin- 1 chain and close connections between the phase diagram of these anyonic spin- 1 chains and the biquadratic $\text{SU}(2)$ spin- 1 chain.

Finally, we explain that the collective states of these anyonic chains correspond to the *edges* between different (non-) Abelian quantum Hall liquids, where a *droplet* of a new, distinct liquid is nucleated by populating a ‘parent’ quantum Hall state with a finite density of interacting anyons.

Anyonic spin-1/2 chains.– We first turn to $\text{su}(2)_k$ quantum deformations of ordinary quantum spin- $1/2$ chains which have been introduced in Ref. 5. These ‘golden chains’ consist of a linear arrangement of non-Abelian anyons with ‘angular mo-

mentum' $\nu = 1/2$ in the $\text{su}(2)_k$ theory. Pairwise interactions between adjacent anyons favor fusion into the 'singlet' $j = 0$ for 'antiferromagnetic' (AFM) exchange and into the 'triplet' $j = 1$ for 'ferromagnetic' (FM) exchange. These $\text{su}(2)_k$ spin-1/2 chains turn out to be gapless for all levels k [5]. The critical theory is a conformal field theory (CFT) closely related to the original $\text{su}(2)_k$ theory, namely a particular coset theory [8]. We first concentrate on AFM couplings for which this CFT description is given by the $(k-1)$ -th unitary minimal model $\mathcal{M}_k := \text{su}(2)_{k-1} \times \text{su}(2)_1 / \text{su}(2)_k$ with central charge $c_k = 1 - 6/[(k+1)(k+2)]$. For $k = 2$ this is the Ising model, for $k = 3$ the tricritical Ising model, and in the limit $k \rightarrow \infty$ becomes the $c = 1$ -theory describing the ordinary Heisenberg chain. The explicit Z_2 symmetry of these models is reflected in a Z_2 sublattice symmetry, e.g. there are two soft modes in the energy spectra at (lattice) momenta $K = 0$ and $K = \pi$. We discuss the assignment of momenta to (primary) fields in more detail below, but already note that there are exactly $k-1$ translational invariant, relevant operators in these critical theories.

For all finite levels k we can identify an unusual symmetry Y in these anyonic chains that is sensitive to a topological flux in the center of a periodic chain [9]. The explicit form of the topological symmetry operator Y is constructed [5] by adding an extra $\nu = 1/2$ -line parallel to the spine of the fusion diagram and merging it with the diagram using the F -matrix (appropriate for the level k) [7]

$$\langle x'_1, \dots, x'_L | Y | x_1, \dots, x_L \rangle = \prod_{i=1}^L \left(F_{\nu x_i \nu}^{x'_{i+1}} \right)_{x_{i+1}}^{x'_i}. \quad (2)$$

The eigenvalues y_μ of the operator Y are given in terms of the modular- S matrix via $S_{\mu\nu}/S_{\mu 0}$ where $\mu = 0, 1/2, 1, \dots, k/2$ is a generalized angular momentum. Thus, there are $(k+1)$ distinct symmetry sectors for the $\text{su}(2)_k$ chain.

This topological symmetry plays a fundamental role in determining whether the observed gaplessness of the $\text{su}(2)_k$ spin-1/2 chains is protected and stable against local perturbations. Small, translationally invariant perturbations preserving this symmetry can only drive the critical system into a gapped phase if there is an operator (at momentum $K = 0$), relevant in the renormalization group sense, which is in the same topological sector as the ground state (typically the trivial sector). Relevant operators in other sectors break the topological symmetry and are thus prohibited from opening a gap. This is exactly the scenario that has been observed for the Fibonacci chain ($k = 3$), where the two translationally invariant relevant operators (the thermal operators ϵ and ϵ' in the tricritical Ising theory) are found to be in the topologically non-trivial sector, while the ground state is in the trivial sector [5]. For higher levels k the number of translationally invariant relevant operators grows as $k-1$, while the number of topological sectors grows as $k+1$. The question thus is whether these two diametrical effects can result in a cancellation and lead to a topological protection for all anyonic systems analogous to the case at $k = 3$.

We explain that this topological protection indeed exists for all finite levels k by observing a powerful connection between the coset theories describing the gapless state and the assignments of topological symmetry sectors to the relevant operators in these theories: The primary fields in these coset theories carry (a pair of) $\text{su}(2)_k$ labels like those of the original anyonic degrees of freedom. This observation allows us to obtain topological sectors for all fields in the gapless theory and identify those operators which for arbitrary k can drive the system into a gapped phase. In the limit $k \rightarrow \infty$ we recover the behavior of the ordinary spin-1/2 chain. We have checked, for the levels $k = 2, 3, 4, 5$, that the so-obtained topological assignments agree with results from exact diagonalization of chains with both AFM and FM interactions with up to $L = 24$ anyons (for $k = 5$) using Eq. (2).

In describing the details of the above topological symmetry assignments we will concentrate for brevity on the case that k is odd. In this case we can restrict ourselves to the 'integer spin' representations $0, 1, \dots, (k-1)/2$ of $\text{su}(2)_k$. The generalized spin-1/2 chains are then based on anyons carrying 'angular momentum' $j = (k-1)/2$ [10]. The fields $\phi_{j_2}^{j_1}$ in the gapless theories of the AFM chains carry two labels j_1, j_2 . In the coset construction of these theories, i.e. $\text{su}(2)_{k-1} \times \text{su}(2)_1 / \text{su}(2)_k$, the label j_2 corresponds to the representations of $\text{su}(2)_{k-1}$, the label i_2 to those of $\text{su}(2)_1$, and j_1 to those of $\text{su}(2)_k$. The label j_1 turns out to determine the topological sector of the fields. In particular, the fields in the topologically trivial sector turn out to be $\phi_{j_2}^0$. We consider the 'character decomposition' $\chi_{i_2}^{(1)} \chi_{j_2}^{(k-1)} = \sum_{j_1} B_{j_2}^{j_1} \chi_{j_1}^{(k)}$, where $\chi_j^{(k)}$ denotes the ('affine') character of $\text{su}(2)_k$ (and $i_2 = j_1 - j_2 \bmod 1$), and the $B_{j_2}^{j_1}$'s the (Virasoro) characters [11] of a unitary minimal model [8]. The Z_2 symmetry of these coset models also allows us to identify the sublattice symmetry of the primary fields. The states at $K = 0$ correspond to fields with integer j_2 , while the states at $K = \pi$ correspond to fields with half-integer j_2 . The scaling dimensions which result from this decomposition are those of the 'Kac-table', $x(k, j_1, j_2) = 2\{1 + j_2(j_2 + 1)/(k+1) - j_1(j_1 + 1)/(k+2)\}$. It easily follows that the ground state at $K = 0$ is in the trivial topological sector. The lowest lying field at $K = 0$ in this sector is ϕ_1^0 , with scaling dimension $2(k+3)/(k+1)$. Thus, we find that for any finite k , there is no relevant operator (with scaling dimension < 2) which can drive the system into a gapped phase. In the limit $k \rightarrow \infty$ of the ordinary spin-1/2 chain, this operator becomes exactly marginal. At momentum $K = \pi$, the most relevant field in the trivial sector is $\phi_{1/2}^0$, with scaling dimension $\frac{1}{2}(k+4)/(k+1)$ which is bounded by one for $k \geq 2$: staggered perturbations can always drive the system into a gapped 'dimerized' phase. We note that the results above hold for k even as well; with the difference that all $k+1$ topological sectors are present.

The critical behavior for FM interactions is described by the Z_k -parafermion coset $\text{su}(2)_k / \text{u}(1)$, whose fields also carry two labels, ψ_m^j . Again $j = 0, 1, \dots, (k-1)/2$ (for k odd) originates from $\text{su}(2)_k$ and determines the topological sector




FIG. 1: (color online) Phase diagrams of the ‘biquadratic’ spin-1 chain for a) ordinary SU(2) spin-1’s and b) the anyonic $su(2)_k$ -theories with $k \geq 5$. Projections onto the triplet ($j = 1$) and quintuplet ($j = 2$) states are parametrized by an angle θ as $J_1 = \sin(\theta)$ and $J_2 = -\cos(\theta)$.

as before, and $m = 0, 1, \dots, k-1$ determines the momentum, $K = 2\pi m/k$. The fields in the trivial sector are the Z_k -parafermion fields, ψ_m^0 , of scaling dimensions $2m(k-m)/k$. The only state whose corresponding scaling dimension is less than two at momentum $K = 0$ in the trivial sector is the ground state, e.g. the identity ψ_0^0 , implying that for all k this critical phase is stable against small perturbations which do not break translational invariance.

Anyonic spin-1 chains.— We now turn to chains formed by a set of anyons with generalized ‘angular momentum’ $\nu = 1$ in the $su(2)_k$ theory for $k \geq 4$. In the limit $k \rightarrow \infty$ we recover the ordinary biquadratic SU(2) spin-1 chain which has a rich phase diagram: There are two gapped phases, the Haldane phase [12] and a dimerized phase [13], and two gapless phases, a ferromagnetic phase and a nematic phase [14].

We have numerically calculated the phase diagrams of anyonic spin-1 chains for the $su(2)_5$ and $su(2)_7$ theories, which allows us to generalize this phase diagram to all levels $k \geq 5$. As illustrated in Fig. 1b), there are four phases that closely mirror their $k \rightarrow \infty$ counterparts: The Haldane phase survives as a gapped phase and includes a generalized AKLT point where the exact ground states are known [15]. It is surrounded by two gapless phases which have the same sublattice symmetries as their SU(2) counterparts, e.g. a Z_3 sublattice symmetry for the phase in the upper wedge which becomes the nematic phase [16], and a Z_2 sublattice symmetry for the phase in the lower wedge which turns into the dimerized phase. For arbitrary $k \geq 5$ the gapless phase in the upper wedge is the k -th member of the family of coset models $su(2)_{k-4} \times su(2)_4 / su(2)_k$. For $k = 5, 7$ we have numerically verified that the low-energy spectra indeed match these theories of central charges $c_k^{(4)} = 2 - 24/(k^2 - 4)$. The gapless phase in the lower wedge is for arbitrary $k \geq 5$ the k -th member of the family of the so-called off-diagonal modular invariants [17] of the unitary minimal models \mathcal{M}_k , a result which we have numerically verified. The critical endpoints of this phase with the Haldane phase correspond

to the $N = 1$ supersymmetric minimal models with central charge $c = 3/2 - 12/[k(k+2)]$. The chain Hamiltonian at this point can be mapped to an exactly integrable model [18]. In the $k \rightarrow \infty$ limit this critical point turns into the $su(2)_2$ Wess-Zumino-Witten (WZW) point of the ordinary biquadratic spin-1 chain.

Quantum Hall Liquids.— Our program of exploring collective states of anyonic spin chains is, at the same time, a tool to systematically study topological phases which can occur inside non-Abelian quantum Hall liquids due to population of such liquids by a macroscopic number of interacting non-Abelian anyons. This also provides us with the properties of the (so far unexplored) edge states appearing at the interfaces between these two liquids. Let us focus for brevity on *bosonic* quantum Hall fluids [19]. Corresponding statements for *fermionic* states with the same non-Abelian statistics [3] involve only differences in (trivial) Abelian factors.

First reconsider the case of a linear arrangement of $j = 1/2$ anyons in a surrounding $su(2)_k$ topological 2D fluid with pairwise interactions favoring singlet pairs. As we have seen, the gapless state is the minimal model \mathcal{M}_k . We can think of this critical state as two non-interacting counterpropagating (neutral) edge states of central charge $c_k = 1 - 6/(k+1)(k+2)$, which are basically located ‘on top of each other’, residing in the surrounding $su(2)_k$ topological bulk fluid. Let us now imagine separating these two edge states slightly in space, so that a narrow strip opens between them, see Fig. 2b). Let us ask if it is possible to place *another* topological quantum Hall fluid X into this narrow strip such that the above pair of counterpropagating edges of central charge c_k are precisely the edge states between the surrounding topological $su(2)_k$ fluid and the new, intervening fluid X in the strip. The answer to this question is ‘yes’: The intervening liquid X is a topological fluid characterized by $su(2)_{k-1} \times su(2)_1$. This can be seen [20] from the coset representation [8] of the minimal model $\mathcal{M}_k = su(2)_{k-1} \times su(2)_1 / su(2)_k$. Recall that the central charges of $X = su(2)_{k-1} \times su(2)_1$ (numerator) and




FIG. 2: Topological liquids and their edge states.

the surrounding fluid $su(2)_k$ (denominator) differ by c_k , thus resulting in an edge state between the two liquids of central charge c_k [21]. To summarize, our results say that the (AFM) interactions between an array of $j = 1/2$ anyons in an $su(2)_k$ topological liquid nucleate a new intervening liquid characterized by the topological properties of $su(2)_{k-1} \times su(2)_1$ [22].

Even though our anyons were initially confined to one dimension, these results will also apply to a macroscopic number of interacting non-Abelian anyons occupying two-dimensional (2D) regions of the surrounding liquid, thereby nucleating larger, 2D regions of the intervening liquid. The simplest example of this phenomenon was observed [25] for the Moore-Read Pfaffian quantum Hall liquid, which, when occupied with a macroscopic number of interacting quasi-hole anyons (even at random positions), turns into the simple Abelian ‘strong-pairing’ state. In general, the CFT of the edge state between the nucleated and the surrounding ‘parent’ liquid is given by the coset construction. For the simpler case of FM anyon interactions the liquid X is Abelian, and a Z_k parafermion CFT edge replaces the neutral \mathcal{M}_k edge CFT. The Pfaffian case above corresponds to $k = 2$.

Even more interesting examples arise when we express our results for the anyonic spin-1 chains in terms of intervening liquids. A family of novel liquids with topological character [23] $su(2)_{k-4} \times su(2)_4$ nucleated within a $su(2)_k$ surrounding liquid corresponds to the family of critical phases in the upper wedge of Fig. 1 b) with corresponding edge states described by the coset $su(2)_{k-4} \times su(2)_4 / su(2)_k$. Interestingly, the critical phases in the lower wedge of Fig. 1 b) describe the appearance of edge states between the $su(2)_k$ bulk fluid and a novel topological quantum liquid of type $su(2)_{k-1} \times su(2)_1$, with the property that the pair of edge states represents the *off-diagonal* modular invariant [17] of the minimal model \mathcal{M}_k . In all cases, the new resulting liquid has *less* anyon types than its $su(2)_k$ ‘parent’ liquid, due to the macroscopic occupation of the latter liquid by interacting anyons. The nucleation process thus amounts to a reduction of the non-Abelian statistics. Recent work in [24] also discusses the coset description of an interface between two different non-Abelian quantum liquids, one spin-polarized and the other spin-singlet. There, the reduction of non-Abelian statistics originates from polarization of spins.

In the context of topological liquids the topological symmetry described earlier acquires a very physical meaning: Local perturbations of the chain Hamiltonians correspond to tunneling events across the intervening liquid. It is precisely the perturbations in the topologically trivial sector which correspond to tunneling processes that are not accompanied by simulta-

neous ejection into the surrounding liquid of anyon particles with non-trivial topological quantum numbers [such processes correspond to $j_1 = 0$ in Fig 2b)].

We thank D. A. Huse for collaboration on a closely related project and M. Freedman, A. Kitaev, M. P. A. Fisher, A. Läuchli and K. Schoutens for discussions. Our numerical work employed the ALPS libraries [26]. A. W. W. L. was supported, in part, by NSF DMR-0706140.

-
- [1] More precisely $U_q[SU(2)]$ with q a root of unity. See, e.g., C. Kassel, *Quantum Groups*, Springer-Verlag, New York (1995).
- [2] N. Read and D. Green, Phys. Rev. B **61**, 10267 (2000).
- [3] N. Read and E. Rezayi, Phys. Rev. B **59**, 8084 (1999).
- [4] C. Nayak, *et al.*, Rev. Mod. Phys. **80**, 1083 (2008).
- [5] A. Feiguin, *et al.*, Phys. Rev. Lett. **98**, 160409 (2007).
- [6] S. Trebst, *et al.*, Phys. Rev. Lett. **101**, 050401 (2008).
- [7] S. Trebst, *et al.*, submitted to Prog. Theor. Phys. Suppl. (2008).
- [8] P. Goddard, *et al.*, Comm. Math. Phys. **103**, 105 (1986).
- [9] A similar symmetry exists for open chains.
- [10] The fusion rules imply that $1/2 \times k/2 = (k-1)/2$ and $(k-1)/2 \times (k-1)/2 = 0+1$. The spin-1/2 chains we consider for odd k are based on the $j = (k-1)/2$ representation.
- [11] Satisfying standard field identifications, e.g.: D. Gepner, Phys. Lett. **B 222**, 207 (1989).
- [12] F. D. M. Haldane, Phys. Lett. **93A**, 464 (1983); Phys. Rev. Lett. **50**, 1153 (1983).
- [13] M.N. Barber and M.T. Batchelor, Phys. Rev. B **40**, 4621 (1989); A. Klümper, Europhys. Lett. **9**, 815 (1989).
- [14] A. Läuchli, *et al.*, Phys. Rev. B **74**, 144426 (2006).
- [15] C. Gils, *et al.*, in preparation.
- [16] G. Fáth and J. Sólyom, Phys. Rev. B **44**, 11836 (1991).
- [17] A. Cappelli, C. Itzykson, and J.-B. Zuber, Nucl. Phys. **B280**, 445 (1987).
- [18] E. Date, *et al.*, Nucl. Phys. **B290**, 231 (1987).
- [19] see e.g., N.R. Cooper, N.K. Wilkin, and J.M.F. Gunn, Phys. Rev. Lett. **87**, 120405 (2001).
- [20] Consider joining $su(2)_k$ and $su(2)_{k-1} \times su(2)_1$ Hall liquids along a line, letting first *vacuum* intervene, see Fig. 2a). The, say, left moving $[su(2)_{k-1} \times su(2)_1]_L$ edge state between the latter fluid and vacuum is the same as the left moving CFT $[\mathcal{M}_k \times su(2)_k]_L$; this is the content of the coset construction [8]. The right moving edge state between the $su(2)_k$ fluid and the intervening vacuum is $[su(2)_k]_R$. Right and left $su(2)_k$ edges interact through the vacuum by a coupling of the two right and left $su(2)$ currents which is known to gap out these gapless modes. What remains is a left moving minimal model $[\mathcal{M}_k]_L$.
- [21] We can also rewrite this statement by bringing the Abelian factor $su(2)_1$, a semion, from the fluid X to the edge. We then arrive at the statement that juxtaposition of the two topological fluids $su(2)_k$ and $su(2)_{k-1}$, namely Read-Rezayi states, along a line yields an edge consisting of the (neutral) \mathcal{M}_k theory and a charged semion $su(2)_1$ moving in the opposite direction. For corresponding fermionic quantum Hall fluids the edge state between parafermionic Z_k and Z_{k-1} Read-Rezayi states is the neutral CFT \mathcal{M}_k and an oppositely moving Abelian $u(1)$ edge state. For $k = 3$, a tricritical Ising model \mathcal{M}_3 neutral edge appears between the Z_3 Read-Rezayi and Z_2 Moore-Read states.
- [22] The wave function of X is the product of the $su(2)_{k-1}$ and $su(2)_1$ wave functions. Symmetrization over the two species of

- particles gives the $su(2)_k$ Read-Rezayi state. (Cf., A. Cappelli, L.S. Georgiev, I.T. Todorov, Nucl. Phys. **B 599**, 499 (2001).)
- [23] Only the maximally Abelian subtheory $su(3)_1$ of $su(2)_4$ enters.
- [24] E. Grosfeld and K. Schoutens, arXiv:0810.1955.
- [25] N. Read and A.W.W. Ludwig, Phys. Rev. B **63**, 024404 (2000).
- [26] A.F. Albuquerque, *et al.*, J. of Magn. and Magn. Materials **310**, 1187 (2007).