

Symmetric Teleparallel Gravity: Some exact solutions and spinor couplings

Muzaffer Adak^{1*}, Özcan Sert^{1†}, Mestan Kalay^{1‡}, Murat Sari^{2§}

¹ Department of Physics, Faculty of Arts and Sciences, Pamukkale University
20017 Denizli, Turkey

² Department of Mathematics, Faculty of Arts and Sciences, Pamukkale University
20017 Denizli, Turkey

12 April 2012, *file CosmicSpeedUpInSTPG14.tex*

Abstract

In this paper we elaborate on the symmetric teleparallel gravity (STPG) written in a non-Riemannian spacetime with nonzero nonmetricity, but zero torsion and zero curvature. Firstly we give a prescription for obtaining the nonmetricity from the metric in a peculiar gauge. Then we state that under a novel prescription of parallel transportation of a tangent vector in this non-Riemannian geometry the autoparallel curves coincides with those of the Riemannian spacetimes. Subsequently we represent the symmetric teleparallel theory of gravity by the most general quadratic and parity conserving lagrangian with lagrange multipliers for vanishing torsion and curvature. We show that our lagrangian is equivalent to the Einstein-Hilbert lagrangian for certain values of coupling coefficients. Thus we arrive at calculating the field equations via independent variations. Then we obtain in turn conformal, spherically symmetric static, cosmological and pp-wave solutions exactly. Finally we discuss the minimal coupling of a spin-1/2 field to STPG.

PACS numbers: 04.50.Kd, 98.80.Jk, 02.40.Yy

Keywords: Non-Riemannian geometry, Modified theories of gravity, Lagrange formulation, Dirac equation

*madak@pau.edu.tr

†osert@pau.edu.tr

‡mkalay@pau.edu.tr

§msari@pau.edu.tr

1 Introduction

Einstein's theory of gravity, the so-called general relativity (GR), is formulated in a Riemannian spacetime. But there are convincing reasons to go beyond the Riemannian spacetime coming from both mathematical and physical points of view. Mathematically both group theoretical [1],[2] and gauge theoretical [3],[4] approaches predict non-Riemannian geometries with especially nonmetricity. Physically unsuccessful affords to quantize gravity with standard field theoretical methods are longstanding reasons for modification of GR. Furthermore the recent observational data received from type-Ia supernovae have showed that the expansion of the Universe is currently undergoing a period of acceleration [5],[6]. Cosmic speed-up can be explained within general relativity by invoking a mysterious cosmic fluid with large negative pressure, the so-called dark energy. Although the simplest possibility for dark energy is a cosmological constant, unfortunately the estimated value for it is too large compared with today's value. Consequently GR must be modified or replaced with new one.

Einstein's equations

$$(Ric)_a - \frac{1}{2}\mathcal{R}e_a + \Lambda e_a = \kappa *t_a \quad (1)$$

are obtained as local extremum of the action $I = \int_M L_0$ where

$$L_0 = -\frac{1}{2\kappa}R^a{}_b \wedge *e_a{}^b + \frac{\Lambda}{\kappa}*1 + \lambda_a \wedge T^a + \mu_{ab} \wedge Q^{ab} - K \quad (2)$$

is the Einstein-Hilbert lagrangian 4-form with the cosmological constant Λ . Here κ is a coupling constant, $t_a = \delta K/\delta e^a$ are the matter energy-momentum 3-forms, λ_a are the Lagrange multiplier 2-forms constraining torsion to zero, μ_{ab} are the Lagrange multiplier 3-forms constraining nonmetricity to zero. Other notations will be introduced in the next section. In one approach of modified gravity models one passes to the non-Riemannian geometries, in general $R^a{}_b \neq 0$, $T^a \neq 0$ and $Q_{ab} \neq 0$, sometimes the so-called the metric-affine gauge theory of gravity (MAG) [4]. Despite a big degrees of freedom of those models it is not an easy matter to handle general cases. Therefore people put some restrictions on their gravity models, see for example [4],[7],[8],[9],[10],[11] and the references therein. In this work we adhere the Yang-Mills manner for a gravitational lagrangian, that is, we consider the most general nonmetricity quadratic and parity conserving lagrangian 4-form

$$L = \sum_{I=1}^4 k_I Q_{ab} \wedge *(^I)Q^{ab} + k_5 \left((^3)Q_{ab} \wedge e^b \right) \wedge * \left((^4)Q^{ac} \wedge e_c \right) + \Lambda *1 + \lambda_a \wedge T^a + R^a{}_b \wedge \rho^b{}_a - K \quad (3)$$

where k_I s are coupling constants, $(^I)Q^{ab}$ are irreducible decompositions of nonmetricity [8],[12], Λ is the cosmological constant, K is the matter Lagrangian 4-form, λ_a are the Lagrange multiplier 2-forms for zero-torsion, $\rho^b{}_a$ are the Lagrange multiplier 2-forms for teleparallelism. In other

words, we have simply replaced the left hand side of (1) with a new one written in terms of only nonmetricity. This model is called symmetric teleparallel gravity (STPG) [13]. Thereby metric $g_{\alpha\beta}$ is seen as gauge potential and the corresponding gauge field strength is nonmetricity $Q_{\alpha\beta} \sim Dg_{\alpha\beta}$ that satisfies the Bianchi identity $DQ_{\alpha\beta} = 0$.

There is another teleparallel approach to gravity, the so-called teleparallel gravity (TPG), formulated only with torsion. Although TPG has attracted a lot of attention in the literature, STPG is infancy. One can consult for example the references [14],[15] for starting to read the wide literature on TPG. On the other hand, a few physically interesting analysis and solutions of STPG can be found in [12],[13],[16],[17]. Our wish here is to fill in a bit this big gap at STPG. Our another motivation is to gain new insights on the nonmetricity. Recently it has been related with the breaking of the Weyl symmetry [7], with the spin-3 gravity field [18] and used in the solar neutrino mixing problem [19].

The outline of the paper is as follows. In the Section 2 after introducing the non-Riemannian geometry and then symmetric teleparallel geometry we define a gauge fixing procedure and a deliberative parallel transportation rule for a tangent vector. From now on we use the computer algebra system REDUCE and its package EXCALC intensively [20],[21]. In the Section 3 we present a systematic treatment of STPG through a lagrange 4-form quadratic in the nonmetricity. Here we show explicitly that for certain values of coupling coefficients our lagrangian corresponds to the Einstein-Hilbert lagrangian in Riemannian spacetime. Then the field equations are obtained consistently by independent variations. By solving $D\lambda_a$ from the connection varied equation and then by inserting it into the coframe varied equation we write down the dynamical field equation (39) in a naive way. In the next four sections we obtain some classes of exact solutions of type: conformal, spherically symmetric static, cosmological and pp-wave, respectively. Finally we analyze a minimal coupling of a Dirac spinor to STPG. The Section 9 is devoted for the concluding remarks.

2 Mathematical preliminaries

Spacetime is denoted by the triple $\{M, g, \nabla\}$ where M is differentiable and orientable 4-dimensional manifold, $g = g_{\alpha\beta}e^\alpha \otimes e^\beta$ is the metric tensor, and ∇ represents connection. $\{X_\alpha\}$ denotes the basis vector set of tangent space $T_p(M)$ at the point p of manifold. Similarly, $\{e^\alpha\}$ denotes the basis co-vector set of cotangent space $T_p^*(M)$ at the point p of manifold. We show duality as

$$e^\alpha(X_\beta) \equiv \iota_{X_\beta}(e^\alpha) = \delta_\beta^\alpha \quad (4)$$

where δ_β^α is Kronecker symbol and $\iota_{X_\beta} \equiv \iota_\beta$ is the interior product. Full connection is determined by local 1-forms, ω^α_β as follows:

$$\nabla_{X_\alpha} X_\beta = \omega^\gamma_\beta(X_\alpha) X_\gamma. \quad (5)$$

From now on we call e^α basis 1-forms and $\omega^\alpha{}_\beta$ full connection 1-forms. Thus, under a general coordinate transformation while basis 1-forms transform as

$$e^{\alpha'} = h^{\alpha'}{}_\alpha e^\alpha \quad (6)$$

full connection 1-forms transform in the following manner

$$\omega^{\alpha'}{}_{\beta'} = h^{\alpha'}{}_\alpha \omega^\alpha{}_\beta h^\beta{}_{\beta'} + h^{\alpha'}{}_\alpha dh^{\alpha}{}_{\beta'} \quad (7)$$

where $h^{\alpha'}{}_\alpha$ are transformation elements. The following shorthand notation for exterior multiplications is adhered $e^\alpha \wedge e^\beta \wedge \dots \equiv e^{\alpha\beta\dots}$. A geometry is determined by the Cartan structure equations

$$Q_{\alpha\beta} := -\frac{1}{2}Dg_{\alpha\beta} = \frac{1}{2}(-dg_{\alpha\beta} + \omega_{\alpha\beta} + \omega_{\beta\alpha}), \quad (8)$$

$$T^\alpha := De^\alpha = de^\alpha + \omega^\alpha{}_\beta \wedge e^\beta, \quad (9)$$

$$R^\alpha{}_\beta := D\omega^\alpha{}_\beta := d\omega^\alpha{}_\beta + \omega^\alpha{}_\gamma \wedge \omega^\gamma{}_\beta, \quad (10)$$

where $Q_{\alpha\beta}$ are the nonmetricity 1-forms, T^α are the torsion 2-forms and $R^\alpha{}_\beta$ are the curvature 2-forms. d and D denote the exterior derivative and the covariant exterior derivative, respectively. These tensors satisfy the Bianchi identities

$$DQ_{\alpha\beta} = \frac{1}{2}(R_{\alpha\beta} + R_{\beta\alpha}), \quad (11)$$

$$DT^\alpha = R^\alpha{}_\beta \wedge e^\beta, \quad (12)$$

$$DR^\alpha{}_\beta = 0. \quad (13)$$

In general, full connection 1-forms can be decomposed as follows [4]-[8]:

$$\omega^\alpha{}_\beta = \underbrace{(g^{\alpha\gamma}dg_{\gamma\beta} + p^\alpha{}_\beta)/2 + \tilde{\omega}^\alpha{}_\beta}_{\text{Riemannian}} + \underbrace{K^\alpha{}_\beta}_{\text{Torsional}} + \underbrace{q^\alpha{}_\beta + Q^\alpha{}_\beta}_{\text{Nonmetrical}} \quad (14)$$

where Levi-Civita connection 1-forms $\tilde{\omega}^\alpha{}_\beta$

$$\tilde{\omega}^\alpha{}_\beta \wedge e^\beta = -de^\alpha, \quad (15)$$

contortion 1-forms $K^\alpha{}_\beta$

$$K^\alpha{}_\beta \wedge e^\beta = T^\alpha, \quad (16)$$

and anti-symmetric 1-forms

$$p_{\alpha\beta} = -(\iota_\alpha dg_{\beta\gamma})e^\gamma + (\iota_\beta dg_{\alpha\gamma})e^\gamma, \quad (17)$$

$$q_{\alpha\beta} = -(\iota_\alpha Q_{\beta\gamma})e^\gamma + (\iota_\beta Q_{\alpha\gamma})e^\gamma. \quad (18)$$

This decomposition is self-consistent. To see that one just has to multiply (14) by $\wedge e^\beta$ and to use above definitions, where special care is needed to raise and lower an index in front of both d and D. Symmetric part of (14) comes from (8)

$$\omega_{(\alpha\beta)} = Q_{\alpha\beta} + \frac{1}{2}dg_{\alpha\beta} \quad (19)$$

and the remainder is anti-symmetric

$$\omega_{[\alpha\beta]} = \frac{1}{2}p_{\alpha\beta} + \tilde{\omega}_{\alpha\beta} + K_{\alpha\beta} + q_{\alpha\beta} . \quad (20)$$

We denote symmetrization and anti-symmetrization by $\omega_{(\alpha\beta)} := \frac{1}{2}(\omega_{\alpha\beta} + \omega_{\beta\alpha})$ and $\omega_{[\alpha\beta]} := \frac{1}{2}(\omega_{\alpha\beta} - \omega_{\beta\alpha})$, respectively.

In general, the triple of nonmetricity, torsion and curvature classifies a geometry. If $Q_{\alpha\beta} = 0$, $T^\alpha \neq 0$ and $R^\alpha_\beta \neq 0$, we have a metric compatible connection in the Riemann-Cartan geometry. If $Q_{\alpha\beta} = 0$, $T^\alpha = 0$ and $R^\alpha_\beta \neq 0$, then we have the Levi-Civita connection of a Riemann geometry. If $Q_{\alpha\beta} = 0$, $T^\alpha \neq 0$ and $R^\alpha_\beta = 0$, it is called the Weitzenböck geometry. Here we work in the symmetric teleparallel geometry defined by $Q_{\alpha\beta} \neq 0$, $T^\alpha = 0$ and $R^\alpha_\beta = 0$.

It is always possible to choose orthonormal basis 1-forms which we denote $\{e^a(x)\}$. Then the metric turns out to be $g = \eta_{ab}e^a(x) \otimes e^b(x)$ where $\eta_{ab} = \text{diag}(-1, +1, +1, +1)$. In this case, the orientation of the manifold can be fixed by the Hodge star $*1 = e^{0123}$ and also the connection 1-forms decompose according to

$$\omega^a_b = \tilde{\omega}^a_b + K^a_b + q^a_b + Q^a_b . \quad (21)$$

A kind of gauge fixing

Suppose we choose a local coordinate system x^α and write the metric $g = g_{\alpha\beta}(x)dx^\alpha \otimes dx^\beta$. Then $e^\alpha = dx^\alpha$ and $de^\alpha = 0$. If we further require that $T^\alpha = 0$ and $R^\alpha_\beta = 0$, then these two conditions can be met provided $\omega^\alpha_\beta = 0$ so that $Q_{\alpha\beta} = -\frac{1}{2}dg_{\alpha\beta} \neq 0$. At this point a choice of local vierbein and its inverse allows us to pass to an orthonormal basis $e^a = h^a_\alpha dx^\alpha$. In the orthonormal frame, one calculates full connection $\omega^a_b = h^a_\alpha \omega^\alpha_\beta h^\beta_b + h^a_\alpha dh^\alpha_b = h^a_\alpha dh^\alpha_b \neq 0$, curvature $R^a_b = h^a_\alpha R^\alpha_\beta h^\beta_b = 0$, torsion $T^a = h^a_\alpha T^\alpha = 0$ and nonmetricity $Q^a_b = h^a_\alpha Q^\alpha_\beta h^\beta_b \neq 0$. This result corresponds to $\tilde{\omega}_{ab} + q_{ab} = 0$ in the orthonormal decomposition (21) which means $\omega_{ab} = Q_{ab}$ (together with $K_{ab} = 0$ since $T^a = 0$). This sequence of operations may be seen as a kind of gauge fixing.

Autoparallel curves in a symmetric teleparallel geometry

In the Riemann geometry one requires that the tangent vector, $V^\alpha = dx^\alpha/d\tau$, to the autoparallel curve, $x^\alpha(\tau)$ with the parameter τ , points in the same direction as itself when parallel

propagated, and demand that it maintains the same length,

$$DV^\alpha = 0. \quad (22)$$

On the other hand, since intuitively autoparallel curves are "those as straight as possible" we do not demand the vector to keep the same length [22] during parallel propagation in STPG. It is known that nonmetricity is related to length and angle standards at parallel transportation [4],[8]. Therefore we prescribe a novel parallel propagation of the tangent vector [23]

$$DV^\alpha = (Q^\alpha{}_\beta + q^\alpha{}_\beta)V^\beta. \quad (23)$$

Then in the gauge of $e^\alpha = dx^\alpha$ and $\omega^\alpha{}_\beta = 0$ the autoparallel curve equation is obtained as

$$\frac{d^2x^\alpha}{d\tau^2} + \frac{1}{2}g^{\alpha\beta}(\partial_\gamma g_{\beta\delta} + \partial_\delta g_{\beta\gamma} - \partial_\beta g_{\gamma\delta})\frac{dx^\gamma}{d\tau}\frac{dx^\delta}{d\tau} = 0. \quad (24)$$

This is the same as the geodesic equation of the Riemann geometry.

3 Field equations of STPG

With the assignments

$$\alpha_a = Q_{ab} \wedge e^b, \quad \beta_a = l^b Q_{ab}, \quad Q = \eta_{ab} Q^{ab}, \quad \gamma_a = \iota_a Q \quad (25)$$

the lagrangian (3) is equivalent to the following

$$\begin{aligned} L = & c_1 Q_{ab} \wedge^* Q^{ab} + c_2 \alpha_a \wedge^* \alpha^a + c_3 \beta_a \wedge^* \beta^a + c_4 Q \wedge^* Q + c_5 \beta_a \wedge^* \gamma^a \\ & + \Lambda^* 1 + \lambda_a \wedge T^a + R^a{}_b \wedge \rho^b{}_a - K \end{aligned} \quad (26)$$

under the redefinitions of the coupling coefficients, see [12]

$$\begin{aligned} c_1 &= k_1, \\ c_2 &= 2(-k_1 + k_2)/3, \\ c_3 &= 2(-k_1 - k_2 + 2k_3)/9, \\ c_4 &= (-8k_1 - 32k_2 + 4k_3 + 36k_4 + 9k_5)/144, \\ c_5 &= (-8k_1 + 16k_2 - 8k_3 - 9k_5)/36. \end{aligned} \quad (27)$$

Since an odd number of Hodge stars occurring in a Lagrangian warrants its parity conservation the Lagrangian (3) or (26) is the most general nonmetricity quadratic and parity conserving lagrangian.

Certain values of c_i s correspond to GR. For determining these values firstly the full curvature 2-form is decomposed as Riemannian curvature plus non-Riemannian terms by substituting (21) (with $T^a = 0$) into $R^a{}_b$;

$$R^a{}_b = \tilde{R}^a{}_b + \tilde{D}\Omega^a{}_b + \Omega^a{}_c \wedge \Omega^c{}_b \quad (28)$$

where $\Omega_{ab} = Q_{ab} + q_{ab}$. Riemannian quantities are marked by a tilde. Then the Einstein-Hilbert lagrangian 4-form is formed

$$R^a{}_b \wedge *e_a{}^b = \tilde{R}^a{}_b \wedge *e_a{}^b + \left(\tilde{D}\Omega^a{}_b \right) \wedge *e_a{}^b + \Omega^a{}_c \wedge \Omega^c{}_b \wedge *e_a{}^b. \quad (29)$$

Here since $\tilde{D}e_a{}^b = 0$, the second term is an exact form, i.e. $\left(\tilde{D}\Omega^a{}_b \right) \wedge *e_a{}^b = d\left(\Omega^a{}_b \wedge *e_a{}^b \right)$. Finally, because of vanishing of the full curvature 2-form in this geometry the left hand is zero. Correspondingly, it may be written

$$-\frac{1}{2}\tilde{R}^a{}_b \wedge *e_a{}^b = \frac{1}{2}\Omega^a{}_c \wedge \Omega^c{}_b \wedge *e_a{}^b \quad (30)$$

from which one computes the GR-equivalent values of c_i s by comparing with (26)

$$c_1 = -1/2, \quad c_2 = 1, \quad c_3 = 0, \quad c_4 = -1/2, \quad c_5 = 1. \quad (31)$$

Now we vary the lagrangian (26) to obtain the field equations by noticing $\delta Q_{ab} = \delta\omega_{(ab)}$ and $(\delta\iota^a)e^b = -\iota^a(\delta e^b)$, and by using the general formula

$$\delta(A \wedge *B) = \delta A \wedge *B + \delta B \wedge *A - \delta e^a \wedge [(\iota_a B) \wedge *A - (-1)^p A \wedge (\iota_a *B)] \quad (32)$$

where A and B are any p -forms on n -dimensional manifold. Thus the variations with respect to λ_a , $\rho^b{}_a$, e^a and $\omega^b{}_a$ yield the following field equations, respectively

$$T^a = 0, \quad R^a{}_b = 0, \quad (33)$$

$$\tau_a + \Lambda^*e_a + D\lambda_a = t_a, \quad (34)$$

$$\Sigma^a{}_b + e^a \wedge \lambda_b + D\rho^a{}_b = \ell^a{}_b, \quad (35)$$

where $t_a = \frac{\delta K}{\delta e^a}$ are the energy-momentum 3-forms of matter, $\ell^a{}_b = \frac{\delta K}{\delta \omega^b{}_a}$ are the angular momentum 3-forms of matter, τ_a are the energy-momentum 3-forms of nonmetricity

$$\begin{aligned} \tau_a &= c_1[\iota_a(Q_{bc} \wedge *Q^{bc}) - 2(\iota_a Q_{bc}) \wedge (*Q^{bc})] + c_2[\iota_a(\alpha_b \wedge *\alpha^b) - 2(\iota_a Q_{bc}) \wedge (e^c \wedge *\alpha^b)] \\ &\quad + c_3[\iota_a(\beta_b \wedge *\beta^b) - 2(\iota_a Q_{bc}) \wedge (\iota^c *\beta^b)] + c_4[\iota_a(Q \wedge *Q) - 2(\iota_a Q_{bc}) \wedge (\eta^{bc} *Q)] \\ &\quad + c_5[\iota_a(\beta_b \wedge *\gamma^b) - (\iota_a Q_{bc}) \wedge (\eta^{bc} \iota_d *\beta^d + \iota^c *\gamma^b)] \end{aligned} \quad (36)$$

and $\Sigma^a{}_b$ are the angular momentum 3-forms of nonmetricity

$$\Sigma^{ab} = 2c_1 *Q^{ab} + 2c_2 e^{(a} \wedge *\alpha^{b)} + 2c_3 \iota^{(a} *\beta^{b)} + 2c_4 \eta^{ab} *Q + c_5 \left(\eta^{ab} \iota_c *\beta^c + \iota^a *\gamma^b \right). \quad (37)$$

We count the number of the field equations under the assumption of given t_a and $\ell^a{}_b$: 24 from $T^a = 0$ plus 96 from $R^a{}_b = 0$ plus 16 from (34) plus 64 from (35) that add to 200. On the other hand, the number of the unknowns is 200 as well: 16 for e^a plus 64 for $\omega^a{}_b$ plus 24 for λ_a plus 96 for $\rho^a{}_b$. Thus the system is closed. In principle, the Lagrangian multipliers are solved from

(35) and the result is substituted into (34). We notice that what we need in the equation (34) is only $D\lambda_a$ rather than λ_a or ρ^a_b . Thus, by multiplying (35) first by D and then by ι_a we obtain

$$D\lambda_a = \iota_b D(\Sigma^b_a - \ell^b_a) \quad (38)$$

where we used $De^a = T^a = 0$ and $D^2\rho^a_b = R^a_c \wedge \rho^c_b - R^c_b \wedge \rho^a_c = 0$. Substitution of this into (34) yields the dynamical field equation of STPG

$$\iota_b D\Sigma^b_a + \tau_a + \Lambda *e_a = t_a + \iota_b D\ell^b_a. \quad (39)$$

4 Conformal solution

We consider the metric

$$g = \exp(2\Phi)\eta_{\alpha\beta}dx^\alpha \otimes dx^\beta \quad (40)$$

where $\eta_{\alpha\beta} = \text{diag}(-1, +1, +1, +1)$ and $\Phi = \Phi(t, x, y, z)$ is the conformal function in the cartesian coordinates, $x^\alpha = (t, x, y, z)$. In the coordinate frame, i.e. $e^\alpha = dx^\alpha$, we choose such a gauge that $\omega^\alpha_\beta = 0$ and then calculate the nonmetricity via $Q_{\alpha\beta} = -\frac{1}{2}dg_{\alpha\beta}$ as

$$Q_{\alpha\beta} = -\eta_{\alpha\beta} \exp(2\Phi)d\Phi. \quad (41)$$

Now we pass to the orthonormal frame

$$g = \eta_{ab}e^a \otimes e^b \quad \text{with} \quad e^a = \exp(\Phi)dx^a. \quad (42)$$

Accordingly the nonmetricity in the orthonormal frame can be computed

$$Q_{ab} = -\eta_{ab}d\Phi. \quad (43)$$

Here we checked that $\tilde{\omega}_{ab} + q_{ab} = 0$. After substituting this into the equation (39) with $t_a = 0$ and $\ell^a_b = 0$ we see that the conformal function is arbitrary under the constraints

$$4c_1 + 5c_2 - c_3 + 16c_4 = 0, \quad 2(c_1 + c_2 + 4c_4) + c_5 = 0 \quad (44)$$

with $\Lambda = 0$. We also remark that GR-values of c_i s (31) do not satisfy these constraints.

5 Solution in the vicinity of spherically symmetric static body

The vicinity of spherically symmetric static body means that $t_a = 0$, $\ell^a_b = 0$ and

$$g = -F^2 dt^2 + G^2 dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) \quad (45)$$

where F and G are functions of the radial coordinate r . Now in the coordinate frame and in the gauge $\omega^\alpha_\beta = 0$ we calculate the nonmetricity, $Q_{\alpha\beta} = -\frac{1}{2}dg_{\alpha\beta}$,

$$\begin{aligned} Q_{tt} &= FF'dr, & Q_{rr} &= -GG'dr, & Q_{\theta\theta} &= -rdr, \\ Q_{\phi\phi} &= -r\sin^2\theta dr - r^2\sin\theta\cos\theta d\theta, & \text{others} &= 0, \end{aligned} \quad (46)$$

where prime denotes derivative with respect to r . Then we pass to the orthonormal frame

$$e^0 = Fdt, \quad e^1 = Gdr, \quad e^2 = rd\theta, \quad e^3 = r\sin\theta d\phi. \quad (47)$$

That gives the chance of computing the nonmetricity in the orthonormal frame

$$Q_{00} = \frac{F'}{FG}e^1, \quad Q_{11} = -\frac{G'}{G^2}e^1, \quad Q_{22} = -\frac{1}{rG}e^1, \quad Q_{33} = -\frac{1}{rG}e^1 - \frac{\cot\theta}{r}e^2, \quad \text{others} = 0. \quad (48)$$

When we substitute these findings to the field equation (39) we obtain a very complicated set of equations. Therefore we try simplifying them firstly by choosing $G = 1/F$. Then under the case of $c_1 \neq 0$, $c_2 = c_5 = -2c_1$, $c_3 = 0$, $c_4 = c_1$ a dynamical solution is obtained

$$F^2 = 1 - \frac{M}{r} + \frac{\Lambda}{6c_1}r^2 \quad (49)$$

where M is an integration constant. For $c_1 = -1/2$ this is exactly the Schwarzschild-de Sitter solution of GR. This result is also a crosscheck of GR-values of c_i s (31). If we set $c_1 = c_2 = c_4 = c_5 = 0$, but $c_3 \neq 0$ we find another solution

$$F = a + br, \quad G = 1 \quad (50)$$

where a and b are constants. These solutions are not conformally invariant. One class of conformally invariant solution is obtained as

$$F = \exp(-m/r), \quad G = kr \quad (51)$$

with $\Lambda = 0$ and the constraints $c_1 \neq 0$, $c_2 = c_3 = -c_1$, $c_4 = c_5 = 0$, where m and k are arbitrary constants.

6 Homogeneous and isotropic cosmology

In spite of the fact that if a model more general than the general relativity is investigated, principally a complete model with any kind matter such as hyperfluid, spinorial etc has to be considered, we assume at this point that the matter lagrangian does not depend explicitly on the connection, i.e. $\ell^a_b = 0$. This assumption is because we will be searching cosmological solutions with a matter source in the form of non-spinning perfect fluid. Otherwise, one can

always drop this restriction for spinning sources. Thus we consider the non-spinning perfect fluid energy-momentum 3-form

$$t_a = [(\rho + p)\eta_{0a}\eta_{0b} + \eta_{ab}p]^* e^b \quad (52)$$

where ρ is matter density, p is pressure. We consider the Robertson-Walker metric in the isotropic coordinates $x^\alpha = (t, x, y, z)$

$$g = -dt^2 + S^2(t) \frac{dx^2 + dy^2 + dz^2}{(1 + kr^2/4)^2} \quad (53)$$

where k is a constant, $r^2 = x^2 + y^2 + z^2$ and $S(t)$ is the expansion function. First, we choose such a frame and a connection that $e^\alpha = dx^\alpha$ and $\omega^\alpha_\beta = 0$, then torsion and curvature are both zero, but $Q_{\alpha\beta} = -\frac{1}{2}dg_{\alpha\beta} \neq 0$,

$$Q_{xx} = Q_{yy} = Q_{zz} = -\frac{S\dot{S}dt}{(1 + kr^2/4)^2} + \frac{kS^2(xdx + ydy + zdz)}{2(1 + kr^2/4)^3}, \quad \text{others} = 0 \quad (54)$$

where dot denotes derivative with respect to t . Now we pass to the orthonormal frame

$$e^0 = dt, \quad e^1 = \frac{S}{1 + kr^2/4} dx, \quad e^2 = \frac{S}{1 + kr^2/4} dy, \quad e^3 = \frac{S}{1 + kr^2/4} dz \quad (55)$$

and correspondingly calculate

$$Q_{11} = Q_{22} = Q_{33} = -\frac{\dot{S}}{S}e^0 + \frac{k}{2S}(xe^1 + ye^2 + ze^3), \quad \text{others} = 0. \quad (56)$$

Now we check $\tilde{\omega}_{ab} + q_{ab} = 0$ which means $\omega_{ab} = Q_{ab}$ together with $K_{ab} = 0$ in this frame according to (21). Here again we crosschecked GR-values of c_i s given by (31). Then we deal with the case with $k = 0$, because cosmic microwave background experiments have indicated that the geometry of the Universe is spatially flat [24],[25]. Thus by setting $k = 0$ from beginning we arrive at the equations

$$3a\ddot{S}/S + 3b\left(\dot{S}/S\right)^2 = -(\Lambda + \rho) \quad (57)$$

$$(b - 2a)\left[2\ddot{S}/S + \left(\dot{S}/S\right)^2\right] = -(\Lambda - p) \quad (58)$$

where $a = 2c_4 + c_5$ and $b = c_1 + c_2 + 7c_4 + 2c_5$. Here note that because of the assumed homogeneity, ρ and p are functions of t alone. Here there are two equations: (57), (58) but three unknowns: S, p, ρ . So by postulating an equation of state $p = w\rho + B$ we close the system where w and B are constants. This equation of state takes place in the modified Chaplygin gas defined as $p = w\rho + B/\rho^\alpha$ where $B < 0$ and $0 \leq \alpha \leq 1$. As a result we obtain a solution

$$S = \left[K_2 e^{-nt} (e^{nt} - K_1)^2\right]^{m/n} \quad (59)$$

where K_1 and K_2 are integration constants, and

$$m = \sqrt{\frac{-B + (1 + w)\Lambda}{3(2a - b - w(a + b))}}, \quad n = \frac{6m(2a - b - w(a + b))}{4a - 2b - 3aw}. \quad (60)$$

7 pp-Wave Solution

We start with the metric

$$g = -dt^2 + p^2 dx^2 + q^2 dy^2 + dz^2 \quad (61)$$

where $p = p(u)$ and $q = q(u)$ are functions of the null coordinate $u = z - t$. This metric represents the plane fronted the gravitational waves propagating in the z -direction with the speed of light. In the gauge $e^\alpha = dx^\alpha$ and $\omega^\alpha_\beta = 0$ we calculate the nonmetricity, $Q_{\alpha\beta} = -\frac{1}{2}dg_{\alpha\beta}$

$$Q_{xx} = -pdp, \quad Q_{yy} = -qdq, \quad \text{others} = 0. \quad (62)$$

Now we pass to orthonormal frame

$$\begin{aligned} e^0 &= dt, & e^1 &= p dx, & e^2 &= dy, & e^3 &= dz, \\ Q_{11} &= -\frac{p'}{p}(e^3 - e^0), & Q_{22} &= -\frac{q'}{q}(e^3 - e^0), & \text{others} &= 0, \end{aligned} \quad (63)$$

where prime denotes derivative with respect to u . When we insert these components of nonmetricity to the Eqn(39) in the vacuum we obtain $\Lambda = 0$. Then the set of equations turns out to be

$$c_5 pq(pq'' + qp'') + 2(c_1 + c_2 + c_4)[(pq')^2 + (qp')^2] + 2(2c_4 + c_5)pq'q' = 0 \quad (64)$$

Here we checked that GR-equivalent values of c_i s gives the Einstein equation, namely $p''/p + q''/q = 0$. There are numerous solutions to this equation in the literature, for example $p = \cos(a_1 u + a_2)$ and $q = \cosh(a_1 u + a_3)$ where a_i s are arbitrary constants. On the other hand if $c_1 + c_2 = 0$, then $pq = \text{constant}$ is a solution. Consequently we see that STPG predicts gravitational waves as well.

8 A Dirac field coupling to STPG

Matter is made up mainly fermionic particles. Therefore we investigate a spin-1/2 particle in STPG. We are using the formalism of Clifford algebra $\mathcal{C}\ell_{1,3}$ -valued exterior forms. The $\mathcal{C}\ell_{1,3}$ algebra is generated by the relation among the orthonormal basis $\{\gamma_0, \gamma_1, \gamma_2, \gamma_3\}$

$$\gamma_{(a}\gamma_{b)} = \eta_{ab}. \quad (65)$$

One particular representation of the γ^a 's is given by the following Dirac matrices

$$\gamma_0 = i \begin{pmatrix} -I & 0 \\ 0 & I \end{pmatrix}, \quad \gamma_1 = i \begin{pmatrix} 0 & \sigma^1 \\ -\sigma^1 & 0 \end{pmatrix}, \quad \gamma_2 = i \begin{pmatrix} 0 & \sigma^2 \\ -\sigma^2 & 0 \end{pmatrix}, \quad \gamma_3 = i \begin{pmatrix} 0 & \sigma^3 \\ -\sigma^3 & 0 \end{pmatrix} \quad (66)$$

where $\sigma^1, \sigma^2, \sigma^3$ are the Pauli matrices. In this case a Dirac spinor Ψ can be represented by a 4-component column matrix. Thus we write explicitly the covariant exterior derivative of Ψ via the Kosmann lift

$$D\Psi = d\Psi + \frac{1}{2}\omega^{ab}(\eta_{ab} + \sigma_{ab})\Psi \quad (67)$$

where $\sigma_{ab} := \frac{1}{2}\gamma_{[a}\gamma_{b]}$ are the generators of the Lorentz group [8],[12],[26]. Correspondingly we calculate $D^2\Psi$

$$D(D\Psi) = \frac{1}{2}R^{ab}(\eta_{ab} + \sigma_{ab})\Psi. \quad (68)$$

Here let us note that in teleparallel geometries as the curvature of cotangent bundle, i.e. R^a_b , is zero, the curvature of the spinor bundle, i.e. $\frac{1}{2}R^{ab}(\eta_{ab} + \sigma_{ab})$, is zero as well. Some relations of the Dirac matrices are

$$\begin{aligned} \sigma_{ab}\gamma_c &= \gamma_c\sigma_{ab} + \eta_{bc}\gamma_a - \eta_{ac}\gamma_b, \\ \sigma_{ab}\gamma_c + \gamma_c\sigma_{ab} &= -\epsilon_{abcd}\gamma^d\gamma_5, \\ \gamma_c\sigma_{ab} &= \frac{1}{2}\eta_{ac}\gamma_b - \frac{1}{2}\eta_{bc}\gamma_a - \frac{1}{2}\epsilon_{abcd}\gamma^d\gamma_5, \\ [\sigma_{ab}, \sigma_{cd}] &= -\eta_{ac}\sigma_{bd} - \eta_{bd}\sigma_{ac} + \eta_{ad}\sigma_{bc} + \eta_{bc}\sigma_{ad}, \end{aligned} \quad (69)$$

where $\gamma_5 := \gamma_0\gamma_1\gamma_2\gamma_3$. In order to obtain the Bjorken-Drell conventions [27] one has to replace $\gamma^a \rightarrow -i\gamma^a$ and $\gamma_5 \rightarrow i\gamma_5$. Now we take the minimally coupled Dirac lagrangian given by the *hermitian* 4-form

$$K = \frac{i}{2}(\bar{\Psi} * \gamma \wedge D\Psi + D\bar{\Psi} \wedge * \gamma \Psi) + im \bar{\Psi} \Psi * 1, \quad (70)$$

where $\gamma := \gamma_a e^a$ is $\mathcal{C}\ell_{1,3}$ -valued 1-form and $\bar{\Psi}$ is the Dirac adjoint spinor, $\bar{\Psi} := \Psi^\dagger \gamma_0$. The coframe e^a necessarily occurs in the Dirac Lagrangian, even in special relativity. The hermiticity of the lagrangian (70) leads to a charge current which admits the usual probabilistic interpretation. Then for a Dirac field we calculate the energy-momentum 3-form

$$t_a = \frac{i}{2}[\bar{\Psi} * (\gamma \wedge e_a) \wedge D\Psi - D\bar{\Psi} \wedge * (\gamma \wedge e_a) \Psi] + im \bar{\Psi} \Psi * e_a, \quad (71)$$

the angular momentum 3-form

$$\ell^a_b = \frac{i}{4}(\bar{\Psi} \gamma \gamma_5 \Psi) \wedge e^a_b. \quad (72)$$

The matter field equation $\delta K / \delta \bar{\Psi} = 0$ yields the Dirac equation in the most general non-Riemannian spacetime with $Q_{ab} \neq 0$, $T^a \neq 0$, $R^a_b \neq 0$;

$$* \gamma \wedge \left(D - \frac{1}{2}T - Q \right) \Psi + \gamma_{(a} * e_{b)} \wedge Q^{ab} \Psi + m \Psi * 1 = 0 \quad (73)$$

where $T = \iota_a T^a$ which is zero in STPG. So, in the gauge, $e^\alpha = dx^\alpha$ and $\omega^{\alpha\beta} = 0$, the Dirac equation turns out to be

$$* \gamma \wedge \left(d - \frac{1}{2}Q \right) \Psi + \gamma_{(a} * e_{b)} \wedge Q^{ab} \Psi + m \Psi * 1 = 0. \quad (74)$$

9 Concluding Remarks

In this paper we studied the general symmetric teleparallel gravity model within the framework of the generic non-Riemannian formulation. A similar but narrow-scoped work was performed in [12]. Generalizing the previous work, we prescribed a gauge, $e^\alpha = dx^\alpha$ and $\omega^a{}_\beta = 0$, for writing down the nonmetricity directly from a metric ansatz, then obtained new solutions strictly with the help of computer and finally considered the minimal Dirac spinor coupling to STPG in a consistent way.

We described a peculiar parallel transportation of a tangent vector. Thus for the fixed gauge the autoparallel curve of the symmetric teleparallel geometry coincides with that of the Riemannian geometry. As a consequence we may cast the orbital equations of the planets in the solar system in a correct way for at least spherically symmetric solution (49). We also showed that some values of the c_i parameters are the GR-equivalent at the level of lagrangian and solutions. It is important to know them because the GR-equivalent STPG, for obvious reasons, is satisfactorily supported by observations. Since the first most important motivation for a new theory of gravity was the quantization problems of GR and the conformally invariant theories predict better UV behavior [28], we investigated that type of solution. But whether our scale-invariant solution really makes sense at the quantum level is an open issue. We could find three classes of the spherically symmetric static solutions. Nevertheless there may be more solutions of this type. The second most important motivation for looking for a new theory of gravity was the cosmic speed-up. Correspondingly, we investigated a cosmological solution for a non-spinning perfect fluid source. Our cosmological solution can predict accelerating universe by some suitable adjustments of coupling parameters. Furthermore we looked at the pp-wave solution and saw that STPG accepts the existence of gravitational waves. Finally because of the fact that matter is mainly made up by fermions it was indispensable to elaborate consistently on a minimal Dirac spinor coupling to STPG. Thus we wrote down naively the Dirac equation in the most general non-Riemannian geometry and then reduced it to the symmetric teleparallel geometry in our special gauge.

Acknowledgment

MA thanks T Dereli for the stimulating discussions, F W Hehl for the fruitful correspondences and B Tekin for the valuable conversations especially on Section 4.

References

- [1] Kirsch I, *Phys. Rev. D* **72** (2005) 024001

- [2] Boulanger N and Kirsch I, *Phys. Rev. D* **73** (2006) 124023
- [3] Benn I M, Dereli T and Tucker R W, *J. Phys. A* **15** (1982) 849
- [4] Hehl F W *et al.*, *Phys. Rep.* **258** (1995) 1
- [5] Riess A G *et al.*, *Astron. J.* **116** (1998) 1009
- [6] Perlmutter S *et al.*, *Astron. J.* **517** (1999) 565
- [7] Tucker R W and Wang C, *Class. Quant. Grav.* **12** (1995) 2587
- [8] Adak M, *Non-Riemannian Theories of Gravitation and the Dirac Equation*, Lambert Academic Pub., 2010, ISBN: 978-3-8383-5139-1
- [9] Dereli T *et al.*, *Class. Quant. Grav.* **13** (1996) L103
- [10] Dereli T and Tucker R W, *JHEP* **03** (2002) 041
- [11] Baekler P *et al.*, *Phys. Rev. D* **83** (2011) 024001
- [12] Adak M *et al.*, *Int. J. Mod. Phys. D* **15** (2006) 619
- [13] Nester J M and Yo H J, *Chinese J. Phys.* **37** (1999) 113
- [14] Obukhov Y N and Pereira, *Phys. Rev. D* **67** (2003) 044016
- [15] Maluf J W and Faria F F, *Phys. Rev. D* **85** (2012) 027502
- [16] Adak M and Sert Ö, *Turk. J. Phys.* **29** (2005) 1
- [17] Adak M and Dereli T, *Europhys. Lett.* **82** (2008) 30008
- [18] Baekler P, Boulanger N and Hehl F W, *Phys. Rev. D* **74** (2006) 125009
- [19] Adak M, Dereli T and Ryder L H, *Phys. Rev. D* **69** (2004) 123002
- [20] Hearn A C, 2004 *REDUCE User's Manual Version 3.8*, <http://www.reduce-algebra.com/docs/reduce.pdf>
- [21] Schrüfer E, 2004 *EXCALC: A System for Doing Calculations in the Calculus of Modern Differential Geometry*, <http://www.reduce-algebra.com/docs/excalc.pdf>
- [22] Wald R M, *General Relativity*, The University of Chicago Press, 1984, ISBN: 0-226-87033-2
- [23] Adak M, *A Note on Parallel Transportation in Symmetric Teleparallel Geometry*, arXiv:1102.1878

- [24] Bernardis P de *et al.*, *Nature* **404** (2000) 955
- [25] Stompor R *et al.*, *Ap. J.* **561** (2001) L7
- [26] Kosmann Y, *C. R. Acad. Sci.* **262** (1966) 289
- [27] Bjorken J D and Drell S D, *Relativistic Quantum Mechanics* (McGraw-Hill, New York, 1964)
- [28] Tanhayi M R, Dengiz S and Tekin B, *Phys. Rev. D* **85** (2012) 064016