

On a class of multidimensional integrable hierarchies and their reductions

L.V. Bogdanov*

Abstract

A class of multidimensional integrable hierarchies connected with commutation of general (unreduced) $(N+1)$ -dimensional vector fields containing derivative over spectral variable is considered. They are represented in the form of generating equation, as well as in the Lax-Sato form. A dressing scheme based on nonlinear vector Riemann problem is presented for this class. The hierarchies connected with Manakov-Santini equation and Dunajski system are considered as illustrative examples.

1 Introduction

Using a construction developed in [1, 2], we describe a structure of general integrable hierarchy connected with commutation of $(N + 1)$ -dimensional vector fields (containing a derivative over spectral variable), which defines $(N + 2)$ -dimensional integrable equations. We consider several special cases of the general hierarchy and its reductions, some of them defining known integrable models. The simplest case of $(1+1)$ -dimensional vector field (one space variable and a spectral variable) corresponds to $(2+1)$ -dimensional integrable system introduced by Manakov and Santini [3] (see also [4, 5]). The case of $(2+1)$ -dimensional vector fields (two space variables and a spectral variable), after a reduction to volume-preserving vector fields, leads to $(2+2)$ -dimensional Dunajski integrable model [6], describing anti-self-dual Einstein-Maxwell equation for signature $(2,2)$ (twistor setting).

*L.D. Landau ITP, Kosygin str. 2, Moscow 119334, Russia, e-mail leonid@landau.ac.ru

2 General $(N + 2)$ -dimensional hierarchy

To define general hierarchy, we consider $N + 1$ formal series, depending on N infinite sets of additional variables ('times')

$$\Psi^0 = \lambda + \sum_{n=1}^{\infty} \Psi_n^0(\mathbf{t}^1, \dots, \mathbf{t}^N) \lambda^{-n}, \quad (1)$$

$$\Psi^k = \sum_{n=0}^{\infty} t_n^k (\Psi^0)^n + \sum_{n=1}^{\infty} \Psi_n^k(\mathbf{t}^1, \dots, \mathbf{t}^N) (\Psi^0)^{-n}. \quad (2)$$

where $0 \leq k \leq N$, $\mathbf{t}^k = (t_0^k, \dots, t_n^k, \dots)$. We denote $\partial_n^k = \frac{\partial}{\partial t_n^k}$ and introduce the projectors $(\sum_{-\infty}^{\infty} u_n \lambda^n)_+ = \sum_{n=0}^{\infty} u_n \lambda^n$, $(\sum_{-\infty}^{\infty} u_n \lambda^n)_- = \sum_{-\infty}^{n=-1} u_n \lambda^n$.

The general hierarchy is defined by the generating relation

$$(J_0^{-1} d\Psi^0 \wedge d\Psi^1 \wedge \dots \wedge d\Psi^N)_- = 0, \quad (3)$$

where the differential includes both times and a spectral variable,

$$df = \sum_{k=1}^N \sum_{n=0}^{\infty} \partial_n^k f dt_n^k + \partial_\lambda f d\lambda,$$

and J_0 is a determinant of Jacobian matrix J_{ij} ,

$$J_{ij} = \partial_i \Psi^j, \quad 0 \leq i, j \leq N, \quad \partial_0 = \frac{\partial}{\partial \lambda}, \quad \partial_k = \frac{\partial}{\partial x^k}, \quad 1 \leq k \leq N,$$

where $x^k = t_0^k$. Equation (3) is a generating relation for Lax-Sato form of the hierarchy.

Proposition 1 *Relation (3) is equivalent to the set of equations*

$$\partial_n^k \Psi = \sum_{i=0}^N (J_{ki}^{-1} (\Psi^0)^n)_+ \partial_i \Psi, \quad 0 \leq n \leq \infty, 1 \leq k \leq N. \quad (4)$$

The proof of this proposition is completely analogous to the similar proof given in [2] for Dunajski hierarchy. The proof of (3) \Rightarrow hierarchy (4) is based on the following statement.

Lemma 1 Given identity (3), for arbitrary first order operator \hat{U} ,

$$\hat{U}\Psi = \left(\sum_{k=1}^N \sum_i u_i^k(\lambda, \mathbf{t}) \partial_i^k + u^0(\lambda, \mathbf{t}^1, \mathbf{t}^2) \partial_\lambda \right) \Psi$$

with ‘plus’ coefficients $((u_i^k)_- = u_-^0 = 0)$, the condition $(\hat{U}\Psi)_+ = 0$ (where for Ψ^k , $k \neq 0$, derivatives of series (2) are taken with fixed Ψ^0) implies that $\hat{U}\Psi = 0$.

The statement (4) \Rightarrow (3) directly follows from the relation

Lemma 2

$$\det\left(\frac{\partial}{\partial \tau_i} \Psi_j\right) = \det(V_{\tau_i+}^j) J_0, \quad (5)$$

where τ_i , $0 \leq i \leq N$, is an arbitrary set of times of the hierarchy, and $V_{\tau_i+}^j$ are the coefficients of corresponding vector fields given by the r.h.s. of equations (4),

$$\partial_{\tau_i} \Psi = \sum_{j=0}^N V_{\tau_i+}^j \partial_j \Psi.$$

Similar to [2], it is also possible to prove that Lax-Sato flows of the hierarchy (4) are compatible.

3 Manakov-Santini hierarchy

A simplest case of general hierarchy (4) with $N = 1$ is connected with the system recently introduced by Manakov and Santini [3]. For this hierarchy we have two series

$$\Psi^0 = \lambda + \sum_{n=1}^{\infty} \Psi_n^0(\mathbf{t}^1, \mathbf{t}^2) \lambda^{-n}, \quad (6)$$

$$\Psi^1 = \sum_{n=0}^{\infty} t_n (\Psi^0)^n + \sum_{n=1}^{\infty} \Psi_n^1(\mathbf{t}) (\Psi^0)^{-n} \quad (7)$$

and generating relation

$$(J_0^{-1} d\Psi^0 \wedge d\Psi^1)_- = 0. \quad (8)$$

Lax-Sato equations for Manakov-Santini hierarchy read

$$\partial_n \Psi = \sum_{i=0,1} (J_{1i}^{-1} (\Psi^0)^n)_+ \partial_i \Psi, \quad 0 \leq n \leq \infty, \quad (9)$$

where

$$J = \begin{pmatrix} \Psi_\lambda^0 & \Psi_\lambda^1 \\ \Psi_x^0 & \Psi_x^1 \end{pmatrix}, \quad (10)$$

and

$$\begin{aligned} J_{10}^{-1} &= -J_0^{-1}\Psi_x^0, \quad J_{11}^{-1} = J_0^{-1}\Psi_\lambda^0, \\ J_0 = \det J &= 1 + \partial_x \Psi_1^1 \lambda^{-1} + (\partial_x \Psi_2^1 - \Psi_1^0) \lambda^{-2} + \dots \end{aligned}$$

Lax-Sato equations for the first two flows are given by

$$\begin{aligned} \partial_y \Psi &= ((\lambda - v_x) \partial_x - u_x \partial_\lambda) \Psi, \\ \partial_t \Psi &= ((\lambda^2 - v_x \lambda + u - v_y) \partial_x - (u_x \lambda + u_y) \partial_\lambda) \Psi, \end{aligned}$$

where $u = \Psi_1^0$, $v = \Psi_1^1$. Compatibility condition for these equations represents a system of equations for u , v ,

$$\begin{aligned} u_{xt} &= u_{yy} + (uu_x)_x + v_x u_{xy} - u_{xx} v_y, \\ v_{xt} &= v_{yy} + uv_{xx} + v_x v_{xy} - v_{xx} v_y. \end{aligned} \quad (11)$$

This system was introduced by Manakov and Santini in [3] (see also [4], [5]). For $v = 0$ the system reduces to the dKP equation

$$u_{xt} = u_{yy} + (uu_x)_x. \quad (12)$$

Respectively, $u = 0$ reduction gives an equation [7] (see also [8, 9, 10])

$$v_{xt} = v_{yy} + v_x v_{xy} - v_{xx} v_y. \quad (13)$$

More generally, to reduce Manakov-Santini hierarchy (9) to dispersionless KP hierarchy, one should consider a reduction $J_0 = 1$ (area preservation or Hamiltonian condition), then from Lax-Sato equations (9) one easily gets Lax-Sato equations for dKP hierarchy.

Respectively, reduction $\Psi^0 = \lambda$ leads to the hierarchy connected with equation (13), which was considered in [9, 10].

4 Dunajski equation hierarchy

This hierarchy is connected with the integrable model introduced by Dunajski [6], which generalizes the famous Plebanski second heavenly equation.

It corresponds to general hierarchy with $N = 2$ with volume-preserving reduction ($J_0 = 1$). To define this hierarchy, we introduce three series

$$\begin{aligned}\Psi^0 &= \lambda + \sum_{n=1}^{\infty} \Psi_n^0(\mathbf{t}^1, \mathbf{t}^2) \lambda^{-n}, \\ \Psi^1 &= \sum_{n=0}^{\infty} t_n^1 (\Psi^0)^n + \sum_{n=1}^{\infty} \Psi_n^1(\mathbf{t}^1, \mathbf{t}^2) (\Psi^0)^{-n} \\ \Psi^2 &= \sum_{n=0}^{\infty} t_n^2 (\Psi^0)^n + \sum_{n=1}^{\infty} \Psi_n^2(\mathbf{t}^1, \mathbf{t}^2) (\Psi^0)^{-n}.\end{aligned}$$

Generating equation for Dunajski equation hierarchy is of the form

$$(d\Psi^0 \wedge d\Psi^1 \wedge d\Psi^2)_- = 0 \quad (14)$$

(taking into account the reduction $J_0 = 1$). Lax-Sato equations of the hierarchy read

$$\begin{aligned}\partial_n^1 \Psi = & + \left((\Psi^0)^n \begin{vmatrix} \Psi_\lambda^0 & \Psi_\lambda^2 \\ \Psi_y^0 & \Psi_y^2 \end{vmatrix} \right)_+ \partial_x \Psi - \left((\Psi^0)^n \begin{vmatrix} \Psi_\lambda^0 & \Psi_\lambda^2 \\ \Psi_x^0 & \Psi_x^2 \end{vmatrix} \right)_+ \partial_y \Psi - \\ & \left((\Psi^0)^n \begin{vmatrix} \Psi_x^0 & \Psi_x^2 \\ \Psi_y^0 & \Psi_y^2 \end{vmatrix} \right)_+ \partial_\lambda \Psi, \quad (15)\end{aligned}$$

$$\begin{aligned}\partial_n^2 \Psi = & - \left((\Psi^0)^n \begin{vmatrix} \Psi_\lambda^0 & \Psi_\lambda^1 \\ \Psi_y^0 & \Psi_y^1 \end{vmatrix} \right)_+ \partial_x \Psi + \left((\Psi^0)^n \begin{vmatrix} \Psi_\lambda^0 & \Psi_\lambda^1 \\ \Psi_x^0 & \Psi_x^1 \end{vmatrix} \right)_+ \partial_y \Psi + \\ & \left((\Psi^0)^n \begin{vmatrix} \Psi_x^0 & \Psi_x^1 \\ \Psi_y^0 & \Psi_y^1 \end{vmatrix} \right)_+ \partial_\lambda \Psi \quad (16)\end{aligned}$$

(plus a condition $J_0 = 1$). It is easy to check that for $\Psi^0 = \lambda$ Dunajski equation hierarchy reduces to heavenly equation hierarchy [12, 13], while for $\Psi^2 = y$ it reduces to dispersionless KP hierarchy. The first two flows of the hierarchy (15), (16) read

$$\partial_1^1 \Psi = (\lambda \partial_x - v_x \partial_x - u_x \partial_y - f_x \partial_\lambda) \Psi, \quad (17)$$

$$\partial_1^2 \Psi = (\lambda \partial_y - v_y \partial_x - u_y \partial_y - f_y \partial_\lambda) \Psi, \quad (18)$$

here

$$u = \Psi_1^2, \quad v = \Psi_1^1, \quad f = \Psi_1^0,$$

condition $J_0 = 1$ implies that $u_y + v_x = 0$, and we can introduce a potential Θ , $v = \Theta_y$, $u = -\Theta_x$. After the identification $z = -t_1^1$, $w = t_1^2$, we get Lax pair for Dunajski system

$$\begin{aligned}\Theta_{wx} + \Theta_{zy} + \Theta_{xx}\Theta_{yy} - \Theta_{xy}^2 &= f, \\ f_{xw} + f_{yz} + \Theta_{yy}f_{xx} + \Theta_{xx}f_{yy} - 2\Theta_{xy}f_{xy} &= 0.\end{aligned}$$

The second flows can be written in the form

$$\begin{aligned}\partial_2^1 \Psi &= (\lambda \partial_1^1 + f \partial_x - (\partial_1^1 v) \partial_x - (\partial_1^1 u) \partial_y - (\partial_1^1 f) \partial_\lambda) \Psi, \\ \partial_2^2 \Psi &= (\lambda \partial_1^2 + f \partial_y - (\partial_1^2 v) \partial_x - (\partial_1^2 u) \partial_y - (\partial_1^2 f) \partial_\lambda) \Psi.\end{aligned}$$

To write down vector fields more explicitly, one should use first flows (17), (18). Commutation relations for any pair of the flows give an equation for Θ , f (with different set of times).

5 Dressing scheme

Let us consider nonlinear vector Riemann problem of the form

$$\Psi_+ = \mathbf{F}(\Psi_-), \quad (19)$$

where Ψ_+ , Ψ_- denote the boundary values of the $(N+1)$ -component vector function on the sides of some oriented curve γ in the complex plane of the variable λ . The problem is to find the function analytic outside the curve with some fixed behavior at infinity (normalization) which satisfies (19). The problem of this type was used by Takasaki [11, 12, 13], who stressed its connection to Penrose nonlinear graviton construction.

The problem (19) is connected with a class of integrable equations, which can be represented as a commutation relation for vector fields containing a derivative on the spectral variable. We give a sketch of the dressing scheme corresponding to the general $(N+2)$ -dimensional integrable hierarchy (3), (4). First, we chose γ as a unit circle. To get solutions of the general hierarchy using the problem (19), one should find solution to this problem with the singularities at $\lambda = \infty$ defined by the series (1), (2) (for simplicity we suggest that only a finite number of times is not equal to zero). If the determinant J_0 for this solution is not equal to zero, we come to the conclusion that the form

$$\Omega = J_0^{-1} d\Psi^0 \wedge d\Psi^1 \wedge \dots \wedge d\Psi^N$$

is analytic in the complex plane, and so it satisfies the generating identity (3). Then the solution of the Riemann problem (19) satisfies Lax-Sato

equations (4) and provides a solution for the general $(N+2)$ -dimensional hierarchy.

Due to the fact that singularities of solution of the Riemann problem are defined by the series (1), (2) using the solution itself, solution of the hierarchy is usually defined in terms of implicit functions.

Acknowledgments

The author was partially supported by Russian Foundation for Basic Research under grants no. 08-01-90104, 07-01-00446 and 06-01-92053.

References

- [1] L. V. Bogdanov, V. S. Dryuma and S. V. Manakov, On the dressing method for Dunajski anti-self-duality equation, *nlin/0612046* (2006).
- [2] L. V. Bogdanov, V. S. Dryuma and S. V. Manakov, Dunajski generalization of the second heavenly equation: dressing method and the hierarchy, *J Phys. A: Math. Theor.* **40** (2007) 14383–14393.
- [3] S. V. Manakov and P. M. Santini, The Cauchy problem on the plane for the dispersionless Kadomtsev-Petviashvili equation, *JETP Lett.* **83** (2006) 462-6.
- [4] S. V. Manakov and P. M. Santini, A hierarchy of integrable PDEs in $2+1$ dimensions associated with 2-dimensional vector fields, *Theor. Math. Phys.* **152** (2007) 1004–1011.
- [5] S. V. Manakov and P. M. Santini, On the solutions of the dKP equation: the nonlinear, Riemann-Hilbert problem, longtime behaviour, implicit solutions and wave breaking, *J Phys. A: Math. Theor.* **41** (2008) 055204.
- [6] M. Dunajski, *Proc. Roy. Soc. Lond. A* **458**, 1205 (2002)
- [7] M.V. Pavlov, Integrable hydrodynamic chains, *J. Math. Phys.* **44**(9) (2003) 4134–4156
- [8] M. Dunajski, A class of Einstein-Weyl spaces associated to an integrable system of hydrodynamic type, *J. Geom. Phys.* **51** (2004) 126–137.
- [9] L. Martínez Alonso and A. B. Shabat, Energy-dependent potentials revisited: a universal hierarchy of hydrodynamic type, *Phys. Lett. A* **300** (2002) 58–64.

- [10] L. Martínez Alonso and A. B. Shabat, Hydrodynamic reductions and solutions of a universal hierarchy, *Theor. Math. Phys.* **140** (2004) 1073–1085
- [11] K. Takasaki, *Phys. Lett. B* **285**, 187–190 (1992)
- [12] K. Takasaki, *J. Math. Phys.* **30**(7), 1515–1521 (1989)
- [13] K. Takasaki, *J. Math. Phys.* **31**(8), 1877–1888 (1989)