

Connecting spin and statistics in quantum mechanics

Arthur Jabs

Alumnus, Technische Universität Berlin
Voßstr. 9, 10117 Berlin, Germany
arthur.jabs@alumni.tu-berlin.de

(14 October 2008)

Abstract. The spin-statistics connection is derived in a simple manner under the postulates that the original and the exchange wave functions are simply added, and that the azimuthal phase angle, which defines the orientation of the spin part of each single-particle spin-component eigenfunction in the plane normal to the spin-quantization axis, is exchanged along with the other parameters. The spin factor $(-1)^{2s}$ belongs to the exchange wave function when this function is constructed so as to get the spinor ambiguity under control. This is achieved by effecting the exchange of the azimuthal angle by means of rotations and admitting only rotations in one sense. The procedure works in Galilean as well as in Lorentz-invariant quantum mechanics. Relativistic quantum field theory is not required.

Keywords: spin, statistics, spinor

PACS: 03.65.Ta

Physics is simple but subtle
Paul Ehrenfest

1 Introduction

The standard method of treating systems of identical particles in quantum mechanics is to require that every wave function or state vector is either symmetric or antisymmetric, that is, multiplied by either $+1$ or -1 when the labels or parameters referring to any two particles are interchanged. There are thus two classes of systems, with different collective behaviour of the particles: systems of bosons and systems of fermions. These two classes are connected with the spins of the particles: all particles which are known to be bosons are empirically found to have integral spin, in units of \hbar , while all known fermions have half-integral (i.e., half-odd-integral) spin.

Within quantum mechanics the connection with spin could not be derived and had to be taken as another postulate. The first derivation was provided by Fierz [1] and Pauli [2], who founded it on relativistic quantum field theory. This also remained the framework for the papers which in subsequent years refined and generalized Pauli's proof [3, 4]. Typically, in these papers it is postulated that no negative-

energy states exist, that the metric in Hilbert space is positive definite, and that the fields either commute or anticommute for spacelike separations (locality, micro-causality). Under these conditions it is shown that integral-spin fields cannot satisfy the (fermionic) anticommutation relations, and half-integral-spin fields cannot satisfy the (bosonic) commutation relations. This does not exclude the possibility that fields exist which satisfy other commutation relations and show statistics that differ from Bose and Fermi statistics.

In 1965 Feynman in his Lectures [5, p. 4-3] objected:

An explanation has been worked out by Pauli from complicated arguments of quantum field theory and relativity. He has shown that the two [spin and statistics] must necessarily go together, but we have not been able to find a way of reproducing his arguments on an elementary level. It appears to be one of the few places in physics where there is a rule which can be stated very simply, but for which no one has found a simple and easy explanation. The explanation is deep down in relativistic quantum mechanics. This probably means that we do not have a complete understanding of the fundamental principle involved.

The aim of the present paper is to propose such a simple and easy explanation.

Actually, since 1965 more than a hundred publications appeared deriving the spin-statistics connection under different sets of conditions [6]. Reviews are contained in [7 - 10]. Many of these publications derive the connection in settings far removed from standard (local) relativistic quantum field theory; and they are also far from simple and easy.

Closest to the present approach are those papers that use only quantum mechanics, relativistic or nonrelativistic, and are written in the spirit of Feynman's demand for simplicity. These papers nevertheless contain one or several of the following restrictions: the wave functions must have special invariance [11], continuity [12] or symmetry [13] properties, or must lie in special spin-component subspaces [14]. The systems considered must be nonrelativistic [12 - 16], have only two spatial dimensions [16], contain only two particles [13, 14], only particles with zero spin [12] or spin $\leq 1/2$ [15], only point particles [17], must admit antiparticles [18], or the exchange must be considered as physical transportation of real objects [11, 17, 19].

The present proposal is not subject to any of these restrictions. It grew out of an attempt to understand the papers by York [20] in the framework of the realist interpretation which I developed some time ago [21]. The premises of the present proposal are seen when the organization of the paper is considered:

In Section 2 we start with Feynman's method of superposing transition amplitudes. This is shown to be equivalent to symmetrizing or antisymmetrizing only the final but not the initial wave function in the transition amplitudes. Moreover, in line with another proposal by Feynman, symmetrizing or antisymmetrizing the final wave function, which for two particles means addition or subtraction of the original and the exchange wave function, is replaced by the "addition postulate", which admits only addition.

The minus sign in the superposition of fermionic wave functions arises from the construction of the exchange function. In Section 3 we point out that it is important to consider the azimuthal spin angle χ , which defines the orientation of the spin part of a single-particle spin-component eigenfunction. The angle χ is also exchanged, but requires a special treatment because it expresses the well known spinor ambiguity when the spin component m is half-integral. This means that we cannot know which of the two possible values of the function with the exchanged χ has to be chosen.

In Section 4 it is shown that the ambiguity can be overcome by effecting the exchange of χ by way of rotations and by admitting only rotations in one direction, either clockwise or counterclockwise. In Section 5 the rotations leading from the original to the exchange function are explicitly carried out, and it is shown that the exchange function thereby acquires the desired spin factor $(-1)^{2s}$. It is thus the rotation group, a subgroup of both the Galilei and the Lorentz group, that determines the type of statistics. In standard quantum mechanics it is the permutation group that does this: its one-dimensional representations are associated with Bose and Fermi statistics, and its other representations with “parastatistics”. The rotation group in our approach leads only to Bose and Fermi statistics, and there is no reason to suggest experiments in search of particles with parastatistics.

In Section 6 the proof is extended to N particles, and in Section 7 to particles with different spin components. Finally, in Section 8, using the properties of helicity functions, it is pointed out that the proof also holds in relativistic quantum mechanics.

2 Adding up transition amplitudes

It is remarkable that wave-function symmetrization (or antisymmetrization) is never mentioned in the Feynman Lectures on Physics [5] and yet the same physical situations as in the other textbooks, which do use symmetrization, are accounted for. Feynman basically considers *transition amplitudes*: when two transitions cannot be distinguished in principle from each other, the amplitudes, rather than the probabilities, have to be superposed [5, pp. 1-10, 3-7...4-3]. The superposition includes the phase factor $\exp(i\delta)$

$$f = f(\theta) + e^{i\delta} f(\pi - \theta) \tag{1}$$

in Feynman’s notation. In line with standard quantum mechanics the phase factor in the Lectures is eventually postulated to be +1 or -1.

What does this mean in terms of (Schrödinger) wave functions? The transition amplitude for the transition from $\Psi_a(1, 2)$ to $\Psi_b(1, 2)$ is given by the scalar product

$$f = (\Psi_b(1, 2), \Psi_a(1, 2)). \tag{2}$$

These scalar products are basic elements of quantum mechanics because any expression of physical significance, that is, an expression which yields the probability of a result of a measurement, is formulated in terms of absolute squares of transition amplitudes

$$|f|^2 = |(\Psi_b, \Psi_a)|^2. \tag{3}$$

Now, in systems of identical particles the transition from $\Psi_a(1, 2)$ to $\Psi_b(1, 2)$ cannot be distinguished in principle from a transition from $\Psi_a(1, 2)$ to $\Psi_b(2, 1)$, where $\Psi_b(2, 1)$ denotes the exchange function, in which the labels or parameters referring to particles 1 and 2 have been exchanged. Thus, in terms of wave functions Eq. (1) takes on the form

$$f = \left(\Psi_b(1, 2), \Psi_a(1, 2) \right) + e^{i\delta} \left(\Psi_b(2, 1), \Psi_a(1, 2) \right) \quad (4)$$

or

$$f = \left(\Psi_b(1, 2) \pm \Psi_b(2, 1), \Psi_a(1, 2) \right). \quad (5)$$

Here only one of the two functions in the scalar product is a superposition of the original and the exchange function, and no normalization factors appear. This is the form that is exclusively used by Feynman.

This is to be compared with the standard method in quantum mechanics, where the amplitude in the above case is written as

$$f = \left(\frac{1}{\sqrt{2}} [\Psi_b(1, 2) \pm \Psi_b(2, 1)], \frac{1}{\sqrt{2}} [\Psi_a(1, 2) \pm \Psi_a(2, 1)] \right). \quad (6)$$

Here both functions in the scalar product are superpositions of an original and an exchange function, and both functions are normalized, by means of the factors $1/\sqrt{2}$.

Feynman's method (5) and the standard method (6) are mathematically equivalent provided the phase factor $\exp(i\delta)$ is restricted from the outset to the values ± 1 (see Appendix). For more general factors this equivalence does not hold. Such factors appear however during our treatment when it comes to one-particle wave functions with different spin components, in Section 7. We thus have to make a choice. It is Feynman's method that we choose because in contrast to the standard method the Feynman method allows us to incorporate an explanation of the spin-statistics connection and yet finally to arrive at the approved expressions of physical significance. Physically, Feynman's method recommends itself because it is really only necessary to consider physically significant expressions, and the scalar products are closer to these ($|f|^2$) than are the wave functions (cf. e.g. [22, 23]). In any case it appears reasonable to number the particles in the first function arbitrarily and, assuming that the numbering gets lost in the transition, to add up all possibilities of numbering in the second function.

In his own attempt to derive the spin-statistics connection Feynman [19] moreover suggests that we may

take the view that the Bose rule is obvious from some kind of understanding that the amplitude[s] in quantum mechanics that correspond to alternatives must be added.

That is, he is proposing to start a priori with $\exp(i\delta) = +1$ everywhere. We follow also this proposal by Feynman and just add up the transition amplitudes. That is, we start from

$$f = \left(\Psi_b(1, 2) + \Psi_b(2, 1), \Psi_a(1, 2) \right)$$

in the case of two particles. In the case of N particles this obviously generalizes to

$$f = \left(\sum_{\alpha} P_{\alpha} \Psi_{\mathbf{b}}(1, 2, \dots, N), \Psi_{\mathbf{a}}(1, 2, \dots, N) \right), \quad (7)$$

where P_{α} is a permutation of the parameters referring to the single particles. The sum extends over the $N!$ possible permutations, including the identity I . The sum $\sum_{\alpha} P_{\alpha} \Psi_{\mathbf{b}} = \Psi_{\mathbf{bS}'}$ is the extension of the sum of the original and the exchange function from two-particle to N -particle systems and corresponds to the symmetrized function in standard quantum mechanics.

3 The special parameter

In order to present the essential points in a simple way we begin by considering a non-relativistic (Galilean) system of two identical particles of spin s ($\mathbf{S}^2 \psi = s(s+1) \psi$) described by a (Schrödinger) wave function which is a product of two normalized one-particle wave functions

$$\Psi = \psi^{(1)}(a, m) \psi^{(2)}(b, m),$$

and the one-particle wave functions are eigenfunctions of the operator of the spin component with respect to an arbitrary but common spin-quantization axis. Moreover, both functions belong to the same eigenvalue m . These restrictions will be removed in Sections 6 to 8.

The single-particle wave functions are functions of the variables x, y, z, t . The mathematical form of the functions is determined by the parameters a, b and m , where a and b stand for *all* parameters that, in addition to m (and χ , below), determine the form, such as mass, charge, total spin, expansion coefficients etc. Charge, mass and total spin are of course the same for identical particles and their exchange has no effect. An alternative notation would be $\psi(a, m, x^{(1)}, y^{(1)}, z^{(1)}, t)$, where the labels in parentheses, (1) and (2), distinguish the particles in the formalism. We have suppressed here the variables and have put the particle labels directly at the function symbols. In Sections 2 to 6 the eigenvalue m is always the same and will also be omitted from the notation. Thus we write

$$\Psi = \psi^{(1)}(a) \psi^{(2)}(b). \quad (8)$$

Among the parameters of the wave function there is one that requires special treatment in the construction of the exchange function. This is the azimuthal angle χ , which defines the orientation of the spin part of each single-particle spin-component eigenfunction in the plane normal to the common spin-quantization axis, counted from some arbitrary reference direction. Therefore we keep this angle out of the set a (and b) and exhibit it explicitly. Each function has its own angle, but the particular values do not matter. The values of χ are restricted to the interval $[0, 2\pi]$.

The specific form of the parametric dependence of the spin-component eigenfunction on χ is given by the factor

$$\exp(im\chi), \quad (9)$$

$$\psi^{(1)}(a, \chi_a) = \exp(im\chi) \psi^{(1)}(a), \quad (10)$$

and the exponential factor expresses the spinor ambiguity: in the case of half-integral m it is $+1$ for $\chi = 0$ and -1 for $\chi = 2\pi$. This is standard quantum mechanics. The angle χ appears “only” in a phase factor, and this factor is usually considered as an overall (global) phase factor without any physical significance. Indeed, every wave function, even after normalization, is only defined to within an arbitrary phase factor which is independent of \mathbf{r} and t , and usually the same experimental predictions result with or without this factor [24, pp. 132, 398].

However, in the present approach the angle χ in the phase factor (9) is given special attention and is exchanged, though in a specific way, along with the other parameters. In this procedure it does have physical significance, for it is then instrumental in determining the *relative* phase when, in Section 5, wave functions with different χ s will be superposed.

The letter χ , rather than the customary φ , is used in order to emphasize that it is the spin part, not the orbital part, which is concerned, that the spin-quantization axis need not coincide with the z -axis, and that the angle χ is not a variable of the one-particle wave function, as \mathbf{r} or φ are. Rather, the dependence on χ is a parametric dependence, like that on m and the other parameters in a and b . Therefore the application of a differential operator like $-i\hbar\partial/\partial\chi$, analogous to the z -component of *orbital* angular momentum, does not make sense for the spin part of the wave function.

Thus, with the explicit appearance of χ Eq. (8) becomes

$$\Psi = \psi^{(1)}(a, \chi_a) \psi^{(2)}(b, \chi_b). \quad (11)$$

In order to obtain the exchange wave function we simply replace a by b and vice versa in the original wave function (11). But because of the spinor ambiguity the exchange of χ_a with χ_b and vice versa cannot be done in such a simple way.

4 Controlling the spinor ambiguity

The special feature with the factor (9) is that χ is an angle, so that we may go from some particular value χ_a to some other value χ_b in two ways, either clockwise or counterclockwise. In the case of half-integral m one way leads to a different wave function at χ_b than the other, the two functions having different signs. In other words, the value of the function at χ_b then depends not only on the value of χ_b but also on the path leading from χ_a to χ_b . This leads to double-valued functions and represents another aspect of the spinor ambiguity.

One may imagine the function $\exp(im\chi)$ with half-integral m to lie on the two-sheeted Riemannian surface of the function \sqrt{z} [25], where one sheet carries only one set of function values. The clockwise path from χ_a to χ_b always ends up in a different sheet than the counterclockwise path. Or one may imagine a Möbius band, where on the first round trip over the band one set of function values is met, and the corresponding other set on the second round trip. In fact, devices like twisted ribbon belts [19, p. 58], contortions of an arm holding a cup [19, p. 30] and others [26] are

similar to the Riemannian surface and the Möbius band in that they construct an indicator of whether we are in the first or in the second turn, and in that they return to the original situation after the second turn.

Now, when adding the original and exchange wave functions the functions must be uniquely defined. This is not the same as the general requirement that wave functions be single-valued. Single-valuedness can only be required for measurable quantities such as transition probabilities or expectation values, but not for the wave functions themselves [27]. In many textbooks it is nevertheless invoked for the wave functions themselves, in particular for justifying integral values of m for orbital angular momentum. The real justification of integral m here rests on group representations and properties of observables [28].

Our case is different because we are concerned with the procedure of constructing one wave function by superposition of others, formally similar to interference. The demand for removing the spinor ambiguity is in line with the demand for removing the ambiguity known as exchange degeneracy, that is, to the fixation of the coefficients of the superposed terms in Eqs. (1) to (7).

Now, according to what has been said above the spinor ambiguity is removed (i.e., kept under control) if we make a choice between the two possible paths from χ_a to χ_b , that is, if we exchange the χ s by way of rotations and decide to make all rotations in one sense only, either clockwise or counterclockwise.

In the language of group theory the clockwise and the counterclockwise way from χ_a to χ_b correspond to paths of different homotopy classes (e.g. [29]). So our choice means that we are admitting only paths of the same homotopy class.

5 Constructing the exchange function

We are now ready to take the decisive step. We want to construct the exchange function from the original function (11), not by simply replacing χ_a by χ_b in the wave function $\psi^{(1)}$ and χ_b by χ_a in the wave function $\psi^{(2)}$ (as is done with the other parameters, a and b), but by continuously rotating the spin part of the functions from χ_a to χ_b and from χ_b to χ_a respectively, with due consideration being given to the paths connecting χ_a and χ_b .

Thus, we start from formula (11) where the a s and b s have already been exchanged, but the χ s have not:

$$\psi^{(1)}(b, \chi_a) \psi^{(2)}(a, \chi_b). \quad (12)$$

We then rotate the function $\psi^{(1)}(b, \chi_a)$ from χ_a to χ_b . We take the counterclockwise sense of the rotations, and we assume $\chi_a < \chi_b$ and $m \geq 0$. In order to get from χ_a to χ_b we then have to run through $\chi_b - \chi_a$. This yields the rotation factor $\exp(im(\chi_b - \chi_a))$ and we obtain

$$\psi^{(1)}(b, \chi_b) = e^{im(\chi_b - \chi_a)} \psi^{(1)}(b, \chi_a). \quad (13)$$

Likewise, rotating the function $\psi^{(2)}(a, \chi_b)$ counterclockwise from χ_b to χ_a means that we have to run through $2\pi - (\chi_b - \chi_a)$. This yields the rotation factor

$\exp(im(2\pi + \chi_a - \chi_b))$ and we obtain

$$\psi^{(2)}(a, \chi_a) = e^{im(2\pi + \chi_a - \chi_b)} \psi^{(2)}(a, \chi_b). \quad (14)$$

Inserting (13) and (14) into (12) then yields the exchange function

$$F \times \psi^{(1)}(b, \chi_b) \psi^{(2)}(a, \chi_a) \quad (15)$$

with

$$F = e^{-im(\chi_b - \chi_a)} e^{-im(2\pi + \chi_a - \chi_b)} = e^{-im2\pi} = (-1)^{2m} = (-1)^{2s}, \quad (16)$$

where for the last equality we have used the fact that s and m are either both integral or both half-integral. Had we chosen the clockwise sense we would have obtained $F = \exp(+im2\pi)$, which is also equal to $(-1)^{2s}$. The same result obtains with the alternative assumption $\chi_a > \chi_b$. The case $\chi_a = \chi_b$ is of statistical weight zero and can be neglected.

In the case of negative m the rotation factor (9) is $\exp(-i|m|\chi)$ and means a rotation in the clockwise rather than in the counterclockwise sense when χ increases. We can, however, reduce this case to that of positive m and counterclockwise rotations: the rotation factor is the same if we consider the minus sign in the exponent to be connected with χ , and $|m|$ to be replaced by m , with $m \geq 0$. We then have negative angles $-|\chi|$. But as we have confined the azimuthal spin angles within $[0, 2\pi]$ we have to replace $-|\chi|$ by $\chi' = -|\chi| + 2\pi$ denoting the same point on the circle, but being positive. Thereby the rotation factor becomes $\exp(im\chi')$, both m and χ' are positive, and the rotations are counterclockwise when χ' increases. As within $[0, 2\pi]$ the values of the angles are arbitrary the cases with negative m are thus reduced to those with positive m . Indeed it is easy to verify that F as given by Eq. (16) is the same whether calculated with χ or with χ' in the single-particle functions.

Finally, adding the original function (11) and the exchange function (15) we arrive at

$$\Psi_S = \psi^{(1)}(a, \chi_a) \psi^{(2)}(b, \chi_b) + (-1)^{2s} \psi^{(1)}(b, \chi_b) \psi^{(2)}(a, \chi_a). \quad (17)$$

Ψ_S need not be normalized in Feynman's method, as emphasized after formula (5). The angle χ and the rotations become effective only in the procedure of exchanging the parameters of the wave functions. In this procedure the angles χ_a and χ_b in the original and the exchange wave function are related in such a way that, although χ_a and χ_b may be randomly distributed in the original function, the resultant factor, $(-1)^{2s}$, is independent of χ_a and χ_b . Thus, once exchange and addition are accomplished, we may, according to formula (9), write the functions in Eq. (17) in the form $\psi^{(1)}(a, \chi_a) = \exp(im\chi_a) \psi^{(1)}(a)$ etc. Either term in Eq. (17) thereby receives the same factor

$$\exp(im[\chi_a + \chi_b]), \quad (18)$$

which can be put as an overall phase factor in front of the parentheses. There it can be omitted, that is, absorbed in the general arbitrary phase factor connected with every wave function.

Thus we are returning to the standard form of the wave functions, which do not exhibit the dependence on χ :

$$\Psi_S = \psi^{(1)}(a) \psi^{(2)}(b) + (-1)^{2s} \psi^{(1)}(b) \psi^{(2)}(a) \quad (19)$$

There is some formal analogy with interference between two parts of a split wave. One part is left unmodified [wave function (11)], the other is subject to a phase shift [exchange, wave function (15)], and then the two are recombined [wave function (17) or (19)].

With formula (19) we have reached our goal for the considered class of functions: we have derived the factor $(-1)^{2s}$ in a simple way from basic principles. This factor yields +1 (bosons) for integral s and -1 (fermions) for half-integral s , and this is the desired connection between spin and statistics.

6 General case. Equal spin components

We begin now to remove the restrictions imposed on the wave function in the previous sections. In the present section we remove the restriction to two particles and to functions of product form. We begin with N -particle functions of product form

$$\Psi_b = \psi^{(1)}(u_{r_1}, \chi_{t_1}) \psi^{(2)}(u_{r_2}, \chi_{t_2}) \cdots \psi^{(N)}(u_{r_N}, \chi_{t_N})$$

with u_{r_1}, u_{r_2}, \dots instead of a, b . The $\psi^{(i)}(u_{r_i}, \chi_{t_i})$ all belong still to the same m , which is therefore dropped from the notation. The symmetrized function is

$$\Psi_{bS'} = \sum_{\alpha} P_{\alpha} \psi^{(1)}(u_{r_1}, \chi_{t_1}) \cdots \psi^{(N)}(u_{r_N}, \chi_{t_N}). \quad (20)$$

The index P' (with prime) is to indicate that P_{α} in (20) permutes the parameter sets $\{u_{r_i}\}$ among the one-particle functions but does not permute the angles χ_{t_i} . The permutation of the angles will be effected separately, by way of rotations. As any permutation can be written as a product of a number of transpositions (interchanges), the term $P_{\alpha} \psi^{(1)}(u_{r_1}, \chi_{t_1}) \cdots \psi^{(N)}(u_{r_N}, \chi_{t_N})$ differs from the term with $P_{\alpha} = I$ by a number k_{α} of transpositions. When the χ rotations are applied, as described in the preceding sections for the case of two particles, every single transposition yields the factor $F = (-1)^{2s}$ in front of the term with interchanged parameters. The angles χ may be different in every such pair, but since the angles are arbitrary and all spin components are equal this does not matter, and every transposition yields the same factor. Hence k_{α} transpositions yield the factor $(-1)^{2sk_{\alpha}}$. The function (20) then changes into the superposition function (symmetric or antisymmetric)

$$\Psi_{bS} = \sum_{\alpha} (-1)^{2sk_{\alpha}} P_{\alpha} \psi^{(1)}(u_{r_1}, \chi_{t_1}) \cdots \psi^{(N)}(u_{r_N}, \chi_{t_N}). \quad (21)$$

The index S (without prime) instead of S' (with prime, as in (20)) is to indicate that the exchange of the angles χ_{t_i} by means of rotation is included. In Eq. (21) P_{α} therefore permutes the pairs $\{u_{r_i}, \chi_{t_i}\}$. The single functions $(-1)^{2sk_{\alpha}} P_{\alpha} \psi^{(1)}(u_{r_1}, \chi_{t_1}) \cdots$

$\psi^{(N)}(u_{r_N}, \chi_{t_N})$ for $P_\alpha \neq I$ are the extensions of the exchange function from two to N particles.

As in the case of two particles, the one-particle functions in (21) may now be written as $\psi^{(i)}(u_{r_i}, \chi_{t_i}) = \exp(im\chi_{t_i}) \psi^{(i)}(u_{r_i})$. This yields the same factor

$$\exp\left(im \sum_{k=1}^N \chi_{t_k}\right)$$

in front of every permutation operator P_α in (21). This factor can thus be drawn out of the permutation sum and becomes again an overall phase factor, which can be omitted. We can thus drop the χ_{t_i} s from the notation in (21) and simply write

$$\Psi_{\text{bS}} = \sum_{\alpha} (-1)^{2sk_\alpha} P_\alpha \psi^{(1)}(u_{r_1}) \cdots \psi^{(N)}(u_{r_N}). \quad (22)$$

Now, if s is an integer, then $(-1)^{2sk_\alpha} = +1$ for any k_α , and Ψ_{bS} is symmetric (bosonic). If s is a half-integer, then $(-1)^{2sk_\alpha} = -1$ for odd k_α , and $+1$ for even k_α , and Ψ_{bS} is antisymmetric (fermionic). And this holds for each one of the $2s+1$ values of m .

In the next step we remove the restriction to wave functions of product form, but still with equal m s. The general N -particle function is

$$\Phi_{\text{b}} = \sum_{r_1, \dots, r_N, t_1, \dots, t_N} a_{r_1 \dots r_N t_1 \dots t_N} \psi^{(1)}(u_{r_1}, \chi_{t_1}) \cdots \psi^{(N)}(u_{r_N}, \chi_{t_N}), \quad (23)$$

where the sum over the r s and t s goes over a possibly infinite number of values. Permuting the parameter sets $\{u_{r_i}\}$ among the one-particle functions and permuting the angles by way of rotations results in

$$\Phi_{\text{bS}} = \sum_{r_1, \dots, r_N, t_1, \dots, t_N} b_{r_1 \dots r_N t_1 \dots t_N} \sum_{\alpha} (-1)^{2sk_\alpha} P_\alpha \psi^{(1)}(u_{r_1}, \chi_{t_1}) \cdots \psi^{(N)}(u_{r_N}, \chi_{t_N}). \quad (24)$$

The sums \sum_{α} may be considered as basis functions in the subspace of wave functions with equal m s. They are symmetric or antisymmetric. The antisymmetric ones are special Slater determinants, on which the Pauli exclusion principle is based.

From the consideration in the preceding section (cf. (18)) it follows that the permutation sum \sum_{α} in (24) can be written as

$$\exp(im[\chi_{t_1} + \cdots + \chi_{t_N}]) \sum_{\alpha} (-1)^{2sk_\alpha} P_\alpha \psi^{(1)}(u_{r_1}) \cdots \psi^{(N)}(u_{r_N}). \quad (25)$$

So

$$\Phi_{\text{bS}} = \sum_{r_1, \dots, r_N, t_1, \dots, t_N} c_{r_1 \dots r_N t_1 \dots t_N} \sum_{\alpha} (-1)^{2sk_\alpha} P_\alpha \psi^{(1)}(u_{r_1}) \cdots \psi^{(N)}(u_{r_N}), \quad (26)$$

where the coefficients $c_{r_1 \dots t_N}$ in (26) are thought to have absorbed the exponentials in (25).

The coefficients include the angles χ_{t_i} . This is no problem because any superposition with definite coefficients means that the arbitrary phase factors, which

accompany every wave function, have been fixed. This includes that the angles χ_{t_i} , which are arbitrary anyway, are fixed when they are part of those arbitrary phases. With this they produce no longer any effect. Note also that the sum over the r s and t s in (26), in contrast to the sum over α , has nothing to do with the identity of the particles in the final state.

The connection between spin and statistics is thus proved for general nonrelativistic N -particle functions composed of one-particle functions with the same spin component.

7 Different spin components

When admitting one-particle wave functions which belong to different spin components m , the factor F in Eq. (16) of Section 5 has to be replaced by a different factor, F_χ . This factor is obtained when the transposition procedure of Section 5 between Eqs. (12) and (16) is repeated with Eq. (12) being replaced by $\psi^{(1)}(b, m_b, \chi_a)$ $\psi^{(2)}(a, m_a, \chi_b)$. The result is

$$F_\chi = (-1)^{2s} \exp(-i(m_a - m_b)(\chi_a - \chi_b)). \quad (27)$$

As any permutation P_α can be written as a product of a number k_α of transpositions, the factor $(-1)^{2sk_\alpha}$ in Eqs. (21) to (26) is then replaced by η_α , which is a product of F_χ s:

$$\eta_\alpha = (-1)^{2sk_\alpha} \prod^{(\alpha)} \exp(-i(m_a - m_b)(\chi_a - \chi_b)(1 - \delta_{k_\alpha 0})). \quad (28)$$

The values that the parameters m_a, χ_a, m_b, χ_b assume in the various factors of the product are those of the particular one-particle functions on which the transpositions in P_α operate. The factor $(1 - \delta_{k_\alpha 0})$ with the Kronecker delta is to make the product 1 when $k_\alpha = 0$, that is, when $P_\alpha = I$. As we shall see, the exact form of η_α does not matter. What matters is first that in the case of equal m s the factor F_χ reduces to F , and η_α to $(-1)^{2sk_\alpha}$, so that we are back to Sections 2 to 6, and second that in any case $|\eta_\alpha| = 1$. What we have to show is that even in the case of different m s the factor η_α is effectively $(-1)^{2sk_\alpha}$ and the angles χ will disappear from the final expressions of physical significance.

The general N -particle wave functions now are (cf. (23))

$$\begin{aligned} \Phi_a(1, \dots, N) &= \\ &= \sum_{\substack{r_1, \dots, r_N \\ s_1, \dots, s_N \\ t_1, \dots, t_N}} a_{r_1 \dots r_N s_1 \dots s_N t_1 \dots t_N} \psi^{(1)}(u_{r_1}, m_{s_1}, \chi_{t_1}) \dots \psi^{(N)}(u_{r_N}, m_{s_N}, \chi_{t_N}) \end{aligned} \quad (29)$$

$$\begin{aligned} \Phi_b(1, \dots, N) &= \\ &= \sum_{\substack{\rho_1, \dots, \rho_N \\ \sigma_1, \dots, \sigma_N \\ \tau_1, \dots, \tau_N}} b_{\rho_1 \dots \rho_N \sigma_1 \dots \sigma_N \tau_1 \dots \tau_N} \psi^{(1)}(u_{\rho_1}, m_{\sigma_1}, \chi_{\tau_1}) \dots \psi^{(N)}(u_{\rho_N}, m_{\sigma_N}, \chi_{\tau_N}), \end{aligned} \quad (30)$$

where the sum over the s_i s and σ_i s goes over the $2s + 1$ possible values of the spin component.

$$\begin{aligned}\Phi_{\text{bS}} &= \sum_{\alpha} \eta_{\alpha} P_{\alpha} \Phi_{\text{b}}(1, \dots, N) = \\ &= \sum_{\rho_1, \dots, \tau_N} b_{\rho_1 \dots \tau_N} \sum_{\alpha} \eta_{\alpha} P_{\alpha} \psi^{(1)}(u_{\rho_1}, m_{\sigma_1}, \chi_{\tau_1}) \cdots \psi^{(N)}(u_{\rho_N}, m_{\sigma_N}, \chi_{\tau_N}).\end{aligned}\quad (31)$$

With a view to an expression of physical significance we consider the transition amplitude

$$f = (\Phi_{\text{bS}}, \Phi_{\text{a}}) = \sum b_{\rho_1 \dots \tau_N}^* \sum a_{r_1 \dots t_N} T_{(rst\rho\sigma\tau)},\quad (32)$$

where

$$\begin{aligned}T_{(rst\rho\sigma\tau)} &= \sum_{\alpha} \eta_{\alpha}^* P_{\alpha} \left(\psi^{(1)}(u_{\rho_1}, m_{\sigma_1}, \chi_{\tau_1}), \psi^{(1)}(u_{r_1}, m_{s_1}, \chi_{t_1}) \right) \cdots \\ &\quad \cdots \left(\psi^{(N)}(u_{\rho_N}, m_{\sigma_N}, \chi_{\tau_N}), \psi^{(N)}(u_{r_N}, m_{s_N}, \chi_{t_N}) \right),\end{aligned}\quad (33)$$

and the P_{α} permute only the sets $\{\rho_i, \sigma_i, \chi_i\}$, not the sets $\{r_i, s_i, t_i\}$, among the one-particle functions. The point is that the one-particle functions with different spin components are mutually orthogonal, irrespective of the u s and χ s. Thus, the scalar products between these functions are all proportional to Kronecker deltas, and the terms (33) become

$$T_{(rst\rho\sigma\tau)} = \sum_{\alpha} \eta_{\alpha}^* P_{\alpha} \kappa_{\sigma_1 s_1} \cdots \kappa_{\sigma_N s_N} \delta_{\sigma_1 s_1} \cdots \delta_{\sigma_N s_N},\quad (34)$$

where

$$\kappa_{\sigma_i s_i} = \left(\psi^{(i)}(u_{\rho_i}, m_{\sigma_i}, \chi_{\tau_i}), \psi^{(i)}(u_{r_i}, m_{s_i}, \chi_{t_i}) \right).$$

The only non-zero terms in the sum \sum_{α} are those where in each Kronecker delta the pair of indices consists of equal numbers, $\sigma_i = s_i$ for each i , although different Kroneckers may have different pairs.

There are three types of these terms:

(i) Terms where all σ_i s, and hence all s_i s, are equal. These cases are those already solved in the preceding sections, yielding the desired result $\eta_{\alpha} = (-1)^{2sk_{\alpha}}$.

(ii) Terms where all σ_i s, and hence all s_i s, are different ($N \leq 2s + 1$). In these terms the sum \sum_{α} over the permutations reduces to one single member, where P_{α} is the identity I , with $k_{\alpha} = 0$ and $\eta_{\alpha} = 1$. We may therefore write $\eta_{\alpha} = (-1)^{2sk_{\alpha}}$, which again conforms with the desired result. The term T then reduces to

$$\begin{aligned}T_{(rst\rho\sigma\tau)} &= \left(\psi^{(1)}(u_{\rho_1}, m_{s_1}, \chi_{\tau_1}) \cdots \psi^{(N)}(u_{\rho_N}, m_{s_N}, \chi_{\tau_N}) \right), \\ &\quad \left(\psi^{(1)}(u_{r_1}, m_{s_1}, \chi_{t_1}) \cdots \psi^{(N)}(u_{r_N}, m_{s_N}, \chi_{t_N}) \right) = (\Psi_{\text{b}}, \Psi_{\text{a}}),\end{aligned}\quad (35)$$

where Ψ_a and Ψ_b are product functions like those in Eqs. (29) and (30) but without the sums, and in the special case $\sigma_i = s_i$.

In the special subcase that the product functions Ψ_a and Ψ_b already represent the total wave functions the transition amplitude (32) is equal to (35), that is, to one term only. The same then holds for the *probability* of the transition, $|(\Psi_b, \Psi_a)|^2$, and there are no interference terms involving different transitions. This is as in a system of distinguishable particles. We have here a generalization of a well known result for two particles [5, p. 3-12], [24, pp. 1407, 1408], [30].

(iii) Terms which consist of two sets with equal σ_i s and hence equal s_i s within each set: $\sigma_1 = \sigma_2 = \dots = \sigma_l \neq \sigma_{l+1} = \sigma_{l+2} = \dots = \sigma_N$. Then those members of the sum (34) where the permutations work only within one set can be treated like those in point (i) and yield the desired result, whereas those members where the permutations exchange parameters of the first set with those of the second are zero. – Cases of more than two such sets are only technically more complicated but add nothing essential. Thus in all expressions of physical significance the general factor η_α of (28) can be replaced by the standard factor $(-1)^{2sk_\alpha}$ (which means $\exp(i\delta) = \pm 1$ in (4)), so that Feynman's method (5) in the end is seen to meet effectively the standard method (6), which thereby is shown to be equally legitimate.

Another point deserves to be mentioned: the term $T_{(rst\rho\sigma\tau)}$ of formula (34) is zero if only one single-particle function, referring to a particular particle, has a different spin component in the first than in the second total wave function. Such terms thus seem to be restricted to spin-independent interactions. Spin-dependent (or whatever dynamical) interactions can, however, be included by considering the possible wave functions that result from such an interaction as intermediate functions Ψ_i between Ψ_a and Ψ_{bS} , where we have to form the sum over the product of the amplitudes $\sum_i (\Psi_{bS}, \Psi_i)(\Psi_i, \Psi_a)$ in the case that the (situations associated with the) Ψ_i are not observed, or over the product of the respective probabilities in the case that the Ψ_i are observed.

8 The relativistic domain

The derivation of the spin-statistics connection presented so far evidently does not require relativity theory. Can it be extended into the relativistic domain? In Lorentz-invariant theory spin and orbital angular momentum are no longer separately conserved quantities, and the two are in general mixed up in a complicated way. There are however functions which are eigenfunctions of the spin-component operator only, with no admixture of orbital angular momentum: the helicity functions [31]. A helicity function describes a free particle with definite non-zero linear momentum and is an eigenfunction of the operator of the spin component with respect to an axis that is parallel or antiparallel to the direction of the particle's momentum. Thus we may replace the previously discussed eigenfunctions of the operator of the spin component along a fixed direction by the helicity functions. Helicities are invariant under ordinary rotations, and the rotation operators commute with the permutation operators, so we may express the momentum eigenfunctions which have their

momenta in arbitrary directions by suitably rotated eigenfunctions with momenta in one common direction (cf. [31, pp. 407, 408]). For these functions we can define a common reference direction for the angles χ , and then construct and add up the functions with the permuted parameters in the previously described way. This works not only for momentum eigenstates, i.e. plane waves, but also for linear superpositions of plane waves, i.e. wave packets.

Appendix. The standard and the Feynman method

In order to make the paper self-contained we here show that Eqs. (5) and (6) of Section 2 are equivalent, provided the phase factor is $\exp(i\delta) = \pm 1$. We mainly follow [24, pp. 1383 - 1385], and we will go into enough detail that it will become clear why the equivalence does not hold in the case of the more general phase factor η_α of Eq. (28).

We at once generalize Eqs. (5) and (6) to N -particle systems and begin with the Fermi case in the standard form of conventional quantum mechanics

$$f = \left(\sqrt{N!} A \Psi_b(1, 2, \dots, N), \sqrt{N!} A \Psi_a(1, 2, \dots, N) \right), \quad (36)$$

where A is the antisymmetrizer

$$A = \frac{1}{N!} \sum_{\alpha} (-1)^{k_{\alpha}} P_{\alpha}. \quad (37)$$

The factor $(-1)^{k_{\alpha}}$ in front of P_{α} is completely determined by P_{α} : k_{α} is the number of transpositions that make up the permutation P_{α} , where it of course only matters whether k_{α} is even or odd.

The numbers $\sqrt{N!}$ accomplish the normalization of the two factor functions in the scalar product, where it is assumed that the single functions in the sums (in A) are mutually orthonormal. The antisymmetrizer satisfies the relations

$$A^{\dagger} = A = A^2, \quad (38)$$

which will be seen to be essential and which will be proved in three steps.

(i) Proof of $A^{\dagger} = A$:

$$A^{\dagger} = \frac{1}{N!} \sum_{\alpha} \left((-1)^{k_{\alpha}} \right)^* P_{\alpha}^{\dagger} = \frac{1}{N!} \sum_{\alpha} \left((-1)^{k_{\alpha}} \right)^* P_{\alpha}^{-1}$$

because P_{α} , being a product of unitary transposition operators, is also unitary. As $(-1)^{k_{\alpha}}$ is a real number we have

$$A^{\dagger} = \frac{1}{N!} \sum_{\alpha} (-1)^{k_{\alpha}} P_{\alpha}^{-1} = \frac{1}{N!} \sum_{\alpha} (-1)^{k_{\alpha}} P_{\alpha} = A$$

because replacing each P_{α}^{-1} by its inverse P_{α} merely changes the order of the terms in the sum since the set $\{P_{\alpha}^{-1}\}$ is again the permutation group. And since the inverse of a permutation operator is equal to the product of the same transposition

operators taken in the opposite order, the operators P_α in A are multiplied by the same factors as the operators P_α^{-1} .

The more general phase factor (28) in place of $(-1)^{k_\alpha}$ is not a real number so that the above proof would not go through. One could replace (28) with a real number by taking the mean of F_χ calculated with counterclockwise rotations only and F_χ calculated with clockwise rotations only. This would result in replacing $\exp(iX)$ by $\cos(X)$ in formulas (27) and (28). Nevertheless there is no essential advantage in this because of the more important feature that both the exponential and the cosine version, in contrast to the factor $(-1)^{k_\alpha}$, depend not only on the permutation P_α but also on the wave function (m, χ) on which P_α operates.

(ii) Proof of $P_{\alpha 0}A = (-1)^{k_{\alpha 0}}A$ (which will be needed in point (iii) below):

$$P_{\alpha 0}A := \frac{1}{N!} \sum_{\alpha} (-1)^{k_\alpha} P_{\alpha 0} P_\alpha. \quad (39)$$

$P_{\alpha 0}P_\alpha =: P_\beta$ is also a permutation operator. If for fixed $P_{\alpha 0}$ we choose successively for P_α all the permutations of the group, we see that each of the P_β coincides with one and only one of these permutation operators, though of course in different order. Therefore the right-hand side of (39) would be proportional to the antisymmetrizer if the factor multiplying P_β in the sum were $(-1)^{k_{\alpha 0} + k_\alpha}$, according to the prescription given after Eq. (37). This is easy to achieve by multiplying every single term in the sum in (39) by $(-1)^{k_{\alpha 0}}$ and neutralizing this by dividing the sum as a whole by $(-1)^{k_{\alpha 0}}$:

$$P_{\alpha 0}A = \frac{1}{N!} \frac{1}{(-1)^{k_{\alpha 0}}} \sum_{\alpha} (-1)^{k_{\alpha 0} + k_\alpha} P_{\alpha 0} P_\alpha. \quad (40)$$

Then we have

$$P_{\alpha 0}A = \frac{1}{(-1)^{k_{\alpha 0}}} A = (-1)^{k_{\alpha 0}} A, \quad (41)$$

where the last equality holds because the factor is equal to its reciprocal value. This is also not true for the factor (28).

(iii) Proof of $A^2 = A$: with $P_\alpha A = (-1)^{k_\alpha} A$ of point (ii) we easily deduce

$$A^2 := \frac{1}{N!} \sum_{\alpha} (-1)^{k_\alpha} P_\alpha A = \frac{1}{N!} \left(\sum_{\alpha} \left((-1)^{k_\alpha} \right)^2 \right) A = \frac{1}{N!} (N!) A = A.$$

This only holds because $((-1)^{k_\alpha})^2 = 1$, which is not true for the factor (28).

Now, with (38) the scalar product in the standard form for N -particle fermion systems (36) may be written as

$$f = N! \left(A^\dagger A \Psi_b, \Psi_a \right) = N! \left(A \Psi_b, \Psi_a \right) = \left(\sum_{\alpha} (-1)^{k_\alpha} P_\alpha \Psi_b, \Psi_a \right), \quad (42)$$

which is the scalar product in Feynman's form for N -particle fermion systems.

In the Bose case the antisymmetrizer A is to be replaced by the symmetrizer $S = (1/N!) \sum_{\alpha} P_\alpha$, which satisfies the relations $S^\dagger = S = S^2$ analogous to (38), and leads to formula (42) with $(-1)^{k_\alpha}$ being replaced by 1.

References

- [1] M. Fierz, Über die relativistische Theorie kräftefreier Teilchen mit beliebigem Spin, *Helv. Phys. Act.* **12**, 3 - 37 (1939).
- [2] W. Pauli, The connection between spin and statistics, *Phys. Rev.* **58**, 716 - 722 (1940).
- [3] R. Jost, Das Pauli-Prinzip und die Lorentz-Gruppe, in: *Theoretical Physics in the Twentieth Century*, edited by M. Fierz and V. F. Weisskopf (Interscience Publishers, New York, 1960), pp. 107 - 136.
- [4] I. Duck and E. C. G. Sudarshan, *Pauli and the Spin-Statistics Theorem* (World Scientific, Singapore, 1997).
- [5] R. P. Feynman, R. B. Leighton, and M. Sands, *The Feynman Lectures on Physics*, Vol. III (Addison-Wesley, Reading MA, 1965).
- [6] About half of these publications are accessible via the internet under [arXiv.org/find](https://arxiv.org/find) [Title: spin AND statistics]; the others can be traced back from these.
- [7] R. C. Hilborn, Answer to Question #7 [“The spin-statistics theorem,” Dwight E. Neuenschwander, *Am. J. Phys.* 62 (11), 972 (1994)], *Am. J. Phys.* **63**, 298 - 299 (1995).
- [8] I. Duck and E. C. G. Sudarshan, Toward an understanding of the spin-statistics theorem, *Am. J. Phys.* **66**, 284 - 303 (1998).
- [9] R. H. Romer, The spin-statistics theorem, *Am. J. Phys.* **70**, 791 (2002).
- [10] J. A. Morgan, Spin and statistics in classical mechanics, *Am. J. Phys.* **72**, 1408 - 1417 (2004) (quant-ph/0401070).
- [11] A. A. Broyles, Derivation of the Pauli exchange principle, quant-ph/9906046; Spin and statistics, *Am. J. Phys.* **44**, 340 - 343 (1976).
- [12] M. Peshkin, Reply to “Non-relativistic proofs of the spin-statistics connection”, by Shaji and Sudarshan., quant-ph/0402118; Reply to “Comment on ‘Spin and statistics in nonrelativistic quantum mechanics: The spin-zero case’ ”, *Phys. Rev. A* **68**, 046102 (2003); Reply to “No spin-statistics connection in nonrelativistic quantum mechanics”, quant-ph/0306189; Spin and statistics in non-relativistic quantum mechanics: The spin-zero case, *Phys. Rev. A* **67**, 042102 (2003); On Spin and Statistics in Quantum Mechanics, quant-ph/0207017.
- [13] J. A. Morgan, Demonstration of the spin-statistics connection in elementary quantum mechanics, arXiv:physics/0702058.

- [14] B. Kuckert, Spin and statistics in nonrelativistic quantum mechanics, I, Phys. Letters A **322**, 47 - 53 (2004).
- [15] E. Donth, Ein einfacher nichtrelativistischer Beweis des Spin-Statistik-Theorems und das Verhältnis von Geometrie und Physik in der Quantenmechanik, Wissenschaftliche Zeitschrift der Technischen Hochschule „Carl Schorlemmer“ Leuna-Merseburg **19**, 602 - 606 (1977); Non-relativistic proof of the spin statistics theorem, Phys. Letters **32A**, 209 - 210 (1970).
- [16] B. Kuckert and J. Mund, Spin & statistics in nonrelativistic quantum mechanics, II, Ann. Physik (Leipzig) **14**, 309 - 311 (2005) (arXiv:quant-ph/0411197).
- [17] H. Bacry, Answer to Question #7 [“The spin-statistics theorem,” Dwight E. Neuenschwander, Am. J. Phys. 62 (11), 972 (1994)], Am. J. Phys. **63**, 297 - 298 (1995); Introduction aux concepts de la physique statistique (Ellipses, Paris 1991), pp. 198 - 200.
- [18] A. P. Balachandran, A. Daughton, Z.-C. Gu, R. D. Sorkin, G. Marmo and A. M. Srivastava, Spin-statistics theorems without relativity or field theory, Int. J. Modern Physics A **8**, 2993 - 3044 (1993).
- [19] R. P. Feynman, The reason for antiparticles, in: Elementary Particles and the Laws of Physics, edited by R. P. Feynman and S. Weinberg (Cambridge University Press, New York 1987), pp. 1 - 59, especially pp. 56 - 59.
- [20] M. York, Symmetrizing The Symmetrization Postulate, in: Spin-Statistics Connection and Commutation Relations, edited by R. C. Hilborn and G. M. Tino (American Institute of Physics, Melville 2000), pp. 104 - 110 (quant-ph/0006101); Identity, Geometry, Permutation, And The Spin-Statistics Theorem, quant-ph/9908078.
- [21] A. Jabs, Quantum mechanics in terms of realism, Physics Essays **9**, 36 - 95, 354 (1996) (quant-ph/9606017); An Interpretation of the Formalism of Quantum Mechanics in Terms of Epistemological Realism, Brit. J. Phil. Sci. **43**, 405 - 421 (1992). In this interpretation, which is opposed to the Copenhagen interpretation, the quantum objects are not point particles but extended objects, represented by the wave functions, like $\psi(\mathbf{r}^{(i)}) \equiv \psi^{(i)}(\mathbf{r})$. The index (i), rather than denoting the particle i in the wave function, denotes the wave function itself. The vector $\mathbf{r}^{(i)}$ does not mean the position of point particle i (where is it when its position is not being measured?) but the position variable of wave function i , and $|\psi(\mathbf{r}^{(i)})|^2 d^3r$ is the probability that the wave function i causes an effect about the point \mathbf{r} . In either interpretation, realist or Copenhagen, these labels are needed to avoid self-interactions or to show the one-particle operators which wave function to operate on. So, the construction of the proof presented here is independent of which of the interpretations is adopted, and the present paper sticks to the traditional formulation.

- [22] J. D. Bjorken, S. D. Drell, *Relativistic Quantum Mechanics* (McGraw-Hill, New York 1964), p. 136.
 - [23] R. P. Feynman, *Quantum Electrodynamics* (W. A. Benjamin, Reading MA, 1961), pp. 124, 125. The argumentation formulated here in quantum electrodynamics carries over to quantum mechanics. This holds also for equivalent formulations in many books on quantum field theory.
 - [24] C. Cohen-Tannoudji, B. Diu and F. Laloë, *Quantum Mechanics, Vols. I, II* (John Wiley & Sons, New York, 1977).
 - [25] K. Knopp, *Funktionentheorie, Zweiter Teil* (de Gruyter, Berlin, 1955), p. 90. English translation: *Theory of Functions, Part II*, translated by F. Bagemihl (Dover Publications, Mineola NY, 1996), p. 101.
 - [26] T. von Foerster, Answer to Question #7 [“The spin-statistics theorem,” Dwight E. Neuenschwander, *Am. J. Phys.* 62 (11), 972 (1994)], *Am. J. Phys.* 64 (5), 526 (1996); R. R. Gould, Answer to Question #7 [“The spin-statistics theorem,” Dwight E. Neuenschwander, *Am. J. Phys.* 62 (11), 972 (1994)], *Am. J. Phys.* 63 (2), 109 (1995); R. Penrose and W. Rindler, *Spinors and Space-Time, Vol. 1* (Cambridge University Press, Cambridge, 1984) p. 43; L. C. Biedenharn and J. D. Louck, *Angular Momentum in Quantum Physics* (Addison-Wesley Publishing Company, Reading, MA, 1981), chap. 2; R. W. Hartung, Pauli principle in Euclidean geometry, *Am. J. Phys.* 47 (10), 900 - 910 (1979); E. Rieflin, Some mechanisms related to Dirac’s strings, *Am. J. Phys.* 47 (4), 378 - 381 (1979); and literature cited in these references.
 - [27] E. Schrödinger, Die Mehrdeutigkeit der Wellenfunktion, *Ann. Physik (Leipzig)* 32 (5), 49 - 55 (1938), reprinted in: E. Schrödinger, *Collected Papers, Vol. 3* (Verlag der Österreichischen Akademie der Wissenschaften, Wien, 1984), pp. 583 - 589.
 - [28] C. van Winter, Orbital Angular Momentum and Group Representations, *Ann. Phys. (New York)* 47, 232 - 274 (1968).
 - [29] S. L. Altmann, *Rotations, Quaternions, and Double Groups* (Clarendon Press, Oxford, 1986), chap. 10.
 - [30] W. Pauli, Die allgemeinen Prinzipien der Wellenmechanik, in: *Handbuch der Physik, Vol. 24, Part 1*, edited by H. Geiger and K. Scheel (Springer-Verlag, Berlin, 1933), pp. 189 - 193. English translation of a 1958 reprint: *General Principles of Quantum Mechanics*, translated by P. Achutan and K. Venkatesan (Springer-Verlag, Berlin, 1980), pp. 116 - 121.
 - [31] M. Jacob and G. C. Wick, On the General Theory of Collisions for Particles with Spin, *Ann. Phys. (New York)* 7, 404 - 428 (1959).
-