

Quantum interference by coherence transfer from spin to orbital angular momentum of photons

Eleonora Nagali,¹ Fabio Sciarrino,^{2,1} Francesco De Martini,^{1,3,*} Lorenzo Marrucci,^{4,5,†} Bruno Piccirillo,^{4,6} Ebrahim Karimi,⁴ and Enrico Santamato^{4,6}

¹*Dipartimento di Fisica dell'Università "La Sapienza" and Consorzio Nazionale Interuniversitario per le Scienze Fisiche della Materia, Roma 00185, Italy*

²*Centro Studi e Ricerche "Enrico Fermi", Via Panisperna 89/A, Compendio del Viminale, Roma 00184, Italy*

³*Accademia Nazionale dei Lincei, via della Lungara 10, Roma 00165, Italy*

⁴*Dipartimento di Scienze Fisiche, Università di Napoli "Federico II", Compl. Univ. di Monte S. Angelo, 80126 Napoli, Italy*

⁵*CNR-INFM Coherentia, Compl. Univ. di Monte S. Angelo, 80126 Napoli, Italy*

⁶*Consorzio Nazionale Interuniversitario per le Scienze Fisiche della Materia, Napoli*

The optical "spin-orbit" coupling occurring in a suitably patterned nonuniform birefringent plate known as 'q-plate' allows entangling the polarization of a single photon with its orbital angular momentum (OAM). This process, in turn, can be exploited for building a reversible "spin-OAM interface", capable of transposing the quantum information from the spin to the OAM degree of freedom of photons and *vice versa*. Here, we experimentally demonstrate this process by single-photon quantum tomographic analysis. Moreover, we show that two-photon quantum correlations such as those resulting from coalescence interference, in the Hong-Ou-Mandel effect, can be successfully transferred into the OAM degree of freedom.

In the last few decades, quantum optics has allowed the implementation of a variety of quantum-information protocols. However, the standard information encoding based on the two-dimensional quantum space of photon polarizations (or "spin" angular momentum) imposes significant limitations to the protocols that may be implemented. To overcome such limitations, more recently the orbital angular momentum (OAM) of light, related to the photon's transverse-mode spatial structure [1], has been recognized as a new promising resource, allowing the implementation of a higher-dimensional quantum space, or a "qu-dit", encoded in a single photon [2, 3]. So far, the generation of OAM-entangled photon pairs has been carried out by exploiting the process of parametric down-conversion [4, 5] and has also been utilized in few quantum information protocols [6, 7, 8, 9]. Despite these successes, the optical tools for generating and controlling the OAM photon states (computer-generated holograms, Dove's prisms, cylindrical lens mode converters, etc.) are rather limited and difficult to use. A convenient way to coherently "interface" the OAM degree of freedom of photons with the more easily manipulated spin/polarization one has been missing so far. In this context, the recent invention of an optical device, the so-called "q-plate" (QP), that couples the photon spin to its orbital angular momentum opens up a number of new and unexplored possibilities [10]. In this work we show, both theoretically and experimentally, that this optical "spin-orbit" coupling can be exploited as an effective interface, completely reversible, between polarization and orbital angular momentum in the quantum regime. The QP also enables the efficient generation of single-photon in which the OAM and polarization degrees of freedom are entangled. Furthermore, photon-photon

quantum correlations can be transferred from the spin to the OAM degree of freedom of a photon pair, as experimentally demonstrated here through the observation of a Hong-Ou-Mandel effect in the OAM.

The QP is a birefringent slab having a suitably patterned transverse optical axis, with a topological singularity at its center. Here, we consider a QP with "charge" of the singularity $q = 1$ and uniform birefringent retardation $\delta = \pi$. Such QP modifies the OAM quantum number l (in units of \hbar) of a light beam crossing it, imposing a variation $\Delta l = \pm 2$ whose sign depends on the input polarization, positive for left-circular and negative for right-circular. The helicity of the output circular polarization is also inverted, i.e. the optical spin is flipped.

Let us now rephrase this behavior in a quantum formalism suitable for describing multi-photon states. Let $\hat{a}_{j,l}^\dagger$ be the operator creating a photon in the polarization state $\vec{\pi}_j$ (where $\vec{\pi}_H$ and $\vec{\pi}_V$ stand for horizontal and vertical linear polarizations, respectively) and with the OAM value l . For simplicity, we omit here the radial quantum number which would be needed for a complete characterization of transverse orbital states, as it will play no significant role in the following [11]. The overall dynamics induced by the QP can be then described by the transformations $\hat{a}_{L,0}^\dagger \Rightarrow \hat{a}_{R,2}^\dagger$ and $\hat{a}_{R,0}^\dagger \Rightarrow \hat{a}_{L,-2}^\dagger$, with $\vec{\pi}_L = 2^{-1/2}(\vec{\pi}_H + i\vec{\pi}_V)$ and $\vec{\pi}_R = \vec{\pi}_L^\perp$ standing for left and right circular polarizations, respectively.

The previous relations describe the coupling of the OAM l and the polarization π degrees of freedom taking place in the QP. Interestingly, a first application of this property is to generate single-particle entanglement of π and l degrees of freedom:

$$\left. \begin{array}{l} |H\rangle_\pi |0\rangle_l \\ |V\rangle_\pi |0\rangle_l \end{array} \right\} \xrightarrow{QP} \frac{1}{\sqrt{2}}(|L\rangle_\pi | - 2\rangle_l \pm |R\rangle_\pi | + 2\rangle_l) \quad (1)$$




FIG. 1: Experimental setup. The main source is a Ti:Sa mode-locked laser converted by second harmonic generation (SHG) into a UV beam with wavelength $\lambda_p = 397.5\text{nm}$. This field pumps a nonlinear crystal of β -barium borate (BBO) which emits photon pairs with H and V polarizations and $\lambda = 795\text{nm}$, filtered by the interference filter (IF) with $\Delta\lambda = 6\text{nm}$ and then coupled to a single mode fiber [12]. The gray dot-dashed box has been optionally inserted to prepare a single photon state triggered by detector D_T . Birefringent quartz crystals (Q) having different thicknesses were used to introduce a controlled temporal delay between the two photons. After setting the input polarization by means of a suitably oriented quarter-wave plate, the photons were sent through the q-plate (QP) and the output OAM states were analyzed with the help of a hologram (Hol) (in OAM-to-spin conversion experiments, Hol and QP were interchanged). To measure (or prepare) OAM states in the basis $l = \pm 2$, a double-fork hologram has been used (upper left panel in red box), so that the OAM state of the first diffracted modes is shifted by $\Delta l = \pm 2$, while the undiffracted 0-order beam has $\Delta l = 0$. The photons on the first diffracted modes are then coupled to single mode fibers which select output states with $l = 0$ and convey them to the detectors D_A and D_B . Hence, the detection of a photon in D_A (D_B) corresponds to a photon incident on the hologram with OAM $l = +2$ ($l = -2$). The 1st-order diffraction efficiency of the hologram was $\sim 10\%$. The measurement (or preparation) of OAM in superposition states (see main text) has been realized by adopting the other holograms shown in the red box. For certain measurements, a polarization analysis set (PA) was also inserted before detection. Due to reflection losses, the transmittance of the q-plate is $T \sim 0.90$.

This is an entangled state between two qubit encoded in different degrees of freedom. In particular $\{|+2\rangle_l, |-2\rangle_l\}$ is the basis for the OAM qubit which lies in the subspace of the infinite dimensional hilbert space of orbital angular momentum. As a first experimental step, we set out to verify how accurately the real QP device performs these transformations in the single photon regime (see Fig.1 for experimental details). First, the QP conversion efficiency η from the input TEM_{00} to the $l = \pm 2$ modes has been




FIG. 2: (a-b) Experimental density matrices for the single photon entangled state. The computational values $\{0,1\}$ are associated to the $\{|R\rangle, |L\rangle\}$ polarization states, and to $\{|+2\rangle, |-2\rangle\}$ for the orbital angular momentum l for the first and the second qubit, respectively. The incoming state on the QP is (a) $|H\rangle_\pi|0\rangle_l$, and (b) $|V\rangle_\pi|0\rangle_l$. The average experimental concurrence is $C = (0.95 \pm 0.02)$, while the average purity of the states is $P = \text{Tr}\rho^2 = (0.94 \pm 0.02)$. (c-d) Experimental Poincaré sphere both for the OAM (c) and π (d) degrees of freedom obtained after the $\pi \rightarrow l$ and the $l \rightarrow \pi$ transferrer, respectively. Experimentally we carried out single-qubit tomography in order to determine the Stoke's parameters for the π and their analogous for l degrees of freedom. The mean fidelity values are (c) $F = (98 \pm 1)\%$ and (d) $F = (97 \pm 1)\%$.

estimated through the coupling efficiency with the single mode fiber. We find $\eta \simeq 85\%$, mainly ascribed to the fact that our QP has a birefringent retardation δ not exactly tuned to π , so that the expected conversion efficiency is $\eta = \sin^2 \frac{\delta}{2}$ [10] (the unconverted component remains $l = 0$ and is therefore filtered out). Next, in order to assess the coherence of the transformations in Eq.(1), single photons in the states $|H\rangle_\pi|0\rangle_l$ or $|V\rangle_\pi|0\rangle_l$ were used as input in the QP. We analyzed the output state through a double-fork hologram and a circular-polarization analysis setup along the two diffracted modes: the intensity of the π_R (π_L) polarization component in the mode corresponding to $l = +2$ ($l = -2$) was measured to be equal to 99.8% (99.6%) of the total, with a high agreement with theory. To demonstrate the realization of the pure states given in Eq.(1), a complete single-photon two-qubit quantum state tomography has been carried out, performing measurements both in π and l degrees of freedom. Besides the normal $\{|+2\rangle_l, |-2\rangle_l\}$ OAM basis, measurements were carried out in the two superposition bases $\{|d_+\rangle_l, |d_-\rangle_l\}$ and $\{|d_R\rangle_l, |d_L\rangle_l\}$, defined by $|d_\pm\rangle_l = \frac{1}{\sqrt{2}}(|+2\rangle_l \pm |-2\rangle_l)$ and $|d_{L,R}\rangle_l = \frac{1}{\sqrt{2}}(|+2\rangle_l \pm i|-2\rangle_l)$. The OAM degree of freedom was analyzed in these bases by means of different holograms (see red box in Fig.(1)) [13]. The experimental results are in very good agreement with theoretical

predictions, as shown in Fig.(2-a,b).

Due to its peculiarities, the q-plate provides a convenient way to coherently "interface" the photon OAM with the more easily manipulated spin degree of freedom. Hence as next step we show that such interface can be considered as a real quantum "transferrer" device, which allows to transfer coherently the information from the polarization π to the OAM l degree of freedom, and *vice versa*. Such processes can be formally expressed as follows:

$$|\varphi\rangle_\pi|0\rangle_l \leftrightarrow |H\rangle_\pi|\varphi\rangle_l \quad (2)$$

where $|\varphi\rangle_\pi = (\alpha|H\rangle + \beta|V\rangle)_\pi$ and $|\varphi\rangle_l = (\alpha|+2\rangle + \beta|-2\rangle)_l$. Here we demonstrate only a probabilistic conversion (with probability $p \cong 50\%$), since some output state contributions are discarded. However, by extending the present scheme, it is possible to achieve a complete deterministic information conversion [14].

I - Transferrer $\pi \rightarrow l$. Through a waveplate $\lambda/4$ and a QP, π and l become entangled (Fig. 3-I-a). Then, the information contained in the polarization is erased by inserting a polarizing beam-splitter (PBS). This conversion has been experimentally verified for a set of maximally polarized states. Any π input state, represented by a vector in the Poincaré sphere, has been converted in another vector in the OAM "Poincaré sphere" (see Fig.2-c), determined by carrying out a quantum state tomography of the OAM state. The results demonstrate a high fidelity of the $\pi \rightarrow l$ transformation. An application of this transferrer is shown in Fig.(3-I-b). The initial information encoded in an input state $|\varphi\rangle_\pi|0\rangle_l$ is coherently transferred to the OAM. New information can then be stored in the polarization degree of freedom, thus allowing the encoding of two qubits of information in the same photon. **II - Transferrer $l \rightarrow \pi$.** Let us show that the coupling between the spinorial and the angular degree of freedom in the q-plate is completely reversible (Fig.3-II-a). The input state $|H\rangle_\pi|\varphi\rangle_l$ is sent through a QP and a $\lambda/4$ waveplate (Fig. 3-II-a). By inserting a single mode fiber on the output, only the states with $l = 0$ are efficiently coupled, leading to a probabilistic process. After the fiber, the state $|\varphi\rangle_\pi|0\rangle_l$ has been obtained, which demonstrates the successful conversion from the OAM




FIG. 3: Schematic representation of the two transferrer devices based on the q-plate (QP). The coupling is between the orbital angular momentum degree of freedom l and the polarization one π .

degree of freedom to the polarization one. Analogously to the previous analysis, in Fig.(2-d) is reported the experimental Poincaré sphere obtained after the $l \rightarrow \pi$ conversion. Such transferrer can be for example exploited within a measurement apparatus for analyzing both π and l degrees of freedom without using holograms, with an advantage in terms of efficiency (Fig.(3-II-b)) and flexibility, since there is no need to change the hologram for each state to be analysed. This bidirectional spin-OAM interface which we have described allows the extension to OAM of many protocols currently only possible with polarization. For example, the realization of a two-photon *C-NOT* gate for OAM states can be obtained by exploiting a *C-NOT* for π states. As a further consideration, the QP allows implementing a fast OAM switcher device, by modulating the polarization degree of freedom through a Pockel's cell and then transferring such modulation to the OAM.

It is natural to ask if the QP is able to preserve the photon-photon correlations too. This would also be a crucial test of the QP potential in the quantum optics field, as multi-photon correlations are extremely sensitive to mode matching and optical quality issues. We address here the generation of a quantum state exhibiting two-photon interference, just as in the classic Hong-Ou-Mandel (HOM) effect [15], but taking place in the OAM degree of freedom of photons. We want to create such a state in a single beam exploiting the QP spin-to-OAM conversion capabilities. Consider first the case of two independent linearly polarized photons, one horizontal and the other vertical, going through the QP: each will undergo the QP transformation given in Eq.(1) and the two outgoing photons will end up having opposite OAM values 50% of the times. If, however, the two photons are undistinguishable, except for their polarization, the field state at the QP input side is $|\Psi_{in}\rangle = |1\rangle_{H,0}|1\rangle_{V,0}$, where we are now using a photon-number ket notation for our multi-photon quantum states. This input state can be rewritten in the circular-polarization basis as $|\Psi_{in}\rangle = \frac{1}{\sqrt{2}}(|2\rangle_{R,0}|0\rangle_{L,0} - |0\rangle_{R,0}|2\rangle_{L,0})$. After the QP, this state evolves into

$$|\Psi_{out}\rangle = \frac{1}{\sqrt{2}}(|2\rangle_{L,-2}|0\rangle_{R,2} - |0\rangle_{L,-2}|2\rangle_{R,2}), \quad (3)$$

in which only photons carrying parallel OAM, either $+2$ or -2 are found. Hence, the QP action can be again interpreted as a mode converter, coherently transferring the HOM-like two-photon quantum correlation from the spin degree of freedom to the OAM one. In the end, what is physically relevant is that a non-classical HOM-like statistics should be finally measured in the OAM degree of freedom *irrespective of the polarization*. Indeed, if the polarization information is not erased, the HOM coalescence is still observable in the spin degree of freedom too. The QP is then able to generate, for the first time, a multi-photon state having non-classical correla-




FIG. 4: (a) Coincidence counts between $[D_A, D_B]$ versus the temporal delay t_d , for the state $|\Psi_{out}\rangle$. The continuous line shows the best fit of the experimental data. (b) Coincidence counts $[D_A, D_{A'}]$ versus t_d . (c) Coincidence counts $[D_A, D_B]$ versus t_d ($V = (0.91 \pm 0.01)$) for the state $|\Phi_{out}\rangle$. (d) Coincidence counts $[D_A, D_{A'}]$ in different OAM bases.

tions in OAM within a single longitudinal mode. Let us appreciate that, since the process is started by photons in the well-defined TEM₀₀ mode, the radial part of the final state $|\Psi_{out}\rangle$ is deterministically fixed [16]. In Fig.(1) is reported the OAM-HOM experimental setup. The coincidence between $[D_A, D_B]$ detects the state contribution of $|1\rangle_{l=2}|1\rangle_{l=-2}$. Due to the coalescence interference, for otherwise identical photons this component is vanishing in the state outcoming from the QP. In order to study the transition from the case of classical behaviour (independent photons) to the case of full quantum interference, a variable temporal delay t_d between the H and V polarizations in the state $|\Psi_{in}\rangle$ has been introduced (Q in Fig.1). The experimental visibility of the quantum interference shown in Fig.(4-a) is $V_{exp} = (0.95 \pm 0.02)$. As a further confirmation, we have measured the contribution of two photons with $l = +2$ by recording the coincidence counts between two detectors $[D_A, D'_{A'}]$ placed on the output modes of a fiber-based beam splitter inserted on the same k_A diffracted mode (not reported in figure). Theoretically, the coalescence of the two photons should lead to a coincidence enhancement by a factor $\Gamma = 2$, and experimentally we found $\Gamma_{exp} = (1.94 \pm 0.02)$.

Furthermore, we verified that even after erasing all information still contained in the polarization degree of freedom, the final state is still coherent and exhibits the same non-classical photon correlations in OAM. To this purpose, we let both the two photons pass through a horizontal linear polarizer set in a common state H . The state of Eq.(3) is then projected into:

$$|\Phi_{out}\rangle = \frac{1}{\sqrt{2}}(|0\rangle_{H,2}|2\rangle_{H,-2} - |2\rangle_{H,2}|0\rangle_{H,-2}) \quad (4)$$

in which the polarization information has been erased and only the OAM degree of freedom preserves the non-classical statistics and two-photon coalescence effect. We verified again the coalescence of the photons by a measurement similar to the previous one, and the experimental data are reported in Fig.4-(c). Next, we set out to verify that we really obtain the pure state given in Eq. (4) and not a statistical mixture having similar OAM correlation properties. To do this, we need to verify the coherence of the two contributions appearing in Eq. (4), and this in turn will also verify the purity of the state given in Eq. (3). This was accomplished by analyzing the photons in the other OAM bases already discussed above. In these bases, we have $|\Phi_{out}\rangle = |1\rangle_{H,d_+}|1\rangle_{H,d_-} = \frac{1}{\sqrt{2}}(|0\rangle_{H,d_L}|2\rangle_{H,d_R} + |2\rangle_{H,d_L}|0\rangle_{H,d_R})$. Therefore, our coherence verification is actually turned into a measurement of two photons in the same OAM state ($|d_+\rangle, |d_-\rangle, |d_R\rangle, |d_L\rangle$). As expected for ($|d_+\rangle, |d_-\rangle$), the events of two photons with the same orbital states are strongly suppressed, while for ($|d_R\rangle, |d_L\rangle$) they are doubled, with an overall correlation visibility $V = (0.86 \pm 0.02)$ (Fig. 4(d)).

In conclusion, we have shown that the q-plate device can be used as a coherent and bidirectional quantum interface allowing the transfer of quantum information between the polarization and the orbital angular momentum degrees of freedom of the photon, both in the case of single photon states and of two-photon correlated states. In the latter case, we demonstrated a Hong-Ou-Mandel kind of quantum interference-coalescence in the orbital angular momentum. The ease of use, flexibility and good conversion efficiency demonstrated by this device makes it a very promising tool for fully exploiting the OAM degree of freedom of photons, in combination with polarization, as a new resource to implement high-dimensional quantum information protocols. We thank G. Maiello for assistance in preparing the q-plates, and P. Mataloni and G. Vallone for useful discussions.

* Electronic address: francesco.demartini@uniroma1.it

† Electronic address: lorenzo.marrucci@na.infn.it

- [1] L. Allen, et al., *Phys. Rev. A* **45**, 8185 (1992)
- [2] G. Molina-Terriza, et al., *Nature Phys.* **3**, 305 (2007)
- [3] S. Franke-Arnold, et al., *Laser & Photon. Rev.*, **4**, 299313 (2008)
- [4] A. Mair, et al., *Nature (London)* **412**, 313 (2001)
- [5] G. Molina-Terriza, et al., *Opt. Commun.* **228**, 115 (2003).
- [6] J. Leach et al., *Phys. Rev. Lett.* **88**, 257901 (2002).
- [7] A. Vaziri, et al., *Phys. Rev. Lett.* **91**, 227902 (2003).
- [8] J.T. Barreiro, et al., *Phys. Rev. Lett.* **95**, 260501 (2005).
- [9] J.T. Barreiro, et al., *Nature Phys.* **4**, 282 (2008).
- [10] L. Marrucci, et al., *Phys. Rev. Lett.* **96**, 163905 (2006).
- [11] The radial profile generated by the QP, as well as by other OAM-generating methods, although not a Laguerre-Gauss mode, is a well defined hypergeometric-gaussian

mode [see, E. Karimi et al., *Opt. Lett.* **32**, 3053 (2007); Karimi et al., arXiv0809.4220v1]. In OAM-based quantum optics, when only states with opposite value of the OAM are involved, the radial profile is not critical, because it is deterministic and not entangled with the other degrees of freedom.

- [12] F. De Martini, et al., *Prog. in Quant. Elect.* **29**, 165 (2005).
[13] N.K. Langford, et al., *Phys. Rev. Lett.* **93**, 053601 (2004).

[14] E. Nagali, et al., to be published in a forthcoming paper.

[15] C.K. Hong, et al., *Phys. Rev. Lett.* **59**, 2044 (1987).

[16] This result should be compared with the case in which photon pairs correlated in the OAM degree of freedom are generated by parametric down conversion [2, 4]. In the latter case, any control on the radial part of the transverse modes is impossible and post-selection of the desired OAM value is mandatory.