

Absolutely Continuous Spectrum for the Anderson Model on Some Tree-like Graphs

Florina Halasan ¹

*Department of Mathematics, University of British Columbia,
1984 Mathematics Road, Vancouver, B.C. V6T 1Z2 Canada*

Abstract

We prove persistence of absolutely continuous spectrum for the Anderson model on a general class of tree-like graphs.

MSC: 81Q10.

Keywords: Random Schrödinger operators; Absolutely continuous spectrum; Anderson Model; Hyperbolic geometry

1 Introduction

Random Schrödinger Operators are used as models for disordered quantum mechanical systems. Their investigation goes back to the late 1950s. The Anderson Model was introduced to describe the motion of a quantum-mechanical electron in a crystal with impurities. For this model, the states corresponding to an absolutely continuous spectrum describe mobile electrons. Thus, an interval of absolutely continuous spectrum is an energy range in which the material is a conductor.

An outstanding open problem, the extended states conjecture, is to prove existence of absolutely continuous spectrum for the lattice \mathbb{Z}^d with $d > 2$. Until now, it is only for the Bethe lattice that this has been established. A first result on the topic was obtained by A. Klein, [5], in 1998; he proved that for weak disorder, on the Bethe lattice, there exists absolutely continuous spectrum for almost all potentials. More recently, Aizenman, Sims and Warzel proved similar results for the Bethe lattice using a different method (see [1]). Their method establishes the persistence of absolutely continuous spectrum under weak disorder and also in the presence of a periodic background potential. During the same time, Froese,

¹Email address: halasan@math.ubc.ca

The author was supported by an NSERC/MITACS Scholarship. This paper is based on part of the author's doctoral thesis.

It is a pleasure to thank Dr. R. Froese for all his guidance and support.




Figure 1: The general tree, \mathbb{T}

Hasler and Spitzer introduced a geometric method for proving the existence of ac spectrum on graphs (see [2]). In their second paper on the topic, [3], they proved delocalization for the Bethe lattice of degree 3 using this geometric approach.

In this work, we provide a simplified version of the geometric method, and its extension to more general trees. We prove the existence of some absolutely continuous spectrum for the Anderson Model on a tree-like graph, \mathbb{T} , defined as follows (see Figure 1). We start with an infinite full binary tree in which each node has degree 3 except for the origin, 0, which has degree 2. On the top edge, coming out of each node in the binary tree, we add m new vertices; subsequently we add n , $m \neq n$, new vertices on each lower edge thus obtaining our new tree. In fact, if we start with a k -nary tree, the results in this paper still hold by similar arguments. We present the binary case, for simplicity. Also, the proof for the $m = n$ case is a straightforward generalization of the one for the Bethe lattice presented in [4].

Using the terminology established in [1], we will use the symbol \mathbb{T} for both our tree graph and its set of vertices. For each $x \in \mathbb{T}$ we have at most one neighbor towards the root and one or more in what we refer to as the forward direction. We say that $y \in \mathbb{T}$ is in the future of $x \in \mathbb{T}$ if the path connecting y and the root runs through x . The subtree consisting of all the vertices in the future of x , with x regarded as its root, is denoted by \mathbb{T}^x .

2 Statement of the Main Result

The Anderson Model on \mathbb{T} is given by the random Hamiltonian, H , on the Hilbert space $\ell^2(\mathbb{T})$. This operator is of the form

$$H = \Delta + q$$

where:

1.

$$\ell^2(\mathbb{T}) = \{ \varphi : \mathbb{T} \rightarrow \mathbb{C}; \sum_{x \in \mathbb{T}} |\varphi(x)|^2 < \infty \}$$

2. The free Laplacian Δ is defined by

$$(\Delta\varphi)(x) = \sum_{y:d(x,y)=1} (\varphi(x) - \varphi(y)), \text{ for all } \varphi \in \ell^2(\mathbb{T}),$$

where d defines the distance between sites.

3. The operator q is a random potential,

$$(q\varphi)(x) = q(x)\varphi(x),$$

where $\{q(x)\}_{x \in \mathbb{T}}$ is a family of independent, identically distributed real random variables with common probability distribution ν .

Our main theorem is the following:

Theorem 1. *Let F be the largest open set included in the spectrum of Δ (this spectrum depends on m and n). For any closed subinterval E , $E \subset F$, there exist $0 < p < 1$, $\eta_1 > 0$ and $\eta_2 > 0$, such that for all ν with*

$$\int_{|q| \geq \eta_1} (1 + |q|^{2(1+p)}) d\nu(q) \leq \eta_2$$

the spectrum of H is purely absolutely continuous in E with probability one.

The following theorem together with the criterion from Section 5 will give us the proof of Theorem 1.

Theorem 2. *Let $\delta_x \in \ell^2(\mathbb{T})$ be the indicator function supported at the site $x \in \mathbb{T}$. Under the hypothesis of the previous theorem, we have*

$$\sup_{\lambda \in R(E, \epsilon)} \mathbb{E} \left(|\langle \delta_x, (H - \lambda)^{-1} \delta_x \rangle|^{1+p} \right) < \infty,$$

for some $\epsilon > 0$ and all $x \in \mathbb{T}$. Here $R(E, \epsilon) = \{z \in \mathbb{H} : \text{Re}(z) \in E, 0 < \text{Im}(z) \leq \epsilon\}$ and $\mathbb{H} = \{x + iy : x, y \in \mathbb{R} \text{ and } y > 0\}$ denotes the complex upper half plane.




Figure 2: The nodes in the recurrence relation for the Green's function.

3 Outline of the Proof

The analysis on trees is generally more accessible than on other graphs since various quantities computed at the tree root satisfy recursion relations. One of these quantities is the Green's function. We write H^x for the restriction of H to \mathbb{T}^x and define the Green's function for the truncated graph as follows:

$$G^x(\lambda) = \langle \delta_x, (H^x - \lambda)^{-1} \delta_x \rangle .$$

We notice that since H is random, at each $x \in \mathbb{T}$ the Green's function, G^x , is an \mathbb{H} -valued random variable. We denote by ρ the probability distribution of G^0 .

The fundamental recurrence relation for these Green's functions is given by

$$G^x(\lambda) = \phi(G^{x_1}(\lambda), G^{x_2}(\lambda), q_1 \dots q_M, \lambda), \quad (1)$$

where

$$\phi : \mathbb{H}^2 \times \mathbb{R}^M \times \mathbb{H} \rightarrow \mathbb{H}$$

is defined by

$$\phi(z_1, z_2, q_1 \dots q_M, \lambda) = \frac{-1}{\phi_n(z_1, q_1 \dots q_n, \lambda) + \phi_m(z_2, q_{n+1}, \dots, q_{n+m}, \lambda) + \lambda - q_M} \quad (2)$$

with

$$\begin{aligned} \phi_0(z, \lambda) &= z \\ \phi_1(z, q_1, \lambda) &= \frac{-1}{z + \lambda - q_1 + 1} \\ \phi_n(z, q_1 \dots q_n, \lambda) &= \frac{-1}{\phi_{n-1}(z, q_1 \dots q_{n-1}, \lambda) + \lambda - q_n + 1} \end{aligned}$$

and $M = m + n + 1$. The nodes x, x_1, x_2 and the potentials q_1, \dots, q_M involved in the recurrence (1) are shown in Figure 2.

If $\text{Im}(\lambda) > 0$, $z = \phi(z, z, 0, \dots, 0, \lambda)$ is a cubic equation with a unique root, z_λ , in the upper half plane, i.e. $\text{Im}(z_\lambda) > 0$. The map $\mathbb{H} \ni \lambda \mapsto z_\lambda$ extends continuously onto the real axis. The set $\{z_\lambda\}_{\lambda \in E}$ is a compact curve strictly contained in \mathbb{H} .

Thus, when λ lies in the strip

$$R(E, \epsilon) = \{z \in \mathbb{H} : \text{Re}(z) \in E, 0 < \text{Im}(z) \leq \epsilon\}$$

with $E \subset F$ closed and ϵ sufficiently small, $\text{Im}(z_\lambda)$ is bounded below and $|z_\lambda|$ is bounded above.

To measure the distance between the Green's function of the free Laplacian and the perturbed Green's function on \mathbb{T} we will use a weight function defined by

$$w(z) = 2 \text{Im}(z_\lambda) (\cosh(\text{dist}_{\mathbb{H}}(z, z_\lambda)) - 1) = \frac{|z - z_\lambda|^2}{\text{Im}(z)}.$$

Up to constants, $w(z)$ is the hyperbolic cosine of the hyperbolic distance from z to z_λ . Note: we have dropped the λ -dependence from the notation.

For the proof of Theorem 2 we need a few lemmas about the following quantity:

$$\mu_{2,p}(z_1, z_2, q_1 \dots q_M, \lambda) = \frac{w^{1+p}(\phi(z_1, z_2, q_1 \dots q_M, \lambda)) + w^{1+p}(\phi(z_2, z_1, q_1 \dots q_M, \lambda))}{w^{1+p}(z_1) + w^{1+p}(z_2)}.$$

Lemma 3. *For any closed subinterval E , $E \subset F$ and some $0 < p < 1$, there exist positive constants $\epsilon, \eta_1, \epsilon_0$ and a compact set $\mathcal{K} \subset \mathbb{H}^2$ such that*

$$\mu_{2,p}|_{\mathcal{K}^c \times [-\eta_1, \eta_1]^M \times R(E, \epsilon_0)} \leq 1 - \epsilon. \quad (3)$$

Here \mathcal{K}^c denotes the complement $\mathbb{H}^2 \setminus \mathcal{K}$.

Lemma 4. *For any closed subinterval E , $E \subset F$ and any $0 < p < 1$, there exist positive constants ϵ_0, C and a compact set $\mathcal{K} \subset \mathbb{H}^2$ such that*

$$\mu_{2,p}|_{\mathcal{K}^c \times \mathbb{R}^M \times R(E, \epsilon_0)} \leq C \prod_{k=1}^M (1 + |q_k|^{2(1+p)}). \quad (4)$$

Given these two Lemmas we can prove that the decay of the probability distribution function of the Green's function at infinity is preserved as $\text{Im}(\lambda)$ becomes small, provided that ν has a finite moment of order $2(1 + p)$.

Theorem 5. For any closed subinterval E , $E \subset F$ and some $0 < p < 1$, there exist $\eta_1 > 0$, $\eta_2 > 0$ and $\epsilon > 0$, such that for all ν satisfying

$$\int_{|q| \geq \eta_1} (1 + |q|^{2(1+p)}) d\nu(q) \leq \eta_2,$$

we have

$$\sup_{\lambda \in R(E, \epsilon)} \mathbb{E} \left(w^{1+p}(G^0(\lambda)) \right) < \infty.$$

Proof. Let η_1 and p be given by Lemma 3, and choose ϵ_0 and \mathcal{K} that work in both Lemma 3 and Lemma 4. For any $(z_1, z_2) \in \mathcal{K}^c$ and $\lambda \in R(E, \epsilon)$, we estimate

$$\begin{aligned} & \int_{\mathbb{R}^M} \mu_{2,p}(z_1, z_2, q_1 \dots q_M, \lambda) d\nu(q_1) \dots d\nu(q_M) \\ & \leq (1 - \epsilon) \int_{[-\eta_1, \eta_1]^M} d\nu(q_1) \dots d\nu(q_M) + C \int_{\mathbb{R}^2 \setminus [-\eta_1, \eta_1]^M} \prod_{i=1}^M (1 + |q_i|^{2(1+p)}) d\nu(q_1) \dots \\ & \hspace{20em} d\nu(q_M) \\ & \leq (1 - \epsilon) + C' \eta_2 \\ & \leq 1 - \epsilon/2 \end{aligned}$$

provided η_2 is sufficiently small. The probability distributions on the hyperbolic plane are defined by

$$\rho(A) = \text{Prob}\{G^x(\lambda) \in A\} = \text{Prob}\{G^0(\lambda) \in A\},$$

where x is any site that had degree 3 in \mathbb{T} .

The recursion formula for the Green's function implies that the distributions $d\rho$ are related by

$$\begin{aligned} \rho(A) &= \text{Prob}\{\phi(z_1, z_2, q_1 \dots q_M, \lambda) \in A\} = \text{Prob}\{(z_1, z_2, q_1 \dots q_M, \lambda) \in \phi^{-1}(A)\} \\ &= \int_{\phi^{-1}(A)} d\rho(z_1) d\rho(z_2) d\nu(q_1) \dots d\nu(q_M) \\ &= \int_{\mathbb{H}^2 \times \mathbb{R}^M} \chi_A(\phi(z_1, z_2, q_1 \dots q_M, \lambda)) d\rho(z_1) d\rho(z_2) d\nu(q_1) \dots d\nu(q_M) \end{aligned}$$

which gives us that for any continuous function $f(z)$

$$\int_{\mathbb{H}} f(z) d\rho(z) = \int_{\mathbb{H}^2 \times \mathbb{R}^M} f(\phi(z_1, z_2, q_1 \dots q_M, \lambda)) d\rho(z_1) d\rho(z_2) dv(q_1) \dots dv(q_M).$$

Using this relation, for $\lambda \in R(E, \epsilon_0)$, we obtain

$$\begin{aligned} \mathbb{E} \left(w^{1+p}(G^0(\lambda)) \right) &= \int_{\mathbb{H}} w^{1+p}(z) d\rho(z) \\ &= \int_{\mathbb{H}^2 \times \mathbb{R}^M} w^{1+p}(\phi(z_1, z_2, q_1 \dots q_M, \lambda)) d\rho(z_1) d\rho(z_2) dv(q_1) \dots dv(q_M) \\ &= \int_{\mathbb{H}^2 \times \mathbb{R}^M} \frac{1}{2} (w^{1+p}(\phi(z_1, z_2, q_1 \dots q_M, \lambda)) + w^{1+p}(\phi(z_2, z_1, q_1 \dots q_M, \lambda))) d\rho(z_1) \\ &\hspace{15em} d\rho(z_2) dv(q_1) \dots dv(q_M) \\ &= \frac{1}{2} \int_{\mathcal{K}^c} \left(\int_{\mathbb{R}^M} \mu_{2,p}(z_1, z_2, q_1 \dots q_M, \lambda) dv(q_1) \dots dv(q_M) \right) \times (w^{1+p}(z_1) + w^{1+p}(z_2)) \\ &\hspace{15em} d\rho(z_1) d\rho(z_2) + C \\ &\leq (1 - \epsilon/2) \int_{\mathbb{H}} w^{1+p}(z) d\rho(z) + C, \end{aligned}$$

where C is some finite constant, only depending on the choice of \mathcal{K} . This implies that for all $\lambda \in R(E, \epsilon_0)$,

$$\mathbb{E} \left(w^{1+p}(G^0(\lambda)) \right) \leq \frac{2C}{\epsilon}.$$

□

Theorem 2. *Under the hypotheses of Theorem 5, we have*

$$\sup_{\lambda \in R(E, \epsilon)} \mathbb{E} \left(\left| \langle \delta_x | (H - \lambda)^{-1} | \delta_x \rangle \right|^{1+p} \right) < \infty$$

for some $\epsilon > 0$ and all $x \in \mathbb{T}$.

Proof. It is an immediate consequence of Theorem 5, Lemma 8 from the Appendix and the following inequality which holds for any two complex numbers z and s in \mathbb{H} :

$$|z| \leq 4 \frac{|z - s|^2}{\text{Im}(z)} + 2|s|. \quad (5)$$

The inequality clearly holds for $|z| \leq 2|s|$. In the complementary case, we have $|z| > 2|s|$ and thus $|z - s| \geq ||z| - |s|| \geq |s|$, implying

$$|z|\operatorname{Im}(z) \leq |z|^2 \leq 2|z - s|^2 + 2|s|^2 \leq 4|z - s|^2$$

and further $|z| \leq 4|z - s|^2/\operatorname{Im}(z)$. This proves (5).

By using (5) with $s = z_\lambda$ we obtain that for $\lambda \in R(E, \epsilon)$

$$|z| \leq 4w(z) + C,$$

where C depends only on E and ϵ . □

Now, the proof of our main theorem follows:

Theorem 1. *Let F be the largest open set included in the spectrum of Δ (this spectrum depends on the values of m and n). For any closed subinterval E , $E \subset F$, there exist $0 < p < 1$, $\eta_1 > 0$ and $\eta_2 > 0$, such that for all ν with*

$$\int_{|q| \geq \eta_1} (1 + |q|^{2(1+p)}) d\nu(q) \leq \eta_2$$

the spectrum of H is purely absolutely continuous in E with probability one.

Proof. Let us consider $\lambda = \alpha + i\beta$. Using Fatou's lemma, Fubini's theorem and Theorem 2, we obtain

$$\begin{aligned} & \mathbb{E} \left(\liminf_{\beta \searrow 0} \int_E |\langle \delta_x, (H - \lambda)^{-1} \delta_x \rangle|^{1+p} d\alpha \right) \\ & \leq \liminf_{\beta \searrow 0} \int_E \mathbb{E} \left(|\langle \delta_x, (H - \lambda)^{-1} \delta_x \rangle|^{1+p} \right) d\alpha < \infty. \end{aligned}$$

Therefore we must have

$$\liminf_{\beta \searrow 0} \int_E |\langle \delta_x, (H - \lambda)^{-1} \delta_x \rangle|^{1+p} d\alpha < \infty,$$

with probability one. Since $\langle \delta_x, (H - \lambda)^{-1} \delta_x \rangle$ is the Stieltjes transform of the measure $d\mu_x$, it follows from Proposition 7, Section 5 that the restriction of μ_x to E is purely absolutely continuous with probability one. In other words, the spectral measure for H corresponding to δ_x , for any $x \in \mathbb{T}$ is purely absolutely continuous in E with probability one. Therefore the operator H has purely absolutely continuous spectrum on E . □

4 Proof of Lemmas 3 and 4

For the proof of Lemma 3 we need the following result:

Proposition 6. For all $z_1, z_2 \in \overline{\mathbb{H}}^2 \setminus (z_\lambda, z_\lambda)$ and $\lambda \in \mathbb{R}$,

$$\mu_{2,0}(z_1, z_2, 0 \dots 0, \lambda) < 1. \quad (6)$$

Proof. For computation purposes we define

$$\begin{aligned} (1 + \lambda + z_1)(1 + \lambda + \phi_1(z_1, 0 \dots 0, \lambda)) \dots (1 + \lambda + \phi_{n-1}(z_1, 0 \dots 0, \lambda)) &= A_n \\ (1 + \lambda + z_2)(1 + \lambda + \phi_1(z_2, 0 \dots 0, \lambda)) \dots (1 + \lambda + \phi_{n-1}(z_2, 0 \dots 0, \lambda)) &= B_n \\ (1 + \lambda + z_\lambda)(1 + \lambda + \phi_1(z_\lambda, 0 \dots 0, \lambda)) \dots (1 + \lambda + \phi_{n-1}(z_\lambda, 0 \dots 0, \lambda)) &= C_n \end{aligned}$$

we also have

$$\begin{aligned} z_1 &= x_1 + iy_1 \\ z_2 &= x_2 + iy_2. \end{aligned}$$

With all these simplifications, $\mu_{2,0}(z_1, z_2, 0 \dots 0, \lambda) \leq N/D$, with

$$\begin{aligned} N = & \left((|z_1 - z_\lambda| |B_m C_m| + |z_2 - z_\lambda| |A_n C_n|)^2 (y_2 |A_m|^2 + y_1 |B_n|^2) + (|z_2 - z_\lambda| |A_m C_m| \right. \\ & \left. + |z_1 - z_\lambda| |B_n C_n|)^2 (y_1 |B_m|^2 + y_2 |A_n|^2) \right) y_1 y_2 \end{aligned}$$

and

$$D = (|C_m|^2 + |C_n|^2)(y_2 |A_m|^2 + y_1 |B_n|^2)(y_1 |B_m|^2 + y_2 |A_n|^2)(|z_1 - z_\lambda|^2 y_2 + |z_2 - z_\lambda|^2 y_1).$$

Proving $\mu_{2,0} < 1$ is equivalent to proving that the polynomial

$$\begin{aligned} P(X, Y) = & X^2 y_2 (y_1 y_2 |A_m B_m C_n|^2 + y_1 y_2 |A_n B_n C_m|^2 + y_2^2 |A_n A_m|^2 (|C_m|^2 + |C_n|^2)) + \\ & Y^2 y_1 (y_1 y_2 |A_m B_m C_n|^2 + y_1 y_2 |A_n B_n C_m|^2 + y_1^2 |B_n B_m|^2 (|C_m|^2 + |C_n|^2)) - \\ & 2XY y_1 y_2 |C_m C_n| (|A_n B_m| (y_2 |A_m|^2 + y_1 |B_n|^2) + |A_m B_n| (y_1 |B_m|^2 + y_2 |A_n|^2)) \end{aligned}$$

is strictly positive for $X = |z_1 - z_\lambda|$ and $Y = |z_2 - z_\lambda|$. This can be easily verified.

For our proof to be complete, we still have to check (6) on the boundary of $\overline{\mathbb{H}}^2$. A few cases are to be considered.

- Both z_1 and z_2 are approaching the real axis. After the first necessary blow-up:

$$\begin{aligned} y_1 &= r_1 \omega_1 \\ y_2 &= r_1 \omega_2 \end{aligned}$$

and

$$\omega_1^2 + \omega_2^2 = 1,$$

$\mu_{2,0}(z_1, z_2, 0 \dots 0, \lambda) \leq N/D$ where N and D became

$$\begin{aligned} N &= \left((|z_1 - z_\lambda| |B_m C_m| + |z_2 - z_\lambda| |A_n C_n|)^2 (\omega_2 |A_m|^2 + \omega_1 |B_n|^2) + \right. \\ &\quad \left. (|z_2 - z_\lambda| |A_m C_m| + |z_1 - z_\lambda| |B_n C_n|)^2 (\omega_1 |B_m|^2 + \omega_2 |A_n|^2) \right) \omega_1 \omega_2, \end{aligned}$$

$$\begin{aligned} D &= (|C_m|^2 + |C_n|^2) (\omega_2 |A_m|^2 + \omega_1 |B_n|^2) (\omega_1 |B_m|^2 + \omega_2 |A_n|^2) \\ &\quad (|z_1 - z_\lambda|^2 \omega_2 + |z_2 - z_\lambda|^2 \omega_1). \end{aligned}$$

Extra blow-ups are needed when $|A_i| \rightarrow 0$ or $|B_i| \rightarrow 0$, for $i = m, n$. Since these four cases are similar, we will just look at one of them:

$$\begin{aligned} |A_n|^2 &= r_2 \sin(\alpha) \\ \omega_1 &= r_2 \cos(\alpha). \end{aligned}$$

In this particular case,

$$\mu_2 \rightarrow \frac{|B_m|^2 |C_m|^2 \cos(\alpha)}{(|C_m|^2 + |C_n|^2) (\omega_2 \sin(\alpha) + |B_m|^2 \cos(\alpha))} < 1.$$

The very special case when both $|A_i|$ and $|B_j| \rightarrow 0$ does not imply any extra difficulty.

The discussion outside these four blow-up points is the same as the one inside \mathbb{H}^2 .

- Both z_1 and z_2 go to $i\infty$. For simplicity we use a change of variables, $u_1 = -\frac{1}{z_1}$ and $u_2 = -\frac{1}{z_2}$. With these new variables, in the condition $\mu_{2,0} \leq N/D$, N and D are given by

$$\begin{aligned} N &= \left((|1 + u_1 z_\lambda| |\tilde{B}_m C_m| + |1 + u_2 z_\lambda| |\tilde{A}_n C_n|)^2 (y_2 |\tilde{A}_m|^2 + y_1 |\tilde{B}_n|^2) + \right. \\ &\quad \left. (|1 + u_2 z_\lambda| |\tilde{A}_m C_m| + |1 + u_1 z_\lambda| |\tilde{B}_n C_n|)^2 (y_1 |\tilde{B}_m|^2 + y_2 |\tilde{A}_n|^2) \right) y_1 y_2, \end{aligned}$$

$$D = (|C_m|^2 + |C_n|^2)(y_2|\tilde{A}_m|^2 + y_1|\tilde{B}_n|^2)(y_1|\tilde{B}_m|^2 + y_2|\tilde{A}_n|^2) \\ (|1 + u_1 z_{,\lambda}|^2 y_2 + |1 + u_2 z_{,\lambda}|^2 y_1)$$

where

$$u_1 = x_1 + iy_1 \\ u_2 = x_2 + iy_2$$

and

$$\tilde{A}_n := u_1 A_n \\ \tilde{B}_n := u_2 B_n .$$

Since both u_1 and u_2 are approaching 0, we can write $u_1 = r \cos(\gamma)e^{ix}$ and $u_2 = r \sin(\gamma)e^{iy}$, with $\gamma, x, y \in [0, \pi/2]$. As $r \rightarrow 0$, $\mu_{2,0} \leq N/D$, with N and D given by

$$N = \left((|\tilde{B}_m C_m| + |\tilde{A}_n C_n|)^2 (\sin(\gamma) \sin(y) |\tilde{A}_m|^2 + \cos(\gamma) \sin(x) |\tilde{B}_n|^2) + \right. \\ \left. (|\tilde{A}_m C_m| + |\tilde{B}_n C_n|)^2 (\sin(\gamma) \sin(y) |\tilde{A}_n|^2 + \cos(\gamma) \sin(x) |\tilde{B}_m|^2) \right) \cdot \\ \cdot \sin(x) \sin(y) \sin(\gamma) \cos(\gamma) ,$$

$$D = (|C_m|^2 + |C_n|^2) \left((\sin(\gamma) \sin(y) |\tilde{A}_m|^2 + \cos(\gamma) \sin(x) |\tilde{B}_n|^2) \right. \\ \left. (\sin(\gamma) \sin(y) |\tilde{A}_n|^2 + \cos(\gamma) \sin(x) |\tilde{B}_m|^2) (\sin(x) \cos(\gamma) + \sin(y) \sin(\gamma)) \right) .$$

Since extra problems can appear, we also have to consider the following blow-up:

$$\sin(\gamma) = r_3 \sin(\beta) \\ \sin(x) = r_3 \cos(\beta) .$$

or similarly for $\cos(\gamma)$ and $\sin(y)$. These type of blow-ups imply $\mu_{2,0} \leq N/D$ with new values for N and D :

$$N = \left((|\tilde{B}_m C_m| + |\tilde{A}_n C_n|)^2 (\sin(\beta) \sin(y) |\tilde{A}_m|^2 + \cos(\gamma) \cos(\beta) |\tilde{B}_n|^2) + \right. \\ \left. (|\tilde{A}_m C_m| + |\tilde{B}_n C_n|)^2 (\sin(\beta) \sin(y) |\tilde{A}_n|^2 + \cos(\gamma) \cos(\beta) |\tilde{B}_m|^2) \right) \cdot \\ \cdot \cos(\beta) \sin(y) \sin(\beta) \cos(\gamma) ,$$

$$D = (|C_m|^2 + |C_n|^2)((\sin(\beta) \sin(\gamma)|\tilde{A}_m|^2 + \cos(\gamma) \cos(\beta)|\tilde{B}_n|^2) \\ (\sin(\beta) \sin(\gamma)|\tilde{A}_n|^2 + \cos(\gamma) \cos(\beta)|\tilde{B}_m|^2)(\cos(\beta) \cos(\gamma) + \sin(\gamma) \sin(\beta)).$$

Now $\mu_{2,0}$ looks similar to the previous case for $|z_1 - z_\lambda| = |z_2 - z_\lambda| = 1$.

- One of them approaches the real axis and the other one approaches $i\infty$. We will use the same substitution as in the previous case and let us assume that $u_1 \rightarrow \mathbb{R}$ and $u_2 \rightarrow 0$. We still need the first blow-up from the previous case and afterwards the discussion is very similar.

A more detailed proof of this proposition can be found in [4]. \square

Proof of Lemma 3. We can extend $\mu_{2,p}$ to an upper semi-continuous function on $\overline{\mathbb{H}^2} \times \mathbb{R}^M \times \overline{R}$ by defining it as

$$\mu_{2,p}(Z_0, Q_0, \lambda_0) = \limsup_{Z \rightarrow Z_0, Q \rightarrow Q_0, \lambda \rightarrow \lambda_0} \mu_{2,p}(Z, Q, \lambda),$$

at points Z_0, Q_0, λ_0 where it is not already defined. Here, the points $Z = (z_1, z_2)$, $Q = (q_1 \dots q_M)$ and λ are approaching their limits in the topology of $\overline{\mathbb{H}^2} \times \mathbb{R}^M \times \overline{R}$. To prove the lemma it is enough to show that

$$\mu_{2,p}(Z, Q, \lambda) < 1$$

for (Z, Q, λ) in the compact set $\partial_\infty(\overline{\mathbb{H}^2}) \times \{0\}^M \times E$, since this implies that for some $\epsilon > 0$, the upper semi-continuous function $\mu_{2,p}(Z, Q, \lambda)$ is bounded by $1 - 2\epsilon$ on the set, and by $1 - \epsilon$ in some neighborhood.

Let us rewrite $\mu_{2,p}$ in terms of $\mu_{2,0}$.

$$\mu_{2,p}(Z, Q, \lambda) = \frac{w^{1+p}(\phi(z_1, z_2, q_1 \dots q_M, \lambda)) + w^{1+p}(\phi(z_2, z_1, q_1 \dots q_M, \lambda))}{w^{1+p}(z_1) + w^{1+p}(z_2)} \\ \leq \left(\frac{w(\phi(z_1, z_2, q_1 \dots q_M, \lambda)) + w(\phi(z_2, z_1, q_1 \dots q_M, \lambda))}{w(z_1) + w(z_2)} \right)^{1+p} \cdot \frac{1}{v_1^{1+p} + v_2^{1+p}},$$

where

$$v_i = \frac{w(z_i)}{w(z_1) + w(z_2)}, \text{ for } i = 1, 2.$$

Since we are concentrating on the boundary of $\overline{\mathbb{H}}^2$, we need the following blow-up

$$\begin{aligned}\chi(z_1) &= \frac{1}{w(z_1)} = R_1 \Omega_1 \\ \chi(z_2) &= \frac{1}{w(z_2)} = R_1 \Omega_2\end{aligned}$$

where R_1 , Ω_1 and Ω_2 are defined as functions of Z with the property $\Omega_1^2 + \Omega_2^2 = 1$. Using the result in Proposition 6 we have

$$\begin{aligned}\mu_{2,p}|_{\partial_\infty(\overline{\mathbb{H}})^2 \times \{0\}^M \times E} &< \left(\mu_{2,0}|_{\partial_\infty(\overline{\mathbb{H}})^2 \times \{0\}^M \times E} \right)^{1+p} \frac{(\Omega_1 + \Omega_2)^{1+p}}{\Omega_1^{1+p} + \Omega_2^{1+p}} \\ &\leq (1 - \epsilon) 2^p \\ &< 1,\end{aligned}$$

for the right choice of p . □

Proof of Lemma 4. Each term in the sum appearing in $\mu_{2,p}$ can be estimated

$$\begin{aligned}\frac{w^{1+p}(\phi(z_1, z_2, q_1 \dots q_M, \lambda))}{w^{1+p}(z_1) + w^{1+p}(z_2)} &= \frac{(w(z_1) + w(z_2))^{1+p}}{w^{1+p}(z_1) + w^{1+p}(z_2)} \left(\frac{w(\phi(z_1, z_2, q_1 \dots q_M, \lambda))}{w(z_1) + w(z_2)} \right)^{1+p} \\ &\leq 2^p \left(\frac{w(\phi(z_1, z_2, q_1 \dots q_M, \lambda))}{w(z_1) + w(z_2)} \right)^{1+p}.\end{aligned}$$

Now it is enough to prove

$$\frac{w(\phi(z_1, z_2, q_1 \dots q_M, \lambda))}{w(z_1) + w(z_2)} \leq C \prod_{k=1}^M (1 + |q_k|^2),$$

since this bounds each term in $\mu_{2,p}$ by the desired quantity.

We have

$$\begin{aligned}
& \frac{w(\phi(z_1, z_2, q_1 \dots q_M, \lambda))}{w(z_1) + w(z_2)} \\
&= \frac{|1 + z_\lambda \phi_n(z_1, q_1 \dots q_n, \lambda) + z_\lambda \phi_m(z_2, q_{n+1} \dots q_{n+m}, \lambda) + z_\lambda(\lambda - q_M)|^2}{\operatorname{Im}(\phi_n(z_1, q_1 \dots q_n, \lambda)) + \operatorname{Im}(\phi_m(z_2, q_{n+1} \dots q_{n+m}, \lambda)) + \operatorname{Im}(\lambda)} \\
& \cdot \frac{1}{\sum_{i=1}^2 \frac{|z_i - z_\lambda|^2}{\operatorname{Im}(z_i)}} \\
&= \frac{|A_n B_m - z_\lambda A_{n-1} B_m - z_\lambda B_{m-1} A_n + A_n B_m z_\lambda(\lambda - q_M)|^2}{\operatorname{Im}(z_1 + \lambda)|B_m|^2 + \operatorname{Im}(z_2 + \lambda)|A_n|^2 + \operatorname{Im}(\lambda)|A_n B_m|^2} \cdot \frac{1}{\sum_{i=1}^2 \frac{|z_i - z_\lambda|^2}{\operatorname{Im}(z_i)}} \\
&\leq (1 + |z_\lambda|^2) \frac{|A_n B_m|^2 + |A_{n-1} B_m + A_n B_{m-1} - A_n B_m(\lambda - q_M)|^2}{\operatorname{Im}(z_1 + \lambda)|B_m|^2 + \operatorname{Im}(z_2 + \lambda)|A_n|^2 + \operatorname{Im}(\lambda)|A_n B_m|^2} \cdot \frac{1}{\sum_{i=1}^2 \frac{|z_i - z_\lambda|^2}{\operatorname{Im}(z_i)}} \\
&\leq (1 + |z_\lambda|^2) \left(\frac{|A_n B_m|^2}{\operatorname{Im}(z_1)|B_m|^2 + \operatorname{Im}(z_2)|A_n|^2} + (|A_n B_m|^2 + |A_{n-1} B_m + A_n B_{m-1}|^2) \cdot \frac{1}{\operatorname{Im}(z_1)|B_m|^2 + \operatorname{Im}(z_2)|A_n|^2} \right) \cdot \frac{1}{\sum_{i=1}^2 \frac{|z_i - z_\lambda|^2}{\operatorname{Im}(z_i)}} \\
&\leq (1 + |z_\lambda|^2) \left(\frac{|A_n B_m|^2(2 + |q_M - \lambda|^2)}{\operatorname{Im}(z_1)|B_m|^2 + \operatorname{Im}(z_2)|A_n|^2} + 2 \left(\frac{|A_{n-1}|^2}{\operatorname{Im}(z_1)} + \frac{|B_{m-1}|^2}{\operatorname{Im}(z_2)} \right) \right) \cdot \frac{1}{\sum_{i=1}^2 \frac{|z_i - z_\lambda|^2}{\operatorname{Im}(z_i)}} \\
&\leq (1 + |z_\lambda|^2) \left(\frac{|A_n|^2(2 + |q_M - \lambda|^2)}{\operatorname{Im}(z_1)} + 2 \left(\frac{|A_{n-1}|^2}{\operatorname{Im}(z_1)} + \frac{|B_{m-1}|^2}{\operatorname{Im}(z_2)} \right) \right) \cdot \frac{1}{\sum_{i=1}^2 \frac{|z_i - z_\lambda|^2}{\operatorname{Im}(z_i)}}
\end{aligned}$$

Using the following inequality (details of its proof can be found in [4]),

$$\begin{aligned}
|A_n|^2 &\leq (n-1)^2 \prod_{\substack{k=1 \\ |1+\lambda-q_k|>1}}^n |1 + \lambda - q_k|^2 |1 + z_1|^2 \\
&\leq (n-1) \prod_{k=1}^n (1 + |q_k|^2)(1 + |z_1|^2)
\end{aligned}$$

we obtain

$$\begin{aligned}
\frac{w(\phi(z_1, z_2, q_1 \dots q_M, \lambda))}{w(z_1) + w(z_2)} &\leq C \left(C_1 (1 + |q_M|^2) \prod_{k=1}^n (1 + |q_k|^2) \frac{1 + |z_1|^2}{\text{Im}(z_1)} \right. \\
&\quad + C_2 (1 + |q_M|^2) \prod_{k=1}^{n-1} (1 + |q_k|^2) \frac{1 + |z_1|^2}{\text{Im}(z_1)} \\
&\quad \left. + C_3 (1 + |q_M|^2) \prod_{k=1}^{m-1} (1 + |q_k|^2) \frac{1 + |z_2|^2}{\text{Im}(z_2)} \right) \cdot \frac{1}{\sum_{i=1}^2 \frac{|z_i - z_\lambda|^2}{\text{Im}(z_i)}}
\end{aligned}$$

Choose the compact set \mathcal{K} such that $\sum_{i=1}^2 |z_i - z_\lambda|^2 / \text{Im}(z_i) \geq C > 0$ for some constant C and $(z_1, z_2) \in \mathcal{K}^c$. Then we can estimate each term depending on whether z_i is close to z_λ . If all z_j are sufficiently close, then $\text{Im}(z_j)$ is bounded below and $|z_j|$ is bounded above by a constant. Thus

$$\text{Im}(z_j) \sum_{i=1}^2 |z_i - z_\lambda|^2 / \text{Im}(z_i) \geq \text{Im}(z_j) C \geq C' > 0$$

and $1 + |z_j|^2 \leq C$, so we are done. If all z_j are far from z_λ ,

$$\text{Im}(z_j) \sum_{i=1}^2 |z_i - z_\lambda|^2 / \text{Im}(z_i) \geq |z_j - z_\lambda|^2 \geq C(1 + |z_j|^2)$$

so $1 + |z_j|^2 / \left(\text{Im}(z_j) \sum_{i=1}^2 |z_i - z_\lambda|^2 / \text{Im}(z_i) \right) \leq C$ again. \square

5 Appendix

This section contains two theorems on absolutely continuous spectrum. The first one gives a sufficient condition for a measure to be absolutely continuous with respect to the Lebesgue measure on an interval and the latter gives a sufficient condition for a random Schrödinger operator to exhibit pure absolutely continuous spectrum on some interval.

5.1 A Criterion for Absolutely Continuous Spectrum

Let μ be a finite measure on \mathbb{R} ; its Stieltjes (or Borel) transform F is given by

$$F(z) = \int \frac{d\mu(t)}{t - z}$$

for $z = x + iy$ with $y > 0$. The following criterion has been proven in [7]. We reproduce the argument here for completeness.

Proposition 7. *Let (a, b) be a finite interval and let $p > 0$. Suppose*

$$\liminf_{y \searrow 0} \int_a^b |F(x + iy)|^{1+p} dx < \infty.$$

Then μ is absolutely continuous with respect to the Lebesgue measure on (a, b) .

Proof. Since $\liminf_{y \searrow 0} \int_a^b |F(x + iy)|^{1+p} dx < \infty$, there exists a sequence $y_n \searrow 0$ such that $\sup_n \int_a^b |F(x + iy_n)|^{1+p} dx < C$, where C is some constant. Define $d\mu_{y_n}(x) = \pi^{-1} \text{Im}(F(x + iy_n)) dx$. Then by [6], $d\mu_{y_n} \rightarrow d\mu$ weakly, as $n \rightarrow \infty$. That is, for f a continuous function of compact support we have $\lim_{n \rightarrow \infty} \int f(x) d\mu_{y_n}(x) = \int f(x) d\mu(x)$. Let f be a continuous function supported on (a, b) , then

$$\begin{aligned} \left| \int_a^b f(x) d\mu(x) \right| &= \lim_{n \rightarrow \infty} \left| \int_a^b f(x) d\mu_{y_n}(x) \right| \\ &= \lim_{n \rightarrow \infty} \pi^{-1} \left| \int_a^b f(x) \text{Im}(F(x + iy_n)) dx \right| \\ &\leq \lim_{n \rightarrow \infty} \left(\|f\|_{1+1/p} \| \text{Im}(F(x + iy_n)) \|_{1+p} \right) \\ &\leq C \|f\|_{1+1/p}. \end{aligned}$$

5.2 Bounds on the Green's function at an arbitrary site.

We show here how to obtain bounds on $G^x(\lambda)$ for an arbitrary site $x \in \mathbb{T}$, using the bound on $G^0(\lambda)$ given by Theorem 5.

Lemma 8. *Let F be the largest open set included in the spectrum of Δ . Suppose that*

$$\sup_{\lambda \in R(E, \epsilon)} \mathbb{E} \left(w^{1+p} \left(\langle \delta_0, (H - \lambda)^{-1} \delta_0 \rangle \right) \right) < \infty,$$

for some closed subinterval $E \subset F$, $\epsilon > 0$ and $0 < p < 1$. Then, for every arbitrary $x \in \mathbb{T}$, we also have

$$\sup_{\lambda \in R(E, \epsilon)} \mathbb{E} \left(w^{1+p} \left(\langle \delta_x, (H - \lambda)^{-1} \delta_x \rangle \right) \right) < \infty .$$

Proof. Suppose we pick an arbitrary vertex x in \mathbb{T} and we consider the Green's function at x for the whole tree \mathbb{T} : $G_x(\lambda) = \langle \delta_x, (H - \lambda)^{-1} \delta_x \rangle$. If we consider x to be the new origin of the tree and we look at the vertices in the future of x , we can see that after a finite number of steps, on each branch, we will run into a copy of the original tree. For all these copies of \mathbb{T} , we know from the hypothesis that

$$\sup_{\lambda \in R(E, \epsilon)} \mathbb{E} \left(w^{1+p} \left(\langle \delta_0, (H - \lambda)^{-1} \delta_0 \rangle \right) \right) < \infty .$$

Starting with these copies of \mathbb{T} , we can go back, step by step, and prove the inequality for the truncated Green's function at each subsequent previous vertex.

When tracking back the Green's function we can have either the Möbius transformation ϕ_1 (see (2)), for the step by step recurrence at a vertex of degree 2, or $\phi'(z_1, z_2, \lambda) = -1/(z_1 + z_2 + \lambda - q)$ at a vertex of degree 3. Therefore, using the proofs of Lemma 3 and Lemma 4, we can easily conclude that our inequality holds step by step:

I. At a vertex y of degree 2:

$$\begin{aligned} \mathbb{E} \left(w^{1+p} (G^y(\lambda)) \right) &= \int_{\mathbb{H}} w^{1+p}(z) d\rho'(z) = \int_{\mathbb{H} \times \mathbb{R}} w^{1+p}(\phi_1(z_1, q_1, \lambda)) d\rho(z_1) dv(q_1) \\ &= \frac{1}{2} \int_{\mathcal{K}^c} \left(\int_{\mathbb{R}} \frac{w^{1+p}(z_1, q_1, \lambda)}{w^{1+p}(z_1)} dv(q_1) \right) \times w^{1+p}(z_1) d\rho(z_1) + C \\ &\leq C' \int_{\mathbb{H}} w^{1+p}(z_1) d\rho(z_1) + C \\ &\leq C' \mathbb{E} \left(w^{1+p} (G(\lambda)) \right) + C , \end{aligned}$$

hence $\sup_{\lambda \in R(E, \epsilon)} \mathbb{E} \left(w^{1+p} (G_y(\lambda)) \right) < \infty$.

2. At a vertex y of degree 3:

$$\begin{aligned}
\mathbb{E}\left(w^{1+p}(G^y(\lambda))\right) &= \int_{\mathbb{H}} w^{1+p}(z) d\rho''(z) \\
&= \int_{\mathbb{H}^2 \times \mathbb{R}} w^{1+p}(\phi'(z_1, z_2, q, \lambda)) d\rho(z_1) d\rho(z_2) d\nu(q_1) \\
&= \int_{\mathbb{H}^2 \times \mathbb{R}} \frac{1}{2} (w^{1+p}(\phi'(z_1, z_2, q_1, \lambda)) + w^{1+p}(\phi'(z_2, z_1, q_1, \lambda))) d\rho(z_1) d\rho(z_2) d\nu(q_1) \\
&= \frac{1}{2} \int_{\mathcal{K}^c} \left(\int_{\mathbb{R}} \frac{w^{1+p}(\phi'(z_1, z_2, q_1, \lambda)) + w^{1+p}(\phi'(z_2, z_1, q_1, \lambda))}{w^{1+p}(z_1) + w^{1+p}(z_2)} d\nu(q_1) \right) \times (w^{1+p}(z_1) + \\
&\quad + w^{1+p}(z_2)) d\rho(z_1) d\rho(z_2) + C \\
&\leq C'' \int_{\mathbb{H}} w^{1+p}(z) d\rho(z) + C \\
&\leq C'' \mathbb{E}\left(w^{1+p}(G(\lambda))\right) + C,
\end{aligned}$$

hence $\sup_{\lambda \in R(\bar{E}, \epsilon)} \mathbb{E}\left(w^{1+p}(G_y(\lambda))\right) < \infty$. □

References

- [1] M. Aizenman and R. Sims and S. Warzel. Stability of the Absolutely Continuous Spectrum of Random Schrödinger Operators on Tree Graphs. *Prob. Theor. Rel. Fields*, (136):363-394, 2006.
- [2] R. Froese and D. Hasler and W. Spitzer. Transfer matrices, hyperbolic geometry and absolutely continuous spectrum for some discrete Schrödinger operators on graphs. *J. Func. Anal.*, (230):184-221, 2006.
- [3] R. Froese and D. Hasler and W. Spitzer. Absolutely Continuous Spectrum for the Anderson Model on a Tree: A Geometric Proof of Klein's Theorem. *Commun. Math. Phys.*, (269):239-257, 2007.
- [4] F. Halasan. On the Anderson Model. University of British Columbia. In preparation.
- [5] F. Klein. Extended States in the Anderson Model on the Bethe Lattice. *Advances in Math.*, (133):163-184, 1998.

- [6] B. Simon. Spectral analysis of rank one perturbations and applications. *Mathematical quantum theory. II. Schrödinger operators (Vancouver, BC, 1993) C.R.M. Proc. Lecture Notes, Amer. Math. Soc., Providence, RI,* (8):109-149, 1995.
- [7] B. Simon. Lp norms of the Borel transform and the decomposition of measures. *Proceedings of the American Mathematical Society*, 123(12):3749-3755, Dec. 1995.