

Everett and the Born Rule

Alastair I. M. Rae

*School of Physics and Astronomy, University of Birmingham
Birmingham B15 2TT*

Abstract

During the last ten years or so, derivations of the Born rule based on decision theory have been proposed and developed, and it is claimed that these are valid in the context of the Everett interpretation. This claim is critically assessed and it is shown that one of its key assumptions, although natural in the context of the Copenhagen interpretation, is not consistent with that of Everett. It is further argued that any interpretation that relates outcome likelihood to the expansion coefficients connecting the wavefunction with the eigenfunctions of the measurement operator must be inconsistent with the Everett interpretation.

Key words: Everett, Born Rule, Quantum Measurement

Everett and the Born Rule

Alastair I. M. Rae

*School of Physics and Astronomy, University of Birmingham
Birmingham B15 2TT*

Introduction

The conventional (“Copenhagen”) interpretation of quantum mechanics states that the state of a system following a measurement is one (and only one) of the eigenvalues belonging to the operator representing the measurement and that, as a result of the measurement, the wavefunction “collapses” to become the corresponding eigenfunction. According to the “Born rule”, the probability of any particular outcome is proportional to the squared modulus of the scalar product of this eigenfunction with the pre-measurement wavefunction. This analysis underlies many of the predictions of quantum mechanics that have been invariably confirmed by experiment. An alternative approach to quantum measurement is the Everett interpretation (also known as the “relative states” or the “many worlds” interpretation) which was proposed by Everett III (1957). The essence of this approach is that it assumes no collapse of the wavefunction associated with a measurement: instead, the time development of the state is everywhere governed by the time-dependent Schrödinger equation. After a “measurement-like” event, this results in a splitting of the wavefunction into a number of branches, which are then incapable of reuniting or communicating with each other in any way. This splitting occurs even when a human observer is part of the measurement chain: the resulting branches then each contain a copy of the observer, who is completely unaware of the existence of the others.

Since its inception, the Everett interpretation has been subject to considerable criticism—e.g. Kent (1990), Squires (1990)—which has three main strands (or branches [sic]). First, there is its metaphysical extravagance. The continual evolution of the universe into a “multiverse” containing an immense number of branches would mean that the universe we observe should be accompanied by an immense number of parallel universes, which we do not

observe and have no awareness of—surely such a postulate must be a gross breach of the principle of Occam’s razor! Everett himself was aware of this criticism and, in a footnote to his original paper, he compares the conceptual difficulties of accepting his interpretation with those encountered by Copernicus when the latter proposed the (in his time revolutionary) idea that the earth moves around the sun. However, the reason that the Occam’s razor argument has not led to the universal rejection of Everett’s ideas is less to do with the strength or otherwise of the Copernican analogy and more to do with the fact that the branching of the universe into the multiverse is claimed to be a direct consequence of the time-dependent Schrödinger equation: no additional postulate, such as the collapse of the wavefunction, is required to explain the phenomenon of quantum measurement and the extravagance with universes may therefore be considered a price worth paying for the economy in postulates.

The second strand in the criticism of Everett is known as the “preferred basis” problem. This is because there is an apparent ambiguity in the way the branches are defined. Thus, if the wavefunction of a system has the form $\psi = A\psi_1 + B\psi_2$, then Everett suggests that a measurement should lead to two sets of branches, one associated with each of the statevectors ψ_1 and ψ_2 . However, the original state could just as well be written as $\psi = C\phi_1 + D\phi_2$ where $\phi_1 = 2^{-1/2}(\psi_1 + \psi_2)$, $\phi_2 = 2^{-1/2}(\psi_1 - \psi_2)$, $C = 2^{-1/2}(A + B)$ and $D = 2^{-1/2}(A - B)$, so why should the branches not be just as well defined by ϕ_1 and ϕ_2 —or indeed any other orthogonal pair of linear combinations of ψ_1 and ψ_2 . This problem has been largely resolved by the appreciation of the importance of the effect of the environment on a quantum system and the associated “decoherence”—Zurek (2008), Wallace (2002), Wallace (2003). A quantum measurement is inevitably accompanied by complex, chaotic processes which act to pick out the particular basis defined by the eigenstates of the measurement operator. This basis is then the one “preferred” by the Everett interpretation and this supervenes on the Schrödinger wavefunction. This result is now generally accepted, although Baker (2006) argues that its derivation uses the Born rule so that there is a danger of circularity if it is then assumed as part of its proof.

The third criticism leveled at Everett is the problem of probabilities. The conventional (Copenhagen) interpretation states that, if the wavefunction before a measurement is $\psi = A\psi_1 + B\psi_2$, and if ψ_1 and ψ_2 are eigenstates of the measurement operator with eigenvalues q_1 and q_2 respectively, then the outcome will be *either* q_1 with probability $|A|^2$ *or* q_2 with probability $|B|^2$,

where these probabilities reflect the frequencies of the corresponding outcomes after a large number of similar measurements. However, according to the Everett approach there is no “either-or” because both outcomes are manifest, albeit in different branches. Instead of a *disjunction* to which we can apply standard probability theory, we have a *conjunction*, where it is hard to see how probabilities can make any sense—Squires (1990), Graham (1973), Lewis (2004). There have been several attempts to resolve this conundrum and to show how probability (or something else that is in practice equivalent to it) can be used in an Everettian context. Wallace (2007) has proposed a principle that he calls “subjective uncertainty” in which he claims that a rational observer should expect to emerge in one branch after a measurement, even although she is also reproduced in the other branches. Greaves (2004) has criticized this approach and suggested an alternative in which we have to take into account the observer’s “descendants” in all the branches, but we should “care” more about some than others; the extent to which we should care is quantified by a “caring measure” that is proportional to the corresponding Born-rule weight. Both these approaches are designed to explain why some branches appear to be favoured over others, but both attempt to do this without altering Everett’s main principle that the quantum state evolves under the influence of the time-dependent Schrödinger equation with nothing else added, so that the Born rule supervenes on this. An alternative approach, which I shall not discuss any further in this paper, is to maintain most of the fundamental ideas of the Everettian interpretation, but add a further layer of “reality” to justify the use of probabilities; an example of this can be found in Lockwood (1989).

Interest in the Everett interpretation has been on the increase recently—particularly during 2007, which was the 50th anniversary of the publication of Everett’s original paper Everett III (1957). Much of the renewed interest has developed from work by Deutsch (1999) some eight years earlier, which was then developed by Saunders (2004) and Wallace (2007). This programme (which I refer to below by the initials DSW) aims to derive the Born rule from minimal postulates that are consistent with the Everett interpretation, as well as with other approaches to the measurement problem. A quite similar approach was developed in parallel with this by Zurek and is set out in a recent review paper Zurek (2008). The present paper aims to show that some of the postulates underlying these arguments do not follow from the Everett interpretation and may well not be consistent with it.

The DSW Proof of the Born Rule

This section sets out the DSW derivation of the Born rule by applying it to a particular example. The argument is deliberately kept as simple as possible and more general treatments can be found in the cited references. Consider the case of a spin-half particle, initially in an eigenstate of an operator representing a component of spin in a direction in the xz plane at an angle θ to the z axis, passing through a Stern-Gerlach apparatus oriented to measure a spin component in the z direction.

Standard quantum mechanics tells us that the initial state α_θ can be written as a linear combination of the eigenstates of \hat{S}_z : α with eigenvalue $+1$ (in units of $\hbar/2$) and β with eigenvalue -1 . We have

$$\alpha_\theta = c\alpha + s\beta \tag{1}$$

where $c = \cos(\theta/2)$ and $s = \sin(\theta/2)$. Particles emerge from the two channels of the Stern-Gerlach apparatus, with the upper and lower channels indicating $S_z = +1$ and -1 respectively and are then detected. After they have entered the detectors, but before any collapse¹ associated with the measurement, the total wavefunction of the system is

$$\psi = c\alpha\chi_+ + s\beta\chi_- \tag{2}$$

where χ_+ (χ_-) is the wavefunction of the detectors, including their environment, when a particle is detected in the positive (negative) channel. According to the Copenhagen interpretation, the corresponding probabilities for a positive or negative outcome are given by the Born rule as c^2 and s^2 respectively. From the Everettian point of view, on the other hand, there is no collapse and the system is always in a state of the form ψ . However, because of the effects of the environment and decoherence, phase coherence between the two terms on the right-hand side of (2) is lost, so they can never in practice interfere. The wavefunction has therefore evolved into two “branches” which then develop independently.

¹At a number of points in this paper, I compare the predictions of the Everett model with those produced by the “Copenhagen interpretation”, by which I mean a model in which the wavefunction collapses into one of the eigenstates of the measurement operator. This is assumed to occur early enough in the process for the outcomes to be the same as would be observed if particles were emerge randomly from one or other output channel, with the relative probabilities of the two outcomes determined by the Born rule.

The principle of the DSW approach is to describe the process being studied as a game, or series of games, where we receive rewards, or pay penalties (i.e. receive negative rewards) depending on the outcomes. The derivation proposed by Zurek (2008) is quite similar to this, although it does not use game theory.

Imagine a game where the player receives a reward depending on the outcome of the experiment. Assume that the value of θ is under our control and that, whenever the experimenter observes a particle emerging from the positive or negative channel of the Stern-Gerlach apparatus, she receives a reward equal to x_+ or x_- respectively; these values can be chosen arbitrarily by the experimenter. In the special cases where $\theta = 0$ or $\theta = \pi$, the initial spin state is an eigenstate of \hat{S}_z with eigenvalues $+1$ and -1 respectively. The particle then definitely emerges from the corresponding channel of the Stern-Gerlach apparatus and the corresponding reward is paid.

In the general case, we define the “value”— $V(\theta)$ —of the game as the minimum payment a rational player would accept not to play the game, and look for an expression for $V(\theta)$ of the form

$$V(\theta) = w_+(\theta)x_+ + w_-(\theta)x_- \quad (3)$$

where the w s are non-negative real numbers that we call “weights” and which are normalized so that their total is unity. We shall find that

$$w_+(\theta) = c^2 \text{ and } w_-(\theta) = s^2 \quad (4)$$

which are the probabilities predicted by the Born rule for this setup.

First consider the effect on the wavefunction of rotating the SG magnet through 180° about the y axis. It follows from the symmetry of the Stern-Gerlach apparatus that spins that were previously directed into the upper channel will now be detected in the lower channel and vice versa. Thus

$$V(\theta + \pi) = w_+(\theta + \pi)x_+ + w_-(\theta + \pi)x_- = w_-(\theta)x_+ + w_+(\theta)x_- \quad (5)$$

From standard quantum mechanics, the effect of this rotation on the wavefunction (2) is to transform it to

$$\psi = -s\alpha\chi_+ + c\beta\chi_- \quad (6)$$

We now proceed by considering a series of particular values of θ .

Case 1 The first case is where $\theta = 0$ so that the initial state, α_θ , is identical with α . As noted above, this state is unaffected by the measurement and the particle is always detected in the positive channel. Thus $V(0) = x_+$, $w_+(0) = 1$ and $w_-(0) = 0$. Similarly, $V(\pi) = x_-$, $w_+(\pi) = 0$ and $w_-(\pi) = 1$.

Case 2 In the second case, $\theta = \pi/2$ so that ψ is as in (2) above, but with $c = s = 2^{-1/2}$. Now consider the effect of rotating the Stern Gerlach apparatus through an angle π . Using (5) and (6), we get the following expressions for V and ψ

$$V(3\pi/2) = w_-(\pi/2)x_+ + w_+(\pi/2)x_- \quad (7)$$

$$\psi = 2^{-1/2}[-\alpha\chi_+ + \beta\chi_-] \quad (8)$$

The only change in the wavefunction is the change of sign in the term involving α . DSW point out that this sign, in common with any other phase factor, should not affect the value, because it can be removed by performing a unitary transformation on this part of the wavefunction only—e.g. by a rotation of the spin through 2π or by introducing an additional path length equal to half a wavelength. Moreover, Zurek (2008) shows that one of the effects of the interaction of the system with the environment is to remove any physical significance from these phase factors. It follows that the value should not be affected by the rotation so that $V(3\pi/2) = V(\pi/2)$, which leads directly to

$$w_+(\pi/2) = w_-(\pi/2) = 1/2 \text{ and } V(\pi/2) = (x_+ + x_-)/2 \quad (9)$$

This result, which might be thought to be an inevitable consequence of the symmetry of the system, is considered by DSW to be the key point of the proof. We should note that, although it agrees with the Born rule, it would also be consistent with any alternative weighting scheme that predicted equal weights in this symmetric situation; in particular it is consistent with a model in which the weights are assumed to be independent of θ .

We now extend the result to the case where the number of output channels is M instead of two and the wavefunction is the sum of M terms, each of which corresponds to a different eigenstate of the measurement operator. In the case where the coefficients of this expansion are all equal, any action that has the effect of exchanging any two output channels (which are numbered 1 and 2) must leave the wavefunction unchanged apart from irrelevant changes in phase. The value is then also unchanged, but the roles of w_1 and w_2 are reversed. Hence

$$w_1x_1 + w_2x_2 = w_1x_2 + w_2x_1 \quad (10)$$

where x_i is the reward associated with the i th output channel. It follows that $w_1 = w_2$; consideration of other permutations immediately extends this result to all i and we have $w_i = N^{-1}$.

Case 3 In the third case, $\theta = \pi/3$ so that $\cos(\theta/2) = \sqrt{3}/2$ and $\sin(\theta/2) = 1/2$. We now assume that the system is modified so that, after emerging from the Stern-Gerlach magnet and before being detected, the outgoing particles interact with a separate quantum system that can exist in one of, or a linear combination of four eigenstates ϕ_i . Following Zurek (2008), this is referred to as an ‘‘ancilla’’ from now on. The ancilla is designed so that, if $\theta = 0$ so that all spins emerge from the positive channel, the ancilla is placed in the state $3^{-1/2} \sum_{i=1,3} \phi_i$; while, if $\theta = \pi$ and all spins are negative, its state becomes ϕ_4 . From linear superposition it follows that if the original spin is in a state of the form (2) with $\theta = \pi/3$, the total wavefunction of the spin plus the ancilla is

$$\begin{aligned} \Psi &= 3^{-1/2}[\phi_1 + \phi_2 + \phi_3] \cos(\pi/6)\alpha + \phi_4 \sin(\pi/6)\beta \\ &= \frac{1}{2}[\phi_1\alpha + \phi_2\alpha + \phi_3\alpha + \phi_4\beta] \end{aligned} \quad (11)$$

As the coefficients of each term in the above expansion are equal, it follows from the earlier discussion of Case 2 that all four weights are equal to 0.25. If we were to measure on the ancilla a quantity whose eigenstates were one of the functions ϕ_1 to ϕ_4 , we should obtain a result equal to one of the corresponding eigenvalues. If the result corresponds to one of the first three eigenfunctions, we can conclude that if, instead, we had measured the spin directly, we would have got a positive result, while a result corresponding to ϕ_4 indicates a negative spin. As this is the only such state, it follows that the weight corresponding to a negative spin is $w_-(\pi/3) = 0.25$ and therefore, from normalization, that $w_+(\pi/3) = 0.75$. (The last step, which follows Zurek (2008), establishes these results without assuming that the weights are additive.) The value of the game therefore equals $0.75x_+ + 0.25x_-$. It can also be shown quite straight forwardly—Zurek (2008)—that, after a number of repeats of the experiment, the predicted distribution of the results is as observed experimentally.

Following DSW, the above argument can be extended to the case of a measurement made in the absence of the ancilla if we make a further assumption,

known as “measurement neutrality”. This states that the outcome of the game is independent of the details of the measurement process—i.e. the presence or absence of the ancilla—so that $w_+(\pi/3) = 0.75$ and $w_-(\pi/3) = 0.25$ in either case. These quantities are identical to $\cos^2(\theta/2)$ and $\sin^2(\theta/2)$ respectively, so the derived weights are the same as those predicted by the Born rule. By choosing an appropriate ancilla, the above argument can be directly extended to examples where the ratio of the weights is any rational number, and then to the general case by assuming that the weights are continuous functions of θ . Hence, the expression for the value is the same as that predicted by the Born-rule:

$$V(\theta) = c^2x_+ + s^2x_- \tag{12}$$

Further generalization to experiments with more than two possible outcomes is reasonably straightforward and does not introduce any major new principles.

Discussion

There have been a number of criticisms of the DSW proof when applied to the Everett model in particular—e.g. Baker (2006), Barnum et al. (2000), Lewis (2005), Lewis (2006), Hemmo and I (2007); some of these even challenge the result (9) for the symmetric case. I shall shortly develop arguments to show that, although the symmetric results appear to be correct, this is not so for the asymmetric case.

First consider how the above translates into predictions of experimental results. The game value is the minimum payment a rational observer would accept in order not to play the game. This means that after playing the game a number of times, a rational observer should expect to receive a set of rewards whose average is equal to the game value. Thus, if we consider a sequence of N such observations in which n_+ and $n_-(= N - n_+)$ particles are detected in the positive and negative channels respectively, the total reward received will be $n_+x_+ + n_-x_-$, and this should equal $N(w_+x_+ + w_-x_-)$ implying that $w_+ = n_+/N$ and $w_- = n_-/N$. This, of course, is just what is observed in a typical experiment provided N is large enough for statistical fluctuations to be negligible. It should be noted that frequencies are *not* being used to *define* probabilities, but the derived weights are used to predict the results of experimental measurement of the frequencies.

The above results are of course consistent with the standard Copenhagen interpretation, whose fundamental mantra was set out by Bohr: “...there is essentially the question of an influence on the very conditions which define the possible types of predictions regarding the future behaviour of the system”. In the present context, this means that, because an experiment designed to demonstrate interference would involve a different experimental arrangement, the experiment can be modelled as a classical stochastic system in which spins emerge from *either* the positive *or* the negative channel of the Stern-Gerlach apparatus. It should be noted that this paper does not aim to justify the Copenhagen interpretation, but employs its results as a comparator with the Everettian case.

Why should an Everettian observer have experiences such as those just described? In the Everettian interpretation, the quantum state evolves deterministically and on first sight, there would appear to be no room for uncertainty. However, after a splitting has occurred, observers in different branches have the same memories of their state before the split, but undergo different experiences after it. Given this, it may be meaningful for an experimenter to have an opinion about the likelihood of becoming a particular one of her successors. This introduces a form of subjective uncertainty, and DSW claim that this plays a role in the Everett interpretation that is equivalent to that played by objective stochastic uncertainty in the Copenhagen case. However, we should note that such subjective uncertainty can only come into play at the point where the experimenter becomes aware of an experimental result, in contrast to the Copenhagen model where the splitting is assumed to occur as the particles emerge from the Stern-Gerlach magnet. I shall shortly proceed to compare and contrast the Copenhagen and Everettian interpretations of the different experiments discussed above. To help focus the discussion, I shall initially assume that in such experiments each particular result is associated with only one branch of the final wavefunction. This assumption has been strongly criticized by DSW and others and I shall return to the question of how it affects our conclusions at a later stage. I now analyze our earlier arguments step by step.

Case 1: Copenhagen As the initial spin state is in an eigenstate of S_z , the result is completely determined. The probability of the result equalling the corresponding eigenvalue is 1 and the probability of the alternative is zero.

Case 1: Everett There is only one branch and this contains the only copy of the observer who invariably records the appropriate eigenvalue.

There is therefore no difference between the observers' experiences in case 1 under the Copenhagen and Everettian interpretations.

Case 2: Copenhagen The probabilities of positive and negative results are both 0.5. After a number of repeats of the experiment, the experimenter will expect to have recorded equal numbers of positive and negative results, so her average reward will be $(x_1 + x_2)/2$, which is the same as the game value.

Case 2: Everett The observer will split into two copies each time a spin is observed and the weights of the two branches are equal for the reasons discussed earlier. After N repeats each observer can therefore expect to have observed $N/2$ positive and $N/2$ negative results and their average rewards will both equal the game value.

There is therefore no difference between the observers' experiences in case 2 under the Copenhagen and Everettian interpretations.

Case 3: Copenhagen As emphasized above, this assumes that the experiment is a stochastic process in which a particle emerges from either the positive and or the negative channel and the relative probabilities of the outcomes are equal to the Born weights. In the presence of the ancilla, a particle is detected in one (and only one) of the equally-weighted states ϕ_1 to ϕ_4 , so all four outcomes have equal probability. To have been observed in any of the first three of these, the spin must have emerged from the Stern-Gerlach experiment through the positive channel, while if the final result was ϕ_4 , it must have come from the negative channel. It follows directly that if the ancilla were absent, three times as many spins would be detected as positive than as negative. Thus, the principle of measurement neutrality, assumed in stage 3 of the earlier derivation, follows from the assumptions underlying the Copenhagen interpretation.

Case 3: Everett We first consider the situation where an ancilla is present so that the state is described by (11); there are therefore four equally-weighted branches, one corresponding to each of the ϕ_i . The observer splits into four equally-weighted copies and should expect her descendants to record an equal number of each of the four possible results and therefore conclude that there are three times as many positive as negative spins. However, in the absence of an ancilla, there are only two branches and the observer is split into two copies each time a result is obtained. To show that a typical Everettian observer should record results that are consistent with the Born weights, we should again apply the principle of measurement neutrality. However, the arguments that led to this in the Copenhagen case, do not now apply as I




Figure 1: In the cop mode a ball is emitted from the first box through either the upper or the lower port and detected either before or after entering the second box; the figure shows one possible outcome. In the eve mode, balls emerge from both ports and one of them is detected either before or after the second box, which releases three balls every time one enters.

shall now show.

We saw above that measurement neutrality holds under the Copenhagen interpretation because it assumes that the particles emerge from *either* the positive *or* the negative channel and then into one, *and only one*, of the states ϕ_i . This is not true in the case of the Everett interpretation, where the system evolves deterministically and the state is described by a linear combination of the wavefunctions associated with the different channels. Apparent stochasticity, or subjective uncertainty, only enters the situation when the experimenter observes the result and splits into a number of descendants—two in the absence of the ancilla and four if it is present. There is no requirement for the frequencies to be the same in both cases—i.e. no reason to apply the principle of measurement neutrality. In the language of decision theory, the values of the two games are not necessarily the same, so a decision on whether or not to accept a payoff will depend on whether the game is being played with or without an ancilla. Indeed, as in the absence of an ancilla there are only two branches, we might expect each observer’s experience to be the same as in case 2, with equal numbers of positive and negative results and an equal reward for each outcome—i.e. the statistical outcomes would be independent of the weights. I shall argue later that this is a natural consequence of the Everettian interpretation, but at present simply emphasize that the principle of measurement neutrality is a direct consequence of assumptions underlying the Copenhagen interpretation that are not contained in that of Everett.

The apparent failure of measurement neutrality in the Everettian context is central to our argument, so I illustrate it further by considering a simple

classical example that consists of a box with two exit ports from each of which a series of balls emerges as in figure 1. The apparatus can be operated in one of two modes that we denote as “C” and “E”. In the C mode, balls emerge one at a time from one of two output ports and, on average, three times as many come out of the upper port as from the lower. An experimenter observes the balls as they emerge and confirms this relative likelihood. Still in the C mode, the experiment is modified so that when a ball emerges from the upper port, it passes into a second, “ancillary” box from which it emerges at random through one of three output channels before being detected. The experimenter now detects a ball either in one of these three channels or emerging from the lower port. Clearly the first of these results is three times as likely as the second, so the observed frequencies are independent of the presence or absence of the second box. Thus the equivalent of measurement neutrality holds in this case.

Now consider the game in the E mode, which is also illustrated in figure 1. In this case two balls emerge from the box simultaneously: a black ball, from the upper port and a white ball from the lower. The two balls fall into a receptacle (not shown in the figure) and an experimenter draws one at random; after repeating the experiment a number of times she sees equal numbers of black and white balls. The experiment is now modified so that the black balls are directed into an ancillary box which now contains a device that releases three identical black balls, one through each of the three output ports, whenever one enters. These three balls along with the white one now fall into the receptacle and the observer again draws one at random; she now sees a black ball three times as often as a white ball. Thus, the relative likelihood of a black or white ball depends on the presence or absence of the second box, and we can conclude that measurement neutrality is not preserved when the game is played in the E mode.

A more whimsical analogy follows the precedent set by Schrödinger’s cat by using animals to illustrate our point. First consider Copenhagen rabbits. These come in two colours—black and white; they are all female and capable of giving birth to one (and only one) baby rabbit which is always of the same colour as its mother. Let us suppose we have four Copenhagen rabbits, three black and one white in a hat and suppose that one, and only one, of them is pregnant. We first play the game of “pick the pregnant rabbit” by putting our hand in the hat, identifying and then pulling out the pregnant rabbit. We are paid different rewards (x_b and x_w) depending on whether the extracted rabbit is black or white. After playing the game a number of times, we find

that we have pulled out three times as many black rabbits as white, so that the game value is $(3x_b + x_w)/4$. The second game is one where we wait until the pregnant rabbit has given birth and then pull out and identify the colour of the baby. Clearly the results and the value are the same as in the first game.

Now consider Everettian rabbits, which are also either black or white. In contrast to the Copenhagen rabbits, they are capable of carrying and giving birth to more than one offspring. Suppose we have two pregnant Everettian rabbits: a white rabbit that is pregnant with a single offspring, and a black rabbit that is expecting triplets. If we draw one of the two pregnant rabbits from the hat at random, the game value will be $(x_b + x_w)/2$. However, if, instead, we wait until after the rabbits have given birth and then draw out one of the offspring at random, the game value will now be $(3x_b + x_w)/4$. Thus Copenhagen rabbits preserve measurement neutrality, but Everettian rabbits do not.

Given the assumptions underlying the Copenhagen interpretation, the first game in the C mode and the game with the Copenhagen rabbits form close parallels with the quantum example discussed earlier. Similarly, the first game in the E mode and the game with Everettian rabbits are closely parallel to the quantum case, provided we accept that random selection at the point where the observer becomes aware of the result is equivalent to subjective uncertainty in the quantum case.

In both these examples as well as in the quantum case, I have shown that measurement neutrality is not a necessary consequence of the principles underlying the Everett interpretation. However, in all the cases where it need not apply, the symmetry is broken in the sense that the weights associated with the different outcomes are not equal. It follows that measurement neutrality (or, indeed, some other quite different principle) could be restored in the classical examples by making additional assumptions: for example, it could be arranged that the ball emerging from the upper channel in the E game is heavier than that coming out of the lower, and that it is three times easier to find and extract a more massive ball when making the selection; similarly, it might be three times easier to catch a rabbit carrying triplets than one pregnant with a single offspring. However, such *ad hoc* rules would have to be built into the physics of the apparatus when it was designed and constructed. Similarly, additional *ad hoc* rules could restore measurement neutrality in the quantum case, but for these to be consistent with the Everett interpretation, they would have to be a necessary consequence of the

Schrödinger equation. I shall shortly argue that this cannot be the case.

Measurement neutrality and an associated principle that he calls “equivalence” have been argued for by David Wallace in a number of papers—Wallace (2002), Wallace (2003), Wallace (2007). He considers games in which the measurement result is erased after it triggers an associated reward and before the experimenter has recorded the outcome. In the symmetric ($\theta = \pi/2$) case, the final states are independent of the pattern of rewards, which reinforces the arguments leading to (9). However, this is not an issue in the present discussion, which challenges the assumption of measurement neutrality only in the asymmetric case. Another point emphasized by Wallace (2007) is that the boundary between what is usually taken as preparation and what is part of the “actual” measurement is essentially arbitrary, particularly in the context of the Everett interpretation. However, the observation and recording of the result by a conscious observer is part of the measurement proper, and it is only at this point that subjective uncertainty or the relevance of a caring measure is introduced into the Everettian treatment of the Born rule.

Up to this point I have argued that the assumptions underlying the derivation of the Born rule, in particular measurement neutrality, are not necessary in an Everettian context. I now intend to go further and show that, there is a fundamental inconsistency between the assumptions underlying the Everett interpretation and any rule that relates the likelihood of a measurement outcome to the amplitudes (c and s in the above example) associated with the branching of the wavefunction. We shall continue to use the example of the measurement of the spin component of a spin-half particle as a focus of the discussion.

After interacting with a detector, the state of the system is given by (2). Let us suppose that there is also an observer present who has not yet observed the state of the counter and whose quantum state is represented by Ω_0 , which becomes Ω_+ or Ω_- , after the observation has been made. The wavefunction of the whole system, including the observer is then

$$\Phi = c\Phi_+ + s\Phi_- \longrightarrow \Psi = c\Psi_+ + s\Psi_- \quad (13)$$

where

$$\begin{aligned} \Phi_+ &= \alpha\chi_+\Omega_0 \longrightarrow \Psi_+ = \alpha\chi_+\Omega_+ \\ &\text{and} \\ \Phi_- &= \beta\chi_-\Omega_0 \longrightarrow \Psi_- = \beta\chi_-\Omega_- \end{aligned} \quad (14)$$

We first assume that the observer (whom we call Bob) knows the value of θ , which determines the orientation of the Stern-Gerlach magnet relative to that of the original prepared state. We assume that both the Everett interpretation and the Born rule apply. Although he knows that he will have descendants in each branch, Bob also knows about subjective uncertainty, so he can use the Born rule to form expectations about the likelihoods of the positive and negative outcomes and, if he is playing a game, he will use this information to place appropriate bets on the outcomes, as discussed earlier. He will therefore be in a “state of expectation”, which must be part of the complete description of his state that is contained in Ω_0 . Thus in this situation, Ω_0 must be a function (or perhaps a functional) of θ .

Now suppose that Bob does not know the value of θ before he makes any measurements. N identically prepared spins are passed through the apparatus and Bob finds M of them to be positive. Using the Born rule, he can estimate the value of θ as $2 \cos^{-1}(M/N)^{1/2}$ and his confidence in this value will be the greater, the larger are M and N . As a result of this experience, Bob’s state of expectation has been changed from one of ignorance to one where he has some knowledge of θ , which he can use to guide his betting strategy for subsequent measurements. This change must therefore have been reflected in Bob’s quantum state, causing a modification to Ω . To further emphasize this point, suppose that the value of θ can be changed without Bob’s direct knowledge by another experimenter (“Alice”) who has control of the Stern-Gerlach apparatus. If she does this and the experiment is repeated a number of times at the new setting, Bob will find that his expectations have been consistently wrong. He may initially attribute this to statistical fluctuation, but eventually he will amend his state of expectation to bring it into line with his experience. Indeed, Bob may know that Alice is able to do this, in which case he will be more likely to amend his state of expectation at an earlier stage. Alice could also send signals to Bob by transmitting sets of N particles using the same value of θ for each set, but changing it between sets. If the Born rule applies, Bob can deduce the values of θ that Alice has used from the relative numbers of positive and negative results, so Alice has again caused changes in Bob’s state of expectation and therefore of Ω .

It is one of the principles of the Everett interpretation that, once branching has occurred and the possibility of interference between branches has been eliminated, the wavefunction associated with a branch describes the “relative state” of the system contained in that branch and this cannot be influenced by the state of any other branch. This implies that the proper-

ties of a system represented by such a relative state are independent of the values of the constants c and s and therefore of θ , the setting angle of the Stern-Gerlach apparatus. This is because, although these constants enter the expressions Φ and Ψ for the total wavefunction (13), they do so only as expansion coefficients, which have no effect on the state of the component branches Φ_+ and Φ_- or on their evolution into Ψ_+ and Ψ_- . In particular, the state of the observer must be independent of the values of θ . This is in direct contradiction to the conclusion reached above, assuming that the Born rule holds. There is therefore a fundamental inconsistency between the Everett interpretation and the Born rule—or indeed any rule that implies a correlation between the apparatus setting and the relative frequencies of the possible outcomes.

Two points should be noted about the above. First, the contradiction does not arise in the Copenhagen interpretation because, as noted earlier, this assumes that the stochasticity arises as the spin emerges from the Stern-Gerlach magnet. The information as to which branch is occupied by the spin is additional to that contained in the wavefunction, so no contradiction arises when this is used by the experimenter to guide his expectations about subsequent measurements. Second, according to the Everett interpretation, the wavefunction contains branches corresponding to all possible sequences of results of the measurements performed so far. Some of these will be the same as predicted by the Born rule, so the corresponding observers may well be led into a state of expectation consistent with the Born rule. However, unless an additional principle (such as subjective uncertainty) applies, this will be analogous to the classical case where an observer incorrectly concludes that a tossed coin is biased because she has observed a long run of heads without seeing any tails. Unless we believe that all our experience of the quantum world to date are also random in the sense that they provide no guide at all to future events, an additional principle of some kind has to be invoked. It is my contention that Schrödinger evolution of the wavefunction alone does not allow the transfer of the information needed to allow an observer such as Bob to invoke such a principle in the cases discussed above.

If we accept the above, it follows that the only way probability can enter the Everett interpretation is if all branches are assigned equal probability. Up to now, we have assumed one branch per outcome, without attempting to justify this. We now turn to the question of “branch counting”, which means the number of branches associated with any given measurement outcome. If we accept the argument that the expansion coefficients play no role

in determining the outcome likelihood in an Everettian context, then an experimenter's expectation of a particular outcome should be proportional to the number of branches associated with it. Such an assumption is similar to that made in statistical thermodynamics, where the ergodic hypothesis states that the result of averaging over an ensemble of systems is the same as the time average for a single system. When applied to the symmetric case, this is an essential part of the arguments leading to (9) and (10). However, branch counting has been strongly criticized by DSW on a number of grounds. Wallace (2007) considers a scenario in which extra branching is introduced into one (say the plus) channel by associating with it a device that displays one of, say a million, random numbers. He argues that this must be irrelevant to an experimenter who sees only the measurement result and is indifferent to the outcome of the randomizing apparatus. This is because "if we divide one outcome into equally-valued suboutcomes, that division is not decision-theoretically relevant". However, this argument does not fully take into account the Everettian context. Referring again to the classical game discussed earlier and illustrated in figure 1, we can consider the additional branching on the right of both setups as due to the presence of a randomizer with three possible outputs. In the case of the C game, these are indeed irrelevant to the expectation of the player, because a ball emerges from only one of the three channels and must have passed through the upper channel. However, in the case of the E game, the chances of observing a black ball are enhanced (tripled) by the splitting and this would have to be taken into account by any rational player, even if the only result she sees is the colour of the ball. Similarly, if we introduce a random number machine as Wallace suggests, then its state will be a linear combination of its million possible outcomes and all these will be associated with a positive value of spin. Given that there is only one branch associated with the alternative outcome, we must expect the subjective likelihood of a positive result to be one million times greater than that for a negative outcome.

A second argument deployed to criticize branch counting is based on the fact that the interaction of a quantum system with its environment leads to an immensely complex branching structure. Indeed it is claimed that the number of branches is not only very large (possibly infinite), but is also subject to very large and rapid fluctuations before, during and after the observation of a result; which may mean that it is not meaningful to talk about even the approximate number of branches that exist at any time. This is adduced as a reason why a rational player should ignore the complexity of the

branching structure and instead expect to observe results consistent with the Born rule. However, if the likelihood of observing a particular result is proportional to the number of associated branches, the complexity introduced by decoherence should actually result in the outcome of a measurement being completely unpredictable. The situation is similar to chaos in classical mechanics or to turbulence in hydrodynamics, whose onset certainly does not lead to increased predictability. In the arguments above, we assumed that each outcome was associated with a single branch, so what would be the likely consequences of a complex branch structure in an Everettian context? First, there may well be situations in which we could expect the number of branches associated with different outcomes to be equal, at least when averaged over a number of measurements, and in this case our earlier discussion would not be affected. However, we might be able to devise a situation (e.g. one in which a detector was placed in the positive output channel only) where the numbers of branches in the two channels would be expected to differ greatly. We should then expect to detect a larger number of (say) positive than negative results. This would be true even if the Stern-Gerlach apparatus were oriented symmetrically—i.e. with $\theta = \pi/2$, so the symmetry on which we based some of our earlier arguments would not hold. The complexity and fluctuations of the branch structure in the Everett case would render even the statistical results of a quantum measurement unpredictable. Such a situation is sometimes described as being “incoherent” and it has been argued that this would mean that the universe would be nothing like the one we experience. However, the obvious conclusion to draw from this is that the Everett assumptions are falsified, rather than that the Everett model is correct and the arguments based on it that lead to incoherence must be wrong.

It might be thought that branch counting would restore the Born rule if the number of branches associated with a particular outcome were proportional to the Born weight. However, not only is there no obvious mechanism to achieve this, it is also inconsistent with the Everett model for the same reasons as were set out earlier. The quantum description of the branch structure is contained within Φ and Ψ in (13) and (14) and therefore cannot depend on the orientation, θ .

Conclusions

I have argued that attempts to prove the Born rule make assumptions that are valid in the context of the Copenhagen interpretation, but are not consistent with the principles underlying the Everett model of measurement. I have further argued that probabilities which are functions of the expansion coefficients cannot be generally included within the Everett interpretation, because these quantities are not generally accessible to an observer in the reduced state associated with a branch. An alternative scheme that could be consistent with Everett is one where each branch has the same probability and the probability of a given outcome depends on the number of branches associated with it. However, this also cannot be made consistent with the Born rule and it leads to predictions of chaotic, unpredictable behaviour, in contrast to the relatively well-ordered behaviour, invariably demonstrated in experiments. I conclude that the Born rule is a vitally important principle in determining quantum behaviour, but that it is additional to, rather than supervening upon, the time-dependent Schrödinger equation.

The debate between the different interpretations of quantum mechanics has often been metaphysical in the sense that they often make the same predictions and cannot therefore be distinguished experimentally. The present paper has argued that this is not so in the case of the Everett interpretation, which predicts results different from those that follow from the Copenhagen interpretation, which in turn are supported by experiment. If this is accepted, the Everett model will have been falsified and the search for a consensual resolution of the quantum measurement problem will have to be focussed elsewhere.

I am grateful to Peter Lewis for comments on an earlier draft of this paper and to Simon Saunders and David Wallace for useful discussions.

References

- Baker, D., 2006. Measurement outcomes and probability in everettian quantum mechanics. Phil Sci archive id2717.
- Barnum, H., Caves, C., Finkelstein, J., Fuchs, C. A., Schack, R., 2000. Quantum probability from decision theory? Proc. Roy. Soc. A456 (1175-1182).
- Deutsch, D., 1999. Quantum theory of probability and decisions. Proc. Roy. Soc. A455, 3129–37.
- Everett III, H., 1957. “relative state” formulation of quantum mechanics. Rev. Mod. Phys. 29, 454–462.
- Graham, N., 1973. The Measurement of Relative Frequency. Princeton University Press, Princeton.
- Greaves, H., 2004. Understanding deutsch’s probability in a deterministic universe. Studies in Hist. Phil. Mod. Phys. 35, 423–56.
- Hemmo, M., I, P., 2007. Quantum probability and many worlds. Studies in History and Philosophy of Modern Physics 38, 333–350.
- Kent, A., 1990. Against many-worlds interpretations. Intl. J. Mod. Phys. A 5, 1745–76.
- Lewis, D., 2004. How many lives has Schrödinger ’s cat? Australasian J. of Phil. 82, 3–22.
- Lewis, P. J., 2005. Probability in everettian quantum mechanics. PhilSCI eprint.
- Lewis, P. J., 2006. Uncertainty and probability for branching selves. Studies in History and Philosophy of Modern Physics forthcoming.
- Lockwood, M., 1989. Mind, Brain and the Quantum. Blackwell, Oxford.
- Saunders, S., 2004. Derivation of the born rule from operational assumptions. Proc. Roy. Soc. A460, 1–18.
- Squires, E. J., 1990. On an alleged “proof” of the quantum probability law. Phys. Lett. A145, 67–68.

- Wallace, D., 2002. Worlds in the everett interpretation. *Studies in Hist. Phil. Mod. Phys.* 33, 637–61.
- Wallace, D., 2003. Everett and structure. *Studies in Hist. Phil. Mod. Phys.* 34, 87–105.
- Wallace, D., 2007. Quantum probability from subjective likelihood: Improving on deutsch’s proof of the probability rule. *Studies in History and Philosophy of Modern Physics* 38, 311–332.
- Zurek, 2008. Relative states and the environment. [quant-ph/0707.2832v1](https://arxiv.org/abs/quant-ph/0707.2832v1).