

Comment on “Application of the Lifshitz theory to poor conductors”

The Letter [1] is devoted to the generalization of the Lifshitz theory to spatially nonlocal conducting materials. Spatial dispersion is taken into account approximately by means of the dielectric permittivities calculated in the random phase approximation. Letter [1] recognizes that these permittivities are rigorously defined only for an infinite medium, but claims that they can be used for two semispaces separated by a gap as well. Additionally, the specular reflection of charge carriers on the boundary planes is assumed. It has been proven, however, that for spatially dispersive materials the scattering of carriers is neither specular nor diffuse [2]. Thus, the developed approach is a crude approximation. It does not contain self-consistent checks of its accuracy and could be justified only because of agreement with experimental data and fundamental physical principles. Below we show that the claims of [1] regarding consistency with the experimental data and Nernst’s theorem are based on irregular comparison with the data and misinterpretation of relevant physical quantities.

According to [1], the experimental data of [3] for the difference Casimir force between an Au sphere and a Si plate in the presence and in the absence of laser light are equally consistent with the nonlocal approach and “local theory with zero conductivity of Si”. (Note that in [3] Si conductivity was disregarded only in the absence of light.) To prove this, in Fig. 1(a) of [1] the experimental data [taken from Fig. 1(a) of [4] without a reference] are shown at a 70% confidence level. In the same figure the theoretical band for the nonlocal approach is obtained from the uncertainty in n , $\Delta n = 0.4 \times 10^{19} \text{ cm}^{-3}$, determined in [3] at a 95% confidence level. The reader of [1] is not informed about the confidence levels used. Such a mismatch comparison of experiment with theory is irregular. In Fig. 1(a) of this Comment the same data are compared with the predictions of the nonlocal approach and the local theory (the bands between the dashed and solid lines, respectively), where the band widths are determined at the same 70% confidence level, as the errors of the data. It is clearly seen that the local theory is consistent with data, whereas the nonlocal approach is excluded by the data at a 70% confidence level. [Fig. 1(b) in [1] shows both the data and the theoretical bands at a 95% confidence level. However, as stated in [4], these data cannot be conclusively compared with the nonlocal approach at such a high confidence.]

According to [1], the nonlocal approach is also applicable to metallic test bodies. In Fig. 1(b) we plot as crosses the mean measured Casimir pressures between two Au plates [5] at a 95% confidence compared with computations using the nonlocal approach (the grey

band) and the generalized plasma-like permittivity (the black band). The widths of both bands are determined at a 95% confidence. It can be seen that the nonlocal approach is excluded by the data at a 95% confidence, whereas the local theory is consistent with them.

According to [1], the nonlocal approach satisfies the Nernst theorem, specifically, for ionic conductors possessing an activation type conductivity. To prove this, the Letter [1] arbitrarily separates the thermal dependence $\sim \exp(-E_a/k_B T)$, where E_a is the activation energy, from the mobility μ and attributes it to the “effective density of charges, which are able to move”. This transfer of the temperature dependence from μ to n is incorrect because the commonly used density of charge carriers n producing the effect of screening in ionic conductors is an independently measured quantity [6], which does not vanish with T . In reality, the screening length depends on the standard charge carrier density n and this results in the violation of Nernst’s theorem in the nonlocal approach to ionic conductors [4].

G. L. Klimchitskaya,¹ U. Mohideen,² and V. M. Mostepanenko¹

¹Institute for Theoretical Physics, Leipzig University, D-04009, Leipzig, Germany

²Department of Physics and Astronomy, University of California, Riverside, CA 92521, USA

PACS numbers: 42.50.Ct, 12.20.Ds, 42.50.Lc, 78.20.Ci

[1] V. B. Svetovoy, Phys. Rev. Lett. **101**, 163603 (2008).

[2] J. T. Foley and A. J. Devaney, Phys. Rev. B **12**, 3104 (1975).

[3] F. Chen *et al.*, Phys. Rev. B **76**, 035338 (2007).

[4] G. L. Klimchitskaya *et al.*, arXiv:0802.2698; J. Phys.A **41**, 432001 (2008).

[5] R. S. Decca *et al.*, Eur. Phys. J. C **51**, 963 (2007).

[6] M. Tomazawa and Dong-Wook Shin, J. Non-Cryst. Sol. **241**, 140 (1998).

Figures




FIG. 1: The measured (a) force differences and (b) pressures are shown as crosses versus separation. The theoretical bands in (a) lie between the solid and dashed lines, respectively, and in (b) are indicated as black and grey stripes. See text for further discussion.