

# Effect of resonant interactions on the stability of a three-state Fermi gas

J. H. Huckans, J. R. Williams, E. L. Hazlett, R. W. Stites, and K. M. O'Hara\*

*Department of Physics, Pennsylvania State University, University Park, Pennsylvania 16802-6300, USA*

(Dated: May 23, 2019)

We investigate the stability of a three spin state mixture of ultracold fermionic  ${}^6\text{Li}$  atoms over a range of magnetic fields encompassing three Feshbach resonances. For most field values, we attribute decay of the atomic population to three-body processes involving one atom from each spin state and find that the three-body loss coefficient varies by over four orders of magnitude. We observe high stability when at least two of the three scattering lengths are small, rapid loss near the Feshbach resonances, and two unexpected resonant loss features. At our highest fields, where all pairwise scattering lengths are approaching  $a_t = -2140a_0$ , we measure a three-body loss coefficient  $L_3 \simeq 5 \times 10^{-22} \text{ cm}^6/\text{s}$  and a trend toward lower decay rates for higher fields indicating that future studies of color superfluidity and trion formation in a  $\text{SU}(3)$  symmetric Fermi gas may be feasible.

PACS numbers: 67.85.Lm, 34.50.-s, 67.85.-d, 03.75.Ss

Multi-component Fermi gases with tunable interactions are exceptionally well suited to the study of few- and many-body quantum physics. Ultracold two-state Fermi gases near a Feshbach resonance have been used to characterize the crossover from Bardeen-Cooper-Schreiffer (BCS) superfluidity to Bose-Einstein condensation (BEC) of diatomic molecules [1, 2, 3, 4, 5, 6, 7]. Further, a normal to superfluid transition and phase separation have been observed in imbalanced two-state spin mixtures [8, 9, 10, 11, 12]. The stability of the two-state mixtures against two- and three-body loss processes was critically important to the success of these experiments.

Recently, there has been considerable interest in the study of three-state Fermi gases with tunable interactions [13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25]. Whereas three-body interactions in ultracold two-state Fermi gases are suppressed by the exclusion principle, the addition of a third spin component allows for the study of three-body phenomena such as the Efimov effect in a Fermi system if pairwise interactions are resonantly enhanced [19, 24]. Further, with this additional spin state there may be competition between multiple pairing states and trion formation [13, 14, 15, 16, 17, 20, 21, 22, 23]. If pairwise interactions are all attractive and of equal magnitude, the system is symmetric under  $\text{SU}(3)$  transformations and is predicted to exhibit a novel superfluid phase with  $\text{SU}(2)$  symmetry and an anomalous number of Goldstone modes [15]. This superfluid phase would be a cold atom analog of the color superfluid phase predicted in quantum chromodynamics (QCD) which undergoes a quantum phase transition to a Fermi liquid of trion states (analogous to baryons) as the ratio of the interaction energy to the kinetic energy is adjusted [18].

Future studies of the above phenomena critically depend on the magnitude of two- and three-body loss rates, particularly when two or more scattering lengths are enhanced by scattering resonances. Two- and three-body loss and heating processes, often enhanced near the resonances [26], can impose stringent limits on the maximum

achievable phase space density. Further, the unambiguous observation of three-body resonances requires negligible two-body loss [27].

Most investigations of Fermi gases with tunable interactions have focussed on two-state mixtures. A small admixture of a third spin component has been used for thermometry [28] and recently the rapid decay of a three-state Fermi gas near a Feshbach resonance was noted during an investigation of Cooper pair size by radio-frequency (RF) spectroscopy in a two-state Fermi gas [29]. Very recently our group reported measurements of three-body loss in thermal and degenerate three-state Fermi gases over a wide range of magnetic fields [30] and Ottenstein *et al.* independently reported measurements of three-body loss rate coefficients in a degenerate three-state Fermi gas for field values away from any pairwise scattering resonances [31].

In this Letter, we report the first measurements of three-body loss coefficients for a three-state Fermi gas over a range of magnetic fields where pairwise interactions are resonantly enhanced. This enhancement is due to three overlapping interspecies Feshbach resonances and a zero-energy resonance in the  ${}^6\text{Li}$  triplet molecular potential. Further, we demonstrate that all two-state mixtures of the three lowest energy hyperfine states of  ${}^6\text{Li}$  are stable against two-body loss processes for almost the entire range of field values between 15 and 953 G, making this three-state mixture well suited to studies of three-body physics. We observe a narrow loss feature at 127 G [32] which may be due to a three-body resonance and an additional state-dependent loss feature at 504 G. Finally, we measure the three-body loss rate coefficient at high fields where all three scattering lengths are asymptoting to the triplet scattering length  $a_t = -2140a_0$ . This measurement and an observed trend toward lower loss at higher fields has important implications for realizing cold atom analogs of color superfluid and baryon phases in QCD.

We study a  ${}^6\text{Li}$  Fermi gas with equal populations in the




FIG. 1: (color online) (a) Energies of the hyperfine ground states of  ${}^6\text{Li}$  in a magnetic field. We create a balanced mixture of the three lowest energy states by applying a RF pulse at a field of 568 G (see inset). (b) The three pairwise  $s$ -wave scattering lengths as a function of magnetic field [33] show Feshbach resonances at 690, 811, and 834 G and all asymptote to  $a_t = -2140a_0$  at high field.

three lowest energy hyperfine states. Figure 1(a) shows the energies of the hyperfine ground states of  ${}^6\text{Li}$  as a function of magnetic field. At zero field, the three states correspond to  $|1\rangle = |\frac{1}{2}, +\frac{1}{2}\rangle$ ,  $|2\rangle = |\frac{1}{2}, -\frac{1}{2}\rangle$  and  $|3\rangle = |\frac{3}{2}, -\frac{3}{2}\rangle$  in the  $|f, m_f\rangle$  basis. For fields above  $\simeq 200$  G, these states become increasingly electron-spin polarized and primarily differ by their nuclear spin projection (i.e. in the  $|m_s, m_i\rangle$  basis  $|1\rangle \simeq |-\frac{1}{2}, 1\rangle$ ,  $|2\rangle \simeq |-\frac{1}{2}, 0\rangle$  and  $|3\rangle \simeq |-\frac{1}{2}, -1\rangle$ ). The pairwise  $s$ -wave scattering lengths between these states ( $a_{12}$ ,  $a_{23}$ , and  $a_{13}$ ) exhibit three broad overlapping Feshbach resonances as shown in Fig. 1(b) [33]. For very large magnetic fields, each of the pairwise scattering lengths asymptote to the triplet scattering length for  ${}^6\text{Li}$ ,  $a_t = -2140a_0$ . Therefore, in the limit of large magnetic fields, this three-state mixture becomes invariant under global  $\text{SU}(3)$  transformations of the spinor wavefunction.

Two-body spin-flip processes are expected to be especially small for this ultracold  ${}^6\text{Li}$  mixture. Spin-exchange collisions become energetically forbidden even in a small magnetic field. Dipolar relaxation is small since (1) the energy released in a spin-flip is not enough to overcome the centrifugal barrier in the  $d$ -wave exit channel and (2) electron spin-flip processes are suppressed at high field as the gas becomes increasingly electron-spin polarized in the lowest energy electron spin state.

We can produce a degenerate Fermi gas (DFG) with an equal mixture of atoms in states  $|1\rangle$  and  $|2\rangle$  once ev-

ery 5 seconds by evaporatively cooling this mixture in a red-detuned optical dipole trap [34]. During the first second,  $\sim 10^8$   ${}^6\text{Li}$  atoms from a Zeeman-slowed atomic beam are collected in a magneto-optical trap (MOT). A crossed optical dipole trap which overlaps the MOT is then turned on and optimally loaded by tuning the MOT laser beams  $\simeq 6$  MHz below resonance and reducing their intensity for 7 ms. The atoms are then optically pumped into states  $|1\rangle$  and  $|2\rangle$  prior to extinguishing the MOT laser beams and field gradient. The two beams which form the optical trap derive from a linearly polarized multi-longitudinal-mode 110 W fiber laser operating at 1064 nm. The beams nearly co-propagate in the vertical direction ( $\hat{y}$ ), have orthogonal polarizations and cross at an angle  $\simeq 11^\circ$ . The beams are elliptical with calculated  $e^{-2}$  waist radii  $\sim 30\mu\text{m}$  and  $\sim 100\mu\text{m}$  at the point of intersection. The maximum trap depth per beam is  $\simeq 1$  mK allowing  $\sim 5 \times 10^6$  atoms to be initially loaded. We apply a bias field and a noisy RF pulse to create a 50-50 mixture of atoms in states  $|1\rangle$  and  $|2\rangle$  [34].

Forced evaporation of the atoms occurs at a field of 330 G where the two-state scattering length  $a_{12} \simeq -280a_0$ . We lower the depth of the optical trap  $U_0$  by a factor of 107 over 3.6 seconds to obtain our final temperature. For the work presented here we study a gas at a temperature  $T \gtrsim 0.5 T_F$  ( $T_F$  is the Fermi temperature) so that it is appropriate to treat the cloud as a thermal gas. To suppress further loss by evaporation for the remainder of the experiment,  $U_0$  is then increased by a factor of 4 increasing the ratio of  $U_0/k_B T$  by a factor  $\simeq 2$ . The final oscillation frequencies of the trap are  $\nu_x = 3.84$  kHz,  $\nu_y = 106$  Hz and  $\nu_z = 965$  Hz [35] with a final trap depth per beam  $\simeq 40 \mu\text{K}$ . The total number of atoms  $\simeq 3.6 \times 10^5$  in a balanced two-state mixture at  $T \simeq 1.9 \mu\text{K}$  for which  $T_F \simeq 3.7 \mu\text{K}$ . The background limited  $1/e$ -lifetime  $\simeq 30$  s.

To create an incoherent three-state mixture we first increase the strength of the magnetic field in 10 ms to 568 G where the mixture is stable. We then apply a noisy RF magnetic field with two frequencies centered on the  $|1\rangle - |2\rangle$  and  $|2\rangle - |3\rangle$  spin-flip transitions, both broadened to a width of 1 MHz by frequency modulation with white noise. A magnetic field gradient of  $\simeq 1$  G/cm in the direction of the bias field ( $\hat{z}$  direction) is simultaneously applied to dephase the internal state coherence of the sample. The noisy RF field remains on for 50 ms to create an equal mixture of  $N \simeq 1.2 \times 10^5$  atoms in each state. At this point,  $T = 1.9 \mu\text{K}$  with  $T_F = 3.2 \mu\text{K}$ .

In our first experiment to explore the field dependence of the loss mechanisms in this three-state Fermi gas, we measured the fraction of trapped atoms remaining after an evolution time of 200 ms at fixed magnetic fields between 15 and 953 G. Figure 2(a) shows the fraction remaining for each of the spin states. For each measurement, the magnetic field was ramped in 10 ms from 568 G to the field of interest where it was held constant for ei-




FIG. 2: (color) (a) The fraction of atoms remaining in each of the three spin states after 200 ms of evolution at the magnetic field of interest. (b) Magnetic field dependence of decay in individual two-state mixtures.  $\blacklozenge$  ( $\bullet$ ) indicates the fraction remaining in state |2> (|3>) of a |2>–|3> (|1>–|3>) mixture after 200 ms. The thick lines guide the eye. The representative error bars indicate the standard deviation in the mean.

ther 1 or 201 ms. The field was then ramped in 10 ms to 953 G and held for 20 ms before a  $300 \mu\text{s}$  time-of-flight expansion and absorption imaging at this field. The 10 ms field ramp to 953 G (where  $a_{12}, a_{23}, a_{13} < 0$ ) ensured that any atoms which had formed weakly bound molecules in the vibrational state associated with the Feshbach resonances (but remained trapped) would be disassociated and measured. Further, each of the spin states could be spectroscopically resolved at this field.

To confirm that two-body loss does not contribute significantly to the measurements above, we measured the fraction of atoms remaining after an evolution time of 200 ms at fixed magnetic field values for each of the three possible two-state mixtures. Figure 2(b) shows the fraction remaining in state |3> for a |1>–|3> mixture and state |2> for a |2>–|3> mixture. The fraction of atoms remaining in the |1>–|2> mixture (not shown) is similar to the |2>–|3> data. With the exception of loss features between 600 and 750 G, each mixture was stable. The loss features are due to three-body recombination to bound molecular states associated with the Feshbach resonance. Since the two-state mixtures are stable for fields  $\leq 600$  G and  $\geq 750$  G, we can interpret loss features in a three-state mixture at these fields as being due to three-body

events involving one atom from each spin state.

In Fig. 2(a), the broad dominant loss feature centered at 720 G occurs in the vicinity of three overlapping interspecies Feshbach resonances. Significant loss due to three-body recombination is expected near these resonances since recombination events involving one atom in each spin state are *not* suppressed by the exclusion principle and one expects a significant increase in the event rate when two or more scattering lengths are resonantly enhanced [36]. Similarly, high stability at zero field and near the zero crossings of the Feshbach resonances is not surprising since at least two of the scattering lengths are small at these fields. At high field where  $a_{12}, a_{23}, a_{13} \rightarrow a_t$ , the stability increases relative to that at 720 G.

Unexpected resonant loss features are observed at 127 and 504 G, where the two-body scattering lengths are not predicted to exhibit any resonances. A possible explanation for the feature at 127 G is that the binding energy of a three-body bound state crosses through zero near this field value. At 504 G, we observe differing degrees of enhanced loss for the three states leading to population imbalance. The feature at 228 G in Fig. 2 is consistent with our estimated location of a |1>–|3>  $p$ -wave Feshbach resonance. We also observed a very narrow but inconsistent loss feature at 259 G (not shown).

A second set of experiments measured three-body loss coefficients at fixed magnetic fields where the decaying populations remained balanced. In these experiments, the magnetic field is swept to the field of interest  $B$  in 10 ms after creation of the three-state mixture and the number  $N(t)$  and temperature  $T(t)$  of atoms in state |3> are then measured by time-of-flight absorption imaging at the field  $B$  for various delay times  $t$ . To extract the three-body loss coefficients, we fit this data assuming one- and three-body loss and heating due to three-body recombination.

When three-body recombination involves one atom from each spin state, the number of trapped atoms in each of the equally populated spin states  $N(t)$  evolves according to  $\dot{N} = -L_1 N - L_3 \langle n^2 \rangle N$ . Here,  $\langle n^2 \rangle$  is the average value of the squared density per spin state and  $L_3$  ( $L_1$ ) is the three-body (one-body) atom-loss rate coefficient. The density distribution in the trap is well described by a thermal distribution so that

$$\frac{dN}{dt} = -L_1 N - \gamma \frac{N^3}{T^3} \quad (1)$$

where  $\gamma = L_3 (m\bar{\omega}^2 / 2\pi k_B)^3 / \sqrt{27}$  and  $\bar{\omega} = (\omega_x \omega_y \omega_z)^{1/3}$ . Following Ref. [37], we model the temperature increase in the gas as arising from “anti-evaporation” (the preferential loss of the highest density atoms) and recombination heating  $T_h$  which causes the temperature of the atoms to evolve as

$$\frac{dT}{dt} = \gamma \frac{N^2 (T + T_h)}{T^3} \quad (2)$$




FIG. 3: (a) Three-body atom-loss rate coefficients,  $L_3$ , at various magnetic field values. Evolution of (b) atom number and (c) temperature at a particular field (180 G). The solid lines show a best-fit to the experimental data which determines  $L_3$ .

For a given set of parameters ( $N(0)$ ,  $T(0)$ ,  $L_1$ ,  $L_3$  and  $T_h$ ), these two coupled differential equations can be numerically integrated to give  $N(t)$  and  $T(t)$ . The best-fit parameters for a data set at a given field are determined by minimizing  $\chi^2$  with respect to these fit parameters. For example, Fig. 3 shows the measured number (b) and temperature (c) as a function of time at the field 180 G with the best-fit curves. The best-fit  $L_3$  values at various fields are displayed in Fig. 3(a).

As shown in Fig. 3(a),  $L_3$  varies by over four orders of magnitude and corroborates the features observed in Fig. 2(a). The error bars (typically comparable to the circles) indicate the uncertainty in the determination of  $L_3$  due to statistical fluctuations and do not include a systematic fractional uncertainty in  $L_3$  of  $\pm 70\%$  arising from our fractional uncertainty in atom number ( $\pm 30\%$ ) and trap frequencies ( $\pm 5\%$ ). In the range  $600 \text{ G} < B < 750 \text{ G}$ , the  $L_3$  values ( $\circ$ ) may be overestimates since the measured decay includes loss events we observe in two-state mixtures at these field values. Near 0 and 568 G, three-body recombination rates are relatively small and very stable three-state Fermi gases can be created. For  $L_3 = 10^{-25} \text{ cm}^6/\text{s}$ , a gas with a density  $n = 5 \times 10^{11} \text{ cm}^{-3}$  per spin state has a calculated lifetime  $\sim 40$  seconds. Consistent with the data shown in Fig. 2(a), a resonant peak in  $L_3$  is observed at  $B \simeq 127 \text{ G}$ . The highest  $L_3$  value we report ( $10^{-21} \text{ cm}^6/\text{s}$ ) occurs at 775 G and is likely unitarity limited (at these fields  $T \simeq 6 \mu\text{K}$  due to heating and  $\sqrt{\hbar^2/mk_B T} = 2200a_0$ ). We observe that  $L_3$  decreases by a factor of  $\simeq 2.5$  as the field is increased from 775 to 953 G, the highest field we currently access. This trend suggests that  $L_3$  may decrease further in the high field limit where  $a_{12}, a_{23}, a_{13} \rightarrow -2140a_0$ .

In conclusion, we can create long-lived three-state DFGs of  $^6\text{Li}$  atoms over a range of fields. Additionally, the minimal two-body loss we observe makes this system promising for future studies of Efimov trimers and other three-body bound states. Indeed, the loss reso-

nance at 127 G may be a signature of such a state. The recombination rates at high fields need not preclude experiments designed to study color superfluidity and trion formation [18]. For instance, by working at low density ( $n \sim 5 \times 10^{10} \text{ cm}^{-3}$ ) and using large period ( $d \simeq 2 \mu\text{m}$ ) optical lattices, long lifetimes ( $\gtrsim 0.1 \text{ s}$ ) and strong interactions can simultaneously be achieved.

We gratefully acknowledge support from the AFOSR, ARO, DARPA and NSF.

\* kohara@phys.psu.edu

- [1] C. A. Regal *et al.*, Phys. Rev. Lett. **92**, 040403 (2004).
- [2] M. W. Zwierlein *et al.*, Phys. Rev. Lett. **92**, 120403 (2004).
- [3] J. Kinast *et al.*, Phys. Rev. Lett. **92**, 150402 (2004).
- [4] C. Chin *et al.*, Science **305**, 1128 (2004).
- [5] T. Bourdel *et al.*, Phys. Rev. Lett. **93**, 050401 (2004).
- [6] M. W. Zwierlein *et al.*, Nature **435**, 1047 (2005).
- [7] G. B. Partridge *et al.*, Phys. Rev. Lett. **95**, 020404 (2005).
- [8] G. B. Partridge *et al.*, Science **311**, 503 (2006).
- [9] G. B. Partridge *et al.*, Phys. Rev. Lett. **97**, 190407 (2006).
- [10] M. W. Zwierlein *et al.*, Science **311**, 492 (2006).
- [11] Y. Shin *et al.*, Phys. Rev. Lett. **97**, 030401 (2006).
- [12] M. W. Zwierlein *et al.*, Nature **442**, 54 (2006).
- [13] A. G. K. Modawi and A. J. Leggett, Journal of Low Temperature Physics **109**, 625 (1997).
- [14] C. Honerkamp and W. Hofstetter, Phys. Rev. B **70**, 094521 (2004).
- [15] L. He *et al.*, Phys. Rev. A **74**, 033604 (2006).
- [16] T. Paananen *et al.*, Phys. Rev. A **73**, 053606 (2006).
- [17] P. F. Bedaque *et al.*, arXiv:cond-mat/0602525 (2006).
- [18] A. Rapp *et al.*, Phys. Rev. Lett. **98**, 160405 (2007).
- [19] E. Braaten and H.-W. Hammer, Ann. Phys. **322**, 120 (2007).
- [20] H. Zhai, Phys. Rev. A **75**, 031603(R) (2007).
- [21] R. W. Cherng *et al.*, Phys. Rev. Lett. **99**, 130406 (2007).
- [22] X. W. Guan *et al.*, Phys. Rev. Lett. **100**, 200401 (2008).
- [23] X.-J. Liu *et al.*, Phys. Rev. A **77**, 013622 (2008).
- [24] T. Luu and A. Schwenk, Phys. Rev. Lett. **98**, 103202 (2007).
- [25] S. Capponi *et al.*, Phys. Rev. A **77**, 013624 (2008).
- [26] J. L. Roberts *et al.*, Phys. Rev. Lett. **85**, 728 (2000).
- [27] T. Kraemer *et al.*, Nature **440**, 315 (2006).
- [28] C. A. Regal, Ph.D. thesis, University of Colorado, Boulder (2005).
- [29] C. H. Schunck *et al.*, Nature **454**, 739 (2008).
- [30] J. Williams *et al.*, Bull. Am. Phys. Soc. **53**, 80 (2008).
- [31] T. B. Ottenstein *et al.*, arXiv:0806.0587 (2008).
- [32] This resonance was first reported in Ref.[31].
- [33] M. Bartenstein *et al.*, Phys. Rev. Lett. **94**, 103201 (2005).
- [34] K. M. O'Hara *et al.*, Science **298**, 2179 (2003).
- [35] We determine trap frequencies by observing dipole oscillations ( $\omega_y$ ) and parametric excitation loss ( $\omega_x, \omega_z$ ).
- [36] Three particles strongly interact even if one of the pairwise scattering lengths is small since the third particle acts as a mediator for the weakly interacting pair.
- [37] T. Weber *et al.*, Phys. Rev. Lett. **91**, 123201 (2003).