A symplectic subgroup of a pseudounitary group as a subset of Clifford algebra

Nikolai Marchuk, Roman Dyabirov

March 31, 2022

Abstract

Let Cl1(1,3) and Cl2(1,3) be the subsets of elements of the Clifford algebra Cl(1,3) of ranks 1 and 2 respectively. Recently it was proved that the subset Cl2(p,q)+iCl1(p,q) of the complex Clifford algebra can be considered as a Lie algebra. In this paper we prove that for p=1, q=3 the Lie algebra Cl2(p,q)+iCl1(p,q) is isomorphic to the well known matrix Lie algebra sp(4,R) of the symplectic Lie group Sp(4,R). Also we define the so called symplectic group of Clifford algebra and prove that this Lie group is isomorphic to the symplectic matrix group Sp(4,R).

Introduction

In [1] was considered the complex Clifford algebras $\mathcal{C}\!\ell(p,q)$ and was found subsets of Clifford algebra closed with respect to the commutator [U, V] = UV - VU. For all positive integer p, q $(p + q \ge 2)$ the subset $\mathcal{C}\!\ell_2^{\mathbb{R}}(p,q) \oplus i \mathcal{C}\!\ell_1^{\mathbb{R}}(p,q)$ of Clifford algebra $\mathcal{C}\!\ell(p,q)$ can be considered as a Lie algebra. In this paper we prove that for p = 1, q = 3 the Lie algebra $\mathcal{C}\!\ell_2^{\mathbb{R}}(p,q) \oplus i \mathcal{C}\!\ell_1^{\mathbb{R}}(p,q)$ is isomorphic to the well known matrix Lie algebra $\mathrm{sp}(4, R)$ of the symplectic Lie group $\mathrm{Sp}(4, R)$. Also we define the so called symplectic group of Clifford algebra and prove that this Lie group is isomprphic to the symplectic matrix group $\mathrm{Sp}(4, R)$.

1 Clifford algebras $\mathcal{C}\ell(p,q)$

Let F be the field of real numbers \mathbb{R} or the field of complex numbers \mathbb{C} and let n be a natural number. Consider 2^n dimensional vector space \mathcal{E} over the field \mathbb{F} with a basis

$$e, e^{a}, e^{a_{1}a_{2}}, \dots, e^{1\dots n}, \quad a_{1} < a_{2} < \cdots,$$

$$(1)$$

with elements numbered dy ordered multi-indices of length from 0 to n. Indices a, a_1, \ldots take values from 1 to n.

Let p, q be nonnegative integer numbers and p + q = n. Consider a diagonal matrix of dimension n

$$\eta = \eta(p, q) = \text{diag}(1, \dots, 1, -1, \dots, -1), \tag{2}$$

with p pieces of 1 and q pieces of -1 on the diagonal. By $\eta^{ab} = \eta_{ab}$ denote elements of η .

Following rules define product $U, V \to UV$ of elements of the vector space \mathcal{E} :

1. For any $U, V, W \in \mathcal{E}$

$$U(V+W) = UV + UW, \quad (U+V)W = UW + VW,$$

$$(UV)W = U(VW);$$

- 2. eU = Ue = U, for any $U \in \mathcal{E}$;
- 3. $e^{a}e^{b} + e^{b}e^{a} = 2\eta^{ab}e$, for a, b = 1, ..., n;
- 4. $e^{a_1} \dots e^{a_k} = e^{a_1 \dots a_k}$, for $1 \le a_1 < \dots < a_k \le n$.

The operation of multiplication converts vector space \mathcal{E} to an algebra. This algebra is called *the Clifford algebra* and denoted by $\mathcal{C}\ell^{\mathbb{F}}(p,q)$ (if $\mathbb{F} = \mathbb{C}$, then $\mathcal{C}\ell(p,q) = \mathcal{C}\ell^{\mathbb{C}}(p,q)$). For p = n, q = 0 we use notation $\mathcal{C}\ell^{\mathbb{F}}(n) = \mathcal{C}\ell^{\mathbb{F}}(n,0)$. Elements e^a are called *generators* of Clifford algebra $\mathcal{C}\ell^{\mathbb{F}}(p,q)$. $\mathcal{C}\ell^{\mathbb{R}}(p,q)$ is real Clifford algebra and $\mathcal{C}\ell(p,q)$ is complex Clifford algebra.

Any element $U \in \mathcal{C}\!\!\mathcal{C}^{\mathbb{F}}(p,q)$ can be written in the form

$$U = ue + u_a e^a + \sum_{a_1 < a_2} u_{a_1 a_2} e^{a_1 a_2} + \dots + u_{1\dots n} e^{1\dots n}$$
(3)

with coefficients $u, u_a, u_{a_1a_2}, \ldots, u_{1...n} \in \mathbb{F}$, which numbered by ordered multi-indices of length from 0 to n.

Denote by $\mathcal{C}_{k}^{\mathbb{F}}(p,q)$, $k = 0, \ldots n$, subspaces of the vector space $\mathcal{C}^{\mathbb{F}}(p,q)$ that span over basis elements $e^{a_1 \ldots a_k}$. Elements of $\mathcal{C}_{k}^{\mathbb{F}}(p,q)$ are called elements of rank k. Sometimes it is suitable to denote $\stackrel{k}{U} \in \mathcal{C}_{k}^{\mathbb{F}}(p,q)$. We have

$$\mathcal{C}\!\ell^{\mathbb{F}}(p,q) = \mathcal{C}\!\ell^{\mathbb{F}}_{0}(p,q) \oplus \ldots \oplus \mathcal{C}\!\ell^{\mathbb{F}}_{n}(p,q) = \mathcal{C}\!\ell^{\mathbb{F}}_{\text{even}}(p,q) \oplus \mathcal{C}\!\ell^{\mathbb{F}}_{\text{odd}}(p,q),$$

where

$$\begin{aligned} \mathcal{C}\!\ell^{\mathbb{F}}_{\text{even}}(p,q) &= \mathcal{C}\!\ell^{\mathbb{F}}_{0}(p,q) \oplus \mathcal{C}\!\ell^{\mathbb{F}}_{2}(p,q) \oplus \dots \\ \mathcal{C}\!\ell^{\mathbb{F}}_{\text{odd}}(p,q) &= \mathcal{C}\!\ell^{\mathbb{F}}_{1}(p,q) \oplus \mathcal{C}\!\ell^{\mathbb{F}}_{3}(p,q) \oplus \dots \end{aligned}$$

and

$$\dim \mathcal{C}\!\ell_k^{\mathbb{F}}(p,q) = C_n^k, \quad \dim \mathcal{C}\!\ell_{\text{even}}^{\mathbb{F}}(p,q) = \dim \mathcal{C}\!\ell_{\text{odd}}^{\mathbb{F}}(p,q) = 2^{n-1},$$

 C_n^k are binomial coefficients. Let us take the antisymmetric coefficients $u_{a_1...a_k} = u_{[a_1...a_k]}$. Consider an element

$$\overset{k}{U} = \sum_{a_1 < \dots < a_k} u_{a_1 \dots a_k} e^{a_1 \dots a_k} = \sum_{a_1 < \dots < a_k} u_{a_1 \dots a_k} e^{a_1} \dots e^{a_k}.$$

We have

$$\overset{k}{U} = \frac{1}{k!} u_{a_1 \dots a_k} e^{a_1} \dots e^{a_k} = \frac{1}{k!} u_{a_1 \dots a_k} e^{a_1} \wedge \dots \wedge e^{a_k}.$$

for U from (3).

2 A Hermitian conjugation operation

Using the projection operation onto the one-dimensional subspace $\mathcal{C}\ell_0(p,q)$, we get a trace of an element $U \in \mathcal{C}\ell(p,q)$

Tr :
$$\mathcal{C}\ell(p,q) \to \mathbb{C}$$
,

where

$$\operatorname{Tr}(ue + u_a e^a + \ldots) = u.$$

This operation has the following properties:

$$\operatorname{Tr}(UV) = \operatorname{Tr}(VU), \quad \operatorname{Tr}([U,V]) = 0.$$

Now we introduce the structure of unitary (Euclidian) space on Clifford algebra [1]. Let $\dagger : \mathcal{C}\!\ell^{\mathbb{F}}(p,q) \to \mathcal{C}\!\ell^{\mathbb{F}}(p,q)$ be a linear Hermitian conjugation operation of the Clifford algebra elements. Following rules define a Hermitian conjugation operation of Clifford algebra elements:

$$(e^{i_1\dots i_k})^{\dagger} = e_{i_k}\dots e_{i_1}, \text{ and } \lambda^{\dagger} = \bar{\lambda}, \quad \forall \lambda \in \mathbb{C},$$

 $(U+V)^{\dagger} = V^{\dagger} + U^{\dagger} \quad \forall U, V \in \mathcal{C}\ell^{\mathbb{F}}(p,q),$

where $e_a = \eta_{ab}e^b$. We say that $\dagger : \mathcal{C}\ell^{\mathbb{F}}(p,q) \to \mathcal{C}\ell^{\mathbb{F}}(p,q)$ is the operation of *Hermitian conjugation* of Clifford algebra elements. It is easy to see that

$$(UV)^{\dagger} = V^{\dagger}U^{\dagger}, \quad U^{\dagger\dagger} = U.$$

Now we can define a Hermitian (Euclidian) scalar product of Clifford algebra elements by the formula

$$(U,V) = \operatorname{Tr}(V^{\dagger}U).$$

In this case we have

$$(e^{i_1\dots i_k}, e^{i_1\dots i_k}) = \operatorname{Tr}(e_{i_k}\dots e_{i_1}e^{i_1}\dots e^{i_k}) = \operatorname{Tr}(e) = 1,$$

(no summation w.r.t. i_1, \ldots, i_k). Basis (1) of $\mathcal{C}\ell^{\mathbb{F}}(p,q)$ is orthonormal with respect to this scalar product and property (7) is valid

$$(U,U) = \sum_{k=0}^{n} \sum_{a_1 < \dots < a_k} |u_{a_1 \dots a_k}|^2 > 0.$$
(4)

For generators e^a formula (6) gives

$$(e^{a})^{\dagger} = e^{a}$$
 for $a = 1, \dots, p;$ (5)
 $(e^{a})^{\dagger} = -e^{a}$ for $a = p + 1, \dots, n.$

3 Pseudounitary groups of Clifford algebras

Let r, s be nonnegative integers, r + s > 0. Consider a Lie group of pseudounitary matrices and a Lie group of special pseudounitary matrices:

$$U(r,s) = \{U \in \operatorname{Mat}(n, \mathbb{C}) : U^{\dagger}\beta U = \beta\},\$$

$$SU(r,s) = \{U \in U(r,s) : \det U = 1\},\$$

where n = r + s and $\beta = \text{diag}(1, \dots, 1, -1, \dots, -1)$ is a diagonal matrix with r pieces of 1 and s pieces of -1 on the diagonal.

Now let us consider the four sets of Clifford algebra elements:

$$\begin{aligned} & \mathcal{W}(\mathcal{C}(p,q)) &= \{ U \in \mathcal{C}(p,q) : U^*U = e \}, \\ & \mathcal{SW}(\mathcal{C}(p,q)) &= \{ U \in \mathcal{W}(\mathcal{C}(p,q)) : \operatorname{Det} U = 1 \}, \\ & \operatorname{w}(\mathcal{C}(p,q)) &= \{ U \in \mathcal{C}(p,q) : U^* = -U \}, \\ & \operatorname{sw}(\mathcal{C}(p,q)) &= \{ U \in \operatorname{w}(\mathcal{C}(p,q)) : \operatorname{Tr} U = 0 \}. \end{aligned}$$

Note that the sets $W(\mathcal{C}\ell(p,q))$ and $SW(\mathcal{C}\ell(p,q))$ are closed with respect to the Clifford product and the sets $w(\mathcal{C}\ell(p,q))$ and $sw(\mathcal{C}\ell(p,q))$ are closed with respect to the commutator. The sets $W(\mathcal{C}\ell(p,q))$ and $SW(\mathcal{C}\ell(p,q))$ can be considered as Lie groups

and the sets $w(\mathcal{C}(p,q))$ and $sw(\mathcal{C}(p,q))$ can be considered as Lie algebras of the Lie groups $W(\mathcal{C}(p,q))$ and $SW(\mathcal{C}(p,q))$.

The group $W(\mathcal{C}\ell(p,q))$ is called the pseudounitary group of Clifford algebra $\mathcal{C}\ell(p,q)$. The group $SW(\mathcal{C}\ell(p,q))$ is called the special pseudounitary group of Clifford algebra.

4 Symplectic Lie groups and their Lie algebras

Let us consider a *real symplectic Lie group* of $n \times n$ -matrices, n = 2m and let us consider the real Lie algebra of this group

$$Sp(n, \mathbb{R}) = \{ U \in Mat(n, \mathbb{R}) : U^T J U = J \}, sp(n, \mathbb{R}) = \{ u \in Mat(n, \mathbb{R}) : u^T J = -Ju \},$$

where U^T is a transpose matrix, J is the matrix

$$J = \left(\begin{array}{cc} 0 & -I_m \\ I_m & 0 \end{array}\right)$$

and I_m is an identity $m \times m$ - matrix. Note that $J^2 = -1$ (1 is the identity 4×4 matrix). Mostafazadeh [4] shows that the symplectic group sp(2n, R) is a subgroup of the pseudounitary group U(n, n).

5 A symplectic group of Clifford algebra

Now define the two sets of Clifford algebra elements

$$\begin{aligned} \operatorname{Sp}(\mathcal{C}(1,3)) &= \{ V \in \mathcal{C}\!\ell^{\mathbb{R}}_{\operatorname{even}}(1,3) \oplus i \mathcal{C}\!\ell^{\mathbb{R}}_{\operatorname{odd}}(1,3) : V^*V = e \}, \\ \operatorname{sp}(\mathcal{C}\!\ell(1,3)) &= \{ v \in i \mathcal{C}\!\ell^{\mathbb{R}}_1(1,3) \oplus \mathcal{C}\!\ell^{\mathbb{R}}_2(1,3) \}. \end{aligned}$$

We have $\operatorname{Sp}(\mathcal{C}(1,3)) \subset W(\mathcal{C}(1,3))$. Suppose that this set is closed with respect to the product; then $\operatorname{Sp}(\mathcal{C}(1,3))$ can be considered as a group (the Lie group)

with respect to the product. This group is called the symplectic group of Clifford algebra $\mathcal{C}\ell(1,3)$. The relation of this group with the matrix symplectic group will be explained in Theorem 1.

In [1] we see that the set $\operatorname{sp}(\mathcal{C}(1,3)) \subset \operatorname{w}(\mathcal{C}(1,3))$ is closed with respect to the commutator [u, v] = uv - vu. So $\operatorname{sp}(\mathcal{C}(1,3))$ is a Lie algebra with respect to the commutator.

Theorem 1. The group $\operatorname{Sp}(\mathcal{C}(1,3))$ is isomorphic to the group $\operatorname{Sp}(4,\mathbb{R})$ and the Lie algebra $\operatorname{sp}(\mathcal{C}(1,3))$ is isomorphic to the Lie algebra $\operatorname{sp}(4,\mathbb{R})$:

$$Sp(\mathcal{C}(1,3)) \simeq Sp(4,\mathbb{R}),$$
(6)
$$sp(\mathcal{C}(1,3)) \simeq sp(4,\mathbb{R}).$$

Proof. It is well known [3] that the complex Clifford algebra $\mathcal{C}\ell(1,3)$ is isomorphic to the algebra of complex matrices $Mat(4,\mathbb{C})$. Let e^a , a = 0, 1, 2, 3 be generators and e be the identity element of Clifford algebra $\mathcal{C}\ell(1,3)$. Consider the matrix representation γ of Clifford algebra elements

$$\gamma : \mathcal{C}(1,3) \to \operatorname{Mat}(4,\mathbb{C}),$$

such that

$$\gamma(A + B) = \gamma(A) + \gamma(B),$$

$$\gamma(\lambda A) = \lambda \gamma(A),$$

$$\gamma(AB) = \gamma(A)\gamma(B),$$

$$\gamma(e) = \mathbf{1},$$

where A, B are arbitrary elements of $\mathcal{C}(1,3)$; λ is a complex number; 1 is the identity

 4×4 -matrix and generators of Clifford algebra represented by the following matrices:

$$\begin{split} \gamma(e^{0}) &= \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & i \\ -i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \end{pmatrix}, \qquad \gamma(e^{1}) &= \begin{pmatrix} 0 & -i & 0 & 0 \\ -i & 0 & 0 & 0 \\ 0 & 0 & 0 & i \\ 0 & 0 & i & 0 \end{pmatrix}, \\ \gamma(e^{2}) &= \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & i \\ i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{pmatrix}, \qquad \gamma(e^{3}) &= \begin{pmatrix} -i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \\ 0 & 0 & i & 0 \\ 0 & 0 & 0 & -i \end{pmatrix}. \end{split}$$

We say that the matrix representation γ is a modified Majorana's representation. Let us remind that the representation γ defines the isomorphism of $\mathcal{C}(1,3)$ and $Mat(4,\mathbb{C})$.

Generators e^a of Clifford algebra satisfy the following formulas

$$e^{a}e^{b} + e^{b}e^{a} = 2\eta^{ab}e, \quad a, b = 0, 1, 2, 3.$$

So the matrices $\gamma^a = \gamma(e^a)$, a = 0, 1, 2, 3 satisfy the conditions

$$\gamma^a \gamma^b + \gamma^b \gamma^a = 2\eta^{ab} \mathbf{1}, \quad a, b = 0, 1, 2, 3.$$

Note that the sixteen matrices

$$\mathbf{1}, \ \gamma^{a}, \ \gamma^{a}\gamma^{b}, \ \gamma^{a}\gamma^{b}\gamma^{c}, \ \gamma^{0}\gamma^{1}\gamma^{2}\gamma^{3}, \quad a < b < \dots$$

are linearly independent and form the basis of the matrix algebra $Mat(4, \mathbb{C})$.

Note also that the matrices γ^a satisfy to the conditions

$$(\gamma^{0})^{\dagger} = \gamma^{0}, \quad (\gamma^{k})^{\dagger} = -\gamma^{k}, \quad k = 1, 2, 3.$$

Therefore the representation γ is consistent with the Hermitian conjugation operation

$$\gamma(A)^{\dagger} = \gamma(A^{\dagger}),$$

where $\gamma(A)^{\dagger}$ is the Hermitian conjugated matrix, and $A^{\dagger} = e^0 A^* e^0$ is the Hermitian conjugated element of Clifford algebra $\mathcal{C}\ell(1,3)$.

Denote by γ^{-1} the inverse map

$$\gamma^{-1}$$
 : Mat $(4, \mathbb{C}) \to \mathcal{C}\ell(1, 3)$.

Matrices $i\gamma^a$ are real, therefore the maps γ, γ^{-1} give the isomorphism

$$\mathcal{C}^{\mathbb{R}}_{\text{even}}(1,3) \oplus i\mathcal{C}^{\mathbb{R}}_{\text{odd}}(1,3) \simeq \text{Mat}(4,\mathbb{R}).$$

Our the representation γ is such that

$$J = i\gamma(e^0),$$

where the matrix J is used in the definition of the group of symplectic matrices.

For any $A \in \mathcal{C}\ell(1,3)$ we have

$$A^{\dagger} = e^{0}A^{*}e^{0}, orA^{*} = e^{0}A^{\dagger}e^{0}.$$

Therefore for any element $A \in \mathcal{C}\!\!\ell^{\mathbb{R}}_{\text{even}}(1,3) \oplus i\mathcal{C}\!\!\ell^{\mathbb{R}}_{\text{odd}}(1,3)$ we have

$$\gamma(A^*) = \gamma(e^0)\gamma(A^\dagger)\gamma(e^0) = -J\gamma(A)^T J.$$

Consider the definition of group $\operatorname{Sp}(\mathcal{C}(1,3))$. The condition $A^*A = e$ leads to the relation

$$\gamma(A^*)\gamma(A) = \gamma(e).$$

Using this relation and conditions

$$\gamma(A^*) = -J\gamma(A)^T J, \quad J^2 = -\mathbf{1}$$

we get

$$\gamma(A)^T J \gamma(A) = J.$$

From the definition of the group $\text{Sp}(4, \mathbb{R})$ it follows that the group $\text{Sp}(\mathcal{C}(1,3))$ is isomorphic to the group $\text{Sp}(4, \mathbb{R})$

$$\operatorname{Sp}(\mathcal{C}\ell(1,3)) \simeq \operatorname{Sp}(4,\mathbb{R}).$$

Let us remind the definition of Lie algebra $w(\mathcal{C}(1,3))$ of the pseudounitary group $W(\mathcal{C}(1,3))$

 $w(\mathcal{C}(1,3)) = \{ w \in \mathcal{C}(1,3) : w^* = -w \}.$

This definition is equivalent to the following:

$$\mathbf{w}(\mathcal{C}\ell(1,3)) = \{ w \in i\mathcal{C}\ell_0^{\mathbb{R}}(1,3) \oplus i\mathcal{C}\ell_1^{\mathbb{R}}(1,3) \oplus \mathcal{C}\ell_2^{\mathbb{R}}(1,3) \oplus \mathcal{C}\ell_3^{\mathbb{R}}(1,3) \oplus i\mathcal{C}\ell_4^{\mathbb{R}}(1,3) \}.$$

Note that the definition of Lie algebra $\operatorname{sp}(\mathcal{C}(1,3))$ can be written in the form

$$\operatorname{sp}(\mathcal{C}\ell(1,3)) = \{ w \in \operatorname{w}(\mathcal{C}\ell(1,3)) : w \in \mathcal{C}\ell^{\mathbb{R}}_{\operatorname{even}}(1,3) \oplus i\mathcal{C}\ell^{\mathbb{R}}_{\operatorname{odd}}(1,3) \}.$$

Hence, $\operatorname{sp}(\mathcal{C}(1,3))$ is the intersection of following sets:

$$\{i\mathcal{C}_{0}^{\mathbb{R}}(1,3) \oplus i\mathcal{C}_{1}^{\mathbb{R}}(1,3) \oplus \mathcal{C}_{2}^{\mathbb{R}}(1,3) \oplus \mathcal{C}_{3}^{\mathbb{R}}(1,3) \oplus i\mathcal{C}_{4}^{\mathbb{R}}(1,3)\} \cap \{\mathcal{C}_{even}^{\mathbb{R}}(1,3) \oplus i\mathcal{C}_{odd}^{\mathbb{R}}(1,3)\}.$$

Any element of the Lie algebra $w \in C\!\ell^{\mathbb{R}}_{even}(1,3) \oplus i\mathcal{C}\!\ell^{\mathbb{R}}_{odd}(1,3)$ represents as a real matrix. Therefore using the matrix representation γ for the expression $w^* = -w$ we get the relations

$$-J\gamma(w)^T J = -\gamma(w), \quad \text{or} \quad \gamma(w)^T J = -J\gamma(w).$$

It is clear that the last expression defines the Lie algebra $sp(4, \mathbb{R})$. Hence the Lie algebra $sp(\mathcal{C}(1,3))$ is isomorphic to the Lie algebra $sp(4, \mathbb{R})$. This completes the proof.

References

- Marchuk N. G., Shirokov D. S. Unitary spaces on Clifford algebras, Adv. appl. Clifford alg., 18, (2008), pp.237-254, arXiv:0705.1641v1 [math-ph], (2007).
- [2] Hestenes D., Space-Time Algebra, Gordon and Breach, New York, (1966).
- [3] Lounesto P., Clifford Algebras and Spinors, Cambridge Univ. Press (1997, 2001)
- [4] Mostafazadeh A., Pseudo-Unitary Operators and Pseudo-Unitary Quantum Dynamics, arXiv:0302050v2 [math-ph], (2003).