Discrete realization of group symmetric LOCC-detection of maximally entangled state

Masahito Hayashi^{1,*}

¹Graduate School of Information Sciences, Tohoku University, Aoba-ku, Sendai, 980-8579, Japan

Group symmetric LOCC measurement for detecting maximally entangled state is considered. Usually, this type measurement has continuous-valued outcomes. However, any realizable measurement has finite-valued outcomes. This paper proposes discrete realizations of such a group symmetric LOCC measurement.

PACS numbers: 03.65.Wj,03.65.Ud,02.20.-a

I. INTRODUCTION

Testing of maximally entangled state is a useful method for guaranteeing the quality of generated maximally entangled states. However, if we require a group symmetric condition for this method, the optimal test often requires infinite-valued measurement. Since any realizable measurement has a finite number of outcomes, it is needed to discretize the optimal measurement.

Now, we focus on the bipartite system $\mathcal{H}_d \otimes \mathcal{H}_d$, in which, the party A and B have the computational bases $\{|i\rangle_A\}_{i=0}^{d-1}$ and $\{|i\rangle_B\}_{i=0}^{d-1}$, respectively. When our target is testing whether the generated state is sufficiently close to the maximal entangled state

$$|\phi^0_{AB}\rangle = \frac{1}{\sqrt{d}}\sum_{i=0}^{d-1}|i\rangle_A\otimes|i\rangle_B$$

under a group symmetric condition, the optimal test can be given by

$$T_{inv}^{1,A\to B} := \int d|\varphi \otimes \overline{\varphi}\rangle \langle \varphi \otimes \overline{\varphi}|\nu(d\varphi)$$
$$= |\phi_{A,B}^{0}\rangle \langle \phi_{A,B}^{0}| + \frac{1}{d+1}(I - |\phi_{A,B}^{0}\rangle \langle \phi_{A,B}^{0}|),$$
(1)

where ν is the group invariant probability measure on the set of pure states, and φ and $\overline{\varphi}$ are given as $\varphi = \sum_{i=0}^{d-1} \varphi_i |i\rangle_A$ and $\overline{\varphi} = \sum_{i=0}^{d-1} \overline{\varphi_i} |i\rangle_B$. This measurement can be realized by the following procedure. In the first step, the system A performs the local group covariant measurement $\int d|\varphi\rangle\langle\varphi|\nu(d\varphi)$, and sends the system Bthe outcome φ . In the second step, the system B performs the two-valued measurement $\{|\overline{\varphi}\rangle\langle\overline{\varphi}|, I - |\overline{\varphi}\rangle\langle\overline{\varphi}|\}$. When Bob obtains the event corresponding to $\{|\overline{\varphi}\rangle\langle\overline{\varphi}|,$ we support the maximal entangled state $|\phi_{AB}^0\rangle$.

This detection procedure can be generalize as follows. First, Alice performs a local measurement:

$$M = \{ p_i | u_i \rangle \langle u_i | \}_i, \quad ||u_i|| = 1, \quad 0 \le p_i \le 1$$

and sends Bob the outcome *i*. Bob perform the twovalued measurement $\{ |\overline{u_i}\rangle \langle \overline{u_i} |, I - |\overline{u_i}\rangle \langle \overline{u_i} | \}$. When Bob obtain the event corresponding to $|\overline{u_i}\rangle\langle\overline{u_i}|$, we support the maximal entangled state $|\phi_{AB}^0\rangle$. This test can be written as the positive semi-definite matrix T(M):

$$T(M) \stackrel{\text{def}}{=} \sum_{i} p_i |u_i \otimes \overline{u_i}\rangle \langle u_i \otimes \overline{u_i}|.$$
(2)

Indeed, when the local dimension d is 2, D'Ariano et al. [1] and Hayashi et al.[2] obtained the discrete ownway LOCC realization of the test $T_{inv}^{1,A\to B}$ as test T(M)with an appropriate choice of the local measurement M. However, its general dimensional case was an open problem. In this paper, employing the concepts of symmetric informationally complete POVM (SIC-POVM) and mutually unbiased bases (MUB), we propose discrete ownway LOCC realizations of $T_{inv}^{1,A\to B}$. Also, the optimality of the proposed realization scheme is shown.

Next, we consider the case when Alice's system (Bob's system) is given as $\mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2}$ ($\mathcal{H}_{B_1} \otimes \mathcal{H}_{B_2}$) and the dimensions of all components coincide, i.e., dim $\mathcal{H}_{A_1} = \dim \mathcal{H}_{A_2} = \dim \mathcal{H}_{B_1} = \dim \mathcal{H}_{B_2} = d$. In this case, we focus on the covariant POVM $M_{cov,u}^2$:

$$\begin{split} & M_{cov,u}^2(\,dg_1\,dg_2) \\ \stackrel{\text{def}}{=} d^2(g_1\otimes g_2)|u\rangle\langle u|(g_1\otimes g_2)^*\nu(\,dg_1)\nu(\,dg_2), \end{split}$$

where the vector u is a maximally entangled state and ν is the group invariant probability measure on SU(d). The optimal test is given as the test $T_{inv}^{2,A\to B} \stackrel{\text{def}}{=} T(M_{cov,u}^2)$, which has the form [3]:

$$T_{inv}^{2,A \to B} = |\phi_{A,B}^{0}\rangle\langle\phi_{A,B}^{0}| \otimes |\phi_{A,B}^{0}\rangle\langle\phi_{A,B}^{0}| + \frac{1}{d^{2} - 1}(I - |\phi_{A,B}^{0}\rangle\langle\phi_{A,B}^{0}|) \otimes (I - |\phi_{A,B}^{0}\rangle\langle\phi_{A,B}^{0}|).$$
(3)

Indeed, The positive semi-definite matrix $T_{inv}^{2,A\to B}$ does not depend on the choice of the maximally entangled state u. In this paper, employing the concept of Clifford group, we provide a discrete own-way LOCC realization of $T_{inv}^{2,A\to B}$ when the local system is given as a composite system of a prime-dimensional system. Also, the optimality of the proposed realization scheme is shown.

^{*}Electronic address: hayashi@math.is.tohoku.ac.jp

II. DISCRETE OWN-WAY LOCC REALIZATION OF $T_{inv}^{1,A \rightarrow B}$

A. Realizing scheme by SIC-POVM

In order to design the test $T_{inv}^{1,A\to B}$, we focus on the concept "symmetric informationally complete POVM (SIC-POVM)". A rank-one POVM $\{p_i|u_i\rangle\langle u_i|\}$ on $\mathcal{H}_A = \mathbb{C}^d$ is called a *symmetric informationally complete POVM* (SIC-POVM), if it satisfies the following conditions:

$$\#\{i\} = d^2,$$

$$p_i = \frac{1}{d}$$

$$|\langle u_i | u_j \rangle|^2 = \frac{1}{d+1} \text{ for } i \neq j$$
(4)

Currently, an SIC-POVM analytically is constructed when the dimension d is 2,3[5, 7],4[4, 7],5[7],6[6],7[8], 8[5], or 19[8]. Also, its existence is numerically verified up to d = 45[4]. As is shown in Appendix A, any SIC-POVM $M_{sic} = \{p_i|u_i\rangle\langle u_i|\}_i$ satisfies

$$T(M_{sic}) = T_{inv}^{1,A \to B},\tag{5}$$

that is, the test $T_{inv}^{1,A\to B}$ can be realized by an SIC-POVM. Moreover, if a POVM $M = \{M_i\}_i$ on \mathcal{H}_A satisfies

$$T(M) = T_{inv}^{1,A \to B},$$

the inequality

$$\#\{i\} \ge d^2$$

holds. This is because the rank of the operator $T_{inv}^{1,A\to B}$ (which equal d^2) is less than the number of the elements of POVM M_i . Hence, we obtain

$$\min\{\#\{i\}|T(\{M_i\}_i) = T_{inv}^{1,A \to B}\} = d^2 \tag{6}$$

if there exists an SIC-POVM on \mathbb{C}^d . That is, the proposed realizing scheme by SIC-POVM is optimal in the sense of (6).

B. Realizing scheme by MUB

However, any SIC-POVM is not a randomized combination of projection valued measures as well as a projection valued measure. Since a projection valued measure (PVM) are more realizable than other POVM, it is more desired to design Alice's POVM as a randomized combination of PVMs. For this purpose, we focus on mutually unbiased bases. d + 1 orthonormal bases $\{\mathcal{B}_1, \ldots, \mathcal{B}_{d+1}\}$ are called mutually unbiased bases (MUB) if

$$|\langle u|v\rangle|^2 = \frac{1}{d}, \forall u \in \mathcal{B}_i, \forall v \in \mathcal{B}_j, i \neq j.$$

The existence of MUB is shown when d is a prime[9] or a prime power[10]. Bandyopadhyay *et al.* gave a more explicit form in these cases [11]. Any mutually unbiased bases $\{\mathcal{B}_1, \ldots, \mathcal{B}_{d+1}\}$ make the POVM $M_{\mathcal{B}_1, \ldots, \mathcal{B}_k}$, *i.e.*,

$$M_{\mathcal{B}_1,\dots,\mathcal{B}_{d+1}} = \left\{ \frac{1}{d+1} |u_{i,j}\rangle \langle u_{i,j}| \right\}_{i,j},$$

where $\mathcal{B}_j = \{u_{1,j}, \dots, u_{d,j}\}$. This POVM always produces the desired test $T_{inv}^{1,A \to B}$ as

$$T(M_{\mathcal{B}_1,\dots,\mathcal{B}_{d+1}}) = T_{inv}^{1,A \to B},\tag{7}$$

which is shown in Appendix B. This construction of the test $T_{inv}^{1,A\to B}$ is optimal in the following sense. Let $\{M^j\}$ be the set of projection-valued measures. A randomized combination of $\{M^j\}$, *i.e.*, $M = \sum_j p_j M_j$ satisfies

$$T(M) = T_{inv}^{1,A \to B}.$$
(8)

Then, as is proven in Appendix C,

$$\#\{j\} \ge d+1,$$
 (9)

which implies the optimality of the POVM consisting of MUB. Hence,

$$\min_{M_j:\text{PVM}} \left\{ \#\{j\} \left| T\left(\sum p_j M_j\right) = T_{inv}^{1,A \to B} \right\} = d+1 \quad (10)$$

if d is a prime or a prime power. That is, the proposed realizing scheme by MUB is optimal in the sense of (10).

III. DISCRETE OWN-WAY LOCC REALIZATION OF $T_{inv}^{2,A \rightarrow B}$

Next, we proceed to the case when both local systems consist of two subsystems. Given a finite group G and its projective representation f on $\mathcal{H}_{A_1} = \mathbb{C}^d$, by regarding \mathcal{H}_{A_2} as the dual space of \mathcal{H}_{A_1} , the matrix f(g) can be regarded as an element $|f(g)\rangle$ of $\mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2}$.

Theorem 1 We assume the two conditions: (1) The representation f is irreducible. (2) The action $f \otimes \overline{f}$ of G to $\mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2}$ has only two irreducible components, i.e., the irreducible subspaces of $\mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2}$ for the action

$$v_1 \otimes v_2 \to f(g)v_1 \otimes \overline{f(g)}v_2$$

are only the one-dimensional space $\langle \phi^0_{A_1,A_2} \rangle$ and its orthogonal space $\langle \phi^0_{A_1,A_2} \rangle^{\perp}$. Then, the resolution $M_f = \left\{ \frac{d^2}{|G|} \left| \frac{1}{\sqrt{d}} f(g) \right\rangle \left\langle \frac{1}{\sqrt{d}} f(g) \right| \right\}_{g \in G}$ satisfies the condition for a POVM, and

$$T(M_f) = T_{inv}^{2,A \to B}.$$
(11)

Its proof is given in Appendix D. This theorem yields a discrete own-way LOCC realization of $T_{inv}^{2,A\to B}$ from the representation f satisfying the above two conditions.

For example, Clifford group satisfies this assumption. For readers' convenience, we give its definition and prove that Clifford group satisfies this assumption. Clifford group C(d) for d-dimensional system is given by

$$C(d) := \{ U \in U(d) | U \operatorname{GP}(d) U^{\dagger} = \operatorname{GP}(d) \}$$
$$\operatorname{GP}(d) := \{ e^{\sqrt{-1}\xi} W(i,j) | \xi \in \mathbb{R}, i, j \in \mathbb{Z} \}$$
$$\operatorname{I}(d) := \{ e^{\sqrt{-1}\xi} | \xi \in \mathbb{R} \},$$

where

$$\begin{split} Z := \sum_{j=0}^{d-1} \omega^j |j\rangle \langle j|, \quad X := \sum_{j=0}^{d-1} |j+1\rangle \langle j| \\ W(i,j) := X^i Z^j \end{split}$$

and ω is the *d*-th root of 1. As is shown in Appendix E, the natural representation of the group C(d) satisfies the conditions (1) and (2). Then, the natural projective representation of the group C(d)/I(d) also satisfies the conditions (1) and (2). As is shown in Lemma 5 in Appleby [8], when *d* is prime, the cardinality |C(d)/I(d)| is $d^3(d^2 - 1)$. In the general case,

$$|C(d)/I(d)| = d^2 \left(\sum_{n=0}^{d-1} \nu(n,d)\nu(n+1,d) \right),$$

where $\nu(n, d)$ is the number of distinct ordered pairs $(x, y) \in \mathbb{Z}_d^2$ such that $xy = n \pmod{d}$.

IV. DISCUSSION

This paper has treated discretization of onw-way LOCC protocols. Using the concepts of symmetric informationally complete POVM (SIC-POVM), mutually unbiased bases (MUB), and Clifford group, we have proposed discrete own-way LOCC realizations of $T_{inv}^{1,A\to B}$ and $T_{inv}^{2,A\to B}$. This result indicates the importance of these concept in discrete mathematics. Since the existence of SIC-POVM and MUB is proven in limited cases, we cannot construct a discrete own-way LOCC realization of $T_{inv}^{1,A\to B}$ in the general case. Thus, further investigation for these concepts are required. While the optimal test is given as $T_{inv}^{3,A\to B}$ when the

While the optimal test is given as $T_{inv}^{S,A\to D}$ when the local system consists of three subsystems by Hayashi [3], its discretization has not been obtained. Since the optimal test $T_{inv}^{3,A\to B}$ is closely related to GHZ state[3], its discretization may be related to GHZ state. Its construction remains as a future research.

Further, the optimal protocol is often given as a protocol with infinite elements in quantum information. In such a case, it is required to discretize this protocol. This kind of discretization is an interesting interdisciplinary topic between quantum information and discrete mathematics.

Acknowledgment

This research was partially supported by a Grant-in-Aid for Scientific Research on Priority Area 'Deepening and Expansion of Statistical Mechanical Informatics (DEX-SMI)', No. 18079014 and a MEXT Grant-in-Aid for Young Scientists (A) No. 20686026.

APPENDIX A: PROOF OF (5)

First, we show that $u_1 \otimes \overline{u_1}, \ldots, u_{d^2} \otimes \overline{u_{d^2}}$ are linearly independent. We choose complex numbers a_1, \ldots, a_{d^2} such that

$$\sum_i a_i u_i \otimes \overline{u_i} = 0$$

Taking trace, we have

$$a_1 + \sum_{i \neq 1} a_i = 0$$

On the other hand,

$$0 = \langle u_1 \otimes \overline{u_1} | \sum_i a_i u_i \otimes \overline{u_i} \rangle = a_1 + \frac{1}{d+1} \sum_{i \neq 1} a_i.$$

Hence, we obtain $a_1 = 0$. Similarly, we can show $a_i = 0$, which implies the linear independence.

Since the dimension of $\mathcal{H}_A \otimes \mathcal{H}_B$ is d^2 , any element of $\mathcal{H}_A \otimes \mathcal{H}_B$ can be expressed as

$$\sum_j a_i u_i \otimes \overline{u_i}.$$

We can calculate

$$\left\langle \sum_{i} a_{i} u_{i} \otimes \overline{u_{i}} \middle| T(M_{sic}) \middle| \sum_{j} a_{j} u_{j} \otimes \overline{u_{j}} \right\rangle$$
$$= \left\langle \sum_{i} a_{i} u_{i} \otimes \overline{u_{i}} \middle| \left(\sum_{k} \frac{1}{d} \left| u_{k} \otimes \overline{u_{k}} \right\rangle \left\langle u_{k} \otimes \overline{u_{k}} \right| \right) \middle| \sum_{j} a_{j} u_{j} \otimes \overline{u_{j}} \right\rangle$$
$$= \frac{d+2}{(d+1)^{2}} \left| \sum_{k} a_{k} \right|^{2} + \frac{d}{(d+1)^{2}} \sum_{k} |a_{k}|^{2}.$$

On the other hand, its norm is calculated as

$$\left\|\sum_{j} a_{j} u_{j} \otimes \overline{u_{j}}\right\| = \frac{1}{d+1} \left|\sum_{k} a_{k}\right|^{2} + \frac{d}{d+1} \sum_{k} |a_{k}|^{2}.$$

Since

$$\langle \phi^0_{A,B} | \sum_j a_j u_j \otimes \overline{u_j} \rangle \bigg|^2 = \frac{1}{d} \left| \sum_j a_j \right|^2,$$

we obtain

$$\begin{split} &\left\langle \sum_{i} a_{i} u_{i} \otimes \overline{u_{i}} \middle| T_{inv}^{1,A \to B} \middle| \sum_{j} a_{j} u_{j} \otimes \overline{u_{j}} \right\rangle \\ &= \left\langle \sum_{i} a_{i} u_{i} \otimes \overline{u_{i}} \middle| \left(\frac{d}{d+1} |\phi_{A,B}^{0}\rangle \langle \phi_{A,B}^{0}| + \frac{1}{d+1} I \right) \right. \\ &\left| \sum_{j} a_{j} u_{j} \otimes \overline{u_{j}} \right\rangle \\ &= \frac{d}{d+1} \frac{1}{d} \left| \sum_{j} a_{j} \right|^{2} \\ &\left. + \frac{1}{d+1} \left(\frac{1}{d+1} \left| \sum_{k} a_{k} \right|^{2} + \frac{d}{d+1} \sum_{k} |a_{k}|^{2} \right) \right. \\ &= \left\langle \sum_{i} a_{i} u_{i} \otimes \overline{u_{i}} \middle| T(M_{sic}) \left| \sum_{j} a_{j} u_{j} \otimes \overline{u_{j}} \right\rangle. \end{split}$$

Therefore, we obtain (5).

APPENDIX B: PROOF OF (7)

We focus on the subspace $\langle \phi_{A,B}^0 \rangle^{\perp}$ orthogonal to $\phi_{A,B}^0$. The subspace $\mathcal{B}'_j = \langle u_{1,j} \otimes \overline{u_{1,j}} - \frac{1}{d}\phi_{A,B}^0, \dots, u_{d-1,j} \otimes \overline{u_{d-1,j}} - \frac{1}{d}\phi_{A,B}^0 \rangle$ belongs to the subspace $\langle \phi_{A,B}^0 \rangle^{\perp}$, and its dimension is d-1. Since

$$\langle u_{i,j} \otimes \overline{u_{i,j}} - \frac{1}{d} \phi^0_{A,B} | u_{i',j'} \otimes \overline{u_{i',j'}} - \frac{1}{d} \phi^0_{A,B} \rangle = 0, \quad j \neq j'$$
(B1)

The spaces $\mathcal{B}'_1, \ldots, \mathcal{B}'_{d+1}$ are orthogonal to each other. Since the dimension of the subspace $\langle \phi^0_{A,B} \rangle^{\perp}$ is $d^2 - 1$, the subspace $\langle \phi^0_{A,B} \rangle^{\perp}$ is spanned by the spaces $\mathcal{B}'_1, \ldots, \mathcal{B}'_{d+1}$. Therefore, any element of the space $\mathcal{H}_A \otimes \mathcal{H}_B$ can be expressed as $\sum_{j=1}^{d+1} \sum_{i=1}^d a_{i,j} u_{i,j} \otimes \overline{u_{i,j}}$. In the following, we abbreviate the sum $\sum_{j=1}^{d+1} \sum_{i=1}^d a \sum_{j=1}^d \sum_{j=1}^d a \sum_{j=1}^d \sum_{j=1}^d a \sum_{j=1}^d \sum_{j=1}^d a \sum_{j=1}^d$ We calculate

$$\left\langle \sum_{j,i} a_{i,j} u_{i,j} \otimes \overline{u_{i,j}} \middle| T(M_{\mathcal{B}_1,\dots,\mathcal{B}_{d+1}}) \middle| \sum_{j',i'} a_{i',j'} u_{i',j'} \otimes \overline{u_{i',j'}} \right\rangle$$
$$= \left\langle \sum_{j,i} a_{i,j} u_{i,j} \otimes \overline{u_{i,j}} \middle| \left(\sum_{l,k} \frac{1}{d+1} |u_{k,l} \otimes \overline{u_{k,l}}\rangle \langle u_{k,l} \otimes \overline{u_{k,l}}| \right) \right\rangle$$
$$\left| \sum_{j',i'} a_{i',j'} u_{i',j'} \otimes \overline{u_{i',j'}} \right\rangle$$
$$= \frac{1}{d+1} \sum_{l,k} \left| \sum_{j,i} \langle u_{k,l} \otimes \overline{u_{k,l}}| |a_{i,j} u_{i,j} \otimes \overline{u_{i,j}}\rangle \right|^2$$
$$= \frac{1}{d} \left| \sum_{j,i} a_{i,j} \right|^2 - \frac{1}{d(d+1)} \sum_j |\sum_i a_{i,j}|^2 + \frac{1}{d+1} \sum_{j,i} |a_{i,j}|^2$$

On the other hand, its norm is calculated as

$$\left\|\sum_{j,i} a_{i,j} u_{i,j} \otimes \overline{u_{i,j}}\right\|$$
$$= \frac{1}{d} \left|\sum_{j,i} a_{i,j}\right|^2 - \frac{1}{d} \sum_j |\sum_i a_{i,j}|^2 + \sum_{j,i} |a_{i,j}|^2$$

Since

$$\left| \left\langle \phi_{A,B}^{0} \left| \sum_{j,i} a_{i,j} u_{i,j} \otimes \overline{u_{i,j}} \right\rangle \right|^{2} = \frac{1}{d} \left| a_{i,j} u_{i,j} \right|^{2},$$

we obtain

$$\left\langle \sum_{j,i} a_{i,j} u_{i,j} \otimes \overline{u_{i,j}} \middle| T_{inv}^{1,A \to B} \middle| \sum_{j',i'} a_{i',j'} u_{i',j'} \otimes \overline{u_{i',j'}} \right\rangle$$

$$= \left\langle \sum_{j,i} a_{i,j} u_{i,j} \otimes \overline{u_{i,j}} \middle| \frac{d}{d+1} |\phi_{A,B}^{0}\rangle \langle \phi_{A,B}^{0}| + \frac{1}{d+1}I \right|$$

$$\left| \sum_{j',i'} a_{i',j'} u_{i',j'} \otimes \overline{u_{i',j'}} \right\rangle$$

$$= \frac{1}{d} \left| \sum_{j,i} a_{i,j} \right|^{2} - \frac{1}{d(d+1)} \sum_{j} |\sum_{i} a_{i,j}|^{2} + \frac{1}{d+1} \sum_{j,i} |a_{i,j}|^{2}$$

$$= \left\langle \sum_{j,i} a_{i,j} u_{i,j} \otimes \overline{u_{i,j}} \middle| T(M_{\mathcal{B}_{1},\dots,\mathcal{B}_{d+1}}) \middle| \sum_{j',i'} a_{i',j'} u_{i',j'} \otimes \overline{u_{i',j'}} \right\rangle$$

Therefore, we obtain (7).

APPENDIX C: PROOF OF (9)

Let $M^j = \{|u_{i,j}\rangle\langle u_{i,j}|\}$. We focus on the projection P to the subspace $\langle \phi^0_{A,B} \rangle^{\perp}$ orthogonal to $\phi^0_{A,B}$ and the subspace $\mathcal{B}''_{j} \stackrel{\text{def}}{=} \langle u_{1,j} \otimes \overline{u_{1,j}}, \dots, u_{d,j} \otimes \overline{u_{d,j}} \rangle$. The image $P\mathcal{B}''_{j}$ is $\langle u_{1,j} \otimes \overline{u_{1,j}} - \frac{1}{d}\phi^0_{A,B}, \dots, u_{d-1,j} \otimes \overline{u_{d-1,j}} - \frac{1}{d}\phi^0_{A,B} \rangle$. The condition (8) implies that the sum of the rank of the space $P\mathcal{B}''_{j}$ is greater than $d^2 - 1$, *i.e.*, the dimension of the space $\langle \phi^0_{A,B} \rangle^{\perp}$. Thus, $\#\{j\}(d-1) \geq d^2 - 1$, which implies the inequality (9).

APPENDIX D: PROOF OF THEOREM 1

First, we prove that M_f satisfies the condition for POVM. The irreducibility of the action f guarantees that

$$\frac{d}{|G|} \sum_{g \in G} \langle k|f(g)|l\rangle \langle l'|f(g)|k'\rangle$$
$$= \langle k| \left(\frac{d}{|G|} \sum_{g \in G} f(g)|l\rangle \langle l'|f(g)\right) |k'\rangle$$
$$= \langle k| \langle l|l'\rangle I|k'\rangle = \delta_{k,k'} \delta_{l,l'}.$$

we obtain

$$\frac{d}{|G|} \sum_{g \in G} \left| \langle f(g) | \left(\sum_{k,l} a_{k,l} | k \rangle \otimes | l \rangle \right) \right|^2$$

= $\frac{d}{|G|} \sum_{g \in G} \sum_{k,l} \sum_{k',l'} a_{k,l} \overline{a_{k',l'}} \overline{\langle k|f(g)|l \rangle} \langle l'|f(g)|k' \rangle$
= $\sum_{k,l} a_{k,l} \overline{a_{k,l}},$

which implies

$$\frac{d^2}{|G|} \sum_{g \in G} \left| \frac{1}{\sqrt{d}} f(g) \right\rangle \left\langle \frac{1}{\sqrt{d}} f(g) \right| = I_{A_1, A_2}.$$

Hence,
$$M_f = \left\{ \frac{d^2}{|G|} \left| \frac{1}{\sqrt{d}} f(g) \right\rangle \left\langle \frac{1}{\sqrt{d}} f(g) \right| \right\}_{g \in G}$$
 is a POVM.

Next, we show (11). We focus on the action of the group $G \times G$ to the total space $\mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2} \otimes \mathcal{H}_{B_1} \otimes \mathcal{H}_{B_2}$ as

$$u_1 \otimes u_2 \otimes v_1 \otimes v_2 \\ \mapsto f(g_1)u_1 \otimes \overline{f(g_2)}u_2 \otimes \overline{f(g_1)}v_1 \otimes f(g_2)v_2$$

for $u_i \in \mathcal{H}_{A_i}$, $v_i \in \mathcal{H}_{B_i}$, and any pair $(g_1, g_2) \in G \times G$. Due to the condition (2), the irreducible decomposition of the space $\mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2} \otimes \mathcal{H}_{B_1} \otimes \mathcal{H}_{B_2}$ is given as $\langle \phi^0_{A_1,B_1} \rangle \otimes \langle \phi^0_{A_2,B_2} \rangle \oplus \langle \phi^0_{A_1,B_1} \rangle \otimes \langle \phi^0_{A_2,B_2} \rangle^{\perp} \oplus \langle \phi^0_{A_1,B_1} \rangle^{\perp} \otimes \langle \phi^0_{A_2,B_2} \rangle \oplus \langle \phi^0_{A_1,B_1} \rangle^{\perp} \otimes \langle \phi^0_{A_2,B_2} \rangle^{\perp}$.

As is checked below, the test $T(M_f)$ is invariant for this action:

$$\begin{split} f(g_1) \otimes \overline{f(g_2)} \otimes \overline{f(g_1)} \otimes f(g_2) T(M_f) \left(f(g_1) \otimes \overline{f(g_2)} \otimes \overline{f(g_1)} \otimes f(g_2) \right)^{\dagger} \\ &= \frac{d^2}{|G|} \sum_{g \in G} \left| \frac{1}{d} f(g_1) f(g) f(g_2)^{\dagger} \otimes \overline{f(g_1)} f(g) f(g_2)^{\dagger} \right\rangle \left\langle \frac{1}{d} f(g_1) f(g) f(g_2)^{\dagger} \otimes \overline{f(g_1)} f(g) f(g_2)^{\dagger} \right\rangle \\ &= \frac{d^2}{|G|} \sum_{g' \in G} \left| \frac{1}{d} f(g') \otimes \overline{f(g')} \right\rangle \left\langle \frac{1}{d} f(g') \otimes \overline{f(g')} \right| = T(M_f), \end{split}$$

where we denote $g_1gg_2^{-1}$ by g'. Hence, the test $T(M_f)$

has the form

$$T(M_{f}) = a|\phi_{A_{1},B_{1}}^{0}\rangle\langle\phi_{A_{1},B_{1}}^{0}|\otimes|\phi_{A_{2},B_{2}}^{0}\rangle\langle\phi_{A_{2},B_{2}}^{0}| + b(I - |\phi_{A_{1},B_{1}}^{0}\rangle\langle\phi_{A_{1},B_{1}}^{0}|)\otimes|\phi_{A_{2},B_{2}}^{0}\rangle\langle\phi_{A_{2},B_{2}}^{0}| + c|\phi_{A_{1},B_{1}}^{0}\rangle\langle\phi_{A_{1},B_{1}}^{0}|\otimes(I - |\phi_{A_{2},B_{2}}^{0}\rangle\langle\phi_{A_{2},B_{2}}^{0}|) + d(I - |\phi_{A_{1},B_{1}}^{0}\rangle\langle\phi_{A_{1},B_{1}}^{0}|)\otimes(I - |\phi_{A_{2},B_{2}}^{0}\rangle\langle\phi_{A_{2},B_{2}}^{0}|).$$

Since f(g) is the unitary matrix, $\frac{1}{\sqrt{d}}f(g)$ is a maximally entangled state on $\mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2}$. Since $\left|\frac{1}{\sqrt{d}}f(g)\right\rangle$ is maximally entangled, Lemma 5 in Hayashi [3] yields that

$$T(M_f) = |\phi^0_{A_1,B_1} \otimes \phi^0_{A_2,B_2}\rangle \langle \phi^0_{A_1,B_1} \otimes \phi^0_{A_2,B_2}| + PT(M_f)P$$
(D1)

where

$$P \stackrel{\text{def}}{=} (I - |\phi^0_{A_2, B_2}\rangle \langle \phi^0_{A_2, B_2}|) \otimes (I - |\phi^0_{A_1, B_1}\rangle \langle \phi^0_{A_1, B_1}|)$$

This relation implies that b = c = 0. Thus, the relation $\operatorname{Tr} T(M_f) = d^2$ yields

$$T(M_f) = |\phi^0_{A_1,B_1}\rangle\langle\phi^0_{A_1,B_1}| \otimes |\phi^0_{A_2,B_2}\rangle\langle\phi^0_{A_2,B_2}| + \frac{1}{d^2 - 1}(I - |\phi^0_{A_1,B_1}\rangle\langle\phi^0_{A_1,B_1}|) \otimes (I - |\phi^0_{A_2,B_2}\rangle\langle\phi^0_{A_2,B_2}|),$$

which implies (11).

APPENDIX E: PROOF OF IRREDUCIBILITY

It is known that the natural representation of the subgroup $GP(d) \subset C(d)$ satisfies the condition (1). Hence,

- G. M. D'Ariano, C. Macchiavello, and M. G. A. Paris, "Local observables for entanglement witnesses," *Phys. Rev. A* 67 042310, 2003.
- [2] M. Hayashi, K. Matsumoto, and Y. Tsuda, "A study of LOCC-detection of a maximally entangled state using hypothesis testing," J. Phys. A: Math. and Gen., 39 14427-14446 (2006).
- [3] M. Hayashi, Group theoretical study of LOCC-detection of maximally entangled state using hypothesis testing, arXiv:0810.3380.
- [4] J. M. Renes, R. Blume-Kohout, A. J. Scott, and C. M. Caves, "Symmetric Informationally Complete Quantum Measurements," *J. Math. Phys.*, 45, 2171-2180 (2004); quant-ph/0310075.
- [5] A. Koldobsky and H. König, "Aspects of the Isometric Theory of Banach Spaces," in *Handbookof Geometry of Banach Spaces*, Vol. 1, edited by W. B. Johnson and J. Lindenstrauss (North-Holland, Dordrecht, 2001), pp.899-939.

it is sufficient to show the condition (2). the irreducible spaces of the subgroup $\operatorname{GP}(d) \subset \operatorname{C}(d)$ are d^2 one-dimensional subspaces generated by $|W(i,j)\rangle$ for i,j. The representation of $\operatorname{GP}(d)$ on each irreducible subspaces is different. Thus, the irreducible subspace of the larger group $\operatorname{C}(d)$ should be represented as the direct sum of these subspaces. As is shown in Lemma 1 in Appleby [8], for any (i,j) and any $F \in \operatorname{SL}(2,\mathbb{Z}_{\overline{d}})$, there exists an element $U \in \operatorname{C}(d)$ such that $f(U) \otimes \overline{f(U)}|W(i,j)\rangle = e^{\sqrt{-1}\delta_{i,j,,F}}|W(F(i,j))\rangle$, where

$$\overline{d} := \begin{cases} d & \text{if} d \text{is odd} \\ 2d & \text{if} d \text{is even.} \end{cases}$$
(E1)

For any pair $(i,j) = \neq (0,0)$, there exists an element $F \in SL(2, \mathbb{Z}_{\overline{d}})$ such that (i,j) = F(1,0). Since any irreducible subspace should be spanned by the subset of $\{|W(i,j)\rangle\}_{i,j}$, the space spanned by $\{|W(i,j)\rangle\}_{(i,j)\neq(0,0)}$ is irreducible. Thus, the condition (2) holds.

- [6] Grassl, M., "On SIC-POVMs and MUBs in dimension 6," *Proceedings of EQIS'04*, pp. 60-61, (2004); quant-ph/0406175.
- [7] Zauner, G., "Quantum designs—foundations of a noncommutative theory of designs," (in German), Ph.D. thesis, University of Vienna, (1999).
- [8] Appleby, D M, "SIC-POVMs and the Extended Clifford Group," J. Math. Phys., 46, 052107 (2005).
- [9] I. D. Ivanovic, "Geometrical description of quantum state determination," J. Phys. A: Math. and Gen., 14, No. 12, 3241-3245, (1981).
- [10] W. K. Wootters and B. D. Fields, "Optimal statedetermination by mutually unbiased measurements," *Annals of Physics*, **191**, No. 2, 363-381, (1989).
- [11] S. Bandyopadhyay, P. O. Boykin, V. Roychowdhury, and F. Vatan, "A New Proof for the Existence of Mutually Unbiased Bases," Algorithmica, 34 (2002), pp. 512-528; quant-ph/0103162.