

QUANTUM GEOMETRODYNAMICS CREATES NEW PROBLEMS

T. P. Shestakova

Department of Theoretical and Computational Physics, Southern Federal University¹,

Sorge St. 5, Rostov-on-Don 344090, Russia

E-mail: shestakova@phys.rsu.ru

Abstract

The development of last years in quantum geometrodynamics highlights new problems which were not obvious in its first formulation proposed by Wheeler and DeWitt. At the first stage the main task was to apply known quantization schemes to gravitational field or a certain cosmological model. This way has led to the realization of the fact that a quantum description of the Universe is impossible without implicit or explicit indication to a reference frame presented by some medium, filling the whole Universe, with its own equation of state and thermodynamical properties.

Thus the questions arise, should one seek for a “privileged” reference frame or consider all the variety of gauge conditions and appropriate solutions for the wave function? It is worth noting that thermodynamical properties of such a quantum Universe would also depend on a chosen reference frame to some extent. So, we need a self-consistent quantum theoretical and thermodynamical description of the Universe.

In this talk I would like to present my point of view on the problems of quantum geometrodynamics, as they can be seen now, forty years after the first significant attempt to construct full quantum theory of gravity made by DeWitt in 1967 [1]. As all of us well know, first attempts to apply quantum theory to gravitational field immediately faced enormous obstacles, and the aim of this my talk is to demonstrate the interrelation between those obstacles and the problems quantum geometrodynamics meets now.

At the first stage the main task was to apply known quantization schemes to gravitational field or a certain cosmological model. It was realized that the main difficulty consisted in the nature of general relativity as a completely covariant theory that ran counter to efforts

¹former Rostov State University

to build a Hamiltonian formulation of it as the first step on the way of its quantization. The difficulty was referred to as “the problem of constraints”. Meanwhile, in 1950s Dirac published his outline of a general Hamiltonian theory [2, 3] which was in principle applicable to any system with constraints, in particular, to gravitational field. The next important step was done by Arnowitt, Deser and Misner [4] who proposed a special parametrization of gravitational variables that made the construction of Hamiltonian formalism easier and admitted a clear interpretation. The third source of DeWitt theory was the ideas of Wheeler concerning a wave functional describing a state of gravitational field [5, 6]. Dirac approach to quantization of systems with constraints, Arnowitt – Deser – Misner (ADM) parametrization and Wheeler ideas are the three cornerstones on which the Wheeler – DeWitt quantum geometrodynamics is based.

It seemed that the initial obstacles had been overcome. However, the Wheeler – DeWitt quantum geometrodynamics encountered a number of fundamental problems which cannot be resolved in its own limits and which have made a way for its strong criticism. So, Isham [7] wrote: “...although it may be heretical to suggest it, the Wheeler - DeWitt equation – elegant though it be – may be completely the wrong way of formulating a quantum theory of gravity”.

What did make Isham to claim it? There exist serious doubts that the Dirac approach can be applied to gravitational field. The central part in the Dirac approach is given to a postulate, according to which each constraint $\varphi_m(q, p) = 0$ after quantization becomes a condition on a state vector, or wave functional, Ψ : $\varphi_m \Psi = 0$. Let us emphasize that it is indeed a postulate, since it cannot be justified by the reference to the correspondence principle. The role prescribed to the constraints could be explained by the fact that at the classical level, the constraints express gauge invariance of the theory. It was initially believed that imposing constraints at the quantum level would also ensure gauge invariance of wave functional. But what grounds do we have to expect it? Strictly speaking, the founders of quantum geometrodynamics have not investigated this issue and gauge invariance of the theory has not been proved. It leads us to the next fundamental problem: Could we consider quantum geometrodynamics as a gauge-invariant theory?

An important role was played by the ADM parametrization: it is the ADM parametrization that enables one to write gravitational constraints in the form independent of gauge variables – the lapse and shift functions N, N_i . It gave rise to an illusion that the theory in which the main equations are those of constraints must not depend on a choice of gauge conditions. At the same time the ADM parametrization introduces in 4-dimensional spacetime a set of 3-dimensional hypersurfaces (the so-called (3+1)-splitting). But fixing (3+1)-splitting

prescribes particular values for the lapse and shift functions [8, 9] that is equivalent to fix a reference frame, and gauge invariance breaks down. Thus, the Hamiltonian constraint loses its sense and, with the latter, so does the whole procedure of quantization.


The third point was the idea by Wheeler that the wave functional must be determined on the superspace of all possible 3-geometries. However, the statement that the wave function must depend only on 3-geometry is just a declaration without any mathematical realization. As we know, the state vector always depends on a concrete form of the metric.

Gauge invariance of the Wheeler – DeWitt theory can hardly be proved or refuted within canonical quantization approach. Path integration approach is more powerful, since in this case gauge invariance of the path integral, and the theory as a whole, is ensured by asymptotic boundary conditions. In first works devoted to derivation of the Wheeler – DeWitt equation from the path integral [10, 11], asymptotic boundary conditions were tacitly adopted without careful consideration if they are justified. In the works of our group in the end of 1990s attention was focused on the circumstance that the Universe is topologically non-trivial system without asymptotic states [12, 13]. It led to the conclusion that the picture of quantum evolution of the Universe cannot be independent of a reference frame in which this evolution is studied.

Our work were not the only works in this trend. I should mention the pioneer paper by Brown and Kuchař [14], the works by the group of Montani and collaborators [8, 9, 15] and others that opened the way to “Evolutionary Quantum Gravity”. In all the approaches a reference frame is presented by some medium, filling the whole Universe, with its own equation of state and thermodynamical properties. The reference frame is introduced in these approaches by different ways. In the approach by Brown and Kuchař the reference frame is related with incoherent dust, In the work by Montani and collaborators it is done by means the so-called kinematical action. In the extended phase space approach it is argued that any gauge-fixing term in Batalin – Vilkovisky (Faddeev – Popov) effective action describes a medium with mentioned above properties.

Returning to the problem of constraints, one should confess that it has not been solved in the sense that we failed to construct a gauge-invariant quantum theory of gravity for the whole Universe. Instead of a gauge-invariant theory, in a mathematically consistent approach one should reject the Wheeler – DeWitt equation as a quantum version of the Hamiltonian constraint, and reestablish the role which Schrödinger equation plays in any quantum theory and which it lost in quantum geometrodynamics. A quantum geometrodynamical Schrödinger equation appears to be gauge-dependent. Any changing a reference frame results in varying gauge conditions and, in its turn, in modifying of the Schrödinger

equation for a wave function of the Universe, as well as its solution. It can be expressed by the following scheme:



Let us try to give an assessment of the modern state of quantum geometrodynamics. The first possibility is to search for arguments in favor of some privileged reference frame in which the picture of the Universe evolution would better correspond with observational data. For example, a significant physical argument could be if quantum evolution of the Early Universe would ensure for inflation stage and further classical evolution of the observable Universe. However, today we do not have available such significant arguments and do not have any grounds to postulate a privileged reference frame. In my opinion, we should consider all possible situations and feel about for relations between classes of gauge transformation of diffeomorphism group and classes of solutions to the Schrödinger equation. This task is very laborious since the structure of diffeomorphism group is known to be very complicated. Nevertheless, one can start, as usual, from well-studied subgroups and try to find the way.

In a full description one should also take into account thermodynamical properties of a quantum Universe filled with a medium playing the role of a reference frame. Indeed, one of possible methods to build thermodynamics of the system under consideration is to write a density matrix through a path integral with Euclidean version of an action (in our case it is a gauged gravitational action), so that thermodynamical properties of the system would depend on a chosen reference frame as well. It must not be surprising for us, since the example of Rindler space teaches us that thermodynamical properties could actually change after going over to another frame. But we yet need a clear interpretation of quantum gravitational phenomena taking place under this transition.

And the question remains: Do we have to appeal to an underlying theory to solve these problems or can we hope to resolve them within the framework of quantum geometrodynamics itself? We are still far from the so-called final theory of Everything. In our attempts

to approach the final theory we rely on our experience in ordinary quantum field theory and quantum gravity, and often face the same problems. So, I do not believe that we should give up the search for better understanding the principles on which quantum geometrodynamics ought to be grounded.

References

- [1] B. S. DeWitt, *Phys. Rev.* **160** (1967), P. 1113–1148.
- [2] P. A. M. Dirac, *Can. J. Math.* **2** (1950), P. 129–148.
- [3] P. A. M. Dirac, *Proc. Roy. Soc.* **A246** (1958), P. 326–332.
- [4] R. Arnowitt, S. Deser and C. W. Misner, in: *Gravitation, an Introduction to Current Research*, ed. by L. Witten, John Wiley & Sons, New York (1963) P. 227–284.
- [5] J. A. Wheeler, in: *Relativity, Groups and Topology*, eds. C. DeWitt, B. S. DeWitt, Gordon & Breach, New York (1964).
- [6] J. A. Wheeler, *Einstein's vision*, Springer Verlag, Berlin–Heidelberg–New York (1968).
- [7] C. Isham, *Canonical quantum gravity and the problem of time*, lectures presented at NATO Advanced Study Institute, Salamanca, June 1992, Preprint gr-qc/9210011.
- [8] S. Mercuri and G. Montani, *Int. J. Mod. Phys.* **D13**, (2004), P. 165–186.
- [9] S. Mercuri and G. Montani, in: *Proceeding of the X Marcel Grossmann meeting, Rio de Janeiro, Brazil, July 2003*, Preprint gr-qc/0401127.
- [10] A. O. Barvinsky and V. N. Ponomariov, *Izvestiya Vuzov. Fizika*, No.3 (1986) P. 37.
- [11] J. J. Halliwell, *Phys. Rev.* **D38** (1988), P. 2468–2481.
- [12] V. A. Savchenko, T. P. Shestakova and G. M. Vereshkov, *Gravitation & Cosmology* **7** (2001), P. 18–28.
- [13] V. A. Savchenko, T. P. Shestakova and G. M. Vereshkov, *Gravitation & Cosmology* **7** (2001), P. 102–116.
- [14] J. D. Brown and K. V. Kuchař, *Phys. Rev.* **D51** (1995), P. 5600–5629.
- [15] M. V. Battisti and G. Montani, *Phys. Lett.* **B637** (2006) P. 203–209.