

CLUSTER EXPANSION FORMULAS AND PERFECT MATCHINGS

GREGG MUSIKER AND RALF SCHIFFLER

ABSTRACT. We study cluster algebras with principal coefficient systems that are associated to unpunctured surfaces. We give a direct formula for the Laurent polynomial expansion of cluster variables in these cluster algebras in terms of perfect matchings of a certain graph $G_{T,\gamma}$ that is constructed from the surface by recursive glueing of elementary pieces that we call tiles. We also give a second formula for these Laurent polynomial expansions in terms of subgraphs of the graph $G_{T,\gamma}$.

1. INTRODUCTION

Cluster algebras, introduced in [FZ1], are commutative algebras equipped with a distinguished set of generators, the *cluster variables*. The cluster variables are grouped into sets of constant cardinality n , the *clusters*, and the integer n is called the *rank* of the cluster algebra. Starting with an initial cluster \mathbf{x} (together with a skew symmetrizable integer $n \times n$ matrix $B = (b_{ij})$ and a coefficient vector $\mathbf{y} = (y_i)$ whose entries are elements of a torsion-free abelian group \mathbb{P}) the set of cluster variables is obtained by repeated application of so called *mutations*. To be more precise, let \mathcal{F} be the field of rational functions in the indeterminates x_1, x_2, \dots, x_n over the quotient field of the integer group ring $\mathbb{Z}\mathbb{P}$. Thus $\mathbf{x} = \{x_1, x_2, \dots, x_n\}$ is a transcendence basis for \mathcal{F} . For every $k = 1, 2, \dots, n$, the mutation $\mu_k(\mathbf{x})$ of the cluster $\mathbf{x} = \{x_1, x_2, \dots, x_n\}$ is a new cluster $\mu_k(\mathbf{x}) = \mathbf{x} \setminus \{x_k\} \cup \{x'_k\}$ obtained from \mathbf{x} by replacing the cluster variable x_k by the new cluster variable

$$(1) \quad x'_k = \frac{1}{x_k} \left(y_k^+ \prod_{b_{ki} > 0} x_i^{b_{ki}} + y_k^- \prod_{b_{ki} < 0} x_i^{-b_{ki}} \right)$$

in \mathcal{F} , where y_k^+, y_k^- are certain monomials in y_1, y_2, \dots, y_n . Mutations also change the attached matrix B as well as the coefficient vector \mathbf{y} , see [FZ1].

The set of all cluster variables is the union of all clusters obtained from an initial cluster \mathbf{x} by repeated mutations. Note that this set may be infinite.

It is clear from the construction that every cluster variable is a rational function in the initial cluster variables x_1, x_2, \dots, x_n . In [FZ1] it is shown that every cluster variable u is actually a Laurent polynomial in the x_i , that is, u can be written as a reduced fraction

$$(2) \quad u = \frac{f(x_1, x_2, \dots, x_n)}{\prod_{i=1}^n x_i^{d_i}},$$

where $f \in \mathbb{Z}\mathbb{P}[x_1, x_2, \dots, x_n]$ and $d_i \geq 0$. The right hand side of equation (2) is called the *cluster expansion* of u in \mathbf{x} .

Date: October 20, 2008.

The cluster algebra is determined by the initial matrix B and the choice of the coefficient system. A canonical choice of coefficients is the *principal* coefficient system, introduced in [FZ2], which means that the coefficient group \mathbb{P} is the free abelian group on n generators y_1, y_2, \dots, y_n , and the initial coefficient tuple $\mathbf{y} = \{y_1, y_2, \dots, y_n\}$ consists of these n generators. In [FZ2], the authors show that knowing the expansion formulas in the case where the cluster algebra has principal coefficients allows one to compute the expansion formulas for arbitrary coefficient systems.

Inspired by the work of Fock and Goncharov [FG1, FG2, FG3] and Gekhtman, Shapiro and Vainshtein [GSV1, GSV2] which discovered cluster structures in the context of Teichmüller theory, Fomin, Shapiro and Thurston [FST, FT] initiated a systematic study of the cluster algebras arising from triangulations of a surface with boundary and marked points. In this approach, cluster variables in the cluster algebra correspond to arcs in the surface, and clusters correspond to triangulations. In [S2], building on earlier results in [S1, ST], this model was used to give a direct expansion formula for cluster variables in cluster algebras associated to unpunctured surfaces, with arbitrary coefficients, in terms of certain paths on the triangulation.

Our first main result in this paper is a new parametrization of this formula in terms of perfect matchings of a certain weighted graph that is constructed from the surface by recursive glueing of elementary pieces that we call tiles. To be more precise, let x_γ be a cluster variable corresponding to an arc γ in the unpunctured surface and let d be the number of crossings between γ and the triangulation T of the surface. Then γ runs through $d + 1$ triangles of T and each pair of consecutive triangles forms a quadrilateral which we call a tile. So we obtain d tiles, each of which is a weighted graph, whose weights are given by the cluster variables x_τ associated to the arcs τ of the triangulation T .

We obtain a weighted graph $G_{T,\gamma}$ by glueing the d tiles in a specific way and then deleting the diagonal in each tile. To any perfect matching M of this graph we associate its weight $w(M)$ which is the product of the weights of its edges, hence a product of cluster variables. We prove the following cluster expansion formula:

Theorem 3.1.

$$x_\gamma = \sum_M \frac{w(M) y(M)}{x_{i_1} x_{i_2} \dots x_{i_d}},$$

where the sum is over all perfect matchings M of $G_{T,\gamma}$, $w(M)$ is the weight of M , and $y(M)$ is a monomial in \mathbf{y} .

We also give a formula for the coefficients $y(M)$ in terms of perfect matchings as follows. The F -polynomial F_γ , introduced in [FZ2] is obtained from the Laurent polynomial x_γ (with principal coefficients) by substituting 1 for each of the cluster variables x_1, x_2, \dots, x_n . By [S2, Theorem 6.2, Corollary 6.4], the F -polynomial has constant term 1 and a unique term of maximal degree that is divisible by all the other occurring monomials. The two corresponding matchings are the unique two matchings that have all their edges on the boundary of the graph $G_{T,\gamma}$. We denote by M_- the one with $y(M_-) = 1$ and the other by M_+ . Now, for an arbitrary perfect matching M , the coefficient $y(M)$ is determined by the set of edges of the symmetric difference $M_- \ominus M = (M_- \cup M) \setminus (M_- \cap M)$ as follows.

Theorem 5.1. *The set $M_- \ominus M$ is the set of boundary edges of a (possibly disconnected) subgraph G_M of $G_{T,\gamma}$ which is a union of tiles $G_M = \cup_{j \in J} S_j$. Moreover,*

$$y(M) = \prod_{j \in J} y_{i_j}$$

As an immediate corollary, we see that the corresponding g -vector, introduced in [FZ2], is

$$g_\gamma = \deg \left(\frac{w(M_-)}{x_{i_1} \cdots x_{i_d}} \right).$$

Our third main result is yet another description of the formula of Theorem 3.1 in terms of the graph $G_{T,\gamma}$ only. In order to state this result, we need some notation. If H is a graph, let $c(H)$ be the number of connected components of H , let $E(H)$ be the set of edges of H , and denote by ∂H the set of boundary edges of H . Define \mathcal{H}_k to be the set of all subgraphs H of $G_{T,\gamma}$ such that H is a union of k tiles $H = S_{j_1} \cup \cdots \cup S_{j_k}$ and such that the number of edges of M_- that are contained in H is equal to $k + c(H)$. For $H \in \mathcal{H}_k$, let

$$y(H) = \prod_{S_{i_j} \text{ tile in } H} y_{i_j}.$$

Theorem 6.1. *The cluster expansion of the cluster variable x_γ is given by*

$$x_\gamma = \sum_{k=0}^d \sum_{H \in \mathcal{H}_k} \frac{w(\partial H \ominus M_-) y(H)}{x_{i_1} x_{i_2} \cdots x_{i_d}}.$$

Theorem 3.1 has interesting intersections with work of other people. In [CCS2], the authors obtained a formula for the denominators of the cluster expansion in types A, D and E , see also [BMR]. In [CC, CK, CK2] an expansion formula was given in the case where the cluster algebra is acyclic and the cluster lies in an acyclic seed. Palu generalized this formula to arbitrary clusters in an acyclic cluster algebra [Pa]. These formulas use the cluster category introduced in [BMRRT], and in [CCS] for type A , and do not give information about the coefficients.

Recently, Fu and Keller generalized this formula further to cluster algebras with principal coefficients that admit a categorification by a 2-Calabi-Yau category [FK], and, combining results of [A] and [ABCP, LF], such a categorification exists in the case of cluster algebras associated to unpunctured surfaces.

In [SZ, CZ, Z, MP] cluster expansions for cluster algebras of rank 2 are given, in [Pr1, CP, FZ3] the case A is considered. In section 4 of [Pr1], Propp describes two constructions of snake graphs, the latter of which are unweighted analogues for the case A of the graphs $G_{T,\gamma}$ that we present in this paper. Propp assigns a snake graph to each arc in the triangulation of an n -gon and shows that the numbers of matchings in these graphs satisfy the Conway-Coxeter frieze pattern induced by the Ptolemy relations on the n -gon. In [M] a cluster expansion for cluster algebras of classical type is given for clusters that lie in a bipartite seed, and the forthcoming work of [MW] will concern cluster expansions for cluster algebras of classical type with principal coefficients, for an arbitrary seed.

The formula for $y(M)$ given in Theorem 5.1 also can be formulated in terms of height functions, as found in literature such as [EKLP] or [Pr2]. We discuss this connection in Remark 5.3 of section 5.

The paper is organized as follows. In section 2, we recall the construction of cluster algebras from surfaces of [FST]. Section 3 contains the construction of the graph $G_{T,\gamma}$ and the statement of the cluster expansion formula. Section 4 is devoted to the proof of the expansion formula. The formula for $y(M)$ and the formula for the g -vectors is given in section 5. In section 6, we present the expansion formula in terms of subgraphs and deduce a formula for the F -polynomials. We give an example in section 7.

Acknowledgements. The authors would like to thank Jim Propp for useful conversations related to this work.

2. CLUSTER ALGEBRAS FROM SURFACES

In this section, we recall the construction of [FST] in the case of surfaces without punctures.

Let S be a connected oriented 2-dimensional Riemann surface with boundary and M a non-empty set of marked points in the closure of S with at least one marked point on each boundary component. The pair (S, M) is called *bordered surface with marked points*. Marked points in the interior of S are called *punctures*.

In this paper we will only consider surfaces (S, M) such that all marked points lie on the boundary of S , and we will refer to (S, M) simply by *unpunctured surface*.

We say that two curves in S *do not cross* if they do not intersect each other except that endpoints may coincide.

Definition 1. An arc γ in (S, M) is a curve in S such that

- (a) the endpoints are in M ,
- (b) γ does not cross itself,
- (c) the relative interior of γ is disjoint from M and from the boundary of S ,
- (d) γ does not cut out a monogon or a digon.

Curves that connect two marked points and lie entirely on the boundary of S without passing through a third marked point are called *boundary arcs*. Hence an arc is a curve between two marked points, which does not intersect itself nor the boundary except possibly at its endpoints and which is not homotopic to a point or a boundary arc.

Each arc is considered up to isotopy inside the class of such curves. Moreover, each arc is considered up to orientation, so if an arc has endpoints $a, b \in M$ then it can be represented by a curve that runs from a to b , as well as by a curve that runs from b to a .

For any two arcs γ, γ' in S , let $e(\gamma, \gamma')$ be the minimal number of crossings of γ and γ' , that is, $e(\gamma, \gamma')$ is the minimum of the numbers of crossings of arcs α and α' , where α is isotopic to γ and α' is isotopic to γ' . Two arcs γ, γ' are called *compatible* if $e(\gamma, \gamma') = 0$. A *triangulation* is a maximal collection of compatible arcs together with all boundary arcs. The arcs of a triangulation cut the surface into *triangles*. Since (S, M) is an unpunctured surface, the three sides of each triangle are distinct (in contrast to the case of surfaces with punctures). Any triangulation has $n + m$ elements, n of which are arcs in S , and the remaining m elements are boundary arcs. Note that the number of boundary arcs is equal to the number of marked points.

b	g	m	surface
1	0	n+3	polygon
1	1	n-3	torus with disk removed
1	2	n-9	genus 2 surface with disk removed
2	0	n	annulus
2	1	n-6	torus with 2 disks removed
2	2	n-12	genus 2 surface with 2 disks removed
3	0	n-3	pair of pants

TABLE 1. Examples of unpunctured surfaces

Proposition 2.1. *The number n of arcs in any triangulation is given by the formula $n = 6g + 3b + m - 6$, where g is the genus of S , b is the number of boundary components and $m = |M|$ is the number of marked points. The number n is called the rank of (S, M) .*

Proof. [FST, 2.10] □

Note that $b > 0$ since the set M is not empty. Table 1 gives some examples of unpunctured surfaces.

Following [FST], we associate a cluster algebra to the unpunctured surface (S, M) as follows. Choose any triangulation T , let $\tau_1, \tau_2, \dots, \tau_n$ be the n interior arcs of T and denote the m boundary arcs of the surface by $\tau_{n+1}, \tau_{n+2}, \dots, \tau_{n+m}$. For any triangle Δ in T define a matrix $B^\Delta = (b_{ij}^\Delta)_{1 \leq i \leq n, 1 \leq j \leq n}$ by



$$b_{ij}^\Delta = \begin{cases} 1 & \text{if } \tau_i \text{ and } \tau_j \text{ are sides of } \Delta \text{ with } \tau_j \text{ following } \tau_i \text{ in the} \\ & \text{counter-clockwise order;} \\ -1 & \text{if } \tau_i \text{ and } \tau_j \text{ are sides of } \Delta \text{ with } \tau_j \text{ following } \tau_i \text{ in the} \\ & \text{clockwise order;} \\ 0 & \text{otherwise.} \end{cases}$$

Then define the matrix $B_T = (b_{ij})_{1 \leq i \leq n, 1 \leq j \leq n}$ by $b_{ij} = \sum_{\Delta} b_{ij}^\Delta$, where the sum is taken over all triangles in T . Note that the boundary arcs of the triangulation are ignored in the definition of B_T . Let $\tilde{B}_T = (b_{ij})_{1 \leq i \leq 2n, 1 \leq j \leq 2n}$ be the $2n \times 2n$ matrix whose upper $n \times n$ part is B_T and whose lower $n \times n$ part is the identity matrix. The matrix B_T is skew-symmetric and each of its entries b_{ij} is either 0, 1, -1, 2, or -2, since every arc τ can be in at most two triangles. An example where $b_{ij} = 2$ is given in Figure 1.

Let $\mathcal{A}(\mathbf{x}_T, \mathbf{y}_T, B_T)$ be the cluster algebra with principal coefficients in the triangulation T , that is, $\mathcal{A}(\mathbf{x}_T, \mathbf{y}_T, B_T)$ is given by the seed $(\mathbf{x}_T, \mathbf{y}_T, B_T)$ where $\mathbf{x}_T = \{x_{\tau_1}, x_{\tau_2}, \dots, x_{\tau_n}\}$ is the cluster associated to the triangulation T , and the initial coefficient vector $\mathbf{y}_T = (y_1, y_2, \dots, y_n)$ is the vector of generators of $\mathbb{P} = \text{Trop}(y_1, y_2, \dots, y_n)$.

For the boundary arcs we define $x_{\tau_k} = 1$, $k = n+1, n+2, \dots, n+m$.

For each $k = 1, 2, \dots, n$, there is a unique quadrilateral in $T \setminus \{\tau_k\}$ in which τ_k is one of the diagonals. Let τ'_k denote the other diagonal in that quadrilateral. Define the *flip* $\mu_k T$ to be the triangulation $T \setminus \{\tau_k\} \cup \{\tau'_k\}$. The mutation μ_k of the seed Σ_T in the cluster algebra \mathcal{A} corresponds to the flip μ_k of the triangulation T in the following sense. The matrix $\mu_k(B_T)$ is the matrix corresponding to the triangulation $\mu_k T$, the cluster $\mu_k(\mathbf{x}_T)$ is $\mathbf{x}_T \setminus \{x_{\tau_k}\} \cup \{x_{\tau'_k}\}$, and the corresponding

FIGURE 1. A triangulation with $b_{23} = 2$ FIGURE 2. The tile \overline{S}_k

exchange relation is given by

$$x_{\tau_k} x_{\tau'_k} = x_{\rho_1} x_{\rho_2} y^+ + x_{\sigma_1} x_{\sigma_2} y^-,$$


where y^+, y^- are some coefficients, and $\rho_1, \sigma_1, \rho_2, \sigma_2$ are the sides of the quadrilateral in which τ_k and τ'_k are the diagonals, such that ρ_1, ρ_2 are opposite sides and σ_1, σ_2 are opposite sides too.

3. EXPANSION FORMULA

In this section, we will present an expansion formula for the cluster variables in terms of perfect matchings of a graph that is constructed recursively using so-called *tiles*.

3.1. Tiles. For the purpose of this paper, a *tile* \overline{S}_k is a planar four vertex graph with five weighted edges having the shape of two equilateral triangles that share one edge, see Figure 2. The weight on each edge of the tile \overline{S}_k is a single variable. The unique interior edge is called *diagonal* and the four exterior edges are called *sides* of \overline{S}_k . We shall use S_k to denote the graph obtained from \overline{S}_k by removing the diagonal.

Now let T be a triangulation of the unpunctured surface (S, M) . If $\tau_k \in T$ is an interior arc, then τ_k lies in precisely two triangles in T , hence τ_k is the diagonal of a unique quadrilateral Q_{τ_k} in T . We associate to this quadrilateral a tile \overline{S}_k by assigning the weight x_k to the diagonal and the weights x_a, x_b, x_c, x_d to the sides of

FIGURE 3. Glueing tiles S_k and S_{k+1} along the edge weighted $x_{[\gamma_k]}$

\bar{S}_k in such a way that there is a homeomorphism $\bar{S}_k \rightarrow Q_{\tau_k}$ which sends the edge with weight x_i to the arc labeled τ_i , $i = a, b, c, d, k$, see Figure 2.

3.2. The graph $\bar{G}_{T,\gamma}$. Let T be a triangulation of an unpunctured surface (S, M) and let γ be an arc in (S, M) which is not in T . Choose an orientation on γ and let $s \in M$ be its starting point, and let $t \in M$ be its endpoint. We denote by

$$p_0 = s, p_1, p_2, \dots, p_{d+1} = t$$

the points of intersection of γ and T in order. Let i_1, i_2, \dots, i_d be such that p_k lies on the arc $\tau_{i_k} \in T$. Note that i_k may be equal to i_j even if $k \neq j$. Let $\tilde{S}_1, \tilde{S}_2, \dots, \tilde{S}_d$ be a sequence of tiles so that \tilde{S}_k is isomorphic to the tile \bar{S}_{i_k} , for $k = 1, 2, \dots, d$.

For k from 0 to d , let γ_k denote the segment of the path γ from the point p_k to the point p_{k+1} . Each γ_k lies in exactly one triangle Δ_k in T , and if $1 \leq k \leq d-1$ then Δ_k is formed by the arcs $\tau_{i_k}, \tau_{i_{k+1}}$, and a third arc that we denote by $\tau_{[\gamma_k]}$.

We will define a graph $\bar{G}_{T,\gamma}$ by recursive glueing of tiles. Start with $\bar{G}_{T,\gamma,1} \cong \tilde{S}_1$, where we orient the tile \tilde{S}_1 so that the diagonal goes from northwest to southeast, and the starting point p_0 of γ is in the southwest corner of \tilde{S}_1 . For all $k = 1, 2, \dots, d-1$ let $\bar{G}_{T,\gamma,k+1}$ be the graph obtained by adjoining the tile \tilde{S}_{k+1} to the tile \tilde{S}_k of the graph $\bar{G}_{T,\gamma,k}$ along the edge weighted $x_{[\gamma_k]}$, see Figure 3. We always orient the tiles so that the diagonals go from northwest to southeast. Note that the edge weighted $x_{[\gamma_k]}$ is either the northern or the eastern edge of the tile \tilde{S}_k . Finally, we define $\bar{G}_{T,\gamma}$ to be $\bar{G}_{T,\gamma,d}$.

Let $G_{T,\gamma}$ be the graph obtained from $\bar{G}_{T,\gamma}$ by removing the diagonal in each tile, that is, $G_{T,\gamma}$ is constructed in the same way as $\bar{G}_{T,\gamma}$ but using tiles S_{i_k} instead of \bar{S}_{i_k} .

A *perfect matching* of a graph is a subset of the edges so that each vertex is covered exactly once. We define the weight $w(M)$ of a perfect matching M to be the product of the weights of all edges in M .

3.3. Cluster expansion formula. Let (S, M) be an unpunctured surface with triangulation T , and let $\mathcal{A} = \mathcal{A}(\mathbf{x}_T, \mathbf{y}_T, B)$ be the cluster algebra with principal coefficients in the initial seed $(\mathbf{x}_T, \mathbf{y}_T, B)$ defined in section 2. Each cluster variable in \mathcal{A} corresponds to an arc in (S, M) . Let x_γ be an arbitrary cluster variable corresponding to an arc γ . Choose an orientation of γ , and let $\tau_{i_1}, \tau_{i_2}, \dots, \tau_{i_d}$ be the arcs of the triangulation that are crossed by γ in this order, with multiplicities possible. Let $G_{T,\gamma}$ be the graph constructed in section 3.2.

Theorem 3.1. *With the above notation*

$$x_\gamma = \sum_M \frac{w(M) y(M)}{x_{i_1} x_{i_2} \cdots x_{i_d}},$$

where the sum is over all perfect matchings M of $G_{T,\gamma}$, $w(M)$ is the weight of M , and $y(M)$ is a monomial in \mathbf{y}_T .

The proof of Theorem 3.1 will be given in Section 4.

4. PROOF OF THEOREM 3.1

Throughout this section, T is a triangulation of an unpunctured surface (S, M) , γ is an arc in S with a fixed orientation, and $s \in M$ is its starting point and $t \in M$ is its endpoint. Moreover, $p_0 = s, p_1, p_2, \dots, p_{d+1} = t$ are the points of intersection of γ and T in order, and i_1, i_2, \dots, i_d are such that p_k lies on the arc $\tau_{i_k} \in T$.

4.1. Complete (T, γ) -paths. Following [ST], we will consider paths α in S that are concatenations of arcs in the triangulation T , more precisely, $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_{\ell(\alpha)})$ with $\alpha_i \in T$, for $i = 1, 2, \dots, \ell(\alpha)$ and the starting point of α_i is the endpoint of α_{i-1} . Such a path is called a T -path.

We call a T -path $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_{\ell(\alpha)})$ a *complete (T, γ) -path* if the following axioms hold:

- (T1) The even arcs are precisely the arcs crossed by γ in order, that is, $\alpha_{2k} = \tau_{i_k}$.
- (T2) For all $k = 0, 1, 2, \dots, d$, the segment γ_k is homotopic to the segment of the path α starting at the point p_k following $\alpha_{2k}, \alpha_{2k+1}$ and α_{2k+2} until the point p_{k+1} .

We define the Laurent monomial $x(\alpha)$ of the complete (T, γ) -path α by

$$x(\alpha) = \prod_{i \text{ odd}} x_{\alpha_i} \prod_{i \text{ even}} x_{\alpha_i}^{-1}.$$

Remark 4.1. • *Every complete (T, γ) -path starts and ends at the same point as γ , because of (T2).*

- *Every complete (T, γ) -path has length $2d + 1$.*
- *For all arcs τ in the triangulation T , the number of times that τ occurs as α_{2k} is exactly the number of crossings between γ and τ .*
- *In contrast to the ordinary (T, γ) -paths defined in [ST], complete (T, γ) -paths allow backtracking.*
- *The denominator of the Laurent monomial $x(\alpha)$ is equal to $x_{i_1} x_{i_2} \cdots x_{i_d}$.*

4.2. Universal cover. Let $\pi : \tilde{S} \rightarrow S$ be a universal cover of the surface S , and let $\tilde{M} = \pi^{-1}(M)$ and $\tilde{T} = \pi^{-1}(T)$.

Choose $\tilde{s} \in \pi^{-1}(s)$. There exists a unique lift $\tilde{\gamma}$ of γ starting at \tilde{s} . Then $\tilde{\gamma}$ is the concatenation of subpaths $\tilde{\gamma}_0, \tilde{\gamma}_1, \dots, \tilde{\gamma}_{d+1}$ where $\tilde{\gamma}_k$ is a path from a point \tilde{p}_k to a point \tilde{p}_{k+1} such that $\tilde{\gamma}_k$ is a lift of γ_k and $\tilde{p}_k \in \pi^{-1}(p_k)$, for $k = 0, 1, \dots, d+1$. Let $\tilde{t} = \tilde{p}_{d+1} \in \pi^{-1}(t)$.

For k from 1 to d , let $\tilde{\tau}_{i_k}$ be the unique lift of τ_{i_k} running through \tilde{p}_k and let $\tilde{\tau}_{[\gamma_k]}$ be the unique lift of $\tau_{[\gamma_k]}$ that is bounding a triangle in \tilde{S} with $\tilde{\tau}_{i_k}$ and $\tilde{\tau}_{i_{k+1}}$. Each $\tilde{\gamma}_k$ lies in exactly one triangle $\tilde{\Delta}_k$ in \tilde{T} . Let $\tilde{S}(\gamma) \subset \tilde{S}$ be the union of the triangles $\tilde{\Delta}_0, \tilde{\Delta}_1, \dots, \tilde{\Delta}_{d+1}$ and let $\tilde{M}(\gamma) = \tilde{M} \cap \tilde{S}(\gamma)$ and $\tilde{T}(\gamma) = \tilde{T} \cap \tilde{S}(\gamma)$. Then $(\tilde{S}(\gamma), \tilde{M}(\gamma))$ is a simply connected unpunctured surface of which $\tilde{T}(\gamma)$ is a

triangulation. This triangulation $\tilde{T}(\gamma)$ consists of arcs $\tilde{\tau}_{i_k}, \tilde{\tau}_{[\gamma_k]}$ with $k = 1, 2, \dots, d$, and two arcs incident to \tilde{s} and two arcs incident to \tilde{t} .

The underlying graph of $\tilde{T}(\gamma)$ is the graph with vertex set $\tilde{M}(\gamma)$ and whose set of edges consists of the (unoriented) arcs in $\tilde{T}(\gamma)$.

By [S2, Section 5.5], we can compute the Laurent expansion of x_γ using complete $(\tilde{T}(\gamma), \tilde{\gamma})$ -paths in $(\tilde{S}(\gamma), \tilde{M}(\gamma))$.

4.3. Folding. The graph $\overline{G}_{T,\gamma}$ was constructed by glueing tiles \tilde{S}_{k+1} to tiles \tilde{S}_k along edges with weight $x_{[\gamma_k]}$, see figure 3. Now we will fold the graph along the edges weighted $x_{[\gamma_k]}$, thereby identifying the two triangles incident to $x_{[\gamma_k]}$, $k = 1, 2, \dots, d-1$.

To be more precise, the edge with weight $x_{[\gamma_k]}$, that lies in the two tiles \tilde{S}_{k+1} and \tilde{S}_k , is contained in precisely two triangles Δ_k and Δ'_k in $\overline{G}_{T,\gamma}$: Δ_k lying inside the tile \tilde{S}_k and Δ'_k lying inside the tile \tilde{S}_{k+1} . Both Δ_k and Δ'_k have weights $x_{[\gamma_k]}$, x_k, x_{k+1} , but opposite orientations. Cutting $\overline{G}_{T,\gamma}$ along the edge with weight $x_{[\gamma_k]}$, one obtains two connected components. Let R_k be the component that contains the tile \tilde{S}_k and R_{k+1} the component that contains \tilde{S}_{k+1} .

The *folding of the graph $\overline{G}_{T,\gamma}$ along $x_{[\gamma_k]}$* is the graph obtained by flipping R_{k+1} and then glueing it to R_k by identifying the two triangles Δ_k and Δ'_k .

The graph obtained by consecutive folding of $\overline{G}_{T,\gamma}$ along all edges with weight $x_{[\gamma_k]}$ for $k = 1, 2, \dots, d-1$, is isomorphic to the underlying graph of the triangulation $\tilde{T}(\gamma)$ of the unpunctured surface $(\tilde{S}(\gamma), \tilde{M}(\gamma))$. Indeed, there clearly is a bijection between the triangles in both graphs, and, in both graphs the way the triangles are glued together is uniquely determined by γ . Note that the two graphs may have opposite orientations.

We obtain a map that we call the *folding map*

$$\begin{array}{ccc} \phi & : & \left\{ \begin{array}{l} \text{perfect matchings} \\ \text{in } G_{T,\gamma} \end{array} \right\} & \rightarrow & \left\{ \begin{array}{l} \text{complete } (\tilde{T}(\gamma), \tilde{\gamma})\text{-paths} \\ \text{in } (\tilde{S}(\gamma), \tilde{M}(\gamma)) \end{array} \right\} \\ & & M & \mapsto & \tilde{\alpha}_M \end{array}$$

as follows. First we associate a path α_M in $\overline{G}_{T,\gamma}$ to the matching M , by inserting a diagonal between any two consecutive edges of the perfect matching. More precisely, α_M is the path starting at s going along the unique edge of M that is incident to s , then going along the diagonal of the first tile \tilde{S}_1 , then along the unique edge of M that is incident to the endpoint of that diagonal, and so forth.

Since M has cardinality $d+1$, the path α_M consists of $2d+1$ edges, thus $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_{2d+1})$. Now we define $\tilde{\alpha}_M = (\tilde{\alpha}_1, \tilde{\alpha}_2, \dots, \tilde{\alpha}_{2d+1})$ by folding the path α_M . Thus, if $M = \{\beta_1, \beta_3, \dots, \beta_{2d-1}, \beta_{2d+1}\}$, where the edges are ordered according to γ , then $\phi(M) = (\tilde{\alpha}_1, \tilde{\alpha}_2, \dots, \tilde{\alpha}_{2d+1})$, where $\tilde{\alpha}_{2k+1}$ is the image of β_{2k+1} under the folding and $\tilde{\alpha}_{2k} = \tau_{i_k}$ is the arc crossing γ at p_k . Then $\phi(M)$ satisfies the axiom (T1) by construction. Moreover, $\phi(M)$ satisfies the axiom (T2), because, for each $k = 0, 1, \dots, d$, the segment of the path $\phi(M)$, which starts at the point p_k following $\tilde{\alpha}_{2k}, \tilde{\alpha}_{2k+1}$ and $\tilde{\alpha}_{2k+2}$ until the point p_{k+1} , is homotopic to the segment γ_k , since both segments lie in the simply connected triangle Δ_k formed by $\tau_{i_k}, \tau_{i_{k+1}}$ and $\tau_{[\gamma_k]}$. Therefore, the folding map ϕ is well defined.

Note that it is possible that $\tilde{\alpha}_k, \tilde{\alpha}_{k+1}$ is backtracking, that is, $\tilde{\alpha}_k$ and $\tilde{\alpha}_{k+1}$ run along the same arc $\tilde{\tau} \in \tilde{T}(\gamma)$.




FIGURE 4. Completion of paths

4.4. Unfolding the surface. Let α be a boundary arc in $(\tilde{S}(\gamma), \tilde{M}(\gamma))$ that is not adjacent to \tilde{s} and not adjacent to \tilde{t} . Then there is a unique triangle Δ in $\tilde{T}(\gamma)$ in which α is a side. The other two sides of Δ are two consecutive diagonals, which we denote by $\tilde{\tau}_j$ and $\tilde{\tau}_{j+1}$, see Figure 4.

By cutting the underlying graph of $\tilde{T}(\gamma)$ along $\tilde{\tau}_j$, we obtain two pieces. Let R_{j+1} denote the piece that contains α , $\tilde{\tau}_{j+1}$ and t . Similarly, cutting $(\tilde{S}(\gamma), \tilde{M}(\gamma))$ along $\tilde{\tau}_{j+1}$, we obtain two pieces, and we denote by R_j the piece that contains s , $\tilde{\tau}_j$ and α .

The graph obtained by unfolding along α is the graph obtained by flipping R_j and then gluing it to R_{j+1} along α . In this new graph, we label the edge of R_j that had the label $\tilde{\tau}_{j+1}$ by $\tilde{\tau}_{j+1}^b$ and the edge of R_{j+1} that had the label $\tilde{\tau}_j$ by $\tilde{\tau}_j^b$, indicating that these edges are on the boundary of the new graph, see Figure 4.

Lemma 4.2. *The graph obtained by repeated unfolding of the underlying graph of $\tilde{T}(\gamma)$ along all boundary edges not adjacent to s or t is isomorphic to the graph $\overline{G}_{T,\gamma}$. Moreover, for each unfolding along an edge α , the edges labeled $\tilde{\tau}_j^b, \tilde{\tau}_{j+1}^b$ are on the boundary of $\overline{G}_{T,\gamma}$ and carry the weights x_j, x_{j+1} respectively, the edges labeled $\tilde{\tau}_j, \tilde{\tau}_{j+1}$ are diagonals in $\overline{G}_{T,\gamma}$ and carry the weights x_j, x_{j+1} respectively, and α is an interior edge of \overline{G}_γ that is not a diagonal and carries the weight $x_{[\gamma_j]}$.*

Proof. This follows from the construction. \square


4.5. Unfolding map. We define a map

$$\begin{aligned} \{\text{complete } (\tilde{T}(\gamma), \tilde{\gamma}) \text{ - paths}\} &\rightarrow \{\text{perfect matchings of } G_{T,\gamma}\} \\ \tilde{\alpha} = (\tilde{\alpha}_1, \tilde{\alpha}_2, \dots, \tilde{\alpha}_{2d+1}) &\mapsto M_{\tilde{\alpha}} = \{\beta_1, \beta_3, \beta_5, \dots, \beta_{2d+1}\} \end{aligned}$$

where $\beta_1 = \tilde{\alpha}_1, \beta_{2d+1} = \tilde{\alpha}_{2d+1}$ and

$$\beta_{2k+1} = \begin{cases} \tilde{\alpha}_{2k+1} & \text{if } \tilde{\alpha}_{2k+1} \text{ is a boundary arc in } \tilde{T}(\gamma), \\ \tilde{\tau}_j^b & \text{if } \tilde{\alpha}_{2k+1} = \tilde{\tau}_j \text{ is a diagonal in } \tilde{T}(\gamma). \end{cases}$$

We will show that this map is well-defined. Suppose β_{2k+1} and $\beta_{2\ell+1}$ have a common endpoint x . Then α_{2k+1} and $\alpha_{2\ell+1}$ have a common endpoint y in $(\tilde{S}(\gamma), \tilde{M}(\gamma))$ and the two edges are not separated in the unfolding described in Lemma 4.2. Consequently, there is no triangle in $\tilde{T}(\gamma)$ that is contained in the subpolygon spanned by α_{2k+1} and $\alpha_{2\ell+1}$, hence α_{2k+1} is equal to $\alpha_{2\ell+1}$. This implies that every arc in the subpath $(\alpha_{2k+1}, \alpha_{2k+2} \dots \alpha_{2\ell+1})$ is equal to the same diagonal $\tilde{\tau}_j$, and the only way this can happen is when $\ell = k + 1$ and $(\alpha_{2k+1}, \alpha_{2k+2} \dots \alpha_{2\ell+1}) = (\tilde{\tau}_j, \tilde{\tau}_j, \tilde{\tau}_j)$ and both endpoints of $\tilde{\tau}_j$ are incident to an interior arc other than $\tilde{\tau}_j$. In this case, $\tilde{\tau}_j$ bounds the two triangles $\tilde{\tau}_{j-1}, \tilde{\tau}_j, \tilde{\tau}_{[\gamma_{j-1}]}$ and $\tilde{\tau}_j, \tilde{\tau}_{j+1}, \tilde{\tau}_{[\gamma_j]}$ in $\tilde{T}(\gamma)$. Unfolding along $\tilde{\tau}_{[\gamma_{j-1}]}$ and $\tilde{\tau}_{[\gamma_j]}$ will produce edges β_{2k+1} and $\beta_{2\ell+1}$ that are not adjacent, see Figure 5. This shows that no vertex of $G_{T,\gamma}$ is covered twice in $M_{\tilde{\alpha}}$.

FIGURE 5. Unfolding along $\tilde{\tau}_{[\gamma_{j-1}]}$ and $\tilde{\tau}_{[\gamma_j]}$

To show that every vertex of $G_{T,\gamma}$ is covered in M_α , we use a counting argument. Indeed, the number of vertices of $G_{T,\gamma}$ is $2(d+1)$, and, on the other hand, $2d+1$ is the length of α , since α is complete, and thus $M_{\tilde{\alpha}}$ has $d+1$ edges. The statement follows since every $\beta_j \in M_{\tilde{\alpha}}$ has two distinct endpoints. This shows that $M_{\tilde{\alpha}}$ is a perfect matching and our map is well-defined.

Lemma 4.3. *The unfolding map $\tilde{\alpha} \mapsto M_{\tilde{\alpha}}$ is the inverse of the folding map $M \mapsto \tilde{\alpha}_M$. In particular, both maps are bijections.*

Proof. Let $\tilde{\alpha} = (\tilde{\alpha}_1, \tilde{\alpha}_2, \dots, \tilde{\alpha}_{2d+1})$ be a complete $(\tilde{T}(\gamma), \tilde{\gamma})$ -path. Then $\tilde{\alpha}_{M_{\tilde{\alpha}}} = (\alpha_1, \alpha_2, \dots, \alpha_{2d+1})$ where α_{2k+1} is the image under folding of the arc $\tilde{\tau}_j^b$ if $\tilde{\alpha}_{2k+1} = \tilde{\tau}_j$ is a diagonal in $\tilde{T}(\gamma)$ or, otherwise, the image under the folding of the arc $\tilde{\alpha}_{2k+1}$. Thus $\alpha_{2k+1} = \tilde{\alpha}_{2k+1}$. Moreover, $\alpha_{2k} = \tau_{i_k} = \tilde{\alpha}_{2k}$, and thus $\tilde{\alpha}_{M_{\tilde{\alpha}}} = \tilde{\alpha}$.

Conversely, let $M = \{\beta_1, \beta_3, \dots, \beta_{2d-1}, \beta_{2d+1}\}$ be a perfect matching of $G_{T,\gamma}$. Then $M_{\tilde{\alpha}_M} = \{\tilde{\beta}_1, \tilde{\beta}_3, \dots, \tilde{\beta}_{2d-1}, \tilde{\beta}_{2d+1}\}$ where

$$\begin{aligned} \tilde{\beta}_{2k+1} &= \begin{cases} \tilde{\alpha}_{2k+1} & \text{if } \tilde{\alpha}_{2k+1} \text{ is a boundary arc,} \\ \tilde{\tau}_j^b & \text{if } \tilde{\alpha}_{2k+1} = \tilde{\tau}_j \text{ is a diagonal} \end{cases} \\ &= \begin{cases} \tilde{\tau}_{[\gamma_j]} & \text{if } \beta_{2k+1} = \tilde{\tau}_{[\gamma_j]}, \\ \tilde{\tau}_j^b & \text{if } \beta_{2k+1} = \tilde{\tau}_j^b. \end{cases} \end{aligned}$$

Hence $M_{\tilde{\alpha}_M} = M$. \square

Combining Lemma 4.3 with the results of [S2], we obtain the following Theorem.

Theorem 4.4. *There is a bijection between the set of perfect matchings of the graph $G_{T,\gamma}$ and the set of complete (T, γ) -paths in (S, M) given by $M \mapsto \pi(\tilde{\alpha}_M)$, where $\tilde{\alpha}_M$ is the image of M under the folding map and π is induced by the universal cover $\pi : \tilde{S} \rightarrow S$. Moreover, the numerator of the Laurent monomial $x(\pi(\tilde{\alpha}_M))$ of the complete (T, γ) -path $\pi(\tilde{\alpha}_M)$ is equal to the weight $w(M)$ of the matching M .*

Proof. The map in the Theorem is a bijection, because it is the composition of the folding map, which is a bijection, by Lemma 4.3, and the map π , which is a bijection, by [S2, Lemma 5.8]. The last statement of the Theorem follows from the construction of the graph $G_{T,\gamma}$. \square

4.6. **Proof of Theorem 3.1.** It has been shown in [S2, Theorem 3.2] that

$$(3) \quad x_\gamma = \sum_{\alpha} x(\alpha) y(\alpha),$$

where the sum is over all complete (T, γ) -paths α in (S, M) , $y(\alpha)$ is a monomial in \mathbf{y}_T , and

$$(4) \quad x(\alpha) = \prod_{k \text{ odd}} x_{\alpha_k} \prod_{k \text{ even}} x_{\alpha_k}^{-1}.$$

Applying Theorem 4.4 to equation (3) yields

$$(5) \quad x_\gamma = \sum_M w(M) y(M) (x_{i_1} x_{i_2} \cdots x_{i_d})^{-1},$$

where the sum is over all perfect matchings M of $G_{T, \gamma}$, $w(M)$ is the weight of the matching and $y(M) = y(\pi(\tilde{\alpha}_M))$. This completes the proof of Theorem 3.1.

5. A FORMULA FOR $y(M)$

In this section, we give a description of the coefficients $y(M)$ in terms of the matching M . First, we need to recall some results from [S2].

Recall that T is a triangulation of the unpunctured surface (S, M) , γ is an arc in (S, M) that crosses T exactly d times, we have fixed an orientation for γ and denote by $s = p_0, p_1, \dots, p_d, p_{d+1} = t$ the intersection points of γ and T in order of occurrence on γ . Let i_1, i_2, \dots, i_d be such that p_k lies on the arc $\tau_{i_k} \in T$, for $k = 1, 2, \dots, d$.

For $k = 0, 1, \dots, d$, let γ_k denote the segment of the path γ from the point p_k to the point p_{k+1} . Each γ_k lies in exactly one triangle Δ_k in T . If $1 \leq k \leq d-1$, the triangle Δ_k is formed by the arcs $\tau_{i_k}, \tau_{i_{k+1}}$ and a third arc that we denote by $\tau_{[\gamma_k]}$.

The orientation of the surface S induces an orientation on each of these triangles in such a way that, whenever two triangles Δ, Δ' share an edge τ , then the orientation of τ in Δ is opposite to the orientation of τ in Δ' . There are precisely two such orientations, we assume without loss of generality that we have the ‘‘clockwise orientation’’, that is, in each triangle Δ , going around the boundary of Δ according to the orientation of Δ is clockwise when looking at it from outside the surface.

Let α be a complete (T, γ) -path. Then $\alpha_{2k} = \tau_{i_k}$ is a common edge of the two triangles Δ_{k-1} and Δ_k . We say that α_{2k} is γ -oriented if the orientation of α_{2k} in the path α is the same as the orientation of τ_{i_k} in the triangle Δ_k , see Figure 6.

It is shown in [S2, Theorem 3.2] that

$$(6) \quad y(\alpha) = \prod_{k: \alpha_{2k} \text{ is } \gamma\text{-oriented}} y_{i_k}.$$

Each perfect matching M of $G_{T, \gamma}$ induces a path α_M in $\overline{G}_{T, \gamma}$ as in the construction of the folding map in section 4.3. The even arcs of α_M are the diagonals of the graph $\overline{G}_{T, \gamma}$. We say that an even arc of α_M has upward orientation if α_M is directed from southeast to northwest on that even arc, otherwise we say that the arc has downward orientation. If going upward on the first even arc of α_M is γ -oriented then we have that the $(2k)$ -th arc of $\pi(\tilde{\alpha}_M)$ is γ -oriented if and only if the $2k$ -th arc of α_M is upward if k is odd, and downward if k is even. If, on the




FIGURE 6. Two examples of the (T, γ) -path segment $(\alpha_{2k-1}, \alpha_{2k}, \alpha_{2k+1})$. On the left, α_{2k} is not γ -oriented and on the right, α_{2k} is γ -oriented.

other hand, going downward on the first even arc of α_M is γ -oriented then we have that the $(2k)$ -th arc of $\pi(\tilde{\alpha}_M)$ is γ -oriented if and only if the $2k$ -th arc of α_M is downward if k is odd, and upward if k is even.

There are precisely two perfect matchings M_+ and M_- of $G_{T,\gamma}$ that contain only boundary edges of $G_{T,\gamma}$. The orientations of the even arcs in both of the induced (T, γ) -paths α_{M_+} and α_{M_-} are alternately upward and downward, thus for one of the two paths, say M_+ , each even arc of $\pi(\tilde{\alpha}_{M_+})$ is γ -oriented, whereas for M_- none of the even arcs of $\pi(\tilde{\alpha}_{M_-})$ is γ -oriented. That is, $y(M_-) = 1$ and $y(M_+) = y_{i_1} y_{i_2} \cdots y_{i_d}$.

For an arbitrary perfect matching M , the coefficient $y(M)$ is determined by the set of edges of the symmetric difference $M_- \ominus M = (M_- \cup M) \setminus (M_- \cap M)$ as follows.

Theorem 5.1. *The set $M_- \ominus M$ is the set of boundary edges of a (possibly disconnected) subgraph G_M of $G_{T,\gamma}$ which is a union of tiles*

$$G_M = \cup_{j \in J} S_j.$$

Moreover,

$$y(M) = \prod_{j \in J} y_{i_j}$$

Proof. Choose any edge e_1 and either endpoint in $M_- \setminus (M_- \cap M)$, and walk along that edge until its other endpoint. Since M is a perfect matching, this endpoint is incident to an edge e_2 in M , which is different from e_1 and, hence, not in M_- . Thus $e_2 \in M \setminus (M_- \cap M)$. Now walk along e_2 until its other endpoint. This endpoint is incident to an edge e_3 in M_- which is different from e_2 , and, hence, not in M . Thus $e_3 \in M_- \setminus (M_- \cap M)$. Continuing this way, we construct a sequence of edges in $M_- \ominus M$. Since $G_{T,\gamma}$ has only finitely many edges, this sequence must become periodic after a certain number of steps; thus there exist p, N such that $e_k = e_{k+p}$ for all $k \geq N$.

We will show that one can take $N = 1$. Suppose to the contrary that $N \geq 2$ is the smallest integer such that $e_k = e_{k+p}$ for all $k \geq N$. Then e_{N-1}, e_N and e_{N+p-1}

share a common endpoint. But e_{N-1}, e_N and e_{N+p-1} are elements of the union of two perfect matchings, hence $e_{N-1} = e_{N+p-1}$, contradicting the minimality of N .

Therefore the sequence e_1, e_2, \dots, e_p in $M \ominus M_-$ is the set of boundary edges of a connected subgraph of $G_{T,\gamma}$ which is a union of tiles.

The graph G_M is the union of these connected subgraphs and, hence, it is a union of tiles. Let H be a connected component of G_M . There are precisely two perfect matchings $M_-(H)$ and $M_+(H)$ of H that consist only of boundary edges of H . Clearly, these two matchings are $M_- \cap E(H)$ and $M \cap E(H)$, where $E(H)$ is the set of edges of the graph H . Therefore, in each tile of H , the orientation of the diagonal in α_{M_-} and α_M are opposite. The restrictions of M_- and M to $E(G_{T,\gamma}) \setminus E(G_M)$ are identical, hence in each tile of $G_{T,\gamma} \setminus G_M$, the orientations of the diagonal in α_{M_-} and α_M are equal. It follows from equation (6) that $y(M) = \prod_{j \in J} y_{i_j}$. \square

It has been shown in [FZ2] that, for any cluster variable x_γ in \mathcal{A} , its Laurent expansion in the initial seed $(\mathbf{x}_T, \mathbf{y}_T, B_T)$ is homogeneous with respect to the grading given by $\deg(x_i) = \mathbf{e}_i$ and $\deg(y_i) = B_T \mathbf{e}_i$, where $\mathbf{e}_i = (0, \dots, 0, 1, 0, \dots, 0)^T \in \mathbb{Z}^n$ with 1 at position i . By definition, the g -vector g_γ of a cluster variable x_γ is the degree of its Laurent expansion with respect to this grading.

Corollary 5.2. *The g -vector g_γ of x_γ is given by*

$$g_\gamma = \deg \frac{w(M_-)}{x_{i_1} x_{i_2} \cdots x_{i_d}}.$$

Proof. This follows from the fact that $y(M_-) = 1$. \square

Remark 5.3. *The formula for $y(M)$ can also be phrased in terms of height functions. As described in section 3 of [Pr2], one way to define the height function on the faces of a bipartite planar graph G , covered by a perfect matching M , is to superimpose each matching with the fixed matching M_\emptyset (the unique matching of minimal height). In the case where G is a snake graph, we take M_\emptyset to be M_- , one of the two matchings of G only involving edges on the boundary. Color the vertices of G black and white so that no two adjacent vertices have the same color. In this superposition, we orient edges of M from black to white, and edges of M_- from white to black. We thereby obtain a spanning set of cycles, and removing the cycles of length two exactly corresponds to taking the symmetric difference $M \ominus M_-$. We can read the resulting graph as a relief-map, in which the altitude changes by $+1$ or -1 as one crosses over a contour line, according to whether the counter-line is directed clockwise or counter-clockwise. By this procedure, we obtain a height function $h_M : F(G) \rightarrow \mathbb{Z}$ which assigns integers to the faces of graph G . When G is a snake graph, the set of faces $F(G)$ is simply the set of tiles $\{S_j\}$ of G . Comparing with the definition of $y(M)$ in Theorem 5.1, we see that*

$$y(M) = \prod_{S_j \in F(G)} y_j^{h_M(j)}.$$

An alternative definition of height functions comes from [EKLP] by translating the matching problem into a domino tiling problem on a region colored as a checkerboard. We imagine an ant starting at an arbitrary vertex at height 0, walking along the boundary of each domino, and changing its height by $+1$ or -1 as it traverses the boundary of a black or white square, respectively. The values of the height function under these two formulations agree up to scaling by four.

6. CLUSTER EXPANSION WITHOUT MATCHINGS

In this section, we give a formula for the cluster expansion of x_γ in terms of the graph $G_{T,\gamma}$ only.

For any graph H , let $c(H)$ be the number of connected components of H . Let $E(H)$ be the set of edges of H , and denote by ∂H the set of boundary edges of H . Define \mathcal{H}_k to be the set of all subgraphs H of $G_{T,\gamma}$ such that H is a union of k tiles $H = S_{j_1} \cup \dots \cup S_{j_k}$ and the number of edges of M_- that are contained in H is equal to $k + c(H)$. For $H \in \mathcal{H}_k$, let

$$y(H) = \prod_{S_{i_j} \text{ tile in } H} y_{i_j}.$$

Theorem 6.1. *The cluster expansion of the cluster variable x_γ is given by*

$$x_\gamma = \sum_{k=0}^d \sum_{H \in \mathcal{H}_k} \frac{w(\partial H \ominus M_-) y(H)}{x_{i_1} x_{i_2} \cdots x_{i_d}},$$

Proof. It follows from the theorems 3.1 and 5.1 that

$$x_\gamma = \sum_{k=1}^d \sum_{M: |y(M)|=k} \frac{w(M) y(G_M)}{x_{i_1} x_{i_2} \cdots x_{i_d}},$$


where $|y(M)|$ is the number of tiles in G_M . We will show that for all k , the map $M \mapsto G_M$ is a bijection between the set of perfect matchings M of $G_{T,\gamma}$ such that $|y(M)| = k$ and the set \mathcal{H}_k .

- The map is well-defined. Clearly, G_M is the union of k tiles. Moreover, $E(G_M) \cap M_-$ is a perfect matching of G_M , since M_- consists of every other boundary edge of $G_{T,\gamma}$. Thus the cardinality of $(E(G_M) \cap M_-)$ is half the number of vertices of G_M , which is equal to $2k + 2 + 2c(G_M)$. Therefore, the cardinality of $(E(G_M) \cap M_-)$ is $k + c(G_M)$ and $G_M \in \mathcal{H}_k$.
- The map is injective, since two graphs $G_M, G_{M'}$ are equal if and only if their boundaries are.
- The map is surjective. Let $H = S_{j_1} \cup \dots \cup S_{j_k}$ such that the cardinality of $E(H) \cap M_-$ equals $k + c(H)$. The boundary of H consists of $2k + 2 + 2c(H)$ edges, half of which lie in M_- . As in the proof of Theorem 5.1, let $M_-(H) = E(H) \cap M_-$ and $M_+(H)$ be the two perfect matchings of H that consist of boundary edges only. Let $M = M_+(H) \cup (M_- \setminus M_-(H))$. Then M is a perfect matching of $G_{T,\gamma}$ such that $G_M = H$, and moreover, $|y(M)|$ is equal to the number of tiles in H , which is k . Thus the map is surjective.

Now the boundary edges of G_M are precisely the elements of $M \ominus M_-$, which implies that $\partial(G_M) \ominus M_- = (M \ominus M_-) \ominus M_- = M \ominus (M_- \ominus M_-) = M$. Therefore $w(M) = w(\partial(G_M) \ominus M_-)$, and this completes the proof. \square

Corollary 6.2. *The F -polynomial of γ is given by*

$$F_\gamma = \sum_{k=0}^d \sum_{H \in \mathcal{H}_k} y(H).$$

FIGURE 7. Triangulated surface with dotted arc γ

7. EXAMPLE

We illustrate Theorem 3.1, Theorem 5.1 and Theorem 6.1 in an example. Let (S, M) be the annulus with two marked points on each of the two boundary components, and let $T = \{\tau_1, \dots, \tau_8\}$ be the triangulation shown in Figure 7. Let γ be the dotted arc in that figure. It has $d = 6$ crossings with the triangulation. The sequence of crossed arcs $\tau_{i_1}, \dots, \tau_{i_6}$ is $\tau_1, \tau_2, \tau_3, \tau_4, \tau_1, \tau_2$, and the corresponding segments $\gamma_0, \dots, \gamma_6$ of the arc γ are labeled in the figure. Moreover, $\tau_{[\gamma_1]} = \tau_6$, $\tau_{[\gamma_2]} = \tau_8$, $\tau_{[\gamma_3]} = \tau_7$, $\tau_{[\gamma_4]} = \tau_5$ and $\tau_{[\gamma_5]} = \tau_6$.

The graph $G_{T, \gamma}$ is obtained by glueing the corresponding six tiles $\tilde{S}_1, \tilde{S}_2, \tilde{S}_3, \tilde{S}_4, \tilde{S}_1$, and \tilde{S}_2 . The result is shown in Figure 8.




FIGURE 8. Construction of the graphs $\overline{G}_{T,\gamma}$ and $G_{T,\gamma}$

Theorems 3.1 and 5.1 imply that $x_\gamma(x_{i_1}x_{i_2} \cdots x_{i_d})$ is equal to

$$\begin{aligned}
 & x_4x_6x_8x_4x_4x_6x_8 y_1y_3y_4y_1 + x_4x_6x_8x_4x_4x_1x_3 y_1y_3y_4y_1y_2 \\
 + & x_4x_6x_8x_4x_5x_2x_8 y_1y_3y_4 + x_4x_6x_2x_3x_1x_2x_8 y_1 \\
 + & x_4x_6x_2x_7x_5x_2x_8 y_1y_4 + x_4x_6x_2x_7x_4x_6x_8 y_1y_4y_1 \\
 + & x_4x_1x_3x_4x_4x_6x_8 y_1y_2y_3y_4y_1 + x_4x_1x_3x_4x_4x_1x_3 y_1y_2y_3y_4y_1y_2 \\
 + & x_4x_1x_3x_4x_5x_2x_8 y_1y_2y_3y_4 + x_5x_2x_8x_4x_4x_6x_8 y_3y_4y_1 \\
 + & x_5x_2x_8x_4x_4x_1x_3 y_3y_4y_1y_2 + x_5x_2x_8x_4x_5x_2x_8 y_3y_4 \\
 + & x_5x_2x_2x_3x_1x_2x_8 + x_5x_2x_2x_7x_4x_6x_8 y_4y_1 \\
 + & x_5x_2x_2x_7x_4x_1x_3 y_4y_1y_2 + x_5x_2x_2x_7x_5x_2x_8 y_4,
 \end{aligned}$$

which is equal to

$$\begin{array}{rcl}
& x_4^3 y_1^2 y_3 y_4 & + x_1 x_3 x_4^3 y_1^2 y_2 y_3 y_4 \\
+ & x_2 x_4^2 y_1 y_3 y_4 & + x_1 x_2^2 x_3 x_4 y_1 \\
+ & x_2^2 x_4 y_1 y_4 & + x_2 x_4^2 y_1^2 y_4 \\
+ & x_1 x_3 x_4^3 y_1^2 y_2 y_3 y_4 & + x_1^2 x_3^2 x_4^3 y_1^2 y_2^2 y_3 y_4 \\
+ & x_1 x_2 x_3 x_4^2 y_1 y_2 y_3 y_4 & + x_2 x_4^2 y_1 y_3 y_4 \\
+ & x_1 x_2 x_3 x_4^2 y_1 y_2 y_3 y_4 & + x_2^2 x_4 y_3 y_4 \\
+ & x_1 x_2^3 x_3 & + x_2^2 x_4 y_1 y_4 \\
+ & x_1 x_2^2 x_3 x_4 y_1 y_2 y_4 & + x_2^3 y_4.
\end{array}$$

For example, the first term corresponds to the matching M consisting of the horizontal edges of the first three tiles and the horizontal edges of the last two tiles. The matching M_- consists in the boundary edges weighted x_5 and x_2 in the first tile, x_2 in the third tile, x_1 and x_3 in the fourth, x_2 in the fifth and x_8 in the sixth tile. Thus $M_- \ominus M = (M_- \cup M) \setminus (M_- \cap M)$ is the union of the first, third, fourth and fifth tile, whence $y(M) = y_{i_1} y_{i_3} y_{i_4} y_{i_5} = y_1 y_3 y_4 y_1$.

To illustrate Theorem 6.1, let $k = 2$. Then \mathcal{H}_k consists of the subgraphs H of $G_{T,\gamma}$ which are unions of two tiles and such that $E(H) \cap M_-$ has three elements if H is connected, respectively four elements if H has two connected components. Thus \mathcal{H}_2 has three elements

$$\mathcal{H}_2 = \{S_{i_3} \cup S_{i_4}, S_{i_4} \cup S_{i_5}, S_{i_1} \cup S_{i_4}\}$$

corresponding to the three terms

$$x_2^2 x_4 y_3 y_4, x_2^2 x_4 y_1 y_4 \text{ and } x_2^2 x_4 y_1 y_4.$$

REFERENCES

- [A] C. Amiot, Cluster categories for algebras of global dimension 2 and quivers with potential, preprint, [arXiv:0805.1035](#).
- [ABCP] I. Assem, T. Brüstle, G. Charbonneau-Jodoin, P.G. Plamondon, Gentle algebras arising from surface triangulations, in preparation.
- [BMRRT] A. Buan, R. Marsh, M. Reineke, I. Reiten and G. Todorov, Tilting theory and cluster combinatorics, *Adv. Math.* **204** (2006), 572–612, [arXiv:math.RT/0402054](#).
- [BMR] A. B. Buan, R. Marsh and I. Reiten, Denominators of cluster variables, preprint, [arXiv:0710.4335](#).
- [CC] P. Caldero and F. Chapoton, Cluster algebras as Hall algebras of quiver representations, *Comment. Math. Helv.* **81** (2006), 595–616, [arXiv:math.RT/0410187](#).
- [CCS] P. Caldero, F. Chapoton and R. Schiffler, Quivers with relations arising from clusters (A_n case), *Trans. Amer. Math. Soc.* **358** (2006), no. 3, 1347–1364, [arXiv:math.RT/0401316](#).
- [CCS2] P. Caldero, F. Chapoton and R. Schiffler, Quivers with relations and cluster tilted algebras, *Algebr. and Represent. Theory* **9**, (2006), no. 4, 359–376, [arXiv:math.RT/0411238](#).
- [CK] P. Caldero and B. Keller, From triangulated categories to cluster algebras, *Invent. Math.* **172** (2008), 169–211, [arXiv:math.RT/0506018](#).
- [CK2] P. Caldero, B. Keller, From triangulated categories to cluster algebras II, *Ann. Sci. École Norm. Sup. (4)* **39** (2006), no. 6, 983–1009, [arXiv:math.RT/0510251](#).
- [CZ] P. Caldero and A. Zelevinsky, Laurent expansions in cluster algebras via quiver representations, *Mosc. Math. J.* **6** (2006), no. 3, 411–429, [arXiv:math.RT/0604054](#).
- [CP] G. Carroll and G. Price, (unpublished result).
- [EKLP] N. Elkies, G. Kuperberg, M. Larsen, J. Propp, Alternating-Sign Matrices and Domino Tilings (Part I), *J. Algebraic Combin.* **1** (1992), no. 2, 11–132, [arXiv:math/9201305](#).
- [FG1] V. Fock and A. Goncharov, Moduli spaces of local systems and higher Teichmüller theory. *Publ. Math. Inst. Hautes Études Sci.* No. 103 (2006), 1–211.

- [FG2] V. Fock and A. Goncharov, Cluster ensembles, quantization and the dilogarithm, preprint (2003), [arXiv:math.AG/0311149](#).
- [FG3] V. Fock and A. Goncharov, Dual Teichmüller and lamination spaces. Handbook of Teichmüller theory. Vol. I, 647–684, IRMA Lect. Math. Theor. Phys., 11, Eur. Math. Soc., Zürich, 2007.
- [FST] S. Fomin, M. Shapiro and D. Thurston, Cluster algebras and triangulated surfaces. Part I: Cluster complexes, preprint (2006), [arXiv:math/0608367](#).
- [FT] S. Fomin and D. Thurston, Cluster algebras and triangulated surfaces. Part II: Lambda lengths, preprint (2008).
- [FZ1] S. Fomin and A. Zelevinsky, Cluster algebras I. Foundations, *J. Amer. Math. Soc.* **15**(2), (2002), 497–529 (electronic), [arXiv:math.RT/0104151](#).
- [FZ2] S. Fomin and A. Zelevinsky, Cluster Algebras IV: Coefficients, *Comp. Math.* **143** (2007), 112–164, [arXiv:math.RA/0602259](#).
- [FZ3] S. Fomin and A. Zelevinsky, (unpublished result).
- [FK] C. Fu and B. Keller, On cluster algebras with coefficients and 2-Calabi-Yau categories, preprint (2008), [arXiv:0710.3152](#).
- [GSV1] M. Gekhtman, M. Shapiro and A. Vainshtein, Cluster algebras and Poisson geometry. *Mosc. Math. J.* **3** (2003), no. 3, 899–934, 1199, [arXiv:math.QA/0208033](#).
- [GSV2] M. Gekhtman, M. Shapiro and A. Vainshtein, Cluster algebras and Weil-Petersson forms. *Duke Math. J.* **127** (2005), no. 2, 291–311, [arXiv:math.QA/0309138](#).
- [LF] D. Labardini-Fragoso, Quivers with potentials associated to triangulated surfaces, preprint (2008), [arXiv:0803.1328](#).
- [MP] G. Musiker and J. Propp, Combinatorial interpretations for rank-two cluster algebras of affine type. *Electron. J. Combin.* **14** (2007), no. 1, Research Paper 15, 23 pp. (electronic). [arXiv:math/0602408](#).
- [M] G. Musiker, A graph theoretic expansion formula for cluster algebras of classical type, to appear in *Ann. Comb.*, [arXiv:0710.3574](#).
- [MW] G. Musiker and L. Williams, Combinatorial formulas for F -polynomials and g -vectors for cluster algebras of classical type, in preparation.
- [Pa] Y. Palu, Cluster characters for triangulated 2-Calabi–Yau categories, to appear in *Annales de l’Institut Fourier*, [arXiv:math/0703540](#).
- [Pr1] J. Propp, The combinatorics of frieze patterns and Markoff numbers, preprint (2005), [arXiv:math.CO/0511633](#).
- [Pr2] J. Propp, Lattice structure for orientations of graphs, preprint (1993), [arXiv:math/0209.5005](#).
- [S1] R. Schiffler, A cluster expansion formula (A_n case), *Electron. J. Combin.* **15** (2008), #R64 1, [arXiv:math.RT/0611956](#).
- [S2] R. Schiffler, On cluster algebras arising from unpunctured surfaces II, preprint (2008), [arXiv:0809.2593](#).
- [ST] R. Schiffler and H. Thomas, On cluster algebras arising from unpunctured surfaces, preprint (2007), [arXiv:0712.4131](#).
- [SZ] P. Sherman and A. Zelevinsky, Positivity and canonical bases in rank 2 cluster algebras of finite and affine types. *Mosc. Math. J.* **4** (2004), no. 4, 947–974, 982, [arXiv:math.RT/0307082](#).
- [Z] A. Zelevinsky, Semicanonical basis generators of the cluster algebra of type $A_1^{(1)}$. *Electron. J. Combin.* **14** (2007), no. 1, Note 4, 5 pp. (electronic), [arXiv:math.RA/0606775](#).

Gregg Musiker
 Department of Mathematics, Room 2-332
 Massachusetts Institute of Technology
 77 Massachusetts Ave.
 Cambridge, MA 02139
 musiker (at) math.mit.edu

Ralf Schiffler
 Department of Mathematics
 University of Connecticut
 196 Auditorium Road
 Storrs, CT 06269-3009
 schiffler (at) math.uconn.edu