Optimal codes in deletion and insertion metric¹

Hyun Kwang Kim², Joon Yop Lee², and Dong Yeol Oh³

Abstract

We improve the upper bound of Levenshtein for the cardinality of a code of length 4 capable of correcting single deletions over an alphabet of even size. We also illustrate that the new upper bound is sharp. Furthermore we will construct an optimal perfect code capable of correcting single deletions for the same parameters.

1 Introduction

Let B_q be a set with q elements. The set B_q is referred to as an alphabet and its elements are referred to as letters. We will assume for definiteness that $B_q = \{0, 1, \ldots, q-1\}$. A sequence $x = (x_1, \ldots, x_n)$ of n letters of B_q is called a word, and the number n is called its length. Together with writing $x = (x_1, \ldots, x_n)$, we will also use the multiplicative writing $x = x_1 \cdots x_n$. Let B_q^n be the set of words over B_q of length n, and put

$$B_q^* = \bigcup_{n=0}^{\infty} B_q^n$$

The deletion and insertion distance $\rho(x, y)$, which was first introduced by Levenshtein [5], between two words x and y in B_q^* is defined by the minimum number of deletions and insertions of letters required to transform x into y. For example, let x = 12243 and y = 14223 be two words in B_5^5 . If we delete the fourth letter and the second letter of x and y respectively, we have derived words x' = 1223, y' = 1223 which are identical, hence $\rho(x, y) = 2$. For a code $C \subseteq B_q^n$, we put

$$\rho(C) = \min\left\{ \rho(x, y) \mid x, y \in C, x \neq y \right\}.$$

Let $N(n, q, d) = \max \{ |C| | C \subseteq B_q^n, \rho(C) > 2d \}$. A code C in B_q^n is called an optimal code if |C| = N(n, q, d). In [6], the author proved an upper bound for N(n, q, 1) for any $n(\geq 2)$ and $q(\geq 2)$, namely

$$N(n,q,1) \le \lfloor \frac{q^{n-1} + (n-2)q^{n-2} + q}{n} \rfloor.$$
 (1)

The upper bound in (1) is sharp when n = 3. Indeed, a code capable of correcting single deletions whose cardinality meets the upper bound in (1) was

¹The material of this paper was presented in part at the 10th International Workshop on Algebraic and Combinatorial Coding Theory, Zvenigorod, Russia, September 2006.

 $^{^{2}}$ This work was supported by the Basic Research Program of the Korea Science and Engineering Foundation (grant # R01-2006-000-11176-0). Department of Mathematics, Pohang University of Science and Technology, Pohang 790-784, The Republic of Korea (hkkim@postech.ac.kr, flutelee@postech.ac.kr).

 $^{^3 \}rm National Institute for Mathematical Sciences, Daejeon 305-340, The Republic of Korea(dyoh@nims.re.kr).$

constructed in [6] when n = 3 and $q \geq 2$ is an arbitrary integer. However this bound is not sharp when $n \geq 4$. In this paper, we improve the upper bound of Levenshtein in the case of n = 4 and q even, and prove that the new upper bound is sharp.

In Section 2, we will derive a new upper bound for the cardinality of a code capable of correcting single deletions when n = 4 and q is even by analyzing the deletion map.

In Section 3, we will construct a code in B_q^4 for $q \equiv 2, 4 \pmod{6}$ capable of correcting single deletions whose cardinalities meet the new upper bound in Section 2. We first introduce the concept of the step property for a Steiner quadruple system, and prove that there is a code in B_q^4 capable of correcting single deletions whose cardinalities meet the upper bound in Section 2 under the assumption that there is a Steiner quadruple system on B_q which satisfies the step property. In the next, we show that there is a Steiner quadruple system on B_q which satisfies the step property for $q \equiv 2, 4 \pmod{6}$.

In Section 4, we will construct a code in B_q^4 for $q \equiv 0 \pmod{6}$ capable of correcting single deletions whose cardinalities meet the upper bound in Section 2. Since there dose not exist a Steiner quadruple system for alphabets of this size, we will use a group divisible system. We divide our construction into two steps. In the first, we prove that there is an optimal code for an alphabet of size q = 6m where m is odd. In the next, we prove that there is an optimal code for an alphabet of size an alphabet of size 2q under the assumption that there is an optimal code for an alphabet of size q.

In Section 5, we modify our construction of optimal codes slightly, and construct an optimal perfect code in B_q^4 when q is even.

2 New upper bound for N(4, q, 1)

In this section, we improve the upper bound of Levenshtein for n = 4 and even q. Our result follows from an analysis of the deletion map which will be introduced in a moment.

We begin with a simple observation. For any word x over B_q and any number $s(\geq 1)$, denote by $\lfloor x \rfloor_s$ the set of words obtained form x by deleting s of its letters. The multi-map $\lfloor \rfloor_s : B_q^n \to B_q^{n-s}$ which sends x to $\lfloor x \rfloor_s$ will be called an s-deletion map or simply a deletion map. For a subset $C \subseteq B_q^n$, denote by $\lfloor C \rfloor_s$ the set $\bigcup_{x \in C} \lfloor x \rfloor_s$. A set $C \subseteq B_q^n$ is called a code capable of correcting s deletions if all the sets $\lfloor x \rfloor_s$ and $\lfloor x \rfloor_s$ for $z \in C$.

deletions if all the sets $\lfloor x \rfloor_s$, $x \in C$, do not pairwise intersect. It follows from the definition that, for a code C of B_q^n , $\rho(C) > 2s$ if and only if C is capable of correcting s deletions.

The following is an immediate consequence of the preceding observation.

Lemma 2.1 (Substitution lemma). Let C be a code of B_q^n such that $\rho(C) > 2s$. If $x \in C$ and $\lfloor y \rfloor_s \subseteq \lfloor x \rfloor_s$, then $\rho((C \setminus \{x\}) \cup \{y\}) > 2s$. More generally, if $A \subset C$, $\lfloor B \rfloor_s \subseteq \lfloor A \rfloor_s$, and $\rho(B) > 2s$, then $\rho((C \setminus A) \cup B) > 2s$.

From now on, C will denote a code in B_q^4 with $\rho(C) > 2$. Let

$$C_i = \{x \in C \mid ||x|_1| = i\} \text{ for } 1 \le i \le 4.$$

By the substitution lemma, we may assume without changing the cardinality of ${\cal C}$ that

 $\left\{ \begin{array}{l} x \in C_1 \Rightarrow x \text{ is of the type } (a, a, a, a); \\ x \in C_2 \Rightarrow x \text{ is of the type } (a, a, b, b); \\ x \in C_3 \Rightarrow x \text{ is of the type } (a, b, b, a), (a, a, b, c), (b, c, a, a), \text{ or } (a, b, b, c); \\ x \in C_4 \Rightarrow x \text{ is of the type } (a, b, c, d), (a, b, a, c), (a, b, c, a), \text{ or } (b, a, c, a). \end{array} \right.$

In the case of i = 3, 4, we analyze further and put

 $\left\{ \begin{array}{l} C_{3,1} = \{x \in C_3 \\ C_{3,2} = \{x \in C_3 \\ C_{4,1} = \{x \in C_4 \\ C_{4,2} = \{x \in C_4 \\ C_{4,3} = \{x \in C_4 \\ x \text{ is of the type } (a, b, c, a)\}, \\ C_{4,3} = \{x \in C_4 \\ x \text{ is of the type } (a, b, c, a)\}, \\ x \text{ is of the type } (a, b, c, a)\}, \\ x \text{ is of the type } (a, b, c, a)\}, \\ x \text{ is of the type } (a, b, c, a)\}, \\ x \text{ is of the type } (a, b, c, a)\}, \\ x \text{ is of the type } (a, b, a, c) \text{ or } (b, a, c, a)\}. \end{array} \right.$

Consider the following subsets of B_q^3 ;

$$\begin{cases} U = \{y \in B_q^3 \mid y \text{ is of the type } (a, a, a)\}, \\ V = \{y \in B_q^3 \mid y \text{ is of the type } (a, a, b) \text{ or } (a, b, b)\}, \\ W = \{y \in B_q^3 \mid y \text{ is of the type } (a, b, a)\}, \\ Z = \{y \in B_q^3 \mid y \text{ is of the type } (a, b, c)\}. \end{cases}$$

Note that we have

$$\begin{cases} |U| = q, |V| = 2q(q-1), \\ |W| = q(q-1), |Z| = q(q-1)(q-2). \end{cases}$$
(2)

The following table counts the contribution of each codeword in C_i (or $C_{i,j}$) to U, V, W, and Z under the deletion map $\lfloor \rfloor_1 : B_q^4 \to B_q^3$.

	C_1	C_2	$C_{3,1}$	$C_{3,2}$	$C_{4,1}$	$C_{4,2}$	$C_{4,3}$
U	1	0	0	0	0	0	0
V	0	2	2	2	0	0	1
W	0	0	0	1	0	2	1
Z	0	0	1	0	4	2	2

Table 1: The contribution of each codeword in C_i (or $C_{i,j}$) to U, V, W, and Z

Let $T \in \{U, V, W, Z\}$ and define $T_i = T \cap \lfloor C_i \rfloor_1$ and $T_{i,j} = T \cap \lfloor C_{i,j} \rfloor_1$ (for example $V_{3,2} = V \cap \lfloor C_{3,2} \rfloor_1$). One can easily obtain the following relations from (2) and Table 1;

 $\begin{cases} |U_1| \leq |U| = q, \\ |V_2| + |V_{3,1}| + |V_{3,2}| + |V_{4,3}| \leq |V| = 2q(q-1), \\ |W_{3,2}| + |W_{4,2}| + |W_{4,3}| \leq |W| = q(q-1), \\ |Z_{3,1}| + |Z_{4,1}| + |Z_{4,2}| + |Z_{4,3}| \leq |Z| = q(q-1)(q-2), \\ |C_1| = |U_1|, |C_2| = \frac{1}{2}|V_2|, |C_{3,1}| = \frac{1}{2}|V_{3,1}| = |Z_{3,1}|, |C_{3,2}| = \frac{1}{2}|V_{3,2}| = |W_{3,2}|, \\ |C_{4,1}| = \frac{1}{4}|Z_{4,1}|, |C_{4,2}| = \frac{1}{2}|W_{4,2}| = \frac{1}{2}|Z_{4,2}|, |C_{4,3}| = |V_{4,3}| = |W_{4,3}| = \frac{1}{2}|Z_{4,3}|. \end{cases}$

We are now in the position to derive the first main result.

Theorem 2.2. Let $C \subseteq B_q^4$ be a code with $\rho(C) > 2$. Then

$$|C| \le \frac{q(q^2 + q + 2\lfloor \frac{q-1}{2} \rfloor + 2)}{4} = \begin{cases} \frac{q^2(q+2)}{4} & \text{if } q \text{ is even,} \\ \frac{q(q+1)^2}{4} & \text{if } q \text{ is odd.} \end{cases}$$

Proof. It follows from the above calculation that

$$\begin{split} |C| &= |C_1| + |C_2| + |C_{3,1}| + |C_{3,2}| + |C_{4,1}| + |C_{4,2}| + |C_{4,3}| \\ &= |U_1| + \frac{1}{2}|V_2| + \frac{1}{2}|V_{3,1}| + \frac{1}{2}|V_{3,2}| + \frac{1}{4}|Z_{4,1}| + \frac{1}{2}|Z_{4,2}| + \frac{1}{2}|Z_{4,3}| \\ &= |U_1| + \frac{1}{2}(|V_2| + |V_{3,1}| + |V_{3,2}| + |V_{4,3}|) - \frac{1}{2}|V_{4,3}| \\ &+ \frac{1}{4}(|Z_{3,1}| + |Z_{4,1}| + |Z_{4,2}| + |Z_{4,3}|) - \frac{1}{4}|Z_{3,1}| + \frac{1}{4}|Z_{4,2}| + \frac{1}{4}|Z_{4,3}| \\ &= |U_1| + \frac{1}{2}(|V_2| + |V_{3,1}| + |V_{3,2}| + |V_{4,3}|) + \frac{1}{4}(|Z_{3,1}| + |Z_{4,1}| + |Z_{4,2}| + |Z_{4,3}|) \\ &+ \frac{1}{2}|C_{4,2}| - \frac{1}{4}|C_{3,1}| \\ &= |U| + \frac{1}{2}|V| + \frac{1}{4}|Z| + \frac{1}{2}|C_{4,2}| - \frac{1}{4}|C_{3,1}|. \end{split}$$

Since $|C_{4,2}| \le q \lfloor \frac{q-1}{2} \rfloor$ and $|C_{3,1}| \ge 0$, the result follows.

Remark. The proof of Theorem 2.2 can be used in the construction of optimal codes as follows: Let C be a code in B_q^4 with $|C| = \frac{1}{4}q(q^2 + q + 2\lfloor \frac{q-1}{2} \rfloor + 2)$. Then we should have

$$\begin{aligned} |U_1| &= |U| = q, |V_2| + |V_{3,1}| + |V_{3,2}| + |V_{4,3}| = |V| = q(q-1), |C_{3,1}| = 0, \\ |Z_{3,1}| + |Z_{4,1}| + |Z_{4,2}| + |Z_{4,3}| = |Z| = q(q-1)(q-2), |C_{4,2}| = q\lfloor \frac{q-1}{2} \rfloor. \end{aligned}$$

Since $|C_{3,1}| = 0$, we have $|V_{3,1}| = |Z_{3,1}| = 0$. From the fact that $|C_{4,2}| = q\lfloor \frac{q-1}{2} \rfloor$, we may assume that $|C_{3,2}| = |C_{4,3}| = 0$. Then $|V_{3,2}| = |V_{4,3}| = |Z_{4,3}| = 0$. In this case $|V_2| = |V|$. Hence

$$\begin{aligned} |C_1| &= q, \ |C_2| = q(q-1), \ |C_3| = 0, \\ |C_{4,1}| &= \frac{1}{4}(q(q-1)(q-2) - 2q\lfloor \frac{q-1}{2} \rfloor), \ |C_{4,2}| = q\lfloor \frac{q-1}{2} \rfloor. \end{aligned}$$

This means that if we include every word of the types (a, a, a, a) and (a, a, b, b), maximal number of words of the type (a, b, c, a), and generate words of the type (a, b, c, d), using all remaining words of length 3 of the type (a, b, c), we can obtain an optimal code. Since $|U_1| = |U|$, and $|V_2| = |V|$ are always possible, to construct an optimal code one only needs to consider combinations of elements in W and Z which satisfy

$$|C_4| = \frac{1}{4}(q(q-1)(q-2) - 2q\lfloor \frac{q-1}{2} \rfloor) + q\lfloor \frac{q-1}{2} \rfloor.$$

From these considerations, we can construct optimal codes in B_q^4 and B_q^6 which coincide with the construction in [4]. We list the codewords of an optimal code in B_q^4 in Table 2, and the codewords in C_4 of an optimal code in B_q^6 in Table 3.

In the next two sections, we will construct an optimal code in B_q^4 when q is even. Let x be a word of length 4 consists of pairwise distinct letters. The basic ingredients in our construction are the codes $\langle x \rangle_{A_q^4}$ and $\langle x \rangle_{B_q^4}$ generated by x in A_q^4 and B_q^4 , which will be defined in the below. The code $\langle x \rangle_{A_q^4}$ was already used in [6]. The essence of our construction is to replace $\langle x \rangle_{A_q^4}$ by

0000	1111	2222	3333
0011	0022	0033	1100
1122	1133	2200	2211
2233	3300	3311	3322
0230	1231	2012	3013
0321	2103	1302	3120

Table 2: An optimal code in B_4^4

0230	1231	2012	3013	4014	5015
0450	1451	2452	3453	4234	5235
0251	1304	2053	3105	4035	5102
0342	1325	2140	3124	4120	5143
0431	1503	2413	3520	4215	5321
0524	1542	2504	3541	4302	5340

Table 3: The C_4 of an optimal code in B_6^4

 $\langle x \rangle_{B^4_q}$ for some words such that our construction satisfies the conditions of the preceding remark.

Let A_q^n be the set of all words in B_q^n with pairwise distinct letters. For a word $x = (a_1, a_2, a_3, a_4)$ in A_q^4 , the codes $\langle x \rangle_{A_q^4}$ and $\langle x \rangle_{B_q^4}$ generated by x in A_q^4 and B_q^4 , respectively, are defined as follows:

$$\begin{split} < x >_{A_q^4} &= \{(a_1, a_2, a_3, a_4), (a_1, a_4, a_3, a_2), (a_2, a_4, a_1, a_3), \\ &\quad (a_3, a_4, a_1, a_2), (a_3, a_2, a_1, a_4), (a_4, a_2, a_3, a_1)\}, \\ < x >_{B_q^4} &= (< x >_{A_q^4} \setminus \{(a_1, a_2, a_3, a_4), (a_3, a_4, a_1, a_2)\}) \\ &\quad \cup \{(a_1, a_3, a_4, a_1), (a_2, a_3, a_4, a_2), (a_3, a_1, a_2, a_3), (a_4, a_1, a_2, a_4)\}. \end{split}$$

The following lemma which describes the basic properties of these codes under the deletion map can be easily verified.

Lemma 2.3. Let $x = (a_1, a_2, a_3, a_4)$ and $y = (b_1, b_2, b_3, b_4)$ be distinct elements of A_q^4 , and T(x) the set of the letters of x. Then we have

- (i) $\langle x \rangle_{A_q^4}$ and $\langle x \rangle_{B_q^4}$ are codes capable of correcting single deletions,
- (ii) if $|T(x) \cap T(y)| \leq 2$, then $\langle x \rangle_{A^4_q} \cup \langle y \rangle_{B^4_q}$ is a code capable of correcting single deletions,
- (iii) if either $\{a_1, a_2\} \cap \{b_1, b_2\} = \emptyset$ or $\{a_3, a_4\} \cap \{b_3, b_4\} = \emptyset$, then $\langle x \rangle_{B_q^4}$ $\cup \langle y \rangle_{B_q^4}$ is a code capable of correcting single deletions.

3 Construction of optimal codes in B_q^4 for $q \equiv 2, 4 \pmod{6}$

In this section, we will construct a code in B_q^4 for $q \equiv 2, 4 \pmod{6}$ capable of correcting single deletions whose cardinality meets the upper bound in Theorem 2.2. Our construction consists of two steps. In the first, we introduce the concept of the step property for a Steiner quadruple system, and prove that there is a code in B_q^4 capable of correcting single deletions whose cardinality meets the upper bound in Theorem 2.2 under the assumption that there is a Steiner quadruple system on B_q which satisfies the step property. In the next, following the construction of Hanani, we show that there is a Steiner quadruple system on B_q which satisfies the step property for $q \equiv 2, 4 \pmod{6}$.

We begin with the definition of the step property.

Definition 3.1. Let S be a Steiner quadruple system on B_q , in short SQS(q), and $\mathcal{O} = \{L_0, L_1, \ldots, L_{q-1}\}$ be a total order on B_q . We say that S satisfies the step property with respect to \mathcal{O} if, for every quadruple of the form $\{L_{2t}, L_{2t+1}, L_a, L_b\}$, we have either a, b < 2t or 2t + 1 < a, b. We say that Ssatisfies the step property if there is a total order \mathcal{O} on B_q such that S satisfies the step property with respect to \mathcal{O} .

Theorem 3.2. Suppose that there is an SQS(q) which satisfies the step property. Then

$$N(4,q,1) = \frac{q(q^2 + q + 2\lfloor \frac{q-1}{2} \rfloor + 2)}{4} = \frac{q^2(q+2)}{4}.$$

Proof. Let S be a SQS(q) which satisfies the step property. Without loss of generality, we may assume that $\{0, 1, \dots, q-1\}$ is the total order which admits the step property for S. Let $\phi : S \to B_q^4$ be the map defined by $\phi(\{x, y, z, w\}) = (x, y, z, w)$ where x < y < z < w, and put $T = \{\{2t, 2t + 1, a, b\} \in S | 2t + 1 < a, b\}$. It follows from the definition of a Steiner quadruple system and Lemma 2.3 that the code

$$C_1 = \bigcup_{x \in \phi(S \setminus T)} < x >_{A_q^4} \bigcup \bigcup_{x \in \phi(T)} < x >_{B_q^4}$$

is capable of correcting single deletions. Since T has $\frac{\frac{q}{2}(\frac{q}{2}-1)}{2} = \frac{q(q-2)}{8}$ elements, we have

$$|C_1| = 6\left\{\frac{q(q-1)(q-2)}{24} - \frac{q(q-2)}{8}\right\} + 8\frac{q(q-2)}{8} = \frac{q^2(q-2)}{4}$$

We obtain a code C from C_1 by adding all words of the types (a, a, a, a) where $a \in B_q$, and (a, a, b, b) where $a, b \in B_q$ and $a \neq b$. It is easy to see that C is capable of correcting single deletions. Finally we have

$$|C| = q + q(q-1) + \frac{q^2(q-2)}{4} = \frac{q^2(q+2)}{4}.$$

In [2], the author has inductively constructed an SQS(q) for $q \equiv 2$ or 4 (mod 6). By following the construction of Hanani, we will prove that there exists an SQS(q) with the step property for $q \equiv 2$ or 4 (mod 6).

We start with some notions and terminology which will be needed in the sequel. First of all, two partitions of unordered pairs of a set with 2m elements are introduced (see [2]).

Given a set of 2m elements $0, 1, \ldots, 2m - 1$, we decompose the m(2m - 1) pairs $\{r, s\}$ from them into 2m - 1 systems $P_{\alpha}(m)$, $\alpha = 0, 1, \ldots, 2m - 2$, each containing m mutually disjoint pairs.

We shall also need another decomposition of the m(2m-1) pairs from 2m elements into 2m systems $\bar{P}_{\xi}(m)$ ($\xi = 0, 1, \ldots, 2m-1$) such that each of the m systems $\bar{P}_{\eta}(m)$ ($\eta = 0, 1, \ldots, m-1$) should contain m-1 mutually disjoint pairs not containing the elements 2η and $2\eta + 1$, and each of the other m systems should contain m mutually disjoint pairs.

Let $F = \{0, 1, \dots, f-1\}$ be a set of f elements. If a system SQS(f) exists, we say that it is possible to form a quadruple system from F and we write $F \in Q$ or $f \in Q$. Similarly, if a system SQS(f) with the step property exists, we write $F \in S$ or $f \in S$.

If $F \in Q$, we shall in the sequel denote by $\{x, y, z, t\} \subset F$ for any quadruple in F, that is, an element of SQS(f). If $f + 1 \in Q$, then $F \cup \{A\} \in Q$ where Ais an additional element. The quadruples which contain A will be denoted by $\{A, u, v, w\}$.

The following was inductively shown by Hanani[2].

Theorem 3.3. If $q \equiv 2$ or $4 \pmod{6}$, then $q \in Q$.

By using a similar argument as in [2], we will prove the following.

Theorem 3.4. If $q \equiv 2$ or $4 \pmod{6}$, then $q \in S$.

Proof. We will proceed by induction on q. Clearly SQS(4) has the step property, hence $4 \in S$. As an induction step, we will prove the following: Let $q \equiv 2 \text{ or } 4 \pmod{6}$. If $g \in S$ for every g < q satisfying $g \equiv 2 \text{ or } 4 \pmod{6}$, then $q \in S$. The proof will be given separately for each of the following cases which evidently exhaust all the possibilities:

 $\begin{array}{l} q \equiv 4 \text{ or } 8 \pmod{12}, \\ q \equiv 4 \text{ or } 10 \pmod{18}, \\ q \equiv 34 \pmod{36}, \\ q \equiv 26 \pmod{36}, \\ q \equiv 2 \text{ or } 10 \pmod{24} \ (q > 2), \\ q \equiv 14 \text{ or } 38 \pmod{72}. \end{array}$

Case $\mathbf{I} : q \equiv 4 \text{ or } 8 \pmod{12}$.

Put q = 2f where $f \equiv 2$ or 4 (mod 6). Since $f \in S$, there is an SQS(f) with the step property. Without loss of generality, we may assume that $F = \{0, 1, \ldots, f - 1\}$ is the total order on B_q which admits the step property for SQS(f). Let $N = \{(i, j) \mid i = 0, 1 \text{ and } j \in F\}$ and $\{x, y, z, t\}$ be any quadruple in F (i.e. an element of SQS(f)). The following quadruples in N form an SQS(q) (see [2]):

$$Q_1 : \{(a_1, x), (a_2, y), (a_3, z), (a_4, t)\}, a_1 + a_2 + a_3 + a_4 \equiv 0 \pmod{2}, Q_2 : \{(0, j), (0, j'), (1, j), (1, j')\}, j \neq j' (j, j' \in F).$$

Rename the letters of N as

$$(0,j) \to j, (1,j) \to f+j \ (0 \le j \le f-1).$$
 (3)

Notice that the solutions of $a_1 + a_2 + a_3 + a_4 \equiv 0 \pmod{2}$ are

$$\begin{aligned} (a_1, a_2, a_3, a_4) &= (0, 0, 0, 0), (0, 0, 1, 1), (0, 1, 0, 1) \\ (0, 1, 1, 0), (1, 0, 0, 1), (1, 0, 1, 0) \\ (1, 1, 0, 0), (1, 1, 1, 1). \end{aligned}$$

To investigate the step property on SQS(q), we only need to consider the following quadruples in each type: for $0 \le t \le \frac{f}{2} - 1$,

$$\begin{array}{rcl} Q_1 &:& \{(0,2t),\,(0,2t+1),\,(0,x),\,(0,y)\},\,\{(0,2t),\,(0,2t+1),\,(1,x),\,(1,y)\},\\ && \{(1,2t),\,(1,2t+1),\,(0,x),\,(0,y)\},\,\{(1,2t),\,(1,2t+1),\,(1,x),\,(1,y)\},\\ Q_2 &:& \{(0,2t),\,(0,2t+1),\,(1,2t),\,(1,2t+1)\}. \end{array}$$

Note that $\{2t, 2t+1, x, y\}$ is a quadruple in SQS(f). Since SQS(f) satisfies the step property with respect to the natural order on F, it follows from (3) that SQS(q) satisfies the step property. Therefore $q \in S$.

Case II : $q \equiv 4$ or 10 (mod 18).

Put q = 3f + 1 where $f \equiv 1$ or 3 (mod 6). Since $f + 1 \in Q$, we have $f + 1 \in S$ by induction. Hence we may assume that SQS(f+1) satisfies the step property with respect to the natural order on $F = \{0, 1, \ldots, f\}$. In accordance with notations in [2], we denote A = f and $N = \{(i, j), A \mid i = 0, 1, 2 \text{ and } j \in F\}$. And let $\{x, y, z, t\}$ and $\{A, u, v, w\}$ be any quadruples in F and $F \cup \{A\}$, respectively. Note that both of them are elements of SQS(f+1). According to [2], the following quadruples in N form an SQS(q):

 $\begin{array}{rcl} Q_1 &:& \{(a_1,x),(a_2,y),(a_3,z),(a_4,t)\}, \, a_1+a_2+a_3+a_4\equiv 0 \pmod{3},\\ Q_2 &:& \{A,(b_1,u),(b_2,v),(b_3,w)\}, \, b_1+b_2+b_3\equiv 0 \pmod{3},\\ Q_3 &:& \{(i,u),(i,v),(i+1,w),(i+2,w)\},\\ Q_4 &:& \{(i,j),(i,j'),(i+1,j),(i+1,j')\}, \, j\neq j' \ (j,\,j'\in F),\\ Q_5 &:& \{A,(0,j),(1,j),(2,j)\}, \, j\in F. \end{array}$

Rename letters in N as follows: for $0 \le t \le \frac{f-3}{2}$,

$$\begin{cases} (0,2t) \to 6t, & (0,2t+1) \to 6t+1, \\ (1,2t) \to 6t+2, & (1,2t+1) \to 6t+3, \\ (2,2t) \to 6t+4, & (2,2t+1) \to 6t+5 \\ (0,f-1) \to 3f-3, & (1,f-1) \to 3f-2, \\ (2,f-1) \to 3f-1, & A=f \to 3f. \end{cases}$$
(4)

To investigate the step property, we only need to consider the following quadruples of each type: for $0 \le t \le \frac{f-3}{2}$ and $0 \le i \le 3$,

$$\begin{array}{rcl} Q_1 &:& \{(0,2t),(0,2t+1),(0,x),(0,y)\},\{(0,2t),(0,2t+1),(1,x),(2,y)\},\\ && \{(1,2t),(1,2t+1),(0,x),(1,y)\},\{(1,2t),(1,2t+1),(2,x),(2,y)\},\\ && \{(2,2t),(2,2t+1),(0,x),(2,y)\},\{(2,2t),(2,2t+1),(1,x),(1,y)\},\\ Q_2 &:& \{(1,2t),(1,2t+1),(1,x),A\},\{(0,x),(1,y),(2,f-1),A\},\\ Q_3 &:& \{(0,2t),(0,2t+1),(1,x),(2,x)\},\{(1,2t),(1,2t+1),(0,x),(2,x)\},\\ && \{(2,2t),(2,2t+1),(0,x),(1,x)\},\{(2,x),(2,y),(0,f-1),(1,f-1)\},\\ Q_4 &:& \{(i,2t),(i,2t+1),(i+1,2t),(i+1,2t+1)\},\\ Q_5 &:& \{(0,f-1),(1,f-1),(2,f-1),A\}. \end{array}$$

Note that $\{2t, 2t + 1, a, b\}$, $\{2t, 2t + 1, a, A\}$, and $\{a, b, f - 1, A\}$ are quadruples in SQS(f + 1). Since SQS(f + 1) satisfies the step property with respect to the natural order on $F \cup \{A\}$, it follows from (4) that SQS(q) satisfies the step property. Therefore $q \in S$.

Case III : $q \equiv 34 \pmod{36}$.

Put q = 3f + 4 where $f \equiv 10 \pmod{12}$, and denote f = 12k + 10. By induction hypothesis, we may assume that $f + 4 \in S$. In accordance with notations in [2], we can assume that there is an SQS(f + 4) on the letters

$$\hat{F} = \{L_0, L_1, \cdots, L_{f-1}, (A, 0), (A, 1), (A, 2), (A, 3)\}$$

where $L_t = 2t$, $L_{\frac{f}{2}+t} = 2t + 1$ $(0 \le t \le \frac{f}{2} - 1)$. W also assume that given SQS(f+4) satisfies the step property with respect to the order

$$L_0 < L_1 < \dots < L_{f-1} < (A,0) < (A,1) < (A,2) < (A,3).$$

For simplicity, put $F = \{L_0, L_1, \dots, L_{f-1}\}$. Let $N = \{(i, j), (A, h) \mid i = 0, 1, 2; j \in F; h = 0, 1, 2, 3\}$ and $\{x, y, z, t\}$ be any quadruple in \tilde{F} . The following quadruples in N form an SQS(q) (see [2]):

 $\begin{array}{rcl} Q_1 &:& \{(A,0),(A,1),(A,2),(A,3)\},\\ Q_2 &:& \{(i,x),(i,y),(i,z),(i,t)\}, \, \text{the quadruple in } Q_1 \, \text{excluded},\\ Q_3 &:& \{(A,a_1),(0,a_2),(1,a_3),(2,a_4)\}, \, a_1+a_2+a_3+a_4\equiv 0 \, (\text{mod } f),\\ Q_4 &:& \{(i+2,b_3),(i,b_1+2k+1+i(4k+2)-d),\\ && (i,b_1+2k+2+i(4k+2)+d),(i+1,b_2)\}\\ && \text{where } b_1+b_2+b_3\equiv 0 \, (\text{mod } f) \, \text{and } 0\leq d\leq 2k,\\ Q_5 &:& \{(i,r_\alpha),(i,s_\alpha),(i+1,r_{\alpha'}),(i+1,s_{\alpha'})\},\\ && \{r_\alpha,s_\alpha\} \, \text{and } \{r_{\alpha'},s_{\alpha'}\} \, \text{are (equal or different) pairs in } P_\alpha(6k+5),\\ && \alpha=4k+2,4k+3,\cdots 12k+8. \end{array}$

Note that if a quadruple in Q_2 contains an element of (i, (A, h)), then we simply denote it by (A, h). Hence quadruples in Q_2 are from the following sets,

$$\{(i, L_j) \mid 0 \le j \le f - 1\} \cup \{(A, h) \mid 0 \le h \le 3\} \ (0 \le i \le 2).$$

Now rename letters in N as follows:

$$\begin{cases} (i, L_t) \to if + t \ (0 \le i \le 2, \ 0 \le t \le f - 1), \\ (A, h) \to 3f + h \ (0 \le h \le 4). \end{cases}$$
(5)

To investigate the step property, we will first consider the following quadruples of each type except Q_5 : for $0 \le t \le \frac{f}{2} - 1$, $0 \le i \le 3$, and $0 \le h \le 1$,

$$\begin{array}{rcl} Q_1 & : & \{(A,0),(A,1),(A,2),(A,3)\}, \\ Q_2 & : & \{(i,L_{2t}),(i,L_{2t+1}),(i,x),(i,y)\}, \{(i,x'),(i,y'),(A,2h),(A,2h+1)\}, \\ Q_3,\,Q_4 & : & \text{no quadruples to consider,} \end{array}$$

Note that $\{(A, 0), (A, 1), (A, 2), (A, 3)\}$, $\{L_{2t}, L_{2t+1}, x, y\}$, and $\{x', y', (A, 2h), (A, 2h+1)\}$ are quadruples in SQS(f+4). Since SQS(f+4) satisfies the step property with respect to the order above on \tilde{F} , it can be easily checked by (5) that the quadruples above satisfy the step property.

In Q_5 , the followings should be considered: for $0 \le i \le 3$

 $\{(i, L_{2t}), (i, L_{2t+1}), (i+1, x), (i+1, y)\}, \{(i, x'), (i, y'), (i+1, L_{2t}), (i+1, L_{2t+1})\}, (i+1, x), (i+1, y)\}, \{(i, x'), (i, y'), (i+1, L_{2t}), (i+1, L_{2t+1})\}, (i+1, y)\}$

where $\{2t, 2t+1\}$ and $\{x, y\}$ are pairs in $P_{\alpha}(6k+5), \alpha = 4k+2, 4k+3, \ldots, 12k+8$. These quadruples also satisfy the step property. Therefore $q \in S$.

Case IV : $q \equiv 26 \pmod{36}$.

This case is similar to Case III. Put q = 3f + 2 where $f \equiv 8 \pmod{12}$, and denote f = 12k + 8. By induction hypothesis, we have $f + 2 \in S$. We may assume that there is an SQS(f + 2) on the alphabet

$$\tilde{F} = \{L_0, L_1, \cdots, L_{f-1}, (A, 0), (A, 1)\}$$

where $L_t = 2t$, $L_{\frac{f}{2}+t} = 2t + 1$ $(0 \le t \le \frac{f}{2} - 1)$. We also assume that given SQS(f+2) satisfies the step property with respect to the order

$$L_0 < L_1 < \dots < L_{f-1} < (A, 0) < (A, 1).$$

For simplicity, put $F = \{L_0, L_1, \dots, L_{f-1}\}$. Let $N = \{(i, j), (A, h) \mid i = 0, 1, 2; j \in F; h = 0, 1\}$ and $\{x, y, z, t\}$ be any quadruple in \tilde{F} . The following quadruples in N form an SQS(q) (see [2]):

$$\begin{array}{rcl} Q_1 &:& \{(i,x),(i,y),(i,z),(i,t)\},\\ Q_2 &:& \{(A,a_1),(0,a_2),(1,a_3),(2,a_4)\},\,a_1+a_2+a_3+a_4\equiv 0 \pmod{f},\\ Q_3 &:& \{(i+2,b_3),(i,b_1+2k+1+i(4k+2)-d),\\ && (i,b_1+2k+2+i(4k+2)+d),(i+1,b_2)\},\\ && \text{where } b_1+b_2+b_3\equiv 0 \pmod{f} \text{ and } 0\leq d\leq 2k,\\ Q_4 &:& \{(i,r_\alpha),(i,s_\alpha),(i+1,r_{\alpha'}),(i+1,s_{\alpha'})\},\\ && \{r_\alpha,s_\alpha\} \text{ and } \{r_{\alpha'},s_{\alpha'}\} \text{ are pairs in } P_\alpha(6k+4),\\ && \alpha=4k+2,4k+3,\cdots,12k+6. \end{array}$$

Note that if a quadruple in Q_1 contains an element of (i, (A, h)), then we simply denote it by (A, h).

Now we rename letters in N as follows:

$$\begin{cases} (i, L_t) \to if + t \ (0 \le i \le 2, 0 \le t \le f - 1), \\ (A, h) \to 3f + h \ (0 \le h \le 1). \end{cases}$$
(6)

To investigate the step property, we will first consider the following quadruples of each type except Q_4 : for $0 \le t \le \frac{f}{2} - 1$ and $0 \le i \le 2$,

$$\begin{array}{rcl} Q_1 & : & \{(i,L_{2t}),(i,L_{2t+1}),(i,a),(i,b)\}, \{(i,a'),(i,b'),(A,0),(A,1)\},\\ Q_2,Q_3 & : & \text{no quadruples to consider}, \end{array}$$

Note that $\{L_{2t}, L_{2t+1}, a, b\}$ and $\{a', b', (A, 0), (A, 1)\}$ are quadruples in SQS(f + 1). Since SQS(f + 2) satisfies the step property with respect to the order above on \tilde{F} , it can be easily checked by (6) that the quadruples above satisfy the step property.

In case of Q_4 , the followings should be considered: for $0 \le i \le 2$,

$$\{(i, L_{2t}), (i, L_{2t+1}), (i+1, a), (i+1, b)\}, \{(i, a), (i, b), (i+1, L_{2t}), (i+1, L_{2t+1})\}, \{(i, L_{2t}), (i, L_{2t+1}), (i+1, L_{2t+1})\}, \{(i, L_{2t+1}), (i, L_$$

where $\{L_{2t}, L_{2t+1}\}$ and $\{a, b\}$ are pairs in $P_{\alpha}(6k+4), \alpha = 4k+2, 4k+3, \ldots, 12k+6$. These quadruples also satisfy the step property. Therefore $q \in S$.

Case $\mathbf{V}: q \equiv 2 \text{ or } 10 \pmod{24}, q > 2.$

Put q = 4f + 2 where $f \equiv 0$ or 2 (mod 6) (f > 0), and denote f = 2k. By induction hypothesis, we have $f + 2 \in S$. Hence we may assume that SQS(f+2)satisfies property with respect to the natural order on $F = \{0, 1, \ldots, f + 1\}$. In accordance with notations in [2], denote (A, l) = f + l (l = 0, 1). Note that $\tilde{F} = F \cup \{(A, l) : h = 0, 1\} \in Q$. Let $N = \{(h, i, j), (A, l) \mid h = 0, 1; i = 0, 1; j \in$ $F; l = 0, 1\}$ and $\{x, y, z, t\}$ be any quadruple in \tilde{F} . The following quadruples in N form an SQS(q) ([2]):

Q_1	:	$\{(h,i,x),(h,i,y),(h,i,z),(h,i,t)\},$
Q_2	:	$\{(A, l), (0, 0, 2c_1), (0, 1, 2c_2 - \epsilon), (1, \epsilon, 2c_3 + l)\},\$
		where $c_1 + c_2 + c_3 \equiv 0 \pmod{k}$ and $0 \le \epsilon \le 1$,
Q_3	:	$\{(A, l), (0, 0, 2c_1 + 1), (0, 1, 2c_2 - 1 - \epsilon), (1, \epsilon, 2c_3 + 1 - l)\},\$
Q_4	:	$\{(A, l), (1, 0, 2c_1), (1, 1, 2c_2 - \epsilon), (0, \epsilon, 2c_3 + 1 - l)\},\$
Q_5	:	$\{(A, l), (1, 0, 2c_1 + 1), (1, 1, 2c_2 - 1 - \epsilon), (0, \epsilon, 2c_3 + l)\},\$
Q_6	:	$\{(h, 0, 2c_1 + \epsilon), (h, 1, 2c_2 - \epsilon), (h + 1, 0, \bar{r}_{c_3}), (h + 1, 0, \bar{s}_{s_3})\},\$
		$\{\bar{r}_{c_3}, \bar{s}_{c_3}\}$ are pairs $\ln \bar{P}_{c_3}(k) \ (0 \le c_3 \le k - 1),$
Q_7	:	$\{(h, 0, 2c_1 - 1 + \epsilon), (h, 1, 2c_2 - \epsilon), (h + 1, 1, \bar{r}_{c_3}), (h + 1, 1, \bar{s}_{c_3})\},\$
Q_8	:	$\{(h, 0, 2c_1 + \epsilon), (h, 1, 2c_2 - \epsilon), (h + 1, 1, \bar{r}_{k+c_3}), (h + 1, 1, \bar{s}_{k+c_3})\},\$
Q_9	:	$\{(h, 0, 2c_1 - 1 + \epsilon), (h, 1, 2c_2 - \epsilon), (h + 1, 0, \bar{r}_{k+c_3}), (h + 1, 0, \bar{s}_{k+c_3})\},\$
Q_{10}	:	$\{(h, 0, r_{\alpha}), (h, 0, s_{\alpha}), (h, 1, r_{\alpha'}), (h, 1, s_{\alpha'})\},\$
		$\{r_{\alpha}, s_{\alpha}\}$ and $\{r_{\alpha'}, s_{\alpha'}\}$ are pairs in $P_{\alpha}(k)$ $(0 \le \alpha \le f - 2)$.

Note that if a quadruple in Q_1 contains an element of the form (h, i, (A, l)), then we simply denote it by (A, l).

Now rename letters in N as follows: for $0 \le j \le f - 1$,

$$\begin{cases} (0,0,j) \to j, & (0,1,j) \to f+j, \\ (1,0,j) \to 2f+j, & (1,1,j) \to 3f+j, \\ (A,0) = f \to 4f, & (A,1) = f+1 \to 4f+1. \end{cases}$$
(7)

To investigate the step property, the following quadruples of each type should be firstly considered, except $Q_6 - Q_{10}$: for $0 \le t \le \frac{f}{2} - 1$ and $0 \le h$, $i \le 1$,

 $\begin{array}{rcl} Q_1 & : & \{(h,i,2t),(h,i,2t+1),(h,i,a),(h,i,b)\}, \,\{(A,0),(A,1),(h,i,a),(h,i,b)\}, \\ Q_k & : & \text{no quadruples to consider for } k=2,3,4,5. \end{array}$

Note that $\{2t, 2t+1, a, b\}$ and $\{(A, 0), (A, 1), a, b\}$ are quadruples in SQS(f+2). Since SQS(f+2) satisfies the step property with respect to the natural order on \tilde{F} , it can be easily checked by (7) that the quadruples above satisfy the step property.

In Q_6 , the followings are to be considered: for $0 \le h \le 1$ and $0 \le \epsilon \le 1$

$$\{(h, 0, 2c_1 + \epsilon), (h, 1, 2c_2 - \epsilon), (h + 1, 0, 2t), (h + 1, 0, 2t + 1)\}.$$

These quadruples also satisfy the step property. The cases $Q_7 - Q_{10}$ are similar to Q_6 . As a result, $q \in S$.

Case VI : $q \equiv 14 \text{ or } 38 \pmod{72}$.

Put q = 12f + 2 where $f \equiv 1$ or 3 (mod 6). As a first step, we will prove that 14, 38 $\in Q$. According to [1], we can construct an SQS(14), which is shown in Table 4. Following the notations in [2], we denote $F = \{0, 1, ..., 11\}$ and

0,1,2,4	13, 9, 12, 11	$12,\!1,\!4,\!3$	12,8,5,1	5, 6, 10, 11	$3,\!5,\!13,\!8$	4,5,7,8
7,10,13,12	$0,\!8,\!9,\!11$	$13,\!2,\!5,\!4$	$13,\!9,\!6,\!2$	6,0,11,12	$4,\!6,\!7,\!9$	$9,\!12,\!4,\!0$
1,2,3,5	$7,\!3,\!6,\!5$	$3,\!4,\!10,\!11$	$7,\!10,\!0,\!3$	$0,\!1,\!12,\!13$	5,0,8,10	$10,\!13,\!5,\!1$
2,3,4,6	$1,\!9,\!10,\!12$	$9,\!12,\!2,\!5$	$8,\!11,\!1,\!4$	1,2,13,7	6,1,9,11	11,7,6,2
3,4,5,0	$2,\!10,\!11,\!13$	6,1,13,8	0,2,7,9	2,3,7,8	$0,\!2,\!10,\!12$	$12,\!8,\!0,\!3$
4,5,6,1	$3,\!11,\!12,\!7$	$4,\!5,\!11,\!12$	$1,\!3,\!8,\!10$	$3,\!4,\!8,\!9$	$5,\!6,\!8,\!9$	$13,\!9,\!1,\!4$
$5,\!6,\!0,\!2$	$4,\!12,\!13,\!8$	$5,\!6,\!12,\!13$	$2,\!4,\!9,\!11$	$13,\!9,\!0,\!3$	$8,\!11,\!3,\!6$	$7,\!10,\!2,\!5$
6,0,1,3	$5,\!13,\!7,\!9$	6,0,13,7	$3,\!5,\!10,\!12$	$7,\!10,\!1,\!4$	$3,\!5,\!9,\!11$	$4,\!6,\!10,\!12$
$8,\!11,\!7,\!13$	6,7,8,10	$0,\!1,\!7,\!8$	$4,\!6,\!11,\!13$	8,11,2,5	6,0,9,10	5,0,11,13
$9,\!12,\!8,\!7$	$8,\!4,\!0,\!6$	1,2,8,9	5,0,12,7	$9,\!12,\!3,\!6$	0,1,10,11	6,1,12,7
$10,\!13,\!9,\!8$	$9,\!5,\!1,\!0$	$2,\!3,\!9,\!10$	4,5,9,10	$10,\!13,\!4,\!0$	$1,\!2,\!11,\!12$	0,2,13,8
11,7,10,9	$10,\!6,\!2,\!1$	$10,\!13,\!3,\!6$	$12,\!8,\!6,\!2$	11,7,5,1	$2,\!3,\!12,\!13$	$1,\!3,\!7,\!9$
12,8,11,10	11,0,3,2	11,7,4,0	$1,\!3,\!11,\!13$	$2,\!4,\!12,\!7$	$3,\!4,\!13,\!7$	$2,\!4,\!8,\!10$

Table 4: SQS(14)

 $(A,h) = 11 + h \ (h = 0,1).$ Note that $F \cup \{(A,h) \mid h = 0,1\} \in Q$. Letting $N' = \{(i,j), (A,h) \mid i = 0,1,2; j \in F; h = 0,1\}$ be a set of 38 elements, we will show that $N' \in Q$. Let $\{x', y', z', t'\}$ be a quadruple in $F \cup \{(A,h) \mid h = 0,1\} = \{j, (A,h) \mid j \in F; h = 0,1\}$. The following quadruples in N' form an SQS(38) [2]:

Q_1	:	$\{(i, x'), (i, y'), (i, z'), (i, t')\},\$
Q_2	:	$\{(A, h), (0, b_1), (1, b_2), (2, b_3 + 3h)\},\$
		$b_1 + b_2 + b_3 \equiv 0 \pmod{12}, \ h = 0, 1,$
Q_3	:	$\{(i, b_1 + 4 + i), (i, b_1 + 7 + i), (i + 1, b_2), (i + 2, b_3)\},\$
Q_4	:	$\{(i,j), (i+1,j+6\epsilon), (i+2,6\epsilon-2j+1), (i+2,6\epsilon-2j-1)\}, \epsilon = 0, 1,$
Q_5	:	$\{(i,j), (i+1,j+6\epsilon), (i+2,6\epsilon-2j+2), (i+2,6\epsilon-2j-2)\},\$
Q_6	:	$\{(i,j), (i+1, j+6\epsilon-3), (i+2, 6\epsilon-2j+1), (i+2, 6\epsilon-2j+2)\},\$
Q_7	:	$\{(i,j), (i+1, j+6\epsilon+3), (i+2, 6\epsilon-2j-1), (i+2, 6\epsilon-2j-2)\},\$
Q_8	:	$\{(i,j), (i,j+6), (i+1,j+3\epsilon), (i+1,j+6+3\epsilon)\},\$
Q_9	:	$\{(i, 2g+3\epsilon), (i, 2g+6+3\epsilon), (i', 2g+1), (i', 2g+5)\},\$
		$i' \neq i, g = 0, 1, 2, 3, 4, 5,$
Q_{10}	:	$\{(i, 2g+3\epsilon), (i, 2g+6+3\epsilon), (i', 2g+2), (i', 2g+4)\},\$
Q_{11}	:	$\{(i,j),(i,j+1),(i+1,j+3e),(i+1,j+3e+1)\}, e = 0,1,2,3,$
Q_{12}	:	$\{(i,j), (i,j+2), (i+1,j+3e), (i+1,j+3e+2)\},\$
Q_{13}	:	$\{(i,j), (i,j+4), (i+1,j+3e), (i+1,j+3e+4)\},\$
Q_{14}	:	$\{(i,r_{\alpha}),(i,s_{\alpha}),(i',r_{\alpha'}),(i',s_{\alpha'})\},$
		$\{r_{\alpha}, s_{\alpha}\}$ and $\{r_{\alpha'}, s_{\alpha'}\}$ are pairs in $P_{\alpha}(6), \alpha = 4, 5.$

Now we construct SQS(q) for $q \equiv 14$ or 38 (mod 72), that is, $f \equiv 1$ or 3 (mod 6). Since $f \equiv 1$ or 3 (mod 6), we can assume that $f+1 \in Q$. Let $N = \{(i, j), (A, h) \mid i = 0, 1, \ldots, f-1; j = 0, 1, \ldots, 11; h = 0, 1\}$ be a set of 12f + 2 elements. The following quadruples in N form an SQS(q) ([2]):

$$R_{1} : \{(i, x'), (i, y'), (i, z'), (i, t')\}, \\R_{2} : \begin{cases} \{(A, h), (u, b_{1}), (v, b_{2}), (w, b_{3} + 3h)\}, b_{1} + b_{2} + b_{3} \equiv 0 \pmod{12} \\ \{(u, \alpha_{1}), (v, \alpha_{2}), (w, \alpha_{3}), (w, \alpha_{4})\}, \\ \{(i, \beta_{1}), (i, \beta_{2}), (i', \beta_{3}), (i', \beta_{4})\}, i' \neq i \ (i', i \in \{u, v, w\}). \end{cases}$$

In this case, α_{ν} and β_{ν} $(1 \leq \nu \leq 4)$ are to be replaced by the second indices of $Q_3 - Q_{14}$, corresponding to the first indices 0, 1, 2 for u, v, w, respectively. It should be noted that i and i' define uniquely a $\{u, v, w\}$ in which they are contained. Therefore they may be considered as two indices from $\{u, v, w\}$.

$$R_3 : \{(x, a_1), (y, a_2), (z, a_3), (t, a_4)\}, a_1 + a_2 + a_3 + a_4 \equiv 0 \pmod{12}.$$

From now on, we will show that $q \in S$ for $q \equiv 14$ or 38 (mod 72). In case of q = 14, rename letters of SQS(14) in Table 4 as follows:

$$\begin{cases} t \to L_{2t} \text{ for } 0 \le t \le 5, \\ t \to L_{2t-13} \text{ for } 7 \le t \le 12, \\ 6 + 7i \to L_{i+12} = (A, i) \text{ for } 0 \le i \le 1. \end{cases}$$

And we give an order on $\{L_0, L_1, \ldots, L_{13}\}$ as $L_0 < L_1 < \cdots < L_{13}$. Then the following quadruples satisfy the step property:

	$\{0,7,1,8\},$	$\{0,7,2,9\},\$	$\{0,7,3,10\},\$	$\{0,7,4,11\},$	$\{0,7,5,12\},\$	$\{0,7,6,13\},\$
$\{1,\!8,\!0,\!7\},$		$\{1, 8, 2, 9\},\$	$\{1, 8, 3, 10\},\$	$\{1, 8, 4, 11\},\$	$\{1, 8, 5, 12\},\$	$\{1, 8, 6, 13\},\$
$\{2,9,0,7\},\$	$\{2,9,1,8\},\$		$\{2,9,3,10\},\$	$\{2,9,4,11\},\$	$\{2,9,5,12\},\$	$\{2, 9, 6, 13\},\$
$\{3,10,0,7\},\$	$\{3,10,1,8\},\$	$\{3,10,2,9\},\$		$\{3,10,4,11\},\$	$\{3,10,5,12\},\$	$\{3,10,6,13\},\$
$\{4,11,0,7\},\$	$\{4,11,1,8\},\$	$\{4,11,2,9\},\$	$\{4,11,3,10\},\$		$\{4,11,5,12\},\$	$\{4,11,6,13\},\$
$\{5,12,0,7\},\$	$\{5,12,1,8\},\$	$\{5,12,2,9\},\$	$\{5,12,3,10\},\$	$\{5,12,4,11\},\$		$\{5,12,6,13\},\$
$\{6,13,0,7\},$	$\{6,13,1,8\},\$	$\{6, 13, 2, 9\},\$	$\{6,13,3,10\},\$	$\{6, 13, 4, 11\},\$	$\{6, 13, 5, 12\},\$	

This shows that SQS(14) satisfies the step property. Therefore $14 \in S$.

In case of q = 38, we rename letters of N' as follows:

$$\begin{cases} (i, L_t) \to 12i + t & (0 \le i \le 2, 0 \le t \le 11), \\ (A, 0) \to 36, & (A, 1) \to 37. \end{cases}$$
(8)

To investigate the step property of SQS(38), we will consider quadruples of each type.

In case of $Q_1,$ the following types of quadruples should be considered: for $0\leq i\leq 2$

$$\{(i, L_{2t}), (i, L_{2t+1}), (i, a), (i, b)\}, \{(i, a), (i, b), (A, 0), (A, 1)\}$$

Since $\{L_{2t}, L_{2t+1}, a, b\}$ and $\{a, b, (A, 0), (A, 1)\}$ in SQS(14) satisfy the step property, these quadruples also satisfy the step property by (8).

In the cases $Q_2 - Q_{13}$, there is no quadruple to consider.

In Q_{14} , the following quadruples should be considered:

$$\{(i, L_{2t}), (i, L_{2t+1}), (i', a), (i', b)\}$$

By (8), these quadruples satisfy the step property. Therefore $38 \in S$.

Finally, consider general cases. At first, rename letters in N as follows:

$$\begin{cases} (i, L_t) \to 12i + t & (0 \le i \le f - 1, \ 0 \le t \le 11), \\ (A, 0) = 12f, & (A, 1) = 12f + 1. \end{cases}$$
(9)

From now on, we will investigate the step property of quadruples of each type.

 $\begin{array}{l} R_1 \} \quad \{(i, L_{2t}), (i, L_{2t+1}), (i, a), (i, b)\}, \ \{(i, a), (i, b), (A, 0), (A, 1)\}.\\ \text{Since } \{L_{2t}, L_{2t+1}, a, b\} \text{ and } \{a, b, (A, 0), (A, 1)\} \text{ satisfy the step property, these} \end{array}$

quadruples also satisfy the step property by (9).

 $R_2) \quad \{(i, L_{2t}), (i, L_{2t+1}), (i', a), (i', b)\}.$

By the step property of constructed SQS(38), these quadruples satisfy the step property by (9).

 R_3) In this case, there is no quadruple to consider.

Therefore given SQS(q) satisfies the step property. This shows $q \in S$.

If we combine Theorems 3.2 and 3.4, we can deduce the following theorem.

Theorem 3.5. If $q \equiv 2, 4 \pmod{6}$, then $N(4, q, 1) = \frac{q^2(q+2)}{4}$.

Remark. The authors of [9] informed us that every Steiner quadruple system in [9] satisfies the step property. It is not known whether there is a Steiner quadruple system which does not satisfy the step property.

4 Construction of optimal codes in B_q^4 for $q \equiv 0 \pmod{6}$

In this section, we will construct a code in B_q^4 for $q \equiv 0 \pmod{6}$ capable of correcting single deletions whose cardinality meets the upper bound in Theorem 2.2. Since there dose not exist a Steiner quadruple system for any alphabet of this size, we will use a group divisible system. We divide our construction into two steps. In the first, we prove that there is an optimal code over an alphabet of size q = 6m, where m is odd. In the next, we prove that there is an optimal code over an alphabet of size 2q under the assumption that there is an optimal code over an alphabet of size q.

We begin with the definition of a group divisible system. By an r-subset of a set X, we mean a subset of X with r elements.

Definition 4.1 ([7]). Let m and r be positive integers. Let \mathcal{T} be a collection $\{T_1, T_2, \ldots, T_m\}$ of disjoint sets with r element whose union is S. An (m, r, k, b) group divisible system or G(m, r, k, b) system on \mathcal{T} is a collection $\mathcal{B} = \{K_1, K_2, \ldots, K_u\}$ of k-subsets of S such that every b-subset in S is either contained in T_i for some $i = 1, 2, \ldots, m$ or it is contained in a unique k-subset in \mathcal{B} but not both.

By a simple counting argument, one can easily show that

$$|G(m,r,k,b)| = \frac{\binom{mr}{b} - m\binom{r}{b}}{\binom{k}{b}}.$$

The following result was proved by Mills in [7].

Theorem 4.2. A G(m, 6, 4, 3) system exists for every positive integer m.

Since we will use a subfamily \mathcal{A}_1 of a system G(m, 6, 4, 3) in our construction of an optimal code, we briefly review the proof of Theorem 4.2 in [7], which introduces \mathcal{A}_1 when m is odd. Let m be odd. For $i \in I_m = \{1, 2, \ldots, m\}$, let T_i be the set of ordered pairs (i, α) with $\alpha \in Z_6$. Let S be the union of these T_i . We partition the elements of Z_6 into pairs in three ways:

$$\begin{cases} P_1 = \{(0,3), (1,5), (2,4)\}, \\ P_2 = \{(1,4), (2,0), (3,5)\}, \\ P_3 = \{(2,5), (3,1), (4,0)\}. \end{cases}$$

For any pair w, x (w < x) of distinct elements of I_m and any λ in $I_3 = \{1, 2, 3\}$, we form the nine quadruples

$$(w, \alpha), (w, \beta), (x, \gamma), (x, \delta)$$

where $(\alpha, \beta), (\gamma, \delta) \in P_{\lambda}$. This gives us a collection of $\frac{27m(m-1)}{2}$ quadruples, denoted by \mathcal{A} , which is contained in G(m, 6, 4, 3). Among the quadruples in \mathcal{A} , choose quadruples of the following types: for any pair w, x of distinct elements of I_m

$$(w, \alpha), (w, \beta), (x, \gamma), (x, \delta),$$

where $(\alpha, \beta), (\gamma, \delta) \in P_1$. We denote these quadruples by \mathcal{A}_1 . Note that $|\mathcal{A}_1| = 9\binom{m}{2}$.

Theorem 4.3. $N(4,q,1) = \frac{q^2(q+2)}{4}$ for q = 6m and m is odd.

Proof. We will keep the notations of the preceding discussion throughout the proof. Let m be odd, and consider G(m, 6, 4, 3) on $\mathcal{T} = \{T_1, T_2, \ldots, T_m\}$. Put $S = \bigcup_{i=1}^{m} T_i$. Recall that our goal is to construct an optimal code in B_q^4 where q = |S|. Define an order relation on S as follows:

$$\begin{cases} (i, \alpha) < (j, \alpha) \text{ if } i < j, \\ (i, \alpha) < (j, \beta) \text{ if } \alpha < \beta. \end{cases}$$

To each quadruple $\{x, y, z, w\}$ in G(m, 6, 4, 3), we associate a word (a_1, a_2, a_3, a_4) of length 4 such that $a_1 < a_2 < a_3 < a_4$ and $a_i \in \{x, y, z, w\}$. From now on, we consider the quadruples in G(m, 6, 4, 3) as words of length 4 defined above.

Put

$$N = \bigcup_{x \in \mathcal{A}_1} \langle x \rangle_{B^4_q} \bigcup \bigcup_{x \in G(m, 6, 4, 3) \setminus \mathcal{A}_1} \langle x \rangle_{A^4_q} .$$

Note that for any two distinct elements $a = (a_1, a_2, a_3, a_4)$, and $b = (b_1, b_2, b_3, b_4)$ of \mathcal{A}_1 , if $|T(a) \cap T(b)| = 2$, then either $\{a_1, a_2\} = \{b_1, b_2\}$ or $\{a_3, a_4\} = \{b_3, b_4\}$. From the structure of \mathcal{A}_1 , the definition of G(m, 6, 4, 3), and Lemma 2.3, we can deduce that N is a code of B_q^4 capable of correcting single deletions. Since $|\mathcal{A}_1| = 9\binom{m}{2}$ and $|G(m, 6, 4, 3)| = \frac{6m(6m-1)(6m-2)}{24} - 5m$, it follows that

$$|N| = 8|\mathcal{A}_1| + 6|G(m, 6, 4, 3) - \mathcal{A}_1| = 54m^3 - 18m^2 - 36m.$$

Since N(4, 6, 1) = 72 (Table (2)) and $|T_i| = 6$ for each *i*, we can choose an optimal code C_i over T_i of cardinality 72 capable of correcting single deletions. Put

$$L = \bigcup_{i=1}^{m} C_i \bigcup N \bigcup \{ (a, a, b, b) \mid a \in T_i, b \in T_j, i \neq j \}.$$

It is easy to check that L is a code of B_q^4 of capable of correcting single deletions. It is clear from a simple calculation that

$$|L| = m|C_1| + |N| + 2 \times 36\binom{m}{2}$$

= 72m + 54m³ - 18m² - 36m + 36m(m - 1)
= 54m³ + 18m² = $\frac{q^2(q+2)}{4}$.

This proves the theorem.

The following has been known since the nineteenth century [8]. Since we need some terminology in our construction of optimal codes, we borrow a sketch of proof from [3].

Lemma 4.4 (Reiss). The n(2n-1) pairs of 2n elements can be partitioned into 2n-1 sets $S_1, S_2, \ldots, S_{2n-1}$ such that each set contains n disjoint pairs.

Proof. Put $l_{ij} = i + j - 1 \pmod{2n - 1}$ where $1 \le i, j, l_{ij} \le 2n - 1$, and define $l_{i,2n} = l_{ii}, i = 1, 2, \dots, 2n - 1$. Take

$$S_q = \{(i,j) \mid i < j \text{ and } l_{ij} = q\}, \ q = 1, 2, \dots, 2n - 1.$$

Then these sets S_q (q = 1, 2, ..., 2n - 1) have the desired property.

Proposition 4.5. If $N(4,q,1) = \frac{q^2(q+2)}{4}$ for q = 6m, then $N(4,2q,1) = \frac{(2q)^2(2q+2)}{4}$.

Proof. Suppose that $N(4, q, 1) = \frac{q^2(q+2)}{4}$ for q = 6m. Let $A = \{1, 2, \dots, q\}$ and put $A_{\alpha} = \{(a, \alpha) \mid 1 \le a \le q\}$ for $\alpha = 1, 2$. We will construct a code capable of correcting single deletions over $A_1 \cup A_2$ of cardinality $\frac{(2q)^2(2q+2)}{4}$. By applying Lemma 4.4 with n = 3m, the pairs of elements of A_{α} can be partitioned into the sets $S_1^{\alpha}, S_2^{\alpha}, \dots, S_{6m-1}^{\alpha}$, namely

the set of pairs of $A_{\alpha} = S_1^{\alpha} \cup S_2^{\alpha} \cup \cdots \cup S_{6m-1}^{\alpha}, \quad \alpha = 1, 2,$

where $S_l^{\alpha} = \{((a, \alpha), (b, \alpha)) \mid a < b, a + b - 1 = l \pmod{6m - 1}\}$ for $\alpha = 1, 2$ and $l = 1, 2, \dots 6m - 1$. Note that $|S_l^{\alpha}| = 3m = \frac{q}{2}$ for α and l. Let us consider the set T of ordered pairs defined as follows:

$$T = \left\{ (x, y) \mid x \in S_i^1, y \in S_i^2, 1 \le i \le 6m - 1 \right\} = \bigcup_{i=1}^{6m-1} S_i^1 \times S_i^2.$$

Since x and y are pairs of elements in A_{α} ($\alpha = 1, 2$), T is a set of quadruples of elements in $A_1 \cup A_2$ whose cardinality is $(6m - 1)|S_1^1||S_1^2| = 9m^2(6m - 1) = 9m^2(6m - 1)$

 $\frac{q^2(q-1)}{2}$. Furthermore, T has the following property: every triple (c_1, i) , (c_2, j) , and (c_3, k) with elements from $A_1 \cup A_2$, except triples with element from only one of A_1 or A_2 , belong to a unique quadruple in T.

Define an order relation on $A_1 \cup A_2$ as follows:

$$\begin{cases} (a, \alpha) < (b, \alpha) \text{ if and only if } a < b, \\ (a, 1) < (b, 2) \text{ for all } a, b. \end{cases}$$

To each quadruple $x = \{(a, 1), (b, 1), (c, 2), (d, 2)\}$ in T where $((a, 1), (b, 1)) \in S_i^1$ and $((c, 2), (d, 2)) \in S_i^2$, we associate a word (a_1, a_2, a_3, a_4) of length 4 such that $a_1 < a_2 < a_3 < a_4$ and $a_i \in \{(a, 1), (b, 1), (c, 2), (d, 2)\}$. From now on, we consider the quadruples in T as words of length 4 over $A_1 \cup A_2$ defined as above. Put

$$N = \bigcup_{x \in S_1^1 \times S_1^2} < x >_{B_{2q}^4} \bigcup \bigcup_{x \in T \setminus (S_1^1 \times S_1^2)} < x >_{A_{2q}^4}.$$

From the construction of T and Lemma 2.3, it is easy to check that N is a code of B_{2q}^4 capable of correcting single deletions. Note that $|N| = 8|S_1^1||S_1^2| + 6(|T| - |S_1^1||S_1^2|) = \frac{2q^2(3q-2)}{4}$. Since $N(4,q,1) = \frac{q^2(q+2)}{4}$ for q = 6m, there exists a code C^{α} over $A_{\alpha}(\alpha = 1,2)$ of B_q^4 capable of correcting single deletions of cardinality $|C^{\alpha}| = N(4,q,1) = \frac{q^2(q+2)}{4}$.

Finally, put

$$L = C^{1} \cup C^{2} \cup N \cup \{(a, a, b, b) \mid a \in A_{1}, b \in A_{2}\} \cup \{(b, b, a, a) \mid a \in A_{1}, b \in A_{2}\}.$$

It is easy to check that L is a code of B_{2q}^4 capable of correcting single deletions with $|L| = \frac{(2q)^2(2q+2)}{4}$. This proves the theorem.

Combining Theorem 4.3 and Proposition 4.5, we obtain the following theorem.

Theorem 4.6. There exists a code C in B_q^4 capable of correcting single deletions with $|C| = N(4, q, 1) = \frac{1}{4}q^2(q+2)$, for q = 6m, m = 1, 2, 3, ...

By combining Theorems 3.5 and 4.6, we finally obtain the following.

Theorem 4.7. For an even q, $N(4, q, 1) = \frac{q^2(q+2)}{4}$.

5 Perfect codes

In this section, we modify our construction of optimal codes slightly, and construct an optimal perfect code in B_q^4 when q is even.

We start with simple definitions. Recall that a code $C \subseteq B_q^n$ is an *s*-covering of B_q^n from below if $\lfloor C \rfloor_s = \lfloor B_q^n \rfloor_s$ (cf. [6]). A code $C \subseteq B_q^n$ capable of correcting *s* deletions is called a perfect code capable of correcting *s* deletions if *C* is an *s*-covering of B_q^n from below. For brevity a perfect code in B_q^n capable of correcting single deletions will be referred to as a perfect code.

In [6], the author showed that there exists a perfect code in B_q^4 of cardinality $q(q-1)(q-2)/4 + q^2$ for any even q. Note that this is not optimal. In previous sections, we have constructed an optimal code C in B_q^4 for any even q. By

counting the cardinality of $\lfloor C \rfloor_1$, one can show that it can not be a perfect code. (For example, the optimal code in Table 1 is not perfect.) However, we can modify the construction of an optimal code to obtain an optimal perfect code as follows.

Suppose that C is an optimal code of B_q^4 for an even q, which is constructed in previous sections. Decompose C into the subcodes C_1 , C_2 , and C_4 , where $C_i = \{x \in C \mid |\lfloor x \rfloor_1 | = i\}$ for i = 1, 2, and 4. From the structure of C, the codes C_1 and C_2 are as follows:

$$\begin{cases} C_1 = \{(a, a, a, a) \mid a \in B_q\}, \\ C_2 = \{(a, a, b, b) \mid a, b \in B_q \text{ and } a \neq b\}. \end{cases}$$

Note that $|C_1| = q$, $|C_2| = q(q-1)$, and $|C_4| = \frac{q^2(q+2)}{4} - (q+q(q-1))$. Now we modify the subcode C_2 to make C'_2 as follows:

$$\begin{array}{rcl} C_2' &=& C_2 \setminus \{(2t,2t,2t+1,2t+1) \mid 0 \leq t \leq \frac{q}{2}-1\} \\ & \cup \{(2t,2t+1,2t,2t+1) \mid 0 \leq t \leq \frac{q}{2}-1\}. \end{array}$$

It can be easily checked that $C' = C_1 \cup C'_2 \cup C_4$ is a code capable of correcting single deletions (i.e. $\rho(C') > 2$) with |C| = |C'|. Note that $|C'_2| = |C_2|$ and $|C'_2 \setminus C_2| = \frac{q}{2}$. For each $x \in C'$, after single deletions, we have the following relations:

$$\begin{cases} |\lfloor x \rfloor_1| = 1 \text{ if } x \in C_1, \\ |\lfloor x \rfloor_1| = 2 \text{ if } x \in C_2, \\ |\lfloor x \rfloor_1| = 4 \text{ if } x \in C'_2 \setminus C_2, \\ |\lfloor x \rfloor_1| = 4 \text{ if } x \in C_4. \end{cases}$$

By computing the cardinality of $\lfloor C' \rfloor_1$, we obtain:

$$| [C']_1 | = | [C_1]_1 | + | [C'_2]_1 | + | [C_4]_1 | = q+2 (q(q-1)-\frac{q}{2}) + 4\frac{q}{2} + 4 (\frac{q^2(q+2)}{4} - q - q(q-1)) = q^3.$$

Hence C' is an optimal perfect code. Therefore we have obtained the following theorem.

Theorem 5.1. For any even q, we can construct an optimal perfect in B_q^4 code capable of correcting single deletions.

Acknowledgement. The authors would like to thank to Dr. Vladimir Levedev for introducing this problem to them.

References

- P.C. Denny, Search and Enumeration Techniques for Incidence Structures, CDMTCS Research Report Series, 085 (1998).
- [2] H. Hanani, On quadruple systems, Canad. J. Math., 12 (1960), 145-157.
- [3] J.G. Kalbfleisch and R.G. Stanton, Maximal and minimal coverings of (k-1)-tuples by k-tuples, Pacif. J. Math. **26**, (1968) 131-140.
- [4] V. Lebedev, personal communication.

- [5] V.I. Levenshtein, Binary codes capable of correcting deletions, insertions and reversals, Soviet Phys. Dokl. 10 (1966), 707 - 710.
- [6] V.I. Levenshtein, On perfect codes in deletion and insertion metric, Discrete Math. Appl. Vol. 2, No. 3, (1992) 241-258.
- [7] W.H. Mills, On the covering of triples by quadruples, Proc. 5th Southeastern Conference on Combin., Graph Th. and Comp., Boca Raton, 1974, 563-581
- [8] M. Reiss, Uber eine Steinersche Combinatorische Aufgabe welche im 45sten Bande diese Journals, Seite 181, gestellt worden ist, J. Reine Angew. Math. 56, (1859) 326-344.
- [9] D.V. Zinoviev and V.A. Zinoviev, On resolvability of Steiner quadruple systems with the first three smallest values of rank, Proceedings of the 10th International Workshop on Algebraic and Combinatorial Coding Theory (ACCT 2006), September 2006, Zvenigorod, Russia, 282-286.