

A SUPERCHARACTER THEORY FOR THE SYLOW p -SUBGROUPS OF THE FINITE SYMPLECTIC AND ORTHOGONAL GROUPS

CARLOS A. M. ANDRÉ & ANA MARGARIDA NETO

ABSTRACT. We define the superclasses for a classical finite unipotent group U of type $B_n(q)$, $C_n(q)$, or $D_n(q)$, and show that, together with the supercharacters defined in [6], they form a supercharacter theory in the sense of [11]. In particular, we prove that the supercharacters take a constant value on each superclass, and evaluate this value. As a consequence, we obtain a factorization of any superclass as a product of elementary superclasses. In addition, we also define the space of superclass functions, and prove that it is spanned by the supercharacters. As a consequence, we (re)obtain the decomposition of the regular character as an orthogonal linear combination of supercharacters. Finally, we define the supercharacter table of U , and prove various orthogonality relations for supercharacters (similar to the well-known orthogonality relations for irreducible characters).

1. INTRODUCTION

This paper is a continuation of the authors' papers [5, 6], and develops a supercharacter theory for the Sylow p -subgroup of the symplectic or orthogonal groups defined over the finite field \mathbb{F}_q with q elements; throughout the paper, p will stand for an arbitrary odd prime number, and $q = p^e$, $e \geq 1$, will be a fixed power of p . The concept of a “supercharacter theory” for an arbitrary finite group was developed by P. Diaconis and I. M. Isaacs in the paper [11]. Roughly, a supercharacter theory replaces irreducible characters by “supercharacters”, and conjugacy classes by “superclasses”, in such a way that

Date: August 2, 2018.

2000 Mathematics Subject Classification. Primary 20C15; Secondary 20G40.

Key words and phrases. Finite unipotent group, Symplectic group, Orthogonal group, Supercharacter, Superclass, Supercharacter theory, Positive root, Basic set of positive roots.

This research was made within the activities of the Centro de Estruturas Lineares e Combinatórias (University of Lisbon, Portugal) and was partially supported by the Fundação para a Ciência e Tecnologia (Lisbon, Portugal) through the project POCTI-ISFL-1-1431. A large part of the research of the first author was made and concluded while he was visiting the University of Stanford (USA) and participating in the special program on “Combinatorial Representation Theory” at the MSRI (Berkeley, USA) whose hospitality is gratefully acknowledged, and was partially supported by the sabbatical research grant 4/2008 of the Fundação Luso-Americana para o Desenvolvimento (Lisbon, Portugal). The first author also expresses his sincere gratitude to Persi Diaconis for his invitation to visit the University of Stanford, and for many enlightening discussions regarding supercharacters and their applications.

a “supercharacter table” can be constructed as an “almost unitary” matrix with similar properties as the usual character table (namely, orthogonality of rows and columns). More precisely, given any finite group G , a *supercharacter theory* for G consists of a partition \mathcal{K} of G and a set \mathcal{X} of (complex) characters of G satisfying the following three axioms:

- (S1) $|\mathcal{K}| = |\mathcal{X}|$;
- (S2) every irreducible character of G is a constituent of a unique $\xi \in \mathcal{X}$;
- (S3) the characters in \mathcal{X} are constant on the member of \mathcal{K} .

The elements of \mathcal{K} will be referred to as *superclasses*, and the elements of \mathcal{X} as *supercharacters* of G . (We observe that, by [11, Lemma 2.1], axiom (S2) is equivalent to requiring that $\{1\} \in \mathcal{K}$.)

Every finite group G has two “trivial” supercharacter theories: the full character theory (where \mathcal{X} consists of all irreducible characters of G , and \mathcal{K} of all the conjugacy classes of G), and the one where $\mathcal{X} = \{1_G, \rho_G - 1_G\}$ and \mathcal{K} consists of the sets $\{1\}$ and $G - \{1\}$; as usual, we denote by 1_G the trivial character and by ρ_G the regular character of G . Although for some groups these are the only possibilities, there are many groups for which nontrivial supercharacter theories exist, and in many cases it may be possible to obtain useful information using some particular supercharacter theory. An illustrating example can be found in the paper [7] where E. Arias-Castro, P. Diaconis and R. Stanley showed that a special supercharacter theory can be applied to study a random walk on upper triangular matrices over finite fields using techniques that traditionally required the knowledge of the full character theory.

Supercharacters theories were initially developed for the upper unitriangular group $U_n(q)$ consisting of all unipotent upper-triangular $n \times n$ matrices over the finite field \mathbb{F}_q with q elements (where q is a power of some prime number p). In his PhD thesis [1], the first author begun the study of the “basic characters” of $U_n(q)$ (under the assumption that $p \geq n$), and was able to show that “clumping” together some of the conjugacy classes and some of the irreducible characters one attains a workable “approximation” to the representation theory of $U_n(q)$. His results were published in a series of papers in the Journal of Algebra, and showed in particular that the “basic characters” determine uniquely the superclasses of a supercharacter theory for $U_n(q)$. The original theory relies on a construction due to D. Kazhdan (see [13]) and is based on Kirillov’s method of coadjoint orbits (see [14] for a description of Kirillov’s method for the unitriangular group). Later, in his PhD thesis [17], N. Yan showed how the “basic characters” can be obtained using more elementary methods which avoid Kazhdan’s construction and the algebraic geometry involved in it. Yan’s approach is valid for an arbitrary prime, and was generalized later by P. Diaconis and M. Isaacs in the paper [11] in order to extend the theory for an arbitrary finite algebra group defined over \mathbb{F}_q .

In the papers [5] (for sufficiently large primes) and [6] (for arbitrary odd primes), the authors started to develop a supercharacter theory for a Sylow p -subgroup U of one the

(non-twisted) Chevalley groups $C_n(q)$, $B_n(q)$, and $D_n(q)$, by defining the supercharacters of U and proving some of their main properties. As in the case of the unitriangular group, the supercharacters of U are parametrized by certain “minimal” subsets of (positive) roots. In fact, it is known that the supercharacters of $U_n(q)$ can be obtained as certain “reduced” products of “elementary characters” which are irreducible characters corresponding to the “matrix entries” (i, j) , for $1 \leq i < j \leq n$, labelled by nonzero elements of \mathbb{F}_q ; in Yan’s thesis, the “elementary characters” were called “primary characters”, and the supercharacters were called “transition characters”. Following Yan’s method, one can show that the supercharacters of $U_n(q)$ are parametrized by certain combinatorial data consisting of a “basic set” D of matrix entries such that no two elements of D agree in, either the first, or the second, coordinate, and of a map ϕ from D to the nonzero elements of \mathbb{F}_q . In the papers [5, 6], the authors defined the supercharacters also as certain “reduced” products of “elementary characters” (which in general are not necessarily irreducible characters) of the given Sylow p -subgroup U . These “reduced” products are parametrized by pairs consisting of a conveniently chosen “basic subset of roots” and of a map to the nonzero elements of \mathbb{F}_q . (We note that the roots in the unitriangular case are in one-to-one correspondence with the matrix entries.) In fact, the group U can be naturally identified with a subgroup of a unitriangular group, and the supercharacters of U can be obtained as constituents of the restriction of a supercharacter of that unitriangular group.

On the other hand, as shown in Yan’s thesis, the same combinatorial data parametrize the superclasses of $U_n(q)$ (see also [3, 4] where the superclasses were called “basic subvarieties”). In fact, a superclass of $U_n(q)$ can be obtained as a “basic” product of “elementary superclasses” which are conjugacy classes corresponding to the “matrix entries”, and labelled by nonzero elements of \mathbb{F}_q . As for supercharacters, we define the superclasses of U by “restricting” the superclasses of the larger unitriangular group, and show that they are indexed by the same combinatorial data consisting of a conveniently chosen “basic set of roots” where each root is labeled by a nonzero element of \mathbb{F}_q .

The paper is organized as follows. In Section 2, we introduce the necessary notation, and recall the definition and main property of the supercharacters of the group U . In Section 3, we define the superclasses of U by intersecting with the superclasses of the unitriangular group which contains U , and show that the set \mathcal{K} of all superclasses gives a partition of U . Then, in Section 4, we define a superclass function on U as a function taking a constant (complex) value on each superclass, and prove that the supercharacters are superclass functions, and form an orthogonal basis for the complex vector space $\text{scf}(U)$ consisting of all superclass functions. As a consequence, we obtain an explicit decomposition of the regular character as a linear combination with positive integers of all the supercharacters (see [6, Theorem 5.2] for a different proof), and (re)prove the main theorem on supercharacters which states that every irreducible character is a constituent of a (unique) supercharacter. In Section 5, we determine the constant value of

a supercharacter on a superclass, and conclude that the superclasses and supercharacters satisfy the axioms of a supercharacter theory for U in the sense of Diaconis and Isaacs. As a consequence, we show that every superclass factorizes uniquely as a product of elementary superclasses. Finally, in Section 6, we define the supercharacter table of U as the square matrix with entries given by the supercharacter values, and prove the main orthogonality relations for supercharacters. As a consequence, we also deduce that the space $\text{scf}(U)$ of superclass functions is a commutative semisimple algebra with respect to the convolution product.

2. SUPERCHARACTERS

Let $p \geq 3$ be a prime number, $q = p^e$ ($e \geq 1$) a power of p , and \mathbb{F}_q the finite field with q elements. For a fixed positive integer n , let G denote one of the following classical finite groups: the symplectic group $Sp_{2n}(q)$, the even orthogonal group $O_{2n}(q)$, or the odd orthogonal group $O_{2n+1}(q)$ (in alternative notation, these are the (non-twisted) Chevalley groups $C_n(q)$, $B_n(q)$, and $D_n(q)$, respectively). Throughout the paper, we set $U = G \cap U_m(q)$ where

$$m = \begin{cases} 2n, & \text{if } G = Sp_{2n}(q), \text{ or } G = O_{2n}(q), \\ 2n + 1, & \text{if } G = O_{2n+1}(q), \end{cases}$$

and $U_m(q)$ denotes the upper unitriangular group consisting of all unipotent upper-triangular $m \times m$ matrices over \mathbb{F}_q . Then, U is a Sylow p -subgroup of G , and it is described as follows. Let $J = J_n$ be the $n \times n$ matrix with 1's along the anti-diagonal and 0's elsewhere. Then, U consists of all (block) matrices of the form

$$(2a) \quad \begin{pmatrix} x & xu & xz \\ 0 & I_r & -u^T J \\ 0 & 0 & Jx^{-T} J \end{pmatrix}$$

where $x \in U_n(q)$, u is an $n \times r$ matrix over \mathbb{F}_q , and

- (i) $r = 0$, and $Jz^T - zJ = 0$, if $U \leq Sp_{2n}(q)$;
- (ii) $r = 0$, and $Jz^T + zJ = 0$, if $U \leq O_{2n}(q)$;
- (iii) $r = 1$, and $Jz^T + zJ = -uu^T$, if $U \leq O_{2n+1}(q)$.

As mentioned in the Introduction, both supercharacters and superclasses of U will be parametrized by certain subsets of (positive) roots. Thus, we introduce some notation and recall some elementary facts concerning roots; for the details, we refer to the books [8, 9] by R. Carter (see also [10, Chapter 8]). Let T be the maximal torus of G consisting of all diagonal matrices, and Σ the root system defined by T . The elements of Σ are described as follows. For each $1 \leq i \leq n$, let $\varepsilon_i: T \rightarrow \mathbb{F}_q^\times$ be the map defined by $\varepsilon_i(t) = t_i$ for all $t \in T$; here, we denote by $t_i \in \mathbb{F}_q^\times$ the (i, i) th entry of the matrix $t \in T$. Then, $\Sigma = \Phi \cup (-\Phi)$ where

$$\Phi = \{\varepsilon_i \pm \varepsilon_j: 1 \leq i < j \leq n\} \cup \Phi'$$

and

$$\Phi' = \begin{cases} \{2\varepsilon_i: 1 \leq i \leq n\}, & \text{if } G = Sp_{2n}(q), \\ \emptyset, & \text{if } G = O_{2n}(q), \\ \{\varepsilon_i: 1 \leq i \leq n\}, & \text{if } G = O_{2n+1}(q). \end{cases}$$

The roots in Φ are said to be *positive*, and the roots in $-\Phi$ are said to be *negative*. Throughout the paper, the word “root” will always stand for “positive root”.

With Φ we associate the subset of “matrix entries” $\mathcal{E} \subseteq \{(i, j): -n \leq i, j \leq n\}$ as follows. For any $\alpha \in \Phi$, we set

$$\mathcal{E}(\alpha) = \begin{cases} \{(i, j), (-j, -i)\}, & \text{if } \alpha = \varepsilon_i - \varepsilon_j \text{ for } 1 \leq i < j \leq n, \\ \{(i, -j), (j, -i)\}, & \text{if } \alpha = \varepsilon_i + \varepsilon_j \text{ for } 1 \leq i < j \leq n, \\ \{(i, -i)\}, & \text{if } G = Sp_{2n}(q) \text{ and } \alpha = 2\varepsilon_i \text{ for } 1 \leq i \leq n, \\ \{(i, 0), (0, -i)\}, & \text{if } G = O_{2n+1}(q) \text{ and } \alpha = \varepsilon_i \text{ for } 1 \leq i \leq n, \end{cases}$$

and we define

$$\mathcal{E} = \bigcup_{\alpha \in \Phi} \mathcal{E}(\alpha).$$

More generally, for each subset $\Psi \subseteq \Phi$, we set

$$\mathcal{E}(\Psi) = \bigcup_{\alpha \in \Psi} \mathcal{E}(\alpha);$$

hence, $\mathcal{E} = \mathcal{E}(\Phi)$.

On the other hand, we consider the mirror order \prec on the set $\{0, \pm 1, \dots, \pm(n+1)\}$ which is defined as

$$1 \prec 2 \prec \dots \prec n+1 \prec 0 \prec -(n+1) \prec \dots \prec -2 \prec -1,$$

and we shall index the rows (from left to right) and columns (from top to bottom) of any $m \times m$ matrix according to this ordering. Hence, the entries of any matrix $x \in U_m(q)$ are indexed by all the pairs $(i, j) \in \mathcal{E}$: for each $(i, j) \in \mathcal{E}$, we shall write $x_{i,j}$ to denote the (i, j) th entry of x (which occurs in the i th row and in the j th column). For our purposes, it is convenient to consider the set

$$\mathcal{E}^+ = \{(i, j) \in \mathcal{E}: 1 \leq i \leq n, i \prec j \preceq -i\},$$

and extend this notation to any subset $\Psi \subseteq \Phi$ by setting

$$\mathcal{E}^+(\Psi) = \mathcal{E}(\Psi) \cap \mathcal{E}^+.$$

We observe that there exists a one-to-one correspondence between Φ and \mathcal{E}^+ .

For any $\alpha \in \Phi$, we define the subgroup U_α of U as follows:

(i) if $\alpha = \varepsilon_i - \varepsilon_j$ for $1 \leq i < j \leq n$, then

$$U_\alpha = \{x \in U: x_{i,k} = 0, i < k < j\};$$

(ii) if $\alpha = \varepsilon_i - \varepsilon_j$ for $1 \leq i < j \leq n$, then

$$U_\alpha = \{x \in U : x_{i,k} = x_{j,l} = 0, i < k \leq n, j < l \leq 0\};$$

(iii) if, either $\alpha = 2\varepsilon_i$ for $1 \leq i \leq n$ (in the case where $U \leq Sp_{2n}(q)$), or $\alpha = \varepsilon_i$ for $1 \leq i \leq n$ (in the case where $U \leq O_{2n+1}(q)$), then

$$U_\alpha = \{x \in U : x_{i,k} = 0, i < k \leq n\}.$$

Let $\vartheta: \mathbb{F}_q \rightarrow \mathbb{C}^\times$ be a non-trivial linear character of the additive group \mathbb{F}_q^+ of \mathbb{F}_q (this character will be kept fixed throughout the paper; moreover, all characters will be taken over the complex field). For any $r \in \mathbb{F}_q^\times$, the mapping $x \mapsto \vartheta(rx_{i,j})$ defines a linear character $\lambda_{\alpha,r}: U_\alpha \rightarrow \mathbb{C}^\times$ of U_α , and we define the *elementary character* $\xi_{\alpha,r}$ to be the induced character

$$\xi_{\alpha,r} = (\lambda_{\alpha,r})^U$$

(see [2] for the corresponding definition in the case of the unitriangular group; see also [11, Corollary 5.11] and the discussion thereon).

We next define the notion of a “basic subset of roots”. To start with, we recall that a subset $\mathcal{D} \subseteq \mathcal{E}$ is said to be *basic* if it contains at most one entry from each row and at most one root from each column; in other words, $\mathcal{D} \subseteq \mathcal{E}$ is basic if

$$|\{j : i < j \preceq -1, (i, j) \in \mathcal{D}\}| \leq 1 \quad \text{and} \quad |\{i : 1 \preceq i < j, (i, j) \in \mathcal{D}\}| \leq 1$$

for all $-n \leq i, j \leq n$. Then, we say that $D \subseteq \Phi$ is a *basic subset* if $\mathcal{D} = \mathcal{E}(D)$ is a basic subset of \mathcal{E} . (We will always use script letters to denote basic subsets of \mathcal{E} , in contrast to basic subsets of Φ which will be mostly denoted by italic letters.)

Given any non-empty basic subset $D \subseteq \Phi$ and any map $\phi: D \rightarrow \mathbb{F}_q^\times$, we define the supercharacter $\xi_{D,\phi}$ to be the product

$$\xi_{D,\phi} = \prod_{\alpha \in D} \xi_{\alpha,\phi(\alpha)}.$$

For convenience, if D is the empty subset of Φ , we consider the empty map $\phi: D \rightarrow \mathbb{F}_q^\times$, and define $\xi_{D,\phi}$ to be the unit character 1_U of U . Let

$$U_D = \bigcap_{\alpha \in D} U_\alpha \quad \text{and} \quad \lambda_{D,\phi} = \prod_{\alpha \in D} (\lambda_{\alpha,\phi(\alpha)})_{U_D}.$$

Then, $\lambda_{D,\phi}$ is clearly a linear character of U_D and, by [5, Proposition 2.2], the supercharacter $\xi_{D,\phi}$ can be obtained as the induced character

$$(2b) \quad \xi_{D,\phi} = (\lambda_{D,\phi})^U.$$

Throughout the paper, we will refer to the pair (D, ϕ) as a *basic pair* for U ; hence, $D \subseteq \Phi$ is a basic subset, and $\phi: D \rightarrow \mathbb{F}_q^\times$ is a map.

The main result of [6] is the following theorem. (Given any finite group G , we denote by $\text{Irr}(G)$ the set of all irreducible characters of G , and by $\langle \cdot, \cdot \rangle$ (or by $\langle \cdot, \cdot \rangle_G$ if necessary)

the Frobenius' scalar product on the complex vector space of all class functions defined on G .)

Theorem 2.1 ([6, Theorem 1.1]). *Let χ be an arbitrary irreducible character of U . Then, χ is a constituent of a unique supercharacter of U ; in other words, there exists a unique basic subset $D \subseteq \Phi$ and a unique map $\phi: D \rightarrow \mathbb{F}_q^\times$ such that $\langle \chi, \xi_{D,\phi} \rangle \neq 0$.*

We note that this theorem establishes axiom (S2) of the definition of a supercharacter theory.

3. SUPERCLASSES

In this section, we define the superclasses of U . This notion depends strongly on certain ‘‘basic subvarieties’’ defined by polynomial equations on the Lie algebra \mathfrak{u} of U . Let \mathfrak{g} denote one of the following classical Lie algebras defined over \mathbb{F}_q : the symplectic Lie algebra $\mathfrak{sp}_{2n}(q)$, the even orthogonal Lie algebra $\mathfrak{o}_{2n}(q)$, or the odd orthogonal Lie algebra $\mathfrak{o}_{2n+1}(q)$. Then, $\mathfrak{u} = \mathfrak{g} \cap \mathfrak{u}_m(q)$ where $\mathfrak{u}_m(q)$ denotes the upper niltriangular Lie algebra consisting of all nilpotent upper-triangular $m \times m$ matrices over \mathbb{F}_q . Thus, \mathfrak{u} consists of all (block) matrices of the form

$$(3a) \quad \begin{pmatrix} a & u & w \\ 0 & 0_r & -u^T J \\ 0 & 0 & -J a^T J \end{pmatrix}$$

where $a \in \mathfrak{u}_n(q)$, u is an $n \times r$ matrix over \mathbb{F}_q , and

- (i) $r = 0$, and $Jw^T - wJ = 0$, if $\mathfrak{u} \leq \mathfrak{sp}_{2n}(q)$;
- (ii) $r = 0$, and $Jw^T + wJ = 0$, if $\mathfrak{u} \leq \mathfrak{o}_{2n}(q)$;
- (iii) $r = 1$, and $Jw^T + wJ = -uu^T$, if $\mathfrak{u} \leq \mathfrak{o}_{2n+1}(q)$.

For any $\alpha \in \Phi$, we will denote by e_α the matrix in \mathfrak{u} defined as follows (as usual, $1 \leq i < j \leq n$):

$$e_\alpha = \begin{cases} e_{i,j} - e_{-j,-i}, & \text{if } \alpha = \varepsilon_i - \varepsilon_j, \\ e_{i,-j} + e_{j,-i}, & \text{if } \alpha = \varepsilon_i + \varepsilon_j \text{ and } \mathfrak{u} \leq \mathfrak{sp}_{2n}(q), \\ e_{i,-j} - e_{j,-i}, & \text{if } \alpha = \varepsilon_i + \varepsilon_j \text{ and } \mathfrak{u} \leq \mathfrak{o}_{2n}(q) \text{ or } \mathfrak{u} = \mathfrak{o}_{2n+1}(q), \\ e_{i,-i}, & \text{if } \mathfrak{u} \leq \mathfrak{sp}_{2n}(q) \text{ and } \alpha = 2\varepsilon_i, \\ e_{i,0} - e_{0,-i}, & \text{if } \mathfrak{u} \leq \mathfrak{o}_{2n+1}(q) \text{ and } \alpha = \varepsilon_i. \end{cases}$$

It is clear that $\{e_\alpha : \alpha \in \Phi\}$ is an \mathbb{F}_q -basis of \mathfrak{u} .

Given be an arbitrary basic pair (D, ϕ) for U with D non-empty basic subset, we define the element

$$e_{D,\phi} = \sum_{\alpha \in D} \phi(\alpha) e_\alpha \in \mathfrak{u}.$$

Since $\mathfrak{u} \subseteq \mathfrak{u}_m(q)$, we may consider the orbit

$$V_{D,\phi} = U_m(q)e_{D,\phi}U_m(q) \subseteq \mathfrak{u}_m(q)$$

for the natural action of $U_m(q) \times U_m(q)$ on $\mathfrak{u}_m(q)$ given by $(x, y) \cdot u = xuy^{-1}$ for all $x, y \in U_m(q)$ and $u \in \mathfrak{u}_m(q)$. By definition (see, for example, [4], [7], [11], or [17]), the superclass of $U_m(q)$ which contains the element $1 + e_{D,\phi} \in U_m(q)$ is the subset

$$1 + V_{D,\phi} = \{1 + a : a \in V_{D,\phi}\} \subseteq U_m(q),$$

and we define the *superclass* $K_{D,\phi}$ of U to be the intersection

$$K_{D,\phi} = U \cap (1 + V_{D,\phi}) = \{x \in U : x - 1 \in V_{D,\phi}\}.$$

We will also consider the intersection

$$O_{D,\phi} = V_{D,\phi} \cap \mathfrak{u},$$

but observe that it is not necessarily true that $K_{D,\phi} = 1 + O_{D,\phi}$; however, there exists a bijection between $K_{D,\phi}$ and $O_{D,\phi}$ (see Lemma 3.2 below).

The following result is a clear consequence of the definition; in the case where D is the empty subset of Φ , we consider the empty map $\phi: D \rightarrow \mathbb{F}_q^\times$, and define $e_{D,\phi} = 0$, so that $V_{D,\phi} = \{0\}$ and $K_{D,\phi} = \{1\}$.

Lemma 3.1. *For every basic pair (D, ϕ) for U , the superclass $K_{D,\phi}$ is invariant under conjugation. In particular, $K_{D,\phi}$ is a union of conjugacy classes.*

On the other hand, by [4, Theorem 2.1], for any basic pairs (D, ϕ) and (D', ϕ') , we have $V_{D,\phi} \cap V_{D',\phi'} \neq \emptyset$ if and only if $(D, \phi) = (D', \phi')$. In fact, by the definition, $\mathcal{E}(D)$ and $\mathcal{E}(D')$ are basic subsets of $\mathcal{E} = \mathcal{E}(\Phi)$ satisfying $\mathcal{E}(D) = \mathcal{E}(D')$ if and only if $D = D'$, and the elements $e_{D,\phi}, e_{D',\phi'} \in \mathfrak{u}_m(q)$ are equal if and only if $D = D'$ and $\phi = \phi'$. Therefore, we obtain a disjoint union $\bigcup_{D,\phi} K_{D,\phi}$ indexed by all basic pairs (D, ϕ) for U . We next show that U equals this union, thus obtaining a partition \mathcal{K} of U which satisfies axiom (S1) for the required supercharacter theory. The proof of the following auxiliary result can be found in [6, Lemma 2.3].

Lemma 3.2. *Let (D, ϕ) be a basic pair for U . Let*

$$z = \begin{pmatrix} x & xv & xw \\ 0 & I_r & -v^T J \\ 0 & 0 & Jx^{-T} J \end{pmatrix} \in U \quad \text{and} \quad a_z = \begin{pmatrix} u & v & w \\ 0 & 0_r & -v^T J \\ 0 & 0 & -Ju^T J \end{pmatrix} \in \mathfrak{u}$$

where $x = 1 + u$. Then, $z \in K_{D,\phi}$ if and only if $a_z \in V_{D,\phi}$. Moreover, the mapping $z \mapsto a_z$ defines a bijection from U to \mathfrak{u} .

Now, let $z \in U$ be arbitrary, and $a_z \in \mathfrak{u}$ be as in the lemma. Then, by [4, Theorem 2.1] (see also [11, Appendix A]), there exists a unique basic subset $\mathcal{D} \subseteq \mathcal{E}$ and a unique map

$\varphi: \mathcal{D} \rightarrow \mathbb{F}_q^\times$ such that $u \in V_{\mathcal{D},\varphi}$ where $V_{\mathcal{D},\varphi} = U_m(q)e_{\mathcal{D},\varphi}U_m(q)$ and

$$e_{\mathcal{D},\varphi} = \sum_{(i,j) \in \mathcal{D}} \varphi(i,j)e_{i,j} \in \mathbf{u}_m(q).$$

We claim that $\mathcal{D} = \mathcal{E}(D)$ for a uniquely determined basic subset $D \subseteq \Phi$, and $e_{\mathcal{D},\varphi} = e_{D,\phi} \in \mathbf{u}$ for a uniquely determined map $\phi: D \rightarrow \mathbb{F}_q^\times$; thus, we must have $\phi(\alpha) = \varphi(i,j)$ for all $\alpha \in D$ with $(i,j) \in \mathcal{E}^+(\alpha)$. To see this, we recall that $V_{\mathcal{D},\varphi}$ is the zero set of a finite family of polynomial equations which are defined as follows.

For each entry $(i,j) \in \mathcal{E}$, let $\mathcal{D}(i,j)$ denote the subset

$$\mathcal{D}(i,j) = \{(k,l) \in \mathcal{D}: i \prec k \preceq -1, 1 \preceq l \prec j\}.$$

Let $\mathcal{D}(i,j) = \{(i_1, j_1), \dots, (i_t, j_t)\}$ where $j_1 \prec j_2 \prec \dots \prec j_t$, and let $\sigma \in S_t$ be the permutation such that $i_{\sigma(1)} \prec i_{\sigma(2)} \prec \dots \prec i_{\sigma(t)}$; as usual, we denote by S_t the symmetric group of degree t . Then, for any $u \in \mathbf{u}_m(q)$, we define $\Delta_{i,j}^{\mathcal{D}}(u)$ to be the determinant

$$\Delta_{i,j}^{\mathcal{D}}(u) = \begin{vmatrix} u_{i,j_1} & \cdots & u_{i,j_t} & u_{i,j} \\ u_{i_{\sigma(1)},j_1} & \cdots & u_{i_{\sigma(1)},j_t} & u_{i_{\sigma(1)},j} \\ \vdots & & \vdots & \vdots \\ u_{i_{\sigma(t)},j_1} & \cdots & u_{i_{\sigma(t)},j_t} & u_{i_{\sigma(t)},j} \end{vmatrix}.$$

We note that $\Delta_{i,j}^{\mathcal{D}}(u) = u_{i,j}$ whenever $\mathcal{D}(i,j) = \emptyset$; in particular, if \mathcal{D} is empty, then $\Delta_{i,j}^{\mathcal{D}}(u) = u_{i,j}$ for all $u \in \mathbf{u}_m(q)$.

By [4, Proposition 2.3], we know that

$$(3b) \quad V_{\mathcal{D},\varphi} = \{u \in \mathbf{u}_m(q): \Delta_{i,j}^{\mathcal{D}}(u) = \Delta_{i,j}^{\mathcal{D}}(e_{\mathcal{D},\varphi}) \text{ for all } (i,j) \in R(\mathcal{D})\}$$

where $R(\mathcal{D}) = \mathcal{E} - S(\mathcal{D})$ and

$$S(\mathcal{D}) = \bigcup_{(i,j) \in \mathcal{D}} (\{(i,s): j \prec s \preceq -1\} \cup \{(r,j): 1 \preceq r \prec i\});$$

we refer to the entries in $S(\mathcal{D})$ as the \mathcal{D} -singular entries, and to those in $R(\mathcal{D})$ as the \mathcal{D} -regular entries. It is easy to show that, for an arbitrary \mathcal{D} -regular entry $(i,j) \in R(\mathcal{D})$, we have

$$(3c) \quad \Delta_{i,j}^{\mathcal{D}}(e_{\mathcal{D},\varphi}) = \begin{cases} (-1)^t \operatorname{sgn}(\sigma) \varphi(i,j) \prod_{s=1}^t \varphi(i_s, j_s), & \text{if } (i,j) \in \mathcal{D}, \\ 0, & \text{if } (i,j) \notin \mathcal{D}, \end{cases}$$

where $\mathcal{D}(i,j) = \{(i_1, j_1), \dots, (i_t, j_t)\}$, $j_1 \prec j_2 \prec \dots \prec j_t$, and $\sigma \in S_t$ is such that $i_{\sigma(1)} \prec i_{\sigma(2)} \prec \dots \prec i_{\sigma(t)}$.

We now prove the following auxiliary result; for simplicity of writing, given any basic subset $D \subseteq \Phi$ and any entry $(i,j) \in \mathcal{E}$, we set $D(i,j) = \mathcal{D}(i,j)$ for $\mathcal{D} = \mathcal{E}(D)$.

Lemma 3.3. *Let $D \subseteq \Phi$ be a basic subset, let $u \in \mathfrak{u}$, and let $(i, j) \in \mathcal{E}^+$. Then,*

$$\Delta_{i,j}^{\mathcal{E}(D)}(u) = (-1)^{r+1} \Delta_{-j,-i}^{\mathcal{E}(D)}(u)$$

where

$$r = \begin{cases} |D(i, j)|, & \text{if, either } \mathfrak{u} \leq \mathfrak{o}_{2n}(q), \text{ or } \mathfrak{u} \leq \mathfrak{o}_{2n+1}(q), \\ |D(i, j)|, & \text{if } \mathfrak{u} \leq \mathfrak{sp}_{2n}(q) \text{ and } j \leq n, \\ |D'(i, j)| - 1, & \text{if } \mathfrak{u} \leq \mathfrak{sp}_{2n}(q) \text{ and } -j \leq n, \end{cases}$$

and $D'(i, j) = D(i, j) \cap \{(i, j) \in \mathcal{E}^+ : j \leq n\}$.

Proof. Let $D(i, j) = \{(i_1, j_1), \dots, (i_t, j_t)\}$ where $j_1 \prec \dots \prec j_t \prec j$, and let $\sigma \in S_t$ be such that $i \prec i_{\sigma(1)} \prec \dots \prec i_{\sigma(t)}$. By the definition of $\mathcal{E}(D)$, we clearly have

$$D(-j, -i) = \{(-j_1, -i_1), \dots, (-j_t, -i_t)\}$$

where $-j \prec -j_t \prec \dots \prec -j_1$ and $-i_{\sigma(t)} \prec \dots \prec -i_{\sigma(1)} \prec -i$. Thus,

$$\Delta_{-j,-i}^{\mathcal{E}(D)}(u) = \begin{vmatrix} u_{-j,-i_{\sigma(t)}} & \cdots & u_{-j,-i_{\sigma(1)}} & u_{-j,-i} \\ u_{-j_t,-i_{\sigma(t)}} & \cdots & u_{-j_t,-i_{\sigma(1)}} & u_{-j_t,-i} \\ \vdots & & \vdots & \vdots \\ u_{-j_1,-i_{\sigma(t)}} & \cdots & u_{-j_1,-i_{\sigma(1)}} & u_{-j_1,-i} \end{vmatrix}.$$

Firstly, we assume that, either $\mathfrak{u} \not\leq \mathfrak{sp}_{2n}(q)$, or $\mathfrak{u} \leq \mathfrak{sp}_{2n}(q)$ and $j \leq n$. Let $C_{j_1}, \dots, C_{j_t}, C_j$ denote the column vectors of $\Delta_{i,j}^{\mathcal{E}(D)}(u)$, and $L_{-j}, L_{-j_t}, \dots, L_{-j_1}$ the row vectors of $\Delta_{-j,-i}^{\mathcal{E}(D)}(u)$. For any $k \in \{j_1, \dots, j_t, j\}$, we have $L_{-k} = -C_k^T J$ where $J = J_{t+1}$ is the $(t+1) \times (t+1)$ matrix with 1's along the anti-diagonal and 0's elsewhere, and thus we deduce that

$$\begin{aligned} \Delta_{-j,-i}^{\mathcal{E}(D)}(u) &= (-1)^{t+1} \begin{vmatrix} u_{i_{\sigma(t),j} } & \cdots & u_{i_{\sigma(1),j} } & u_{i,j} \\ u_{i_{\sigma(t),j_t} } & \cdots & u_{i_{\sigma(1),j_t} } & u_{i,j_t} \\ \vdots & & \vdots & \vdots \\ u_{i_{\sigma(t),j_1} } & \cdots & u_{i_{\sigma(1),j_1} } & u_{i,j_1} \end{vmatrix} = (-1)^{t+1} \begin{vmatrix} u_{i_{\sigma(t),j} } & u_{i_{\sigma(t),j_t} } & \cdots & u_{i_{\sigma(t),j_1} } \\ \vdots & \vdots & & \vdots \\ u_{i_{\sigma(1),j} } & u_{i_{\sigma(1),j_t} } & \cdots & u_{i_{\sigma(1),j_1} } \\ u_{i,j} & u_{i,j_t} & \cdots & u_{i,j_1} \end{vmatrix} \\ &= (-1)^{t+1} \det(J)^2 \begin{vmatrix} u_{i,j_1} & \cdots & u_{i,j_t} & u_{i,j} \\ u_{i_{\sigma(1),j_1} } & \cdots & u_{i_{\sigma(1),j_t} } & u_{i_{\sigma(1),j} } \\ \vdots & & \vdots & \vdots \\ u_{i_{\sigma(t),j_1} } & \cdots & u_{i_{\sigma(t),j_t} } & u_{i_{\sigma(t),j} } \end{vmatrix} = (-1)^{t+1} \Delta_{i,j}^{\mathcal{E}(D)}(u), \end{aligned}$$

as required.

On the other hand, suppose that $\mathfrak{u} \leq \mathfrak{sp}_{2n}(q)$ and $j = -k$ for some $1 \leq k \leq n$. By setting $i_{\sigma(0)} = i$ and $j_{t+1} = j$, let $0 \leq s \leq t$ and $1 \leq s' \leq t$ be such that $i_{\sigma(s)} \preceq n \prec i_{\sigma(s+1)}$ and $j_{s'} \preceq n \prec j_{s'+1}$. Since $j_1 \prec j_2 \prec \dots \prec j_{s'} \preceq n \prec i_{\sigma(s+1)} \prec \dots \prec i_{\sigma(t)}$, we have

$u_{-j_b, -i_{\sigma(a)}} = u_{i_{\sigma(a)}, j_b} = 0$ for all $s < a \leq t$ and all $1 \leq b \leq s'$, and thus

$$\Delta_{-j, -i}^{\mathcal{E}(D)}(u) = \begin{vmatrix} u_{-j, -i_{\sigma(t)}} & \cdots & u_{-j, -i_{\sigma(s+1)}} & u_{-j, -i_{\sigma(s)}} & \cdots & u_{-j, -i_{\sigma(1)}} & u_{-j, -i} \\ u_{-j_t, -i_{\sigma(t)}} & \cdots & u_{-j_t, -i_{\sigma(s+1)}} & u_{-j_t, -i_{\sigma(s)}} & \cdots & u_{-j_t, -i_{\sigma(1)}} & u_{-j_t, -i} \\ \vdots & & \vdots & \vdots & & \vdots & \vdots \\ u_{-j_{s'+1}, -i_{\sigma(t)}} & \cdots & u_{-j_{s'+1}, -i_{\sigma(s+1)}} & u_{-j_{s'+1}, -i_{\sigma(s)}} & \cdots & u_{-j_{s'+1}, -i_{\sigma(1)}} & u_{-j_{s'+1}, -i} \\ 0 & \cdots & 0 & u_{-j_{s'}, -i_{\sigma(s)}} & \cdots & u_{-j_{s'}, -i_{\sigma(1)}} & u_{-j_{s'}, -i} \\ \vdots & & \vdots & \vdots & & \vdots & \vdots \\ 0 & \cdots & 0 & u_{-j_1, -i_{\sigma(s)}} & \cdots & u_{-j_1, -i_{\sigma(1)}} & u_{-j_1, -i} \end{vmatrix}.$$

Since $\mathbf{u} \leq \mathfrak{sp}_{2n}(q)$, we deduce that

$$\begin{aligned} \Delta_{-j, -i}^{\mathcal{E}(D)}(u) &= \begin{vmatrix} -u_{i_{\sigma(t)}, j} & \cdots & -u_{i_{\sigma(s+1)}, j} & u_{\sigma(s), j} & \cdots & u_{i_{\sigma(1)}, j} & u_{i, j} \\ -u_{i_{\sigma(t)}, j_t} & \cdots & -u_{i_{\sigma(s+1)}, j_t} & u_{i_{\sigma(s)}, j_t} & \cdots & u_{i_{\sigma(1)}, j_t} & u_{i, j_t} \\ \vdots & & \vdots & \vdots & & \vdots & \vdots \\ -u_{i_{\sigma(t)}, j_{s'+1}} & \cdots & -u_{i_{\sigma(s+1)}, j_{s'+1}} & u_{i_{\sigma(s)}, j_{s'+1}} & \cdots & u_{i_{\sigma(1)}, j_{s'+1}} & u_{i, j_{s'+1}} \\ 0 & \cdots & 0 & -u_{i_{\sigma(s)}, j_{s'}} & \cdots & -u_{i_{\sigma(1)}, j_{s'}} & -u_{i, j_{s'}} \\ \vdots & & \vdots & \vdots & & \vdots & \vdots \\ 0 & \cdots & 0 & -u_{i_{\sigma(s)}, j_1} & \cdots & -u_{i_{\sigma(1)}, j_1} & -u_{i, j_1} \end{vmatrix} \\ &= (-1)^{t-s+s'} \begin{vmatrix} u_{i_{\sigma(t)}, j} & \cdots & u_{i_{\sigma(s+1)}, j} & u_{\sigma(s), j} & \cdots & u_{i_{\sigma(1)}, j} & u_{i, j} \\ u_{i_{\sigma(t)}, j_t} & \cdots & u_{i_{\sigma(s+1)}, j_t} & u_{i_{\sigma(s)}, j_t} & \cdots & u_{i_{\sigma(1)}, j_t} & u_{i, j_t} \\ \vdots & & \vdots & \vdots & & \vdots & \vdots \\ u_{i_{\sigma(t)}, j_{s'+1}} & \cdots & u_{i_{\sigma(s+1)}, j_{s'+1}} & u_{i_{\sigma(s)}, j_{s'+1}} & \cdots & u_{i_{\sigma(1)}, j_{s'+1}} & u_{i, j_{s'+1}} \\ 0 & \cdots & 0 & u_{i_{\sigma(s)}, j_{s'}} & \cdots & u_{i_{\sigma(1)}, j_{s'}} & u_{i, j_{s'}} \\ \vdots & & \vdots & \vdots & & \vdots & \vdots \\ 0 & \cdots & 0 & u_{i_{\sigma(s)}, j_1} & \cdots & u_{i_{\sigma(1)}, j_1} & u_{i, j_1} \end{vmatrix}. \end{aligned}$$

Arguing as above (transposing and conjugating by the matrix $J = J_{t+1}$), we conclude that

$$\Delta_{-j, -i}^{\mathcal{E}(D)}(u) = (-1)^{t-s+s'} \Delta_{i, j}^{\mathcal{E}(D)}(u),$$

and the result follows because $t - s + s' = |D'(i, j)|$. \square

We are now able to prove the following result.

Proposition 3.4. *Let $u \in \mathbf{u}$ be arbitrary, and (\mathcal{D}, φ) be the (unique) basic pair for $U_m(q)$ such that $u \in V_{\mathcal{D}, \varphi}$. Then, $e_{\mathcal{D}, \varphi} \in \mathbf{u}$; in particular, there exists a unique basic pair (D, ϕ) for U such that $\mathcal{D} = \mathcal{E}(D)$ and $e_{\mathcal{D}, \varphi} = e_{D, \phi}$.*

Proof. It is enough to show that, for all $(i, j) \in \mathcal{D} \cap \mathcal{E}^+$, we have

$$\varphi(i, j) = (-1)^{\varepsilon_j} \varphi(-j, -i)$$

where

$$\varepsilon_j = \begin{cases} 1, & \text{if, either } \mathbf{u} \not\leq \mathfrak{sp}_{2n}(q), \text{ or } \mathbf{u} \leq \mathfrak{sp}_{2n}(q) \text{ and } j \leq n, \\ 0, & \text{if } \mathbf{u} \leq \mathfrak{sp}_{2n}(q) \text{ and } -j \leq n. \end{cases}$$

To prove this, we proceed by induction on $|\mathcal{D}|$. We consider the total order \preceq on the set of entries \mathcal{E} defined as follows: for all $(i, j), (k, l) \in \mathcal{E}$, we set

$$(3d) \quad (i, j) \prec (k, l) \iff \text{either } j \prec l, \text{ or } j = l \text{ and } k \prec i.$$

Let $(i, j) \in \mathcal{D} \cap \mathcal{E}^+$ be the smallest entry satisfying $\varphi(i, j) \neq (-1)^{\varepsilon_j} \varphi(-j, -i)$ (if it exists), let

$$\mathcal{D}(i, j) = \{(i_1, j_1), \dots, (i_t, j_t)\}$$

with $j_1 \prec \dots \prec j_t$, and let $\sigma \in S_t$ such that $i_{\sigma(1)} \prec \dots \prec i_{\sigma(t)}$. Since $(r, s) \prec (i, j)$, we have $\varphi(r, s) = (-1)^{\varepsilon_s} \varphi(-s, -r)$ for all $(r, s) \in \mathcal{D}(i, j) \cap \mathcal{E}^+$. Using (3a) and (3b), it is easy to conclude that

$$\mathcal{D}(-j, -i) = \{(-j_1, -i_1), \dots, (-j_t, -i_t)\}.$$

Let $\mathcal{D}_0 = \mathcal{D}(i, j) \cup \mathcal{D}(-j, -i)$, and $\varphi_0: \mathcal{D}_0 \rightarrow \mathbb{F}_q^\times$ be the restriction of φ to \mathcal{D}_0 . Then, by induction, the element

$$e_{\mathcal{D}_0, \varphi_0} = \sum_{(r, s) \in \mathcal{D}_0} \varphi(r, s) e_{r, s}$$

lies in \mathbf{u} .

On the other hand, let $c_{-j, -i} \in \mathbb{F}_q$ denote the (i, j) th coefficient of $e_{\mathcal{D}, \varphi}$; hence,

$$c_{-j, -i} = \begin{cases} \varphi(-j, -i), & \text{if } (-j, -i) \in \mathcal{D}, \\ 0, & \text{if } (-j, -i) \notin \mathcal{D}. \end{cases}$$

By (3b), we have

$$\Delta_{-j, -i}^{\mathcal{D}}(e_{\mathcal{D}, \varphi}) = (-1)^t \operatorname{sgn}(\sigma) c_{-j, -i} \prod_{s=1}^t \varphi(-j_s, -i_s).$$

It is easy to show that

$$\prod_{s=1}^t \varphi(-j_s, -i_s) = (-1)^{r'} \prod_{s=1}^t \varphi(i_s, j_s)$$

where

$$r' = \begin{cases} |\mathcal{D}(i, j)|, & \text{if, either } \mathbf{u} \leq \mathfrak{o}_{2n}(q), \text{ or } \mathbf{u} \leq \mathfrak{o}_{2n+1}(q), \\ |\mathcal{D}'(i, j)|, & \text{if } \mathbf{u} \leq \mathfrak{sp}_{2n}(q), \end{cases}$$

and $\mathcal{D}'(i, j) = \mathcal{D}(i, j) \cap \{(i, j) \in \mathcal{E}^+ : j \leq n\}$. Thus, we deduce that

$$\Delta_{-j, -i}^{\mathcal{D}}(e_{\mathcal{D}, \varphi}) = (-1)^{r'} c_{-j, -i} \varphi(i, j)^{-1} \Delta_{i, j}^{\mathcal{D}}(e_{\mathcal{D}, \varphi}).$$

Since $u \in V_{\mathcal{D},\varphi}$, we have

$$\Delta_{i,j}^{\mathcal{D}}(u) = \Delta_{i,j}^{\mathcal{D}}(e_{\mathcal{D},\varphi}) = \Delta_{i,j}^{\mathcal{D}_0}(e_{\mathcal{D},\varphi})$$

and

$$\Delta_{-j,-i}^{\mathcal{D}}(u) = \Delta_{-j,-i}^{\mathcal{D}}(e_{\mathcal{D},\varphi}) = \Delta_{-j,-i}^{\mathcal{D}_0}(e_{\mathcal{D},\varphi});$$

we note that $(-j, -i) \in R(\mathcal{D})$ (by induction and by the choice of (i, j)). By the previous lemma, we conclude that

$$\Delta_{i,j}^{\mathcal{D}}(u) = (-1)^{r+1} \Delta_{-j,-i}^{\mathcal{D}}(e_{\mathcal{D},\varphi})$$

where

$$r = \begin{cases} r' - 1, & \text{if } \mathbf{u} \leq \mathfrak{sp}_{2n}(q) \text{ and } -j \leq n, \\ r', & \text{otherwise.} \end{cases}$$

It follows that

$$(-1)^{r'} c_{-j,-i} \varphi(i, j)^{-1} \Delta_{i,j}^{\mathcal{D}}(e_{\mathcal{D},\varphi}) = (-1)^{r'+1} \Delta_{i,j}^{\mathcal{D}}(e_{\mathcal{D},\varphi}).$$

Since $\Delta_{i,j}^{\mathcal{D}}(u) \neq 0$ (because $(i, j) \in \mathcal{D}$), we obtain

$$c_{-j,-i} = \begin{cases} \varphi(i, j), & \text{if } \mathbf{u} \leq \mathfrak{sp}_{2n}(q) \text{ and } -j \leq n, \\ -\varphi(i, j), & \text{otherwise.} \end{cases}$$

It follows that $(-j, -i) \in \mathcal{D}$ and that $c_{-j,-i} = \varphi(-j, -i)$. Moreover, we note that, in the orthogonal case, if $j = i$, we obtain $\varphi(i, -i) = -\varphi(i, -i)$, hence $(i, -i) \notin \mathcal{D}$.

The above contradicts the minimal choice of (i, j) , and thus we conclude that

$$\varphi(i, j) = (-1)^{\varepsilon_j} \phi_{\varepsilon}(-j, -i)$$

for all $(i, j) \in \mathcal{D} \cap \mathcal{E}^+$. The proof is complete. \square

As observed above, this concludes the proof of the following result; we recall that $O_{D,\phi} = \mathbf{u} \cap V_{D,\phi}$ (by the definition).

Theorem 3.5. *Let $u \in \mathbf{u}$ be arbitrary. Then, there exists a unique basic subset $D \subseteq \Phi$ and a unique map $\phi: D \rightarrow \mathbb{F}_q^\times$ such that $u \in O_{D,\phi}$. Thus, \mathbf{u} is the disjoint union*

$$\mathbf{u} = \bigcup_{D,\phi} O_{D,\phi}$$

where the union runs over all basic pairs (D, ϕ) for U . Moreover, we have

$$O_{D,\phi} = \{a \in \mathbf{u}: \Delta_{i,j}^{\mathcal{D}}(a) = \Delta_{i,j}^{\mathcal{D}}(e_{D,\phi}) \text{ for all } (i, j) \in R(D)\}$$

for every basic pair (D, ϕ) for U .

As a consequence of Theorem 3.5 and Lemma 3.2, we obtain the following main theorem.

Theorem 3.6. *Let $z \in U$ be arbitrary. Then, there exists a unique basic subset $D \subseteq \Phi$ and a unique map $\phi: D \rightarrow \mathbb{F}_q^\times$ such that $z \in K_{D,\phi}$. Thus, U is the disjoint union of all its superclasses; that is,*

$$U = \bigcup_{D,\phi} K_{D,\phi}$$

where the union runs over all basic pairs (D, ϕ) for U .

4. SUPERCLASS FUNCTIONS

In this section, we prove that every supercharacter is a ‘‘superclass function’’ of U ; by definition, a function $\eta: U \rightarrow \mathbb{C}^\times$ is said to be a *superclass function* if it takes a constant value on each superclass of U . In fact, we shall prove the following result (cf. [6, Theorem 3.1]).

Theorem 4.1. *Every supercharacter of U is a superclass function. Moreover, every superclass function on U is a linear combination of supercharacters; hence, the supercharacters of U form a basis for the complex vector space $\text{scf}(U)$ consisting of all superclass functions on U .*

Since every supercharacter is a product of elementary characters (by definition), it is enough to show that every elementary character of U takes a constant value on each superclass of U . In fact, the theorem above will follow from the following result. (Henceforth, for each $z \in U$, we denote by a_z the element of \mathfrak{u} given by Lemma 3.2.)

Proposition 4.2. *Let $\alpha \in \Phi$ and $r \in \mathbb{F}_q^\times$ be arbitrary. Let (D, ϕ) be a basic pair for U , and denote by $z_{D,\phi}$ the unique element of $z \in U$ with $a_z = e_{D,\phi}$. Then,*

$$\xi_{\alpha,r}(z) = \xi_{\alpha,r}(z_{D,\phi})$$

for all $z \in K_{D,\phi}$.

Proof (first part). Let $(i, j) \in \mathcal{E}^+(\alpha)$; hence, $1 \leq i \leq n$ and $i \prec j \preceq -i$. In this first part of the proof, we shall assume that $j \neq -i$ (in the case where $U \leq Sp_{2n}(q)$). Let $\zeta_{i,j,r}$ be the elementary character of $U_m(q)$ associated with (i, j) and r . We recall its definition (see [2, Lemma 3]). We consider the subgroup $U_{i,j} = \{x \in U_m(q) : x_{i,k} = 0 \text{ for all } i \prec k \prec j\}$ of $U_m(q)$, and the linear character $\mu_{i,j,r}: U_{i,j} \rightarrow \mathbb{C}^\times$ defined by $\mu_{i,j,r}(x) = \vartheta(rx_{i,j})$ for all $x \in U_{i,j}$. Then, $\zeta_{i,j,r}$ is defined to be the induced character $\zeta_{i,j,r} = (\mu_{i,j,r})^{U_m(q)}$.

By [6, Proposition 3.2], we have $\xi_{\alpha,r} = (\zeta_{i,j,r})_U$, and so $\xi_{\alpha,r}(z) = \zeta_{i,j,r}(z)$ for all $z \in U$. By [4, Proposition 5.1] (see also [11, Theorem 5.8], or [7, Theorem 2.2]), we have $\zeta_{i,j,r}(x) = \zeta_{i,j,r}(1 + e_{D,\phi})$ for all $x \in 1 + V_{D,\phi}$. In particular, we deduce that

$$\xi_{\alpha,r}(z) = \zeta_{i,j,r}(z) = \zeta_{i,j,r}(1 + e_{D,\phi}) = \xi_{\alpha,r}(z_{D,\phi})$$

for all $z \in K_{D,\phi}$; we recall that $z \in K_{D,\phi}$ if and only if $a_z \in V_{D,\phi}$ (by Lemma 3.2). \square

In order to complete the proof of Proposition 4.2, it remains to consider the case where $U \leq Sp_{2n}(q)$ and $\alpha = 2\varepsilon_i$ for some $1 \leq i \leq n$. In what follows, we will always assume that this is the case; moreover, the basic pair (D, ϕ) will be kept fixed. We prove some elementary auxiliary lemmas; we mention that similar results are valid in the general case (a proof can be found in the second's author PhD thesis [16]). The proof of the first lemma is straightforward.

Lemma 4.3. *Let $u \in O_{D,\phi}$ be arbitrary, and $(k, l) \in \mathcal{E}(D)$ be the smallest entry of $\mathcal{E}(D)$ (with respect to the total order \preceq on \mathcal{E} as defined in (3d)); hence, $1 \leq k \leq n$ and $k \prec l \preceq -k$. Then, there exists $x \in U$ such that $v = xux^{-1} \in O_{D,\phi}$ satisfies $v_{k',l} = v_{k,l'} = 0$ for all $1 \preceq k', l' \preceq -1$ with $k' \neq k$ and $l' \neq l$.*

As a consequence, we obtain the following result.

Corollary 4.4. *Let $\beta \in \Phi$ and $s \in \mathbb{F}_q^\times$ be arbitrary. Then, $O_{\beta,s} = \{x(se_\beta)x^{-1} : x \in U\}$ is the adjoint U -orbit which contains $se_\beta \in \mathfrak{u}$. Moreover, $K_{\beta,s} = \{xz_{\beta,s}x^{-1} : x \in U\}$ is the conjugacy class which contains the element $z_{\beta,s} = 1 + se_\beta \in U$. (In particular, we have $\xi_{\alpha,r}(z) = \xi_{\alpha,r}(z_{\beta,s})$ for all $z \in K_{\beta,s}$.)*

Proof. The first assertion is an immediate consequence of the previous lemma. For the second, we note that $xz_{\beta,s}x^{-1} = 1 + x(se_\beta)x^{-1} \in U \cap (1 + V_{\beta,s}) = K_{\beta,s}$. On the other hand, if $z \in K_{\beta,s}$, then $a_z \in O_{\beta,s}$ (by Lemma 3.2), and thus the mapping $z \mapsto a_z$ defines a bijection from $K_{\beta,s}$ to $O_{\beta,s}$. Therefore, $|K_{\beta,s}| = |O_{\beta,s}| = |\{1 + x(se_\beta)x^{-1} : x \in U\}|$, and the result follows. \square

We observe that, in the notation of the corollary, we have $z_{\beta,s} = z_{D,\phi}$ for $D = \{\beta\}$ and $\phi: D \rightarrow \mathbb{F}_q^\times$ defined by $\phi(\beta) = s$; hence, Proposition 4.2 is true whenever the basic subset D has a unique element. Therefore, we will assume that $|D| > 1$, and $\xi_{\alpha,r}(z) = \xi_{\alpha,r}(z_{D',\phi'})$ for all $z \in K_{D',\phi'}$ and every basic pair (D', ϕ') with $|D'| < |D|$. Proposition 4.2 will then follow by induction. However, we need a concrete formula for the values of an elementary character on any superclass.

As usual, we denote by \mathfrak{u}^* the dual vector space of \mathfrak{u} , and let $\{e_\alpha^* : \alpha \in \Phi\}$ be the \mathbb{F}_q -basis of \mathfrak{u}^* dual to the basis $\{e_\alpha : \alpha \in \Phi\}$ of \mathfrak{u} ; hence, $e_\alpha^*(e_\beta) = \delta_{\alpha,\beta}$ for all $\alpha, \beta \in \Phi$. For each $f \in \mathfrak{u}^*$, we define

$$u(f) = \sum_{\beta \in \Phi} u_\beta e_\beta^* \in \mathfrak{u}_{2n}(q)$$

where

$$u_\beta = \begin{cases} \frac{1}{2} f(e_\beta), & \text{if } \beta = \varepsilon_k \pm \varepsilon_l \text{ for } 1 \leq k < l \leq n, \\ f(e_\beta), & \text{if } \beta = 2\varepsilon_k \text{ for } 1 \leq k \leq n. \end{cases}$$

It is easy to see that $f(v) = \text{Tr}(u(f)^T v)$ for all $v \in \mathfrak{u}$, and that the mapping $f \mapsto u(f)$ defines a vector space isomorphism from \mathfrak{u}^* to \mathfrak{u} . Finally, we define the linear function $\hat{f} \in \mathfrak{u}_{2n}(q)^*$ by

$$\hat{f}(v) = \text{Tr}(u(f)^T v)$$

for all $v \in \mathfrak{u}_{2n}(q)$, and set

$$O_{\alpha,r}^* = \{f \in \mathfrak{u}^* : \hat{f} \in U_{2n}(q)(re_{i,-i}^*)U_{2n}(q)\}$$

where $(xgy)(a) = g(x^{-1}ay^{-1})$ for all $x, y \in U_{2n}(q)$, $g \in \mathfrak{u}_{2n}(q)^*$ and $a \in \mathfrak{u}_{2n}(q)$. By [6, Proposition 5.2], we know that

$$\xi_{\alpha,r}(z) = \frac{\xi_{\alpha,r}(1)}{|O_{\alpha,r}^*|} \sum_{f \in O_{\alpha,r}^*} \vartheta_f(a_z)$$

for all $z \in U$. Given any \mathbb{F}_q -vector space V and any linear map $f \in V^*$, we denote by ϑ_f the composite map $\vartheta \circ f: V \rightarrow \mathbb{C}^\times$; it is straightforward to check that ϑ_f is a linear character of the additive group V^+ and that $\text{Irr}(V^+) = \{\vartheta_f : f \in V^*\}$.

For our purposes, it is convenient to describe the subset $O_{\alpha,r}^* \subseteq \mathfrak{u}^*$ (and the elementary character $\xi_{\alpha,r}$) as follows. By [5, Corollary 5.3], we have $f \in O_{\alpha,r}^*$ if and only if the following holds:

- (i) $\hat{f}(e_{a,-b}) = 0$ for all $(a, -b) \in \mathcal{E}$ with $1 \leq a < i$ or $1 \leq b < i$;
- (ii) $\hat{f}(e_{i,-i}) = r$;
- (iii) $\begin{vmatrix} \hat{f}(e_{i,b}) & \hat{f}(e_{i,-i}) \\ \hat{f}(e_{a,b}) & \hat{f}(e_{a,-i}) \end{vmatrix} = 0$ for all $(a, b) \in \mathcal{E}$ with $1 \prec a \prec b \prec -i$.

On the other hand, for $u = u(f)$, we have

$$\hat{f}(e_{a,b}) = \text{Tr}(u^T e_{a,b}) = \text{Tr}(e_{a,b}^T u) = e_{a,b}^*(u) = u_{a,b}$$

for all $(a, b) \in \mathcal{E}$. Therefore, for any $(a, b) \in \mathcal{E}^+$, we deduce that

$$\hat{f}(e_{-b,-a}) = \begin{cases} \hat{f}(e_{a,b}), & \text{if } (a, b) \in \mathcal{E}^+(\varepsilon_a - \varepsilon_b), \\ -\hat{f}(e_{a,b}), & \text{if } (a, b) \in \mathcal{E}^+(\varepsilon_a + \varepsilon_{-b}). \end{cases}$$

In particular, for $i \prec a \prec -i$, we get

$$\hat{f}(e_{a,-i}) = \begin{cases} \hat{f}(e_{i,-a}), & \text{if } -n \preceq a \prec -i, \\ -\hat{f}(e_{i,-a}), & \text{if } i \prec a \preceq n, \end{cases}$$

and so the elements of $O_{\alpha,r}^*$ can be parametrized by the set \mathcal{C} consisting of all functions $c: \{a: i \prec a \prec -i\} \rightarrow \mathbb{F}_q$. In fact, for each $c \in \mathcal{C}$, there exists a unique linear function $f_c \in \mathfrak{u}^*$ such that $\hat{f}_c \in \mathfrak{u}_{2n}(q)^*$ satisfies $\hat{f}_c(e_{i,a}) = c_a$ for all $i \prec a \prec -i$; here, we write $c_a = c(a)$ for all $i \prec a \prec -i$. This concludes the proof of the following result.

Lemma 4.5. *Let $\alpha = 2\varepsilon_i \in \Phi$ for some $1 \leq i \leq n$, and $r \in \mathbb{F}_q^\times$. Then, in the notation as above, we have*

$$O_{\alpha,r}^* = \{f_c : c \in \mathcal{C}\}.$$

Moreover, the mapping $c \mapsto f_c$ defines a bijection from \mathcal{C} to $O_{\alpha,r}$, and

$$\xi_{\alpha,r}(z) = \frac{1}{q^{n-i}} \sum_{c \in \mathcal{C}} \vartheta_{f_c}(a_z)$$

for all $z \in U$.

Henceforth, we set $\vartheta_c = \vartheta_{f_c}$ for all $c \in \mathcal{C}$, and consider the function $\vartheta: \mathbf{u} \rightarrow \mathbb{C}$ defined by

$$\vartheta(u) = \sum_{c \in \mathcal{C}} \vartheta_c(u)$$

for all $u \in \mathbf{u}$. Since $O_{\alpha,r}^* \subseteq \mathbf{u}^*$ is U -invariant (for the natural action given by conjugation), we clearly have $\vartheta(xux^{-1}) = \vartheta(u)$ for all $u \in \mathbf{u}$. Let $u \in O_{D,\phi}$ be arbitrary, and $(k, l) \in \mathcal{E}$ be the smallest entry of $\mathcal{E}(D)$ (with respect to the order \preceq defined in (3d)); hence, we must have $1 \leq k \leq n$ and $k \prec l \preceq -k$. Let $\beta \in \Phi$ be such that $(k, l) \in \mathcal{E}(\beta)$ (hence, $\beta \in D$), and $s = \phi(\beta)$. By Lemma 3.3, there exists $x \in U$ such that $v = xux^{-1}$ satisfies $v_{a,l} = v_{k,b} = 0$ for all $1 \preceq a, b \preceq -1$ with $a \neq k$ and $b \neq l$, and thus

$$\vartheta(u) = \vartheta(v) = \sum_{c \in \mathcal{C}} \vartheta_c(se_\beta + w) = \sum_{c \in \mathcal{C}} \vartheta_c(se_\beta) \vartheta_c(w)$$

where $w = v - se_\beta \in \mathbf{u}$. Next, we consider the relative positions of the entries (k, l) and $(i, -i)$. There are four distinct cases.

CASE 1 ($k = i$ and $l \prec -i$). In this case, $f_c(e_\beta) = \hat{f}_c(e_{i,l} \pm e_{-l,-i}) = 2\hat{f}(e_{i,l}) = 2c_l$ for all $c \in \mathcal{C}$, and so

$$\vartheta(u) = \sum_{c \in \mathcal{C}} \vartheta(2c_l s) \vartheta_c(w).$$

On the other hand, we have $w_\gamma = w_{a,b}$ whenever $\gamma \in \Phi$ and $(a, b) \in \mathcal{E}^+(\gamma)$, and thus

$$(4a) \quad f_c(w) = \sum_{\gamma \in \Phi} w_\gamma f_c(e_\gamma) = rw_{i,-i} + \sum_{i \prec b \prec -i} c_b w_{i,b} + \sum_{i \prec b \prec -i} \sum_{i \prec a \preceq -b} r^{-1} c_{-a} c_b w_{a,b}.$$

In fact, for all $c \in \mathcal{C}$ and all $\beta \in \Phi$, we have

$$f_c(e_\beta) = \hat{f}_c(e_\beta) = \begin{cases} r, & \text{if } \beta = 2\varepsilon_i, \\ 2c_b, & \text{if } (i, b) \in \mathcal{E}^+(\beta) \text{ and } b \neq -i, \\ 2r^{-1}c_{-a}c_b, & \text{if } (a, b) \in \mathcal{E}^+(\beta) \text{ for } i < a, \\ 0, & \text{otherwise.} \end{cases}$$

Since $w_{a,l} = v_{a,l} = 0$ for all $i \prec a \preceq -i$, the coordinate c_l of any $c \in \mathcal{C}$ does not occur in the expression of $f_c(w)$ given by (4a), and so

$$\vartheta(u) = \left(\sum_{t \in \mathbb{F}_q} \vartheta(2ts) \right) \left(\sum_{c \in \mathcal{C}_1} \vartheta_c(w) \right) = \left(\sum_{t \in \mathbb{F}_q} \vartheta_t(2s) \right) \left(\sum_{c \in \mathcal{C}_1} \vartheta_c(w) \right)$$

where $\mathcal{C}_1 = \{c \in \mathcal{C} : c_l = 0\}$, and ϑ_t , for $t \in \mathbb{F}_q$, denotes the character of \mathbb{F}_q^+ defined by $\vartheta_t(a) = \vartheta(ta)$ for all $a \in \mathbb{F}_q$. Since $\sum_{t \in \mathbb{F}_q} \vartheta_t$ is the regular character of \mathbb{F}_q^+ , we conclude that $\vartheta(u) = 0$. Furthermore, we note that

$$\sum_{c \in \mathcal{C}_1} \vartheta_c(w) = q^{-1} \sum_{c \in \mathcal{C}} \vartheta_c(w)$$

(by the same reason as above).

CASE 2 ($k = i$ and $l = -i$). We have $f_c(e_\beta) = f_c(e_{i,-i}) = r$ for all $c \in \mathcal{C}$, and so

$$\vartheta(u) = \vartheta(rs) \sum_{c \in \mathcal{C}} \vartheta_c(w).$$

CASE 3 ($k < i$). We have $f_c(e_\beta) = 0$ for all $c \in \mathcal{C}$, and so

$$\vartheta(u) = \sum_{c \in \mathcal{C}} \vartheta_c(w).$$

CASE 4 ($k > i$). In this case, we have

$$f_c(e_\beta) = \begin{cases} r^{-1}c_{-k}c_l, & \text{if } l \neq -k, \\ r^{-1}c_{-k}^2, & \text{if } l = -k, \end{cases}$$

for all $c \in \mathcal{C}$.

On the one hand, suppose that $l \neq -k$. Then, the entries c_{-k} and c_l of any $c \in \mathcal{C}$ do not occur in $f_c(w)$ (see the argument in case 1), hence we get

$$\vartheta(u) = \left(\sum_{t, t' \in \mathbb{F}_q} \vartheta(r^{-1}stt') \right) \left(q^{-2} \sum_{c \in \mathcal{C}} \vartheta_c(w) \right).$$

Since $\sum_{t \in \mathbb{F}_q} \vartheta_t$ is the regular character of \mathbb{F}_q^+ , we conclude that

$$\sum_{t, t' \in \mathbb{F}_q} \vartheta(r^{-1}stt') = \sum_{t, t' \in \mathbb{F}_q} \vartheta_t(r^{-1}st') = q,$$

and thus

$$\vartheta(u) = q^{-1} \sum_{c \in \mathcal{C}} \vartheta_c(w).$$

On the other hand, suppose that $l = -k$. Then, the entry c_{-k} of any $c \in \mathcal{C}$ do not occur in $f_c(w)$, hence we get

$$\vartheta(u) = \left(\sum_{t \in \mathbb{F}_q} \vartheta(r^{-1}st^2) \right) \left(q^{-1} \sum_{c \in \mathcal{C}} \vartheta_c(w) \right).$$

Now, we recall that the *quadratic character* of \mathbb{F}_q is, by definition, the linear character η of the multiplicative group \mathbb{F}_q^\times defined by

$$\eta(c) = \begin{cases} 1, & \text{if } c \in (\mathbb{F}_q^\times)^2, \\ -1, & \text{otherwise,} \end{cases}$$

for all $c \in \mathbb{F}_q^\times$. Moreover, given any linear character ν of \mathbb{F}_q^\times and any linear character ϑ of \mathbb{F}_q^+ , the *Gauss sum* of ν and ϑ is defined by

$$G(\nu, \vartheta) = \sum_{c \in \mathbb{F}_q^\times} \nu(c)\vartheta(c).$$

The following result is Theorem 5.3.3 of the book [15].

Theorem 4.6. *Let ϑ be a non-trivial linear character of \mathbb{F}_q^+ , and*

$$h(T) = a_2T^2 + a_1T + a_0 \in \mathbb{F}_q[T]$$

be a polynomial over \mathbb{F}_q with $a_2 \neq 0$. Suppose that q is odd. Then,

$$\sum_{c \in \mathbb{F}_q} \vartheta(h(c)) = \vartheta(a_0 - a_1^2(4a_2)^{-1})\eta(a_2)G(\eta, \vartheta)$$

where η is the quadratic character of \mathbb{F}_q .

Applying this result to our situation (with $h(T) = r^{-1}sT^2$), we obtain

$$\vartheta(u) = q^{-1}\eta(r^{-1}s)G(\eta, \vartheta) \sum_{c \in \mathcal{C}} \vartheta_c(w).$$

It follows that, in any case, we have

$$\vartheta(u) = c_{\beta,s} \sum_{c \in \mathcal{C}} \vartheta_c(w)$$

for some constant $c_{\beta,s} \in \mathbb{C}$ depending only on the root $\beta \in D$ and on the value $s = \phi(\beta)$; in fact,

$$(4b) \quad c_{\beta,s} = \begin{cases} 0, & \text{if } k = i \text{ and } l \prec -i, \\ \vartheta(rs), & \text{if } k = i \text{ and } l = -i, \\ 1, & \text{if } k < i, \\ q^{-1}, & \text{if } k > i \text{ and } l \neq -k, \\ q^{-1}\eta(r^{-1}s)G(\eta, \vartheta), & \text{if } k > i \text{ and } l = -k. \end{cases}$$

We are now able to conclude the proof of Proposition 4.2.

Proof of Proposition 4.2 (second part). Let the notation be as above, and let $z' \in U$ be such that $a_{z'} = w$. Let $D' = D - \{\beta\}$, and $\phi' : D' \rightarrow \mathbb{F}_q^\times$ be the restriction on ϕ to D' . Then, it is easy to check that $w = v - se_\beta \in O_{D',\phi'}$, hence $z' \in K_{D',\phi'}$ (by Lemma 3.2). By induction, we have $\xi_{\alpha,r}(z') = \xi_{\alpha,r}(z_{D',\phi'})$. Since

$$\xi_{\alpha,r}(z') = \frac{1}{q^{n-i}} \sum_{c \in \mathcal{C}} \vartheta_c(w)$$

(by Lemma 4.5), we conclude that

$$\xi_{\alpha,r}(z) = \frac{c_{\beta,s}}{q^{n-i}} \sum_{c \in \mathcal{C}} \vartheta_c(w) = c_{\beta,s} \xi_{\alpha,r}(z') = c_{\beta,s} \xi_{\alpha,r}(z_{D',\phi'}).$$

Therefore, the value $\xi_{\alpha,r}(z)$ does not depend on $z \in K_{D,\phi}$, hence $\xi_{\alpha,r}(z) = \xi_{\alpha,r}(z_{D,\phi})$ for all $z \in K_{D,\phi}$. \square

We next proceed with the proof of Theorem 4.1; a slightly different proof will be given later without reference to the results of [6].

Proof of Theorem 4.1. Since every supercharacter is a product of elementary characters, Proposition 4.2 implies that every supercharacter is a superclass function. By [6, Theorem 4.2], the supercharacters are orthogonal, hence they are linearly independent functions of $\text{scf}(U)$. Since the dimension of the vector space $\text{scf}(U)$ equals the number of basic pairs (D, ϕ) for U , we conclude that the supercharacters form a basis of $\text{scf}(U)$, and this completes the proof. \square

We now observe that, since the regular character of U is clearly a superclass function, Theorem 4.1 implies that it is a linear combination of supercharacters. In particular, we obtain the following result (and also an alternative proof of [6, Theorem 3.2]).

Theorem 4.7. *Every irreducible character is a constituent of a (unique) supercharacter.*

Proof. It is enough to observe that every irreducible character of U is a constituent of the regular character. (The unicity follows by the orthogonality of supercharacters; see [6, Theorem 4.2].) \square

Finally, an easy calculation proves the following result (and gives an alternative proof of [6, Theorem 5.2]).

Theorem 4.8. *Let ρ_U be the regular character of U . Then,*

$$\rho_U = \sum_{D,\phi} \frac{\xi_{D,\phi}(1)}{\langle \xi_{D,\phi}, \xi_{D,\phi} \rangle} \xi_{D,\phi}$$

where the sum is over all basic pairs (D, ϕ) .

Proof. Let $\rho_U = \sum_{D,\phi} m_{D,\phi} \xi_{D,\phi}$ where $m_{D,\phi} \in \mathbb{C}$ for all basic pairs (D, ϕ) . Since supercharacters are orthogonal, we obtain $\langle \rho_U, \xi_{D,\phi} \rangle = m_{D,\phi} \langle \xi_{D,\phi}, \xi_{D,\phi} \rangle$. On the other hand, let $\text{Irr}_{D,\phi}(U)$ denote the subset of $\text{Irr}(U)$ consisting of all irreducible constituents of the supercharacter $\xi_{D,\phi}$; hence, we have a disjoint union $\text{Irr}(U) = \bigcup_{D,\phi} \text{Irr}_{D,\phi}(U)$, and

$$\xi_{D,\phi} = \sum_{\chi \in \text{Irr}_{D,\phi}(U)} \langle \chi, \xi_{D,\phi} \rangle \chi.$$

Since $\rho_U = \sum_{\chi \in \text{Irr}(U)} \chi(1) \chi$, we deduce that

$$m_{D,\phi} \langle \xi_{D,\phi}, \xi_{D,\phi} \rangle = \langle \rho_U, \xi_{D,\phi} \rangle = \sum_{\chi \in \text{Irr}_{D,\phi}(U)} \chi(1) \langle \chi, \xi_{D,\phi} \rangle = \xi_{D,\phi}(1),$$

and the result follows. \square

5. SUPERCHARACTER VALUES

In this section, we obtain explicit formulae that allows to determine the constant value $\xi_{D,\phi}(z_{D',\phi'})$ of the supercharacter $\xi_{D,\phi}$ on the superclass $K_{D',\phi'}$. Since $\xi_{D,\phi}$ is a product of elementary characters, it is enough to determine the value of an arbitrary elementary character $\xi_{\alpha,r}$, for $\alpha \in \Phi$ and $r \in \mathbb{F}_q^\times$, on any superclass. Let $(i, j) \in \mathcal{E}$, and consider be the elementary character $\zeta_{i,j,r}$ of $U_m(q)$. By [4, Proposition 5.1] (see also [7, Theorem 2.2]), $\zeta_{i,j,r}$ is constant on the superclasses of $U_m(q)$, and its value on the superclass associated with a basic pair (\mathcal{D}, φ) equals

$$(5a) \quad \zeta_{i,j,r}(1 + e_{\mathcal{D},\varphi}) = \begin{cases} q^{-t} \zeta_{i,j,r}(1) \vartheta(r\varphi(i, j)), & \text{if } (i, j) \in \mathcal{D}, \\ q^{-t} \zeta_{i,j,r}(1), & \text{if } (i, j) \in R(\mathcal{D}) - \mathcal{D}, \\ 0, & \text{otherwise,} \end{cases}$$

where $t = |\{(k, l) \in \mathcal{D} : i \prec k \prec l \prec j\}|$. (We note that $q^{-t} \zeta_{i,j,r}(1) = q^{t'}$ where t' is the number of \mathcal{D} -regular entries which are directly below the entry (i, j) .)

Using [6, Proposition 3.2], we easily deduce the following result. For simplicity of writing, for any basic subset $D \subseteq \Phi$, we define

$$R(D) = \{\beta \in \Phi : \mathcal{E}(\beta) \subseteq R(\mathcal{E}(D))\},$$

and observe that, for any root $\beta \in \Phi$, we have

$$\mathcal{E}(\beta) \subseteq R(\mathcal{E}(D)) \iff \mathcal{E}(\beta) \cap R(\mathcal{E}(D)) \neq \emptyset;$$

in fact, an entry $(k, l) \in \mathcal{E}$ is D -regular if and only if $(-l, -k)$ is also D -regular. Further, for any root $\alpha \in \Phi$, we set

$$D(\alpha) = \{(k, l) \in \mathcal{E}(D) : i \prec k \prec l \prec j\}$$

where $(i, j) \in \mathcal{E}^+(\alpha)$.

Proposition 5.1. *Let $\alpha \in \Phi$, and suppose that $\alpha \neq 2\varepsilon_i$ for $1 \leq i \leq n$ (in the case where $U \leq Sp_{2n}(q)$). Let $r \in \mathbb{F}_q^\times$, and (D', ϕ') be a basic pair for U . Then,*

$$\xi_{\alpha,r}(z_{D',\phi'}) = \begin{cases} q^{-t(\alpha,D')} \xi_{\alpha,r}(1) \vartheta(r\phi'(\alpha)), & \text{if } \alpha \in D', \\ q^{-t(\alpha,D')} \xi_{\alpha,r}(1), & \text{if } \alpha \in R(D') - D', \\ 0, & \text{otherwise,} \end{cases}$$

where $t(\alpha, D') = |D'(\alpha)|$.

Proof. It is enough to observe that $\xi_{\alpha,r} = (\zeta_{i,j,r})_U$ for $(i, j) \in \mathcal{E}(\alpha)$ (by [6, Proposition 3.2]), and thus

$$\xi_{\alpha,r}(z_{D,\phi}) = \zeta_{i,j,r}(z_{D,\phi}) = \zeta_{i,j,r}(1 + e_{D,\phi}).$$

The result follows by (5a) because $e_{D,\phi} = e_{\mathcal{E}(D),\varphi}$ for a (uniquely determined) map $\varphi: \mathcal{E}(D) \rightarrow \mathbb{F}_q^\times$ (by Proposition 3.4). \square

Next, we consider the case where $U \leq Sp_{2n}(q)$ and $\alpha = 2\varepsilon_i$ for some $1 \leq i \leq n$. Let (D, ϕ) be a basic pair for U . Let $(k, l) \in \mathcal{E}$ be the smallest entry of $\mathcal{E}(D)$ (with respect to the order \preceq defined in (3d)), and $\beta \in D$ be such that $(k, l) \in \mathcal{E}(\beta)$. Let $s = \phi(\beta)$, $D' = D - \{\beta\}$, and $\phi': D' \rightarrow \mathbb{F}_q^\times$ be the restriction of ϕ to D' . We recall from the proof of Proposition 4.2 that $\xi_{\alpha,r}(z_{D,\phi}) = c_{\beta,s} \xi_{\alpha,r}(z_{D',\phi'})$ where $c_{\beta,s} \in \mathbb{C}$ is given by (4b). In particular, for $D = \{\beta\}$, we have $z_{D',\phi'} = 1$, hence $\xi_{\alpha,r}(z_{\beta,s}) = c_{\beta,s} \xi_{\alpha,r}(1) = q^{n-i} c_{\beta,s}$. In the general situation, we get

$$\xi_{\alpha,r}(z_{D,\phi}) = \xi_{\alpha,r}(1) \prod_{\beta \in D} c_{\beta,\phi(\beta)} = q^{n-i} \prod_{\beta \in D} c_{\beta,\phi(\beta)},$$

and so we obtain the following formulae.

Proposition 5.2. *Suppose that $U \leq Sp_{2n}(q)$, let $\alpha = 2\varepsilon_i$ for some $1 \leq i \leq n$, and let (D', ϕ') be a basic pair for U . Moreover, let η be the quadratic character of \mathbb{F}_q , let $G(\eta, \vartheta)$ be the Gauss sum of η and ϑ . Then,*

$$\xi_{\alpha,r}(z_{D',\phi'}) = \begin{cases} q^{-t(\alpha,D')} \xi_{\alpha,r}(1) c_{\alpha,r}^{D',\phi'} \vartheta(r\phi'(\alpha)), & \text{if } \alpha \in D', \\ q^{-t(\alpha,D')} \xi_{\alpha,r}(1) c_{\alpha,r}^{D',\phi'}, & \text{if } \alpha \in R(D') - D', \\ 0, & \text{otherwise,} \end{cases}$$

where $t(\alpha, D') = |D'(\alpha)|$, and

$$c_{\alpha,r}^{D',\phi'} = q^{\frac{1}{2}(t(\alpha,D') - t_0(\alpha,D'))} G(\eta, \vartheta)^{t_0(\alpha,D')} \prod_{\beta \in D'_0(\alpha)} \eta(r^{-1}\phi'(\beta))$$

for $D'_0(\alpha) = D' \cap \{2\varepsilon_k: i < k \leq n\}$ and $t_0(\alpha, D') = |D'_0(\alpha)|$.

As an immediate consequence of Propositions 5.1 and 5.2, we obtain following general formula for the constant value $\xi_{D,\phi}^{D',\phi'} = \xi_{D,\phi}(z_{D',\phi'})$ of the supercharacter $\xi_{D,\phi}$ on the superclass $K_{D',\phi'}$ (see [7, Theorem 2.2] for the corresponding result in the case of the

unitriangular group). As in the previous proposition, given any basic subset $D \subseteq \Phi$, we define

$$D_0 = D \cap \{2\varepsilon_i : 1 \leq i \leq n\}, \quad \text{and} \quad D_0(\alpha) = D \cap \{2\varepsilon_k : i < k \leq n\}$$

whenever $\alpha = 2\varepsilon_i \in \Phi$ for $1 \leq i \leq n$.

Theorem 5.3. *Let (D, ϕ) and (D', ϕ') be basic pairs for U . Then,*

$$\xi_{D, \phi}(z_{D', \phi'}) = \begin{cases} q^{-t(D, D')} \xi_{D, \phi}(1) c_{D, \phi}^{D', \phi'} \prod_{\alpha \in D \cap D'} \vartheta(\phi(\alpha) \phi'(\alpha)), & \text{if } D \subseteq R(D'), \\ 0, & \text{otherwise,} \end{cases}$$

where $t(D, D') = \sum_{\alpha \in D} |D'(\alpha)|$, and

$$c_{D, \phi}^{D', \phi'} = q^{\frac{1}{2}(t(D, D') - t_0(D, D'))} G(\eta, \vartheta)^{t_0(D, D')} \prod_{\alpha \in D_0} \prod_{\beta \in D'_0(\alpha)} \eta(\phi(\alpha)^{-1} \phi'(\beta))$$

for $t_0(D, D') = \sum_{\alpha \in D_0} |D'_0(\alpha)|$.

We observe that, in the case where, either $U \leq O_{2n}(q)$, or $U \leq O_{2n+1}(q)$, the set D_0 is empty, and thus $c_{D, \phi}^{D', \phi'} = 1$. Moreover, in any case, we have $q^{-t(D, D')} \xi_{D, \phi}(1) = q^{t'(D, D')}$ where $t(D, D')$ is the number of D' -regular entries which are directly below the entries in $\mathcal{E}^+(D)$.

We conclude this section with a consequence of Theorem 5.3, proving that every superclass factorizes uniquely as a product (in any order) of ‘‘elementary’’ superclasses; we recall that a similar factorization is valid for supercharacters.

Theorem 5.4. *Let (D, ϕ) be a basic pair for U . Then,*

$$K_{D, \phi} = \prod_{\alpha \in D} K_{\alpha, \phi(\alpha)}$$

where the product can be taken in any order.

Proof. We order the roots according to the total order \preceq defined as follows: given $\alpha, \beta \in \Phi$, let $(i, j) \in \mathcal{E}^+(\alpha)$ and $(k, l) \in \mathcal{E}^+(\beta)$, and define $\alpha \preceq \beta$ if and only if $(i, j) \preceq (k, l)$ (with respect to the order on \mathcal{E} as defined in (3d)). On the one hand, let $z \in \prod_{\alpha \in D} K_{\alpha, \phi(\alpha)}$ be arbitrary, and (D', ϕ') be the unique basic pair with $z \in K_{D', \phi'}$. Suppose that $(D', \phi') \neq (D, \phi)$, and let $\alpha \in \Phi$ be the smallest root in $D \cup D'$ such that, either $\alpha \notin D \cap D'$, or $\phi(\alpha) \neq \phi'(\alpha)$. We consider the elementary character $\xi_{\alpha, r}$ for any $r \in \mathbb{F}_q^\times$. By Propositions 5.1 and 5.2, we have

$$\xi_{\alpha, r}(z) = \begin{cases} c_{D, \phi} \vartheta(r \phi(\alpha)), & \text{if } \alpha \in D, \\ c_{D', \phi'} \vartheta(r \phi'(\alpha)), & \text{if } \alpha \in D', \end{cases}$$

where $c_{D, \phi}, c_{D', \phi'} \in \mathbb{C}$ are non-zero constants depending only on roots $\gamma \in D \cup D'$ with $\gamma \prec \alpha$; moreover, by the choice of α , we have $c_{D, \phi} = c_{D', \phi'}$. Since $\vartheta(s) \neq 0$ for all $s \in \mathbb{F}_q$, we conclude that $\xi_{\alpha, r}(z) \neq 0$, and thus $\alpha \in R(D) \cap R(D')$ (again by Propositions 5.1 and

5.2). Now, suppose that $\alpha \in D - D'$. Then, since $\alpha \in R(D')$, we have $\xi_{\alpha,r}(z) = c_{D',\phi'} = c_{D,\phi}$, and thus $\vartheta(r\phi(\alpha)) = 1$. Since $r \in \mathbb{F}_q^\times$ is arbitrary, we conclude that

$$q = \sum_{r \in \mathbb{F}_q} \vartheta(r\phi(\alpha)) = \rho(\phi(\alpha))$$

where $\rho = \sum_{r \in \mathbb{F}_q} \vartheta_r$ is the regular character of \mathbb{F}_q^+ . It follows that $\phi(\alpha) = 0$, a contradiction. Similarly, we obtain a contradiction assuming that $\alpha \in D' - D$, and thus $\alpha \in D \cap D'$. Thus, we get $\vartheta(r\phi(\alpha)) = \vartheta(r\phi'(\alpha))$ for all $r \in \mathbb{F}_q^\times$, and the argument used above shows that $\phi(\alpha) = \phi'(\alpha)$. This final contradiction implies that $(D', \phi') = (D, \phi)$, and thus

$$\prod_{\alpha \in D} K_{\alpha, \phi(\alpha)} \subseteq K_{D, \phi}.$$

On the other hand, for the reverse inclusion, we consider the complex group algebra $\mathbb{C}U$ of U , and the superclass sum

$$\hat{K} = \sum_{z \in K} z \in \mathbb{C}U$$

associated with a superclass $K \subseteq U$. By [11, Corollary 2.3], the product $\prod_{\alpha \in D} \hat{K}_{\alpha, \phi(\alpha)}$ is a linear combination with nonnegative integer coefficients of the superclass sums of U . By the above, we easily conclude that $\prod_{\alpha \in D} \hat{K}_{\alpha, \phi(\alpha)}$ is an integer multiple of $\hat{K}_{D, \phi}$, and so

$$K_{D, \phi} \subseteq \prod_{\alpha \in D} K_{\alpha, \phi(\alpha)},$$

as required. □

6. THE SUPERCHARACTER TABLE

Theorem 4.1 allows the definition of the *supercharacter table* of U as the square (complex) matrix \mathcal{T} having rows and columns indexed by all the basic pairs for U , and where the coefficient corresponding to the basic pairs (D, ϕ) and (D', ϕ') is the constant value $\xi_{D, \phi}(z_{D', \phi'})$ of the supercharacter $\xi_{D, \phi}$ on the superclass $K_{D', \phi'}$. Since supercharacters are orthogonal (by [6, Theorem 5.4]), the rows of \mathcal{T} are orthogonal (for the usual inner product). In fact, in what follows, we prove various orthogonality relations (which are similar to the well-known for irreducible characters and conjugacy classes).

We start by considering the convolution product of supercharacters; we recall that, for any functions $\zeta, \eta: U \rightarrow \mathbb{C}$, the convolution product of ζ and η is the function $\zeta \star \eta: U \rightarrow \mathbb{C}$ defined by

$$(\zeta \star \eta)(z) = \sum_{x \in U} \zeta(x)\eta(zx^{-1})$$

for all $z \in U$. As an example, it is well-know that

$$(6a) \quad \chi \star \chi' = \delta_{\chi, \chi'} \frac{|U|}{\chi(1)} \chi$$

for all $\chi, \chi' \in \text{Irr}(U)$ (see, for example, [12, Theorem 2.13]); we observe that this corresponds to the generalized orthogonality relations for irreducible characters:

$$\frac{1}{|U|} \sum_{x \in U} \chi(x) \chi'(zx^{-1}) = \delta_{\chi, \chi'} \frac{\chi(z)}{\chi(1)}$$

for all $\chi, \chi' \in \text{Irr}(U)$ and all $z \in U$. For supercharacters, we obtain the following similar result.

Theorem 6.1 (Generalized orthogonality relations for supercharacters). *Let (D, ϕ) and (D', ϕ') be basic pairs for U . Then*

$$\xi_{D, \phi} \star \xi_{D', \phi'} = \delta_{D, D'} \delta_{\phi, \phi'} \frac{|U| \langle \xi_{D, \phi}, \xi_{D, \phi} \rangle}{\xi_{D, \phi}(1)} \xi_{D, \phi}.$$

In other words, we have

$$\frac{1}{|U|} \sum_{x \in U} \xi_{D, \phi}(x) \xi_{D', \phi'}(zx^{-1}) = \delta_{D, D'} \delta_{\phi, \phi'} \frac{\langle \xi_{D, \phi}, \xi_{D, \phi} \rangle \xi_{D, \phi}(z)}{\xi_{D, \phi}(1)}$$

for all $z \in U$.

Proof. By Theorem 5.4, we deduce that

$$\chi(1) = \langle \chi, \rho_U \rangle = \frac{\xi_{D, \phi}(1)}{\langle \xi_{D, \phi}, \xi_{D, \phi} \rangle} \langle \chi, \xi_{D, \phi} \rangle,$$

and thus

$$\xi_{D, \phi} = \frac{\langle \xi_{D, \phi}, \xi_{D, \phi} \rangle}{\xi_{D, \phi}(1)} \sum_{\chi \in \text{Irr}_{D, \phi}(U)} \chi(1) \chi$$

for all basic pairs (D, ϕ) . Now, since supercharacters are orthogonal, (6a) implies that $\xi_{D, \phi} \star \xi_{D', \phi'} = 0$ for all basic pairs (D, ϕ) and (D', ϕ') with $(D, \phi) \neq (D', \phi')$. On the other hand, we obtain

$$\xi_{D, \phi} \star \xi_{D, \phi} = \frac{|U| \langle \xi_{D, \phi}, \xi_{D, \phi} \rangle^2}{\xi_{D, \phi}(1)^2} \sum_{\chi \in \text{Irr}_{D, \phi}} \chi(1) \chi = \frac{|U| \langle \xi_{D, \phi}, \xi_{D, \phi} \rangle}{\xi_{D, \phi}(1)} \xi_{D, \phi},$$

as required. \square

In particular, we deduce that the rows of the supercharacter table \mathcal{T} are orthogonal (an alternative proof can be easily obtained by evaluating the Frobenius scalar product).

Theorem 6.2 (First orthogonality relation for supercharacters). *Let (D', ϕ') and (D'', ϕ'') be basic pairs for U . Then,*

$$\sum_{D, \phi} \frac{|K_{D, \phi}|}{|U|} \xi_{D', \phi'}(z_{D, \phi}) \overline{\xi_{D'', \phi''}(z_{D, \phi})} = \delta_{D', D''} \delta_{\phi', \phi''} \langle \xi_{D', \phi'}, \xi_{D', \phi'} \rangle$$

where the sum is over all basic pairs (D, ϕ) for U .

Proof. Since $\xi_{D'', \phi''}(x^{-1}) = \overline{\xi_{D'', \phi''}(x)}$, the previous theorem gives

$$\sum_{x \in U} \xi_{D', \phi'}(x) \overline{\xi_{D'', \phi''}(x)} = \delta_{D', D''} \delta_{\phi', \phi''} |U| \langle \xi_{D', \phi'}, \xi_{D', \phi'} \rangle,$$

and the result now follows by Theorem 4.1. \square

Another consequence of the generalized orthogonality relations is the following result.

Theorem 6.3. *The space $\text{scf}(U)$ of superclass functions is a commutative semisimple algebra with respect to the convolution product.*

Proof. By Theorem 4.7, the convolution product of supercharacters is a superclass function (in fact, it is a multiple of a supercharacter), hence $\text{scf}(U)$ is a commutative algebra. Furthermore, $\text{scf}(U)$ has a basis of orthogonal idempotents, namely the functions

$$\zeta_{D, \phi} = \frac{\xi_{D, \phi}(1)}{|U| \langle \xi_{D, \phi}, \xi_{D, \phi} \rangle} \xi_{D, \phi}$$

for the basic pairs (D, ϕ) , and thus it is semisimple. \square

Finally, we deduce that the columns of the supercharacter table \mathcal{T} are also orthogonal.

Theorem 6.4 (Second orthogonality relation for supercharacters). *Let (D', ϕ') and (D'', ϕ'') be basic pairs for U . Then,*

$$\sum_{D, \phi} \frac{1}{\langle \xi_{D, \phi}, \xi_{D, \phi} \rangle} \xi_{D, \phi}(z_{D', \phi'}) \overline{\xi_{D, \phi}(z_{D'', \phi''})} = \delta_{D', D''} \delta_{\phi', \phi''} \frac{|U|}{|K_{D', \phi'}|}$$

where the sum is over all basic pairs (D, ϕ) for U .

Proof. For any basic pairs (D, ϕ) and (D', ϕ') , let

$$h_{D', \phi'}^{D, \phi} = \sqrt{\frac{|K_{D, \phi}|}{|U| \langle \xi_{D', \phi'}, \xi_{D', \phi'} \rangle}} \xi_{D', \phi'}(z_{D, \phi}),$$

and consider the square matrix $H = (h_{D', \phi'}^{D, \phi})$ with rows and columns indexed by the basic pairs. By the previous theorem, we have $U\bar{U}^T = I$, and thus $\bar{U}^T U = I$. It follows that

$$\sum_{D, \phi} h_{D', \phi'}^{D, \phi} \overline{h_{D'', \phi''}^{D, \phi}} = \delta_{D', D''} \delta_{\phi', \phi''},$$

as required. \square

REFERENCES

- [1] ANDRÉ, C. A. M. *Irreducible characters of the unitriangular group and coadjoint orbits*. PhD thesis, University of Warwick, 1992.
- [2] ANDRÉ, C. A. M. Basic characters of the unitriangular group. *J. Algebra* 175, 1 (1995), 287–319.
- [3] ANDRÉ, C. A. M. Basic sums of coadjoint orbits of the unitriangular group. *J. Algebra* 176, 3 (1995), 959–1000.
- [4] ANDRÉ, C. A. M. The basic character table of the unitriangular group. *J. Algebra* 241, 1 (2001), 437–471.
- [5] ANDRÉ, C. A. M., AND NETO, A. M. Super-characters of finite unipotent groups of types B_n , C_n and D_n . *J. Algebra* 305, 1 (2006), 394–429.
- [6] ANDRÉ, C. A. M., AND NETO, A. M. Supercharacters of the Sylow p -subgroups of the finite symplectic and orthogonal groups. Preprint, 2008. Available at <http://arxiv.org/abs/0804.4285>.
- [7] ARIAS-CASTRO, E., DIACONIS, P., AND STANLEY, R. A super-class walk on upper-triangular matrices. *J. Algebra* 278, 2 (2004), 739–765.
- [8] CARTER, R. W. *Simple groups of Lie type*. Pure and Applied Mathematics. Wiley, London, 1972.
- [9] CARTER, R. W. *Finite Groups of Lie type (conjugacy classes and complex characters)*. Pure and Applied Mathematics. Wiley, New York, 1985.
- [10] CURTIS, C. W., AND REINER, I. *Methods of representation theory (with applications to finite groups and orders)*. Vol. II. Pure and Applied Mathematics. Wiley, New York, 1987.
- [11] DIACONIS, P., AND ISAACS, I. M. Supercharacters and superclasses for algebra groups. *Trans. Amer. Math. Soc.* 360, 5 (2008), 2359–2392.
- [12] ISAACS, I. M. *Character theory of finite groups*. Dover, New York, 1994.
- [13] KAZHDAN, D. Proof of Springer’s hypothesis. *Israel J. Math.* 28 (1977), 272–286.
- [14] KIRILLOV, A. A. Variations on the triangular theme. In *Lie groups and Lie algebras: E. B. Dynkin’s seminar*, vol. 169 of *Amer. Math. Soc. Transl. Ser. 2*. Amer. Math. Soc., Providence RI, 1995, pp. 43–73.
- [15] LIDL, R., AND NIEDERREITER, H. *Finite fields*, 2nd ed., vol. 20 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1997.
- [16] NETO, A. M. *Caracteres e órbitas coadjuntas de p -grupos lineares finitos*. PhD thesis, University of Lisbon, 2006.
- [17] YAN, N. *Representation theory of the finite unipotent linear groups*. PhD thesis, University of Pennsylvania, 2001.

(C. A. M. André) DEPARTAMENTO DE MATEMÁTICA, FACULDADE DE CIÊNCIAS DA UNIVERSIDADE DE LISBOA, CAMPO GRANDE, EDIFÍCIO C6, PISO 2, 1749-016 LISBOA, PORTUGAL

(A. M. Neto) INSTITUTO SUPERIOR DE ECONOMIA E GESTÃO, UNIVERSIDADE TÉCNICA DE LISBOA, RUA DO QUELHAS 6, 1200-781 LISBOA, PORTUGAL

(C. A. M. André & A. M. Neto) CENTRO DE ESTRUTURAS LINEARES E COMBINATÓRIAS, COMPLEXO INTERDISCIPLINAR DA UNIVERSIDADE DE LISBOA, AV. PROF. GAMA PINTO 2, 1649-003 LISBOA, PORTUGAL

E-mail address: `caandre@fc.ul.pt`

E-mail address: `ananeto@iseg.utl.pt`