

# Study of the discontinuity of the exchange-correlation potential in an exactly soluble case

Paola Gori-Giorgi<sup>1,2</sup> and Andreas Savin<sup>1</sup>

<sup>1</sup> *Laboratoire de Chimie Théorique,  
CNRS UMR76116, Université Pierre et Marie Curie,  
4 Place Jussieu, F-75252 Paris, France*

<sup>2</sup> *Afdeling Theoretische Chemie, Vrije Universiteit,  
De Boelelaan 1083, 1081 HV Amsterdam, The Netherlands*

(Dated: April 1, 2022)

## Abstract

It was found by Perdew, Parr, Levy, and Balduz [Phys. Rev. Lett. **49**, 1691 (1982)] and by Sham and Schlüter [Phys. Rev. Lett. **51**, 1884 (1983)] that the exact Kohn-Sham exchange-correlation potential of an open system may jump discontinuously as the particle number crosses an integer, with important physical consequences. Recently, Sagvolden and Perdew [Phys. Rev. A **77**, 012517 (2008)] have analyzed the discontinuity of the exchange-correlation potential as the particle number crosses one, with an illustration that uses a model density for the  $H^-$  ion. In this work, we extend their analysis to the case in which the external potential is the simple harmonic confinement, choosing spring-constant values for which the two-electron hamiltonian has an analytic solution. This way, we can obtain the exact, analytic, exchange and correlation potentials for particle number fluctuating between zero and two, illustrating the discontinuity as the particle number crosses one without introducing any model or approximation. We also discuss exchange and correlation separately.

## I. INTRODUCTION

Kohn-Sham (KS) density functional theory (DFT) (see, e.g., [1]) is a successful method for electronic structure calculations, thanks to its unique combination of low computational cost and reasonable accuracy. In the Kohn-Sham formalism, the total energy of a many-electron system in the external potential  $\hat{V}_{\text{ext}} = \sum_i v_{\text{ext}}(\mathbf{r}_i)$  is rewritten as a functional of the one-electron density  $\rho(\mathbf{r})$ ,

$$E[\rho] = T_s[\rho] + \int d\mathbf{r} v_{\text{ext}}(\mathbf{r}) \rho(\mathbf{r}) + U[\rho] + E_{\text{xc}}[\rho]. \quad (1)$$

In Eq. (1),  $T_s[\rho]$  is the kinetic energy of a non-interacting system of fermions (usually called KS system) having the same one-electron density  $\rho$  of the physical, interacting, system. The Hartree energy  $U[\rho]$  is the classical repulsion energy,  $U[\rho] = \frac{1}{2} \int d\mathbf{r} \int d\mathbf{r}' \rho(\mathbf{r}) \rho(\mathbf{r}') |\mathbf{r} - \mathbf{r}'|^{-1}$ , and the exchange-correlation functional  $E_{\text{xc}}[\rho]$  must be approximated. Minimization of Eq. (1) with respect to the orbitals forming the KS determinant lead to the KS equations. Thus, instead of the physical problem, in KS DFT we solve the hamiltonian of a model system of non-interacting fermions in the one-body local potential  $\hat{V}_{\text{KS}} = \sum_i v_{\text{KS}}(\mathbf{r}_i)$ , with

$$v_{\text{KS}}(\mathbf{r}) = v_{\text{ext}}(\mathbf{r}) + v_{\text{H}}(\mathbf{r}) + v_{\text{xc}}(\mathbf{r}) \quad (2)$$

$$v_{\text{H}}(\mathbf{r}) = \int d\mathbf{r}' \frac{\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} \quad (3)$$

$$v_{\text{xc}}(\mathbf{r}) = \frac{\delta E_{\text{xc}}[\rho]}{\delta \rho(\mathbf{r})}, \quad (4)$$

and we recover the energy of the physical, interacting system, through the sum of the two functionals  $U[\rho] + E_{\text{xc}}[\rho]$ .

In Refs. [2, 3, 4] an analysis of KS theory for systems with fluctuating particle number lead to the conclusion that the exact exchange-correlation potential  $v_{\text{xc}}(\mathbf{r})$  may jump discontinuously by a spatial-independent constant as the particle number crosses an integer, with important physical consequences. In the last years there has been new interest in the derivative discontinuity of  $E_{\text{xc}}[\rho]$  (see, e.g., [5, 6, 7, 8, 9, 10, 11]), and its existence has even been questioned in Ref. [12]. In a recent paper, Sagvolden and Perdew [13] have given further support to the assumptions used to find the exchange-correlation potential discontinuity, and they have illustrated the discontinuity when the particle number crosses one, using a model density for the  $\text{H}^-$  ion. They have also rigorously proved that the von Weizsäcker

functional,

$$T_{\text{vW}}[\rho] = \frac{1}{2} \int d\mathbf{r} |\nabla \sqrt{\rho(\mathbf{r})}|^2 \quad (5)$$

is the correct  $T_s[\rho]$  for a system with particle number  $N \leq 2$ .

In this work we repeat a similar analysis for electronic systems with particle number fluctuating between zero and two when the external potential is harmonic,  $v_{\text{ext}}(\mathbf{r}) = \frac{1}{2}\omega^2 r^2$ . Taut [14] has shown that the corresponding hamiltonian for  $N = 2$  electrons is analytically soluble for some special values of  $\omega$ , which means that, in such special cases, we can calculate analytically the exact interacting density and the exact exchange-correlation potential [15]. This way, we can illustrate the derivative discontinuity without relying on any approximation. The paper is organized as follows. In Sec. II we report the equations used to extract the exact exchange and correlation potentials from Taut's [14] analytical solutions. The corresponding results are shown and analyzed in Sec. III, and the last Sec. IV is devoted to concluding remarks.

## II. THEORY

We consider a system with particle number fluctuating between 0 and 2, in the harmonic external potential  $v_{\text{ext}}(\mathbf{r}) = \frac{1}{2}\omega^2 r^2$ . Such a system can be thought as arising from identical quantum traps whose centers are separated by a very large distance [16]. If the energy of the physical system is convex as a function of the integer particle number  $M$ , i.e., if  $E(M) \leq (E(M+1) + E(M-1))/2$ , then the system with non-integer particle number  $N$  arises from an ensemble of only the two systems with integer-particle number  $M$  and  $M+1$  such that  $M < N < M+1$  [2]. In the case treated here,  $v_{\text{ext}}(\mathbf{r}) = \frac{1}{2}\omega^2 r^2$ , the energy of the non-interacting system is convex, and the electron-electron interaction seems to make it strictly convex (see, e.g., the results for two-dimensional harmonic traps of Refs. [17] or the three-dimensional case treated in Ref. [18]). The density of the system with particle number  $0 \leq N \leq 2$ ,  $\rho_N(\mathbf{r}) = \rho_N(r)$ , is then equal to [2, 13]

$$\rho_N(\mathbf{r}) = \begin{cases} N\rho_1(\mathbf{r}) & 0 \leq N \leq 1 \\ (2-N)\rho_1(\mathbf{r}) + (N-1)\rho_2(\mathbf{r}) & 1 < N \leq 2 \end{cases} \quad (6)$$

The exact Kohn-Sham potential is, up to a constant, given by the functional derivative of the von Weizsäcker functional of Eq. (5),

$$v_{\text{KS}}(r) = \frac{\nabla^2 \sqrt{\rho_N(r)}}{2\sqrt{\rho_N(r)}} + \text{const.} \quad (7)$$

The Hartree potential  $v_{\text{H}}(r)$  can be easily calculated by plugging Eq. (6) into Eq. (3). The exchange and correlation potential  $v_{\text{xc}}(r)$  is then calculated from Eq. (4),  $v_{\text{xc}}(r) = v_{\text{KS}}(r) - v_{\text{H}}(r) - v_{\text{ext}}(r)$ . Setting  $v_{\text{xc}}(r \rightarrow \infty) = 0$  for any fixed  $N$  determines the arbitrary constant in the exchange-correlation potential.

The potential  $v_{\text{xc}}(r)$  can be further divided into exchange and correlation by noticing that for  $0 \leq N \leq 1$   $v_x(r)$  must exactly cancel the Hartree potential and that for  $1 \leq N \leq 2$  the sum  $U[\rho_N] + E_x[\rho_N]$  must equal the expectation of the electron-electron repulsion  $\hat{V}_{ee}$  over the non-interacting ensemble density matrix that yields  $\rho_N(r)$ . The non-interacting density matrix that yields  $\rho_N(r)$  and corresponds to the kinetic energy functional of Eq. (5) is

$$\Gamma_0 = (2 - N)|\Phi_1\rangle\langle\Phi_1| + (N - 1)|\Phi_2\rangle\langle\Phi_2|, \quad (8)$$

with

$$\Phi_1(r) = \sqrt{\frac{\rho_N(r)}{N}} \quad (9)$$

$$\Phi_2(r_1, r_2) = \sqrt{\frac{\rho_N(r_1)}{N}} \sqrt{\frac{\rho_N(r_2)}{N}}. \quad (10)$$

Thus

$$U[\rho_N] + E_x[\rho_N] = \text{Tr}(\Gamma_0 \hat{V}_{ee}) = \frac{2(N - 1)}{N^2} U[\rho_N] \quad (11)$$

By taking the functional derivative  $\delta/\delta\rho(\mathbf{r})$  of both sides of Eq. (11) (and by considering that  $N = \int d\mathbf{r} \rho_N(\mathbf{r})$ ), we finally obtain, for the whole range  $0 \leq N \leq 2$ ,

$$v_x(r) = \begin{cases} -v_{\text{H}}(r) & 0 \leq N \leq 1 \\ -\frac{(N^2 - 2N + 2)}{N^2} v_{\text{H}}(r) + \left(\frac{2}{N^2} - \frac{4(N-1)}{N^3}\right) U[\rho_N] & 1 < N \leq 2 \end{cases} \quad (12)$$

Equation (12) shows that, within the definition of exchange of Eq. (11), the discontinuity of the exchange potential at  $N = 1$  is always a spatially-independent constant equal to

$$v_x(r)|_{N \rightarrow 1^+} - v_x(r)|_{N \rightarrow 1^-} = 2U[\rho_1]. \quad (13)$$

Exact exchange in an open system of fluctuating electron number as also been widely discussed in Ref. [19], where only the energy, and not the potential, has been analyzed.

When the external potential is harmonic,  $v_{\text{ext}}(r) = \frac{1}{2}\omega^2 r^2$ , we have  $\rho_1(r) = \omega^{3/2}/\pi^{3/2} e^{-\omega r^2}$ . The ground-state wavefunction of the hamiltonian with  $N = 2$  electrons has the form

$$\Psi(\mathbf{r}_1, \mathbf{r}_2) = \eta(R)\psi(r_{12}). \quad R = \frac{|\mathbf{r}_1 + \mathbf{r}_2|}{2}, \quad r_{12} = |\mathbf{r}_2 - \mathbf{r}_1|. \quad (14)$$

The center-of-mass wavefunction  $\eta(R)$  is a simple ground-state three-dimensional harmonic oscillator state. Taut [14] has shown that for some special values of  $\omega$  also the relative wavefunction  $\psi(r_{12})$  has an analytic form. For some of these special  $\omega$ -values we have calculated the corresponding one electron density  $\rho_2(r)$ ,

$$\rho_2(r) = C \frac{e^{-\omega r^2}}{2\omega r} \int_0^\infty dr_{12} r_{12} t(r_{12})^2 \left( e^{-\omega(r-r_{12})^2} - e^{-\omega(r+r_{12})^2} \right), \quad (15)$$

where  $t(x) = e^{\frac{1}{2}\omega x^2} \psi(x)$ , and  $C$  is a normalization constant. The function  $t(x)$  is a polynomial of some finite order  $n$  (depending on  $\omega$ ), so that the corresponding  $\rho_2(r)$  is completely analytical. This allows us to obtain the exact exchange-correlation potentials up to any large  $r$  exactly, without introducing any approximation. The large- $r$  part of the potentials is crucial in order to illustrate the derivative discontinuity.

Notice that similar calculations of the exact exchange-correlation potential for the  $N = 2$  system have been carried out in the past, mainly aimed at comparing the exact potentials with current approximations (see e.g., [15]). Here our aim is to calculate the exact potentials for  $0 \leq N \leq 2$  in order to obtain exactly their discontinuity as the particle number crosses one.

### III. RESULTS

We considered three cases for which  $\psi(r_{12})$  of Eq. (14) is analytical:  $\omega = 1/2$ ,  $\omega = 1/10$ , and  $\omega = (35 - 2\sqrt{57})/712 \approx 0.0173$ . The exact exchange-correlation potentials for  $N$  slightly below and slightly above 1 are reported in Fig. 1. The qualitative behavior is similar to the one reported by Sagvolden and Perdew [13] obtained from the model density of the  $\text{H}^-$  ion: as  $N \rightarrow 1^+$  the exchange-correlation potential is, over a larger and larger range, more and more equal to the  $N \rightarrow 1^-$  potential plus a constant. The magnitude of the constant is, as expected [2, 3, 4],  $I - A$ , where  $I$  is the ionization potential and  $A$  the electron affinity. For closed-shell two-electron systems this amounts to  $I - A = E_2 - 2E_1$ , where  $E_2$  is the energy




FIG. 1: The exact exchange-correlation potential for a system with particle number  $N$  fluctuating between zero and two in the external potential  $\frac{1}{2}\omega r^2$ , for three values of  $\omega$ . The magnitude of the discontinuity at  $N = 1$  is equal to  $E_2 - 2E_1$ . We have for  $\omega = \frac{1}{2}$   $E_2 - 2E_1 = \frac{1}{2}$ , for  $\omega = \frac{1}{10}$   $E_2 - 2E_1 = \frac{1}{5}$ , and for  $\omega \approx 0.0173$   $E_2 - 2E_1 = 4\omega \approx 0.06938$ .

of the system with  $N = 2$  electrons and  $E_1$  is the energy for  $N = 1$ . Figure 1 clearly shows that, as argued in Refs. [2, 3, 4, 13], for any fixed given  $N$  (above or below 1)

$$\lim_{r \rightarrow \infty} v_{\text{xc}}(r) = 0, \quad (16)$$

but

$$\lim_{r \rightarrow \infty} \lim_{N \rightarrow 1^+} v_{\text{xc}}(r) = I - A. \quad (17)$$

#### IV. CONCLUSIONS

The discontinuity of the exact Kohn-Sham exchange-correlation potential was argued in Refs. [2, 3, 4], and is known to have very important physical consequences. In particular, the absence or the underestimation of this discontinuity in local and semilocal functionals explains why they often produce a qualitatively incorrect dissociation limit for non-equilibrium nuclear positions and why they underestimate charge-transfer excitation energies in time-dependent DFT. In this work we have calculated the exact Kohn-Sham exchange and correlation potentials as the particle number crosses 1 for a system which is analytically soluble. This way, we have illustrated, at least in one case, the existence of this discontinuity without relying on any approximation, thus providing further support to the assumptions that were used for its prediction.

#### Acknowledgements

It is our pleasure to dedicate this illustration of some simple principles of density functional theory to Istavan Mayer who always liked simple theorems, proofs, and derivations in quantum chemistry. We are indebted to John P. Perdew for pointing out an error in the previous version, concerning the exchange potential of Eq. (12). This work was supported by the ANR (National French Research Agency) under Grant n. ANR-07-BLAN-0271.

- 
- [1] W. Kohn, Rev. Mod. Phys. **71**, 1253 (1999).
  - [2] J. P. Perdew, R. G. Parr, M. Levy, and J. L. Balduz, Phys. Rev. Lett. **49**, 1691 (1982).
  - [3] L. J. Sham and M. Schlüter, Phys. Rev. Lett. **51**, 1888 (1983).

- [4] J. P. Perdew and M. Levy, Phys. Rev. Lett. **51**, 1884 (1983).
- [5] M. Grüning, A. Marini, and A. Rubio, Phys. Rev. B **74**, 161103 (2006).
- [6] M. Grüning, A. Marini, and A. Rubio, J. Chem. Phys. **124**, 154108 (2006).
- [7] A. Ruzsinszky, J. P. Perdew, G. I. Csonka, O. A. Vydrov, and G. E. Scuseria, J. Chem. Phys. **125**, 194112 (2006).
- [8] D. J. Tozer, J. Chem. Phys. **119**, 12697 (2003).
- [9] A. M. Teale, F. D. Proft, and D. J. Tozer, J. Chem. Phys. **129**, 044110 (2008).
- [10] P. Mori-Sanchez, A. J. Cohen, and W. T. Yang, J. Chem. Phys. **125**, 201102 (2006).
- [11] A. J. Cohen, P. Mori-Sanchez, and W. T. Yang, Science **321**, 792 (2008).
- [12] F. E. Zahariev and Y. A. Wang, Phys. Rev. A **70**, 042503 (2004).
- [13] E. Sagvolden and J. P. Perdew, Phys. Rev. A **77**, 012517 (2008).
- [14] M. Taut, Phys. Rev. A **48**, 3561 (1993).
- [15] C. Filippi, C. J. Umrigar, and M. Taut, J. Chem. Phys. **100**, 1290 (1994).
- [16] W. Yang, Y. Zhang, and P. W. Ayers, Phys. Rev. Lett. **84**, 5172 (2000).
- [17] M. Rontani, C. Cavazzoni, D. Bellucci, and G. Goldoni, J. Chem. Phys. **124**, 124102 (2006).
- [18] M. Taut, K. Pernal, J. Cioslowski, and V. Staemmier, J. Chem. Phys. **118**, 4861 (2003).
- [19] J. P. Perdew, A. Ruzsinszky, G. I. Csonka, O. A. Vydrov, G. E. Scuseria, V. N. Staroverov, and J. Tao, Phys. Rev. A **76**, 040501 (2007).