

# Addendum to “Experimental demonstration of a quantum protocol for Byzantine agreement and liar detection”

Sascha Gaertner,<sup>1,2</sup> Mohamed Bourennane,<sup>3</sup> Christian Kurtsiefer,<sup>4</sup> Adán Cabello,<sup>5,\*</sup> and Harald Weinfurter<sup>1,2</sup>

<sup>1</sup>Max-Planck-Institut für Quantenoptik, D-85748 Garching, Germany

<sup>2</sup>Fakultät für Physik, Ludwig-Maximilians-Universität, D-80799 München, Germany

<sup>3</sup>Department of Physics, Stockholm University, SE-10691 Stockholm, Sweden

<sup>4</sup>Department of Physics, National University of Singapore, 117542 Singapore, Singapore

<sup>5</sup>Departamento de Física Aplicada II, Universidad de Sevilla, E-41012 Sevilla, Spain

(Dated: March 13, 2019)

Gao *et al.* has described a possible intercept-resend attack for the quantum protocol for detectable Byzantine agreement described in Phys. Rev. Lett. **100**, 070504 (2008). Here we describe an extension of the protocol which defeats such attacks.

PACS numbers: 03.67.Hk, 03.67.Pp, 42.50.Dv

Recently [1], Gao *et al.* have pointed out that there exists an intercept-resend attack for the quantum protocol for detectable Byzantine agreement described in [2]. Here we describe how such attacks alter the entanglement of the distributed multipartite state and thus can be revealed by further analysis of the acquired data.

The protocol uses the four-photon state

$$|\Psi^{(4)}\rangle_{abcd} = \frac{1}{2\sqrt{3}}(2|0011\rangle - |0101\rangle - |0110\rangle - |1001\rangle - |1010\rangle + 2|1100\rangle)_{abcd}, \quad (1)$$

to distribute lists securely to three parties which then enable them to achieve the detectable Byzantine agreement.

In [2], step (iii) stated: “*C* randomly chooses a position from his list and asks *A* and *B* to inform him about their results on the same position. If all parties have measured in the same basis, their results must be suitably correlated.” In the attack proposed in [1], the traitor intercepts the qubits sent to one of the loyal generals and performs measurements on them. These measurements are single-qubit measurements in the  $\{|0\rangle, |1\rangle\}$  basis of eigenstates of the Pauli matrix  $\sigma_z$  or in the  $\{|\bar{0}\rangle, |\bar{1}\rangle\}$  basis [where  $|\bar{0}\rangle = (|0\rangle + |1\rangle)/\sqrt{2}$  and  $|\bar{1}\rangle = (|0\rangle - |1\rangle)/\sqrt{2}$ ] of eigenstates of the Pauli matrix  $\sigma_x$ . Then, the traitor resends the qubits in the state resulting from the previous measurements to the loyal general. With this method, the traitor can obtain the other generals’ secret lists which he can use to mislead the loyal generals into following different plans.

Since the roles of *B* and *C* in the protocol in [2] are symmetrical, there are only three cases to be dealt with:

(I) *A* (the commanding general) is the traitor and intercepts, measures, and resends the qubit sent to *B* (qubit *c*).

(II) *B* is the traitor and intercepts, measures, and resends the two qubits sent to *A* (qubits *a* and *b*).

(III) *B* is the traitor and intercepts, measures, and resends the qubit sent to *C* (qubit *d*).

These attacks indeed do not alter the perfect correlations utilized in the protocol so far. However, they obviously change the entanglement of the four photon state and thus the correlations between the results if the parties measured along different directions. This can be used to detect the attack. There are several ways to foil these attacks. Here we present a simple method which does not require additional measurements and in which the traitor does not need to announce his actual results. In the following, we present an example of the method for each of the three cases.

(I) Suppose that *A* is the traitor and intercepts and measures the qubit sent to *B* in the basis of eigenstates of  $\sigma_z$ . In half of the cases, *A* obtains the result corresponding to  $|0\rangle_c$  and resends the state  $|0\rangle_c$  to *B*. Therefore, in these cases the state of the four qubits is

$$|\Psi'\rangle_{abcd} = \frac{1}{\sqrt{6}}(-|0101\rangle - |1001\rangle + 2|1100\rangle)_{abcd}. \quad (2)$$

In the other half of the cases, *A* obtains the result corresponding to  $|1\rangle_c$  and resends *B* the state  $|1\rangle_c$ . Therefore, in these cases the state of the four qubits is

$$|\Psi''\rangle_{abcd} = \frac{1}{\sqrt{6}}(2|0011\rangle - |0110\rangle - |1010\rangle)_{abcd}. \quad (3)$$

Therefore, after *A*’s attack, the state of the four qubits is the mixed state

$$\rho_{abcd} = \frac{1}{2}(|\Psi'\rangle\langle\Psi'| + |\Psi''\rangle\langle\Psi''|)_{abcd}. \quad (4)$$

Suppose that *B* and *C* measure their qubits in the basis of eigenstates of  $\sigma_x$ . Then, *C* asks *A* and *B* for their results. When *C* checks them, *C* expects that  $\langle\sigma_x^{(c)} \otimes \sigma_x^{(d)}\rangle_{\Psi^{(4)}} = \frac{1}{3}$ , which is the prediction for the state  $|\Psi^{(4)}\rangle_{abcd}$ . However, for the mixed state  $\rho_{abcd}$ , *C* finds  $\langle\sigma_x^{(c)} \otimes \sigma_x^{(d)}\rangle_{\rho_{abcd}} = 0$ .

(II) If *B* is the traitor and intercepts and measures qubits *a* and *b* in the basis of eigenstates of  $\sigma_z$  (resulting

\*Electronic address: adan@us.es

TABLE I: Experimental data showing that an intercept-resend attack [1] has not occurred in the experiment described in [2].

	Experimental result	Result for $ \Psi^{(4)}\rangle_{abcd}$
$\langle \sigma_x^{(a)} \otimes \sigma_x^{(b)} \rangle$	$0.262 \pm 0.025$	$1/3$
$\langle \sigma_z^{(a)} \otimes \sigma_z^{(b)} \rangle$	$0.273 \pm 0.025$	$1/3$
$\langle \sigma_x^{(a)} \otimes \sigma_x^{(c)} \rangle$	$-0.602 \pm 0.021$	$-2/3$
$\langle \sigma_z^{(a)} \otimes \sigma_z^{(c)} \rangle$	$-0.631 \pm 0.02$	$-2/3$
$\langle \sigma_x^{(a)} \otimes \sigma_x^{(d)} \rangle$	$-0.612 \pm 0.02$	$-2/3$
$\langle \sigma_z^{(a)} \otimes \sigma_z^{(d)} \rangle$	$-0.663 \pm 0.019$	$-2/3$
$\langle \sigma_x^{(b)} \otimes \sigma_x^{(c)} \rangle$	$-0.603 \pm 0.021$	$-2/3$
$\langle \sigma_z^{(b)} \otimes \sigma_z^{(c)} \rangle$	$-0.615 \pm 0.02$	$-2/3$
$\langle \sigma_x^{(b)} \otimes \sigma_x^{(d)} \rangle$	$-0.61 \pm 0.02$	$-2/3$
$\langle \sigma_z^{(b)} \otimes \sigma_z^{(d)} \rangle$	$-0.621 \pm 0.02$	$-2/3$
$\langle \sigma_x^{(c)} \otimes \sigma_x^{(d)} \rangle$	$0.334 \pm 0.025$	$1/3$
$\langle \sigma_z^{(c)} \otimes \sigma_z^{(d)} \rangle$	$0.326 \pm 0.024$	$1/3$

in some mixed state  $\rho'_{abcd}$ ), and  $A$  measures qubits  $a$  and

$b$  in the basis of eigenstates of  $\sigma_x$ , then  $C$  expects that  $\langle \sigma_x^{(a)} \otimes \sigma_x^{(b)} \rangle = \frac{1}{3}$  but finds  $\langle \sigma_x^{(a)} \otimes \sigma_x^{(b)} \rangle_{\rho'_{abcd}} = 0$ .

(III) If  $B$  is the traitor and intercepts and measures qubit  $d$  in the basis of eigenstates of  $\sigma_z$ , and  $A$  and  $C$  measure their qubits in the basis of eigenstates of  $\sigma_x$ , then  $C$  expects that  $\langle \sigma_x^{(a)} \otimes \sigma_x^{(d)} \rangle_{\Psi^{(4)}} = \langle \sigma_x^{(b)} \otimes \sigma_x^{(d)} \rangle_{\Psi^{(4)}} = -\frac{2}{3}$  but finds  $\langle \sigma_x^{(a)} \otimes \sigma_x^{(d)} \rangle_{\rho'_{abcd}} = \langle \sigma_x^{(b)} \otimes \sigma_x^{(d)} \rangle_{\rho'_{abcd}} = 0$ .

If  $C$  (or the other generals, since they exchange their roles in the next step) obtains results that do not satisfy the quantum predictions for the state  $|\Psi^{(4)}\rangle_{abcd}$ , then the loyal generals decide to abort the protocol; otherwise, the loyal generals can reach an agreement using the lists.

We show that in the experiment described in [2], actual experimental data prove that the attack [1] has not occurred. Table I contains all the tests described above and those obtained from them by changing  $\sigma_x^{(i)}$  by  $\sigma_z^{(i)}$  for all qubits, exploiting the fact that the state (1) is invariant under the same unitary transformation applied to the four qubits.

In conclusion, there is a simple method that makes the protocol in [2] secure against the attack proposed in [1].

[1] F. Gao, F.-Z. Guo, Q.-Y. Wen, and F.-C. Zhu, to be published in Phys. Rev. Lett.

[2] S. Gaertner, M. Bourennane, C. Kurtsiefer, A. Cabello,

and H. Weinfurter, Phys. Rev. Lett. **100**, 070504 (2008).