

# On an extension of Lesche stability

A. El Kaabouchi<sup>1</sup>, A. Le Méhauté<sup>1</sup>, L. Nivanen<sup>1</sup>, C.J. Ou<sup>1,2</sup> and A. Q. Wang<sup>1</sup>

<sup>1</sup>Institut Supérieur des Matériaux et Mécaniques Avancés du Mans  
44 Avenue Bartholdi, 72000 Le Mans, France

<sup>2</sup>College of Information Science and Engineering,  
Huaqiao University, Quanzhou 362021, China

## Abstract

In this paper, we give a new method for proving the Lesche stability of several functionals (Incomplete entropy, Tsallis entropy,  $\kappa$ -entropy, Quantum-Group entropy). We prove also that the Incomplete  $q$ -expectation value and Rényi entropy for  $(0 < q < 1)$  are  $\alpha$ -stable for all  $(0 < \alpha \leq q)$ . Finally, we prove that the Incomplete  $q$ -expectation value is  $\alpha$ -stable for all  $0 < \alpha \leq 1$ .

### Notations and definitions

We denote by  $E_f = \bigcup_{n \in \mathbf{N}^*} \mathbf{R}^n$ . For  $N \in \mathbf{N}^*$  and  $\alpha > 0$ , we define the function  $d_{\alpha, N} : \mathbf{R}^N \times \mathbf{R}^N \rightarrow \mathbf{R}_+$  by

$$d_{\alpha, N}(p, p') = \left( \sum_{i=1}^N |p_i - p'_i|^\alpha \right)^{1/\alpha}.$$

We recall that

for  $\alpha \geq 1$ ,  $d_{\alpha, N}$  is a *distance* on  $\mathbf{R}^N$

$$\forall p, p' \in \mathbf{R}^N, d_{\alpha, N}(p, p') = 0 \Leftrightarrow p = p'$$

$$\forall p, p' \in \mathbf{R}^N, d_{\alpha, N}(p, p') = d_{\alpha, N}(p', p)$$

$$\forall p, p', p'' \in \mathbf{R}^N, d_{\alpha, N}(p, p'') \leq d_{\alpha, N}(p, p') + d_{\alpha, N}(p', p'')$$

for  $\alpha < 1$ ,  $d_{\alpha, N}$  is a *quasi-distance* on  $\mathbf{R}^N$ .

$$\forall p, p' \in \mathbf{R}^N, d_{\alpha, N}(p, p') = 0 \Leftrightarrow p = p'$$

$$\forall p, p' \in \mathbf{R}^N, d_{\alpha, N}(p, p') = d_{\alpha, N}(p', p)$$

$$\forall p, p', p'' \in \mathbf{R}^N, d_{\alpha, N}(p, p'') \leq 2^{\frac{1}{\alpha}-1} (d_{\alpha, N}(p, p') + d_{\alpha, N}(p', p''))$$

Let  $C$  be a function defined on  $A$  a subset of  $E_f$ . For  $N \in \mathbf{N}^*$  and  $\alpha > 0$ , we denote by  $C_{N, \max}$  the number defined by (when it exists)  $C_{N, \max} = \sup \{C(p), p \in A \cap \mathbf{R}^N\}$ .

### Definition 1

A Function  $C$  defined on  $A$  a subset of  $E_f$  is said to be  $\alpha$ -stable if  $C$  posses the following property

$$\forall \varepsilon > 0, \exists \delta > 0, \forall N \in \mathbf{N}^*, \forall p, p' \in A \cap \mathbf{R}^N, d_{\alpha, N}(p, p') < \delta \Rightarrow \left| \frac{C(p) - C(p')}{C_{N, \max}} \right| < \varepsilon.$$

### Remark

The Lesche stability [1] correspond to  $\alpha$ -stability for  $\alpha = 1$ .

### Proposition 2

Let  $0 < \beta \leq \alpha$  and  $C$  be a function defined on a subset of  $E_f$ .

If  $C$  is  $\alpha$ -stable then  $C$  is  $\beta$ -stable.

*Proof.*

It's easy to see that for all  $x \in ]0, 1[$ ,  $x^\alpha \leq x^\beta$  and thus if  $d_{\beta, N}(p, p') < \delta$  then  $d_{\alpha, N}(p, p') < \delta^{\beta/\alpha}$ .

### Proposition 3

If  $C_1, C_2$  are  $\alpha$ -stable with  $C_1 \geq 0, C_2 \geq 0$  and  $\lambda, \mu \in \mathbf{R}_+$ , then  $\lambda C_1 + \mu C_2$  is  $\alpha$ -stable.

*Proof.*

We can suppose that  $C_1 \neq 0, C_2 \neq 0$  and  $\lambda, \mu \in \mathbf{R}_+^*$ .

We remark that for all  $N \in \mathbf{N}^*$  and  $\alpha > 0$ ,

$$\begin{aligned} \left| \frac{(\lambda C_1 + \mu C_2)(p) - (\lambda C_1 + \mu C_2)(p')}{(\lambda C_1 + \mu C_2)_{N, \max}} \right| &= \frac{|\lambda(C_1(p) - C_1(p')) + \mu(C_2(p) - C_2(p'))|}{(\lambda C_1 + \mu C_2)_{N, \max}} \\ &\leq \lambda \frac{|(C_1(p) - C_1(p'))|}{(\lambda C_1 + \mu C_2)_{N, \max}} + \mu \frac{|(C_2(p) - C_2(p'))|}{(\lambda C_1 + \mu C_2)_{N, \max}} \\ &\leq \lambda \frac{|(C_1(p) - C_1(p'))|}{\lambda (C_1)_{N, \max}} + \mu \frac{|(C_2(p) - C_2(p'))|}{\mu (C_2)_{N, \max}} \\ &= \frac{|(C_1(p) - C_1(p'))|}{(C_1)_{N, \max}} + \frac{|(C_2(p) - C_2(p'))|}{(C_2)_{N, \max}} \end{aligned}$$

Let  $\varepsilon > 0$ , there exists  $\delta_1 > 0$ ,  $\delta_2 > 0$ , such that  $\forall N \in \mathbf{N}^*$ ,  $\forall p, p' \in A \cap \mathbf{R}^N$ ,

$$d_{\alpha, N}(p, p') < \delta_1 \Rightarrow \left| \frac{C_1(p) - C_1(p')}{(C_1)_{N, \max}} \right| < \frac{\varepsilon}{2} \text{ and } d_{\alpha, N}(p, p') < \delta_2 \Rightarrow \left| \frac{C_2(p) - C_2(p')}{(C_2)_{N, \max}} \right| < \frac{\varepsilon}{2}.$$

We put  $\delta = \min(\delta_1, \delta_2) > 0$ ,  $\forall p, p' \in A \cap \mathbf{R}^N$ ,

$$d_{\alpha, N}(p, p') < \delta \Rightarrow (d_{\alpha, N}(p, p') < \delta_1 \text{ and } d_{\alpha, N}(p, p') < \delta_2)$$

Consequently 
$$\left| \frac{(\lambda C_1 + \mu C_2)(p) - (\lambda C_1 + \mu C_2)(p')}{(\lambda C_1 + \mu C_2)_{N, \max}} \right| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

**Definition 4**

Let  $q \in ]0, +\infty[ \setminus \{1\}$ . The incomplete entropy  $S_q^I$  [2] is defined on  $A = \bigcup_{n \in \mathbf{N}^*} \left\{ p \in [0, 1]^n, \sum_{i=1}^n (p_i)^q = 1 \right\}$  by

$$S_q^I(p) = \frac{1 - \sum_{i=1}^N p_i}{1 - q} \text{ for all } p \in [0, 1]^N.$$

**Theorem 5**

For all  $q \in ]0, +\infty[ \setminus \{1\}$ , the incomplete entropy  $S_q^I$  is  $\alpha$ -stable for all  $\alpha \leq 1$ .

*Proof.*

It's easy to see that

i) 
$$\left| (S_q^I)_{N, \max} \right| = \left| S_q^I \left( (1/N)^{1/q}, \dots, (1/N)^{1/q} \right) \right| = \left| \frac{1 - N^{-\frac{1}{q}}}{1 - q} \right|.$$

ii) For all  $q \in ]0, +\infty[ \setminus \{1\}$ , the function  $x \mapsto \frac{1}{1 - x^{-\frac{1}{q}}}$  is bounded on  $[2, +\infty[$

Indeed if  $q \in ]0, 1[$ , then  $\lim_{x \rightarrow +\infty} \frac{1}{1 - x^{-\frac{1}{q}}} = 1$  and if  $q > 1$ , then  $\lim_{x \rightarrow +\infty} \frac{1}{1 - x^{-\frac{1}{q}}} = 0$ .

Let  $M > 0$ , such that for all  $N \in \mathbf{N}^* \setminus \{1\}$ , 
$$\left| \frac{1}{1 - N^{-\frac{1}{q}}} \right| \leq M.$$

For all  $\varepsilon > 0$ , there exists  $\delta = \left( \frac{\varepsilon}{M} \right)^{1/\alpha} > 0$ , such that for all  $N \in \mathbf{N}^*$ ,  $p, p' \in A \cap \mathbf{R}^N$ ,  $d_{\alpha, N}(p, p') < \delta$

implies

$$\begin{aligned} \left| \frac{S_q^I(p) - S_q^I(p')}{(S_q^I)_{N, \max}} \right| &= \left| \frac{\sum_{i=1}^N (p_i' - p_i)}{1 - N^{-\frac{1}{q}}} \right| \leq \frac{\sum_{i=1}^N |p_i' - p_i|}{\left| 1 - N^{-\frac{1}{q}} \right|} \\ &\leq M \sum_{i=1}^N |p_i' - p_i| \leq M \sum_{i=1}^N |p_i' - p_i|^\alpha \\ &= M (d_{\alpha, N}(p, p'))^\alpha < M \delta^\alpha = \varepsilon \end{aligned}$$

**Definition 6**

Let  $q \in ]0, +\infty[ \setminus \{1\}$ . The Tsallis entropy  $S_q$  [3] is defined on  $A = \bigcup_{n \in \mathbf{N}^*} \left\{ p \in [0, 1]^n, \sum_{i=1}^n p_i = 1 \right\}$  by

$$S_q(p) = \frac{1 - \sum_{i=1}^N (p_i)^q}{q - 1} \text{ for all } p \in [0, 1]^N.$$

**Theorem 7**

For all  $q \in ]0, +\infty[ \setminus \{1\}$ , the Tsallis entropy  $S_q$  is  $\alpha$ -stable for all  $\alpha \leq 1$ .

**Corollary 8**

a) For all  $\kappa \in ]-1, 1[$ , the  $\kappa$ -entropy is  $\alpha$ -stable for all  $\alpha \leq 1$ .

b) For all  $q \in ]0, +\infty[ \setminus \{1\}$ , the Quantum-Group entropy [4] is  $\alpha$ -stable for all  $\alpha \leq 1$ .

*Proof.*

a) Let  $\kappa \in ]-1, 1[$ . The  $\kappa$ -entropy  $S^\kappa$  is defined on  $A = \bigcup_{n \in \mathbf{N}^*} \left\{ p \in [0, 1]^n, \sum_{i=1}^n p_i = 1 \right\}$  by

$$S^\kappa(p) = - \frac{\sum_{i=1}^N p_i \left( (p_i)^\kappa - (p_i)^{-\kappa} \right)}{2\kappa} \text{ for all } p \in [0, 1]^N.$$

It's easy to see that  $S^\kappa(p) = \frac{1}{2} (S_{\kappa+1}(p) + S_{1-\kappa}(p))$  and we conclude by proposition 3.

b) Let  $q \in ]0, +\infty[ \setminus \{1\}$ . The Quantum-Group entropy is defined on  $A = \bigcup_{n \in \mathbf{N}^*} \left\{ p \in [0, 1]^n, \sum_{i=1}^n p_i = 1 \right\}$  is defined by

$$S_q^{QG}(p) = - \frac{\sum_{i=1}^N \left( (p_i)^q - (p_i)^{1/q} \right)}{q - \frac{1}{q}} \text{ for all } p \in [0, 1]^N.$$

It's easy to see that  $S_q^{QG}(p) = \frac{q}{q+1} S_q(p) + \frac{q}{\frac{1}{q}+1} S_{\frac{1}{q}}(p) = \frac{q}{q+1} S_q(p) + \frac{1}{q+1} S_{\frac{1}{q}}(p)$  and we conclude by

proposition 3.

For the proof of the theorem 7, we have need of the following lemmas.

**Lemma 9**

a) For all  $q \in ]0, 1[$ , the function  $x \mapsto \frac{x^{1-q}}{1-x^{1-q}}$  is bounded on  $[2, +\infty[$  by  $M_1 > 0$ ;

b) For all  $q > 1$ , the function  $x \mapsto \frac{1}{1-x^{1-q}}$  is bounded on  $[2, +\infty[$  by  $M_2 > 0$ .

Indeed if  $q \in ]0, 1[$ , then  $\lim_{x \rightarrow +\infty} \frac{x^{1-q}}{1-x^{1-q}} = 1$  and if  $q > 1$ , then  $\lim_{x \rightarrow +\infty} \frac{1}{1-x^{1-q}} = 0$ .

**Lemma 10**

Let  $\alpha > 0$ ,  $N \in \mathbf{N}^*$ ,  $(p_i), (p'_i) \in [0, 1]^N$ .

a) If  $\alpha < 1$ , then  $\sum_{i=1}^N \left| p_i^\alpha - (p'_i)^\alpha \right| \leq (d_{\alpha, N}(p, p'))^\alpha$ ;

b) If  $\alpha < 1$ , then  $\sum_{i=1}^N \left| p_i^\alpha - (p'_i)^\alpha \right| \leq N^{1-\alpha} (d_{1, N}(p, p'))^\alpha$ ;

c) If  $\alpha > 1$ , then  $\sum_{i=1}^N \left| p_i^\alpha - (p'_i)^\alpha \right| \leq \alpha (d_{1, N}(p, p'))^\alpha$ ;

d) If  $\alpha > 1$ , then  $\sum_{i=1}^N \left| p_i^\alpha - (p'_i)^\alpha \right| \leq \alpha N^{1-\frac{1}{\alpha}} (d_{\alpha, N}(p, p'))^\alpha$ .

*Proof.*

a) We put  $I_1 = \{i \in \{1, \dots, N\}, p_i < p_i'\}$  and  $I_2 = \{i \in \{1, \dots, N\}, p_i > p_i'\}$ .

For all  $i \in I_1$ , we have:  $1 = \left(\frac{p_i}{p_i'}\right) + \left(1 - \frac{p_i}{p_i'}\right) \leq \left(\frac{p_i}{p_i'}\right)^\alpha + \left(1 - \frac{p_i}{p_i'}\right)^\alpha = \left(\frac{p_i}{p_i'}\right)^\alpha + \left|1 - \frac{p_i}{p_i'}\right|^\alpha$ , we deduce that

$(p_i)^\alpha - (p_i')^\alpha \leq |p_i - p_i'|^\alpha$ . And by symmetry we deduce that all  $i \in I_2$ , we have:  $(p_i')^\alpha - (p_i)^\alpha \leq |p_i - p_i'|^\alpha$ .

We conclude that for  $i \in \{1, \dots, N\}$ ,  $|(p_i)^\alpha - (p_i')^\alpha| \leq |p_i - p_i'|^\alpha$ , and thus a) is proved.

b) By a) we have,  $\sum_{i=1}^N |(p_i)^\alpha - (p_i')^\alpha| \leq \sum_{i=1}^N |p_i - p_i'|^\alpha = \sum_{i=1}^N |p_i - p_i'|^\alpha \cdot 1$ , and by Holder's Inequality,

$\left(\frac{1}{1/\alpha} + \frac{1}{1/(1-\alpha)} = 1\right)$ , we obtain

$$\sum_{i=1}^N |p_i - p_i'|^\alpha \cdot 1 \leq \left(\sum_{i=1}^N (|p_i - p_i'|^\alpha)^{1/\alpha}\right)^\alpha \left(\sum_{i=1}^N 1^{1/(1-\alpha)}\right)^{1-\alpha} = \left(\sum_{i=1}^N |p_i - p_i'|^\alpha\right)^\alpha N^{1-\alpha}.$$

c) The function  $x \mapsto x^\alpha$  is continuous on  $[0,1]$ , differentiable on  $]0,1[$ , by the Mean-Value Theorem, we have

$$\forall i \in \{1, \dots, N\}, \exists c_i \in (p_i, p_i'), \text{ such that } \left((p_i)^\alpha - (p_i')^\alpha\right) = (p_i - p_i')^\alpha \alpha (c_i)^{\alpha-1},$$

We deduce that

$$\forall i \in \{1, \dots, N\}, |(p_i)^\alpha - (p_i')^\alpha| \leq |p_i - p_i'|^\alpha \alpha.$$

Consequently,  $\sum_{i=1}^N |p_i^\alpha - (p_i')^\alpha| \leq \alpha \left(\sum_{i=1}^N |p_i - p_i'|^\alpha\right) = \alpha d_{1,N}(p, p')$ .

d) By c) we have  $\sum_{i=1}^N |p_i^\alpha - (p_i')^\alpha| \leq \alpha \left(\sum_{i=1}^N |p_i - p_i'|^\alpha\right) = \alpha \sum_{i=1}^N |p_i - p_i'|^\alpha \cdot 1$ , and by Holder's inequality we obtain

$$\sum_{i=1}^N |p_i - p_i'|^\alpha \cdot 1 \leq \left(\sum_{i=1}^N |p_i - p_i'|^\alpha\right)^{1/\alpha} \left(\sum_{i=1}^N 1^{\alpha/(\alpha-1)}\right)^{1-\frac{1}{\alpha}} = N^{1-\frac{1}{\alpha}} d_{\alpha,N}(p, p').$$

Consequently,  $\sum_{i=1}^N |p_i^\alpha - (p_i')^\alpha| \leq N^{1-\frac{1}{\alpha}} d_{\alpha,N}(p, p')$ .

*Proof of theorem 7.*

It's easy to see that  $\left|(S_q)_{\max}\right| = |S_q((1/N), \dots, (1/N))| = \left|\frac{1 - N^{1-q}}{1 - q}\right|$ .

For  $q \in ]0,1[$ , let  $\varepsilon > 0$ , there exists  $\delta = \left(\frac{\varepsilon}{M_1}\right)^{1/(\alpha q)} > 0$ , such that for all  $N \in \mathbf{N}^* \setminus \{1\}$ ,  $p, p' \in A \cap \mathbf{R}^N$ ,

$d_{1,N}(p, p') < \delta$  implies

$$\begin{aligned}
\left| \frac{S_q(p) - S_q(p')}{(S_q)_{\max}} \right| &= \frac{\left| \sum_{i=1}^N (p_i^q - (p'_i)^q) \right|}{|1 - N^{1-q}|} \leq \frac{\sum_{i=1}^N |p_i^q - (p'_i)^q|}{|1 - N^{1-q}|} \leq \frac{N^{1-q} \left( \sum_{i=1}^N |p_i - p'_i| \right)^q}{|1 - N^{1-q}|} \\
&\leq \frac{N^{1-q} \left( d_{\alpha, N}(p, p') \right)^\alpha}{|1 - N^{1-q}|} \\
&= \frac{N^{1-q} \left( d_{\alpha, N}(p, p') \right)^{\alpha q}}{|1 - N^{1-q}|} < M_1 \delta^{\alpha q} = \varepsilon
\end{aligned}$$

For  $q > 1$ , let  $\varepsilon > 0$ , there exists  $\delta = \left( \frac{\varepsilon}{qM_2} \right)^{1/\alpha} > 0$ , such that for all  $N \in \mathbf{N}^*$ ,  $p, p' \in A \cap \mathbf{R}^N$ ,  $d_{1, N}(p, p') < \delta$  implies

$$\begin{aligned}
\left| \frac{S_q(p) - S_q(p')}{(S_q)_{\max}} \right| &\leq \frac{\sum_{i=1}^N |p_i^q - (p'_i)^q|}{|1 - N^{1-q}|} \leq \frac{q \left( \sum_{i=1}^N |p_i - p'_i| \right)^q}{|1 - N^{1-q}|} \\
&\leq \frac{q \left( d_{\alpha, N}(p, p') \right)^\alpha}{|1 - N^{1-q}|} \\
&< qM_2 \delta^\alpha = \varepsilon
\end{aligned}$$

Thus, the theorem 6 is proved.

### The $\alpha$ – stability of the Incomplete $q$ – expectation

#### Definition 11

Let  $q \in ]0, +\infty[ \setminus \{1\}$  et  $C$  a element of  $E_f$

The Incomplete  $q$  – expectation of  $C$  is defined by  $\langle C \rangle_q(p) = \sum_{i=1}^N (p_i)^q C_i$  for all

$$p \in \left\{ p \in [0, 1]^N, \sum_{i=1}^N (p_i)^q = 1 \right\} \text{ where } N \in \mathbf{N}^*.$$

#### Theorem 12

- For  $q > 1$ , the Incomplete  $q$  – expectation of  $C$  is  $\alpha$  – stable for all  $0 < \alpha \leq 1$ .
- For  $q \in ]0, 1[$ , the Incomplete  $q$  – expectation of  $C$  is  $\alpha$  – stable for all  $0 < \alpha \leq q$ .
- For  $q \in ]0, 1[$ , the Incomplete  $q$  – expectation of  $C$  is 1 – unstable.

*Proof*

We have

$$\begin{aligned}
\left| \langle C \rangle_{N, \max} \right| &= \max \left( \left| \langle C \rangle_q(p) \right|, p \in \left\{ p \in [0, 1]^N, \sum_{i=1}^N (p_i)^q = 1 \right\} \right) \\
&= \max \left( \left| \sum_{i=1}^N (p_i)^q C_i \right|, \text{ where } p \text{ satisfies } \sum_{i=1}^N (p_i)^q = 1 \right) \\
&\geq \max_{i=1, \dots, N} |C_i| \cdot 1^q = \max_{i=1, \dots, N} |C_i|
\end{aligned}$$

- For  $q > 1$ , let  $\varepsilon > 0$ , there exists  $\delta = \left( \frac{\varepsilon}{q} \right)^{1/\alpha} > 0$ , such that for all  $N \in \mathbf{N}^*$ ,  $p, p' \in A \cap \mathbf{R}^N$ ,  $d_{\alpha, N}(p, p') < \delta$  implies

$$\begin{aligned}
\left| \frac{\langle C \rangle_q(p) - \langle C \rangle_q(p')}{(\langle C \rangle_q)_{N, \max}} \right| &\leq \frac{\sum_{i=1}^N |p_i^q - (p'_i)^q| |C_i|}{\left| (\langle C \rangle_q)_{N, \max} \right|} \leq \frac{\sum_{i=1}^N |p_i^q - (p'_i)^q| |C_i|}{\max_{i=1, \dots, N} |C_i| \cdot 1^q} \\
&\leq \sum_{i=1}^N |p_i^q - (p'_i)^q| \leq q \left( \sum_{i=1}^N |p_i - p'_i| \right) \\
&= q(d_{1,N}(p, p')) \leq q(d_{\alpha, N}(p, p'))^\alpha < q\delta^\alpha = \varepsilon
\end{aligned}$$

b) For  $q \in ]0, 1[$ , let  $\varepsilon > 0$ , there exists  $\delta = (\varepsilon)^{1/\alpha} > 0$ , such that for all  $N \in \mathbf{N}^*$ ,  $p, p' \in A \cap \mathbf{R}^N$ ,  $d_{\alpha, N}(p, p') < \delta$  implies

$$\begin{aligned}
\left| \frac{\langle C \rangle_q(p) - \langle C \rangle_q(p')}{(\langle C \rangle_q)_{N, \max}} \right| &\leq \frac{\sum_{i=1}^N |p_i^q - (p'_i)^q| |C_i|}{\left| (\langle C \rangle_q)_{N, \max} \right|} \leq \frac{\sum_{i=1}^N |p_i^q - (p'_i)^q| |C_i|}{\max_{i=1, \dots, N} |C_i| \cdot 1^q} \\
&\leq \sum_{i=1}^N |p_i^q - (p'_i)^q| \leq \left( \sum_{i=1}^N |p_i - p'_i|^q \right) \\
&= (d_{\alpha, N}(p, p'))^\alpha \leq (d_{\alpha, N}(p, p'))^\alpha < \delta^\alpha = \varepsilon
\end{aligned}$$

c) Let  $q \in ]0, 1[$ , and  $\varepsilon = \frac{1}{2} > 0$ , for all  $\delta > 0$ , there is  $N = 2W = 2E \left( \left( \frac{2}{\delta} \right)^{\frac{q}{1-q}} \right) + 2 \in \mathbf{N}^*$ , let  $p, p'$  defined by

$$p = (p_i)_{i=1, \dots, N}, \quad p_i = \left( \frac{1}{W} \right)^{1/q} \text{ if } i \text{ is odd and } p_i = 0 \text{ if } i \text{ is even}$$

$$p' = (p'_i)_{i=1, \dots, N}, \quad p'_i = \left( \frac{1}{W} \right)^{1/q} - p_i$$

We have

$$\sum_{i=1}^N (p_i)^q = \sum_{i=1}^W \left( \left( \frac{1}{W} \right)^{1/q} \right)^q = W \cdot \frac{1}{W} = 1, \quad \sum_{i=1}^N (p'_i)^q = \sum_{i=1}^W \left( \left( \frac{1}{W} \right)^{1/q} \right)^q = W \cdot \frac{1}{W} = 1$$

$$d_{1, N}(p, p') = \sum_{i=1}^N |p_i - p'_i| = N \left( \frac{1}{W} \right)^{1/q} = 2W^{1-1/q} < \delta$$

$$C = (C_i)_{i=1, \dots, N}, \quad C_i = 1 \text{ if } i \text{ is odd and } C_i = 0 \text{ if } i \text{ is even}$$

$$\left| (\langle C \rangle_q)_{N, \max} \right| = \sum_{i=1}^W \left( \left( \frac{1}{W} \right)^{1/q} \right) = W^{1-1/q}$$

$$\text{And } \left| \frac{\langle C \rangle_q(p) - \langle C \rangle_q(p')}{(\langle C \rangle_q)_{N, \max}} \right| = \left| \frac{W \left( \frac{1}{W} - 0 \right)}{W^{1-1/q}} \right| = W^{\frac{1}{q}-1} > 1 > \varepsilon.$$

Consequently, the Incomplete  $q$ -expectation is 1-unstable.

### The $\alpha$ -stability of the Rényi entropy

#### Definition 13

Let  $q \in ]0, +\infty[ \setminus \{1\}$ . The Rényi entropy  $R_q$  is defined on  $A = \bigcup_{n \in \mathbf{N}^*} \left\{ p \in [0, 1]^n, \sum_{i=1}^n p_i = 1 \right\}$  by

$$R_q(p) = \frac{\ln \left( \sum_{i=1}^N (p_i)^q \right)}{1-q} \text{ for all } p \in [0, 1]^N.$$

**Theorem 14**

For  $q \in ]0,1[$ , the Rényi entropy  $R_q$  is  $\alpha$ -stable for all  $0 < \alpha \leq q$ .

*Proof.*

It's easy to see that  $\left| (R_q)_{\max} \right| = \left| R_q((1/N), \dots, (1/N)) \right| = \left| \frac{\ln(N^{1-q})}{1-q} \right| = \ln N$ .

For  $q \in ]0,1[$ , let  $\varepsilon > 0$ , there exists  $\delta = (\varepsilon)^{1/\alpha} > 0$ , such that for all  $N \in \mathbf{N}^* \setminus \{1,2\}$ ,  $p, p' \in A \cap \mathbf{R}^N$ ,  $d_{\alpha,N}(p, p') < \delta$  implies

$$\left| \frac{R_q(p) - R_q(p')}{(R_q)_{\max}} \right| = \frac{\left| \ln \left( \sum_{i=1}^N p_i^q \right) - \ln \left( \sum_{i=1}^N (p'_i)^q \right) \right|}{|\ln N|}$$

By the Mean-Value Theorem, there exists  $c_{q,N} \in \left( \sum_{i=1}^N p_i^q, \sum_{i=1}^N (p'_i)^q \right)$  such that

$$\left| \ln \left( \sum_{i=1}^N p_i^q \right) - \ln \left( \sum_{i=1}^N (p'_i)^q \right) \right| = \left| \sum_{i=1}^N p_i^q - \sum_{i=1}^N (p'_i)^q \right| \frac{1}{c_{q,N}}$$

Because  $\sum_{i=1}^N p_i^q \geq 1$  and  $\sum_{i=1}^N (p'_i)^q \geq 1$ , then  $\frac{1}{c_{q,N}} \leq 1$ , consequently

$$\left| \frac{\ln \left( \sum_{i=1}^N p_i^q \right) - \ln \left( \sum_{i=1}^N (p'_i)^q \right)}{\ln N} \right| = \frac{\left| \sum_{i=1}^N p_i^q - \sum_{i=1}^N (p'_i)^q \right|}{\ln N} \frac{1}{c_{q,N}} \leq \sum_{i=1}^N \left| p_i^q - (p'_i)^q \right|$$

By a) of lemma 10, we obtain  $\left| \frac{R_q(p) - R_q(p')}{(R_q)_{\max}} \right| \leq (d_{q,N}(p, p'))^q \leq (d_{\alpha,N}(p, p'))^\alpha < \delta^\alpha = \varepsilon$ .

**Acknowledgements**

The authors would like to thank Prof. P. Lescot for discussions on the subject.

**References**

- [1] B Lesche, Instabilities of Rényi Entropies. Journal of Statistical Physics, Vol. 27, No. 2, 1982
- [2] Q. A. Wang, Euro. Phys. J. B, **26** (2002)357;  
Q. A. Wang, A. Le Méhauté, L. Nivanen, M. Pezeril, Physica A, **340** (2004)117
- [3] C. Tsallis, J. Stat. Phys. **52** (1988)479
- [4] S. Abe, G. Kaniadakis, A. M. Scarfone; Stability of generalized entropies. Physica A, Math. Gen. **37** (2004) 10513-10519
- [5] S. Abe, Stability of Tsallis entropy and instabilities of Rényi and normalized Tsallis entropies: A basis for  $q$ -exponential distributions. Phy. Rev. E. **66** (2002) 0461134
- [6] G. Kaniadakis, A. M. Scarfone, Lesche Stability of  $\kappa$ -Entropy. arXiv:cond-mat/0310728v2
- [7] M F. Curado, F. D. Nobre, On the stability of analytic entropic form. Physica A 335 (2004) 94-106