

A Quantum Fluid Description of the Free Electron Laser

R. Bonifacio¹, N. Piovella², G.R.M Robb³ & A. Serbeto⁴

¹*I.N.F.N. Sezione di Milano, Via Celoria 16, Milano I-20133,Italy*

²*Dipartimento di Fisica, Universita degli Studi di Milano, via Celoria 16, 20133 Milano*

³*Scottish Universities Physics Alliance (SUPA),*

Department of Physics, University of Strathclyde,

107 Rottenrow, Glasgow, G4 0NG, United Kingdom

⁴*Instituto de Física, Universidade Federal Fluminense, 24210-340 Rio de Janeiro, Brazil*

(Dated: November 2, 2018)

Abstract

Using the Madelung transformation we show that in a quantum Free Electron Laser (QFEL) the beam obeys the equations of a quantum fluid in which the potential is the classical potential plus a quantum potential. The classical limit is shown explicitly.

INTRODUCTION

In the quantum FEL model, the electron beam is described as a macroscopic matter-wave [1, 2, 3, 4]. When slippage due to the difference between the light and electron velocities is neglected, the electron beam-wave interaction is described by the following equations for the dimensionless radiation amplitude $A(\bar{z})$ and the matter wave field $\Psi(\theta, \bar{z})$ [5]:

$$i \frac{\partial \Psi(\theta, \bar{z})}{\partial \bar{z}} = -\frac{1}{2\bar{\rho}} \frac{\partial^2}{\partial \theta^2} \Psi(\theta, \bar{z}) - i\bar{\rho} [A(\bar{z})e^{i\theta} - \text{c.c.}] \Psi(\theta, \bar{z}) \quad (1)$$

$$\frac{dA(\bar{z})}{d\bar{z}} = \int_0^{2\pi} d\theta |\Psi(\theta, \bar{z})|^2 e^{-i\theta} + i\delta A(\bar{z}). \quad (2)$$

The electron beam is therefore described by a Schrödinger equation for a matter-wave field Ψ in a self-consistent pendulum potential proportional to A , where $|A|^2 = |a|^2/(N\bar{\rho})$, $|a|^2$ is the average number of photons in the interaction volume V , and $|\Psi|^2$ is the space-time dependent electron density, normalized to unity. In Eqs. (1) and (2) we have adopted the universal scaling used in the classical FEL theory [6, 7, 8], *i.e.* $\theta = (k + k_w)z - ckt$ is the electron phase, where $k_w = 2\pi/\lambda_w$ and $k = \omega/c = 2\pi/\lambda$ are the wiggler and radiation wavenumbers, $\bar{z} = z/L_g$ is the dimensionless wiggler length, $L_g = \lambda_w/4\pi\rho$ is the gain length, $\rho = \gamma_r^{-1}(a_w/4ck_w)^{2/3}(e^2n/m\epsilon_0)^{1/3}$ is the classical FEL parameter, $\gamma_r = \sqrt{(\lambda/2\lambda_w)(1 + a_w^2)}$ is the resonant energy in mc^2 units, a_w is the wiggler parameter and n is the electron density. Finally, $\bar{p} = (\gamma - \gamma_0)/\rho\gamma_0$ is the dimensionless electron momentum and $\delta = (\gamma_0 - \gamma_r)/\rho\gamma_0$ is the detuning parameter, where $\gamma_0 \approx \gamma_r$ is the initial electron energy in mc^2 units.

Whereas the classical FEL equations in the above universal scaling do not contain any explicit parameter (see ref. [8]), the quantum FEL equations (1) and (2) depend on the quantum FEL parameter

$$\bar{\rho} = \left(\frac{mc\gamma_r}{\hbar k} \right) \rho. \quad (3)$$

From the definition of A , it follows that $\bar{\rho}|A|^2 = |a|^2/N$ is the average number of photons emitted per electron. Hence, since in the classical steady-state high-gain FEL A reaches a maximum value of the order of unity, $\bar{\rho}$ represents the maximum number of photons emitted per electron, and the classical regime occurs for $\bar{\rho} \gg 1$. Note also that in Eq. (1) $\bar{\rho}$ appears as a “mass” term, so one expects a classical limit when the mass is large. As we shall see, when $\bar{\rho} < 1$ the dynamical behavior of the system changes substantially from a classical to a quantum regime.

QUANTUM FLUID DESCRIPTION

We now perform a Madelung-like transformation [9], writing the wavefunction as

$$\Psi = R \exp(i\bar{\rho}S)$$

which allows us to rewrite the Maxwell-Schrodinger equations, eq. (1) and (2), as a system of quantum fluid equations

$$\frac{\partial R}{\partial \bar{z}} = -\frac{\partial R}{\partial \theta} \frac{\partial S}{\partial \theta} - \frac{R}{2} \frac{\partial^2 S}{\partial \theta^2} \quad (4)$$

$$\frac{\partial S}{\partial \bar{z}} = -\frac{1}{2} \left(\frac{\partial S}{\partial \theta} \right)^2 - V(\theta, \bar{z}) \quad (5)$$

$$\frac{dA}{d\bar{z}} = \int_0^{2\pi} R^2 e^{-i\theta} d\theta + i\delta A \quad (6)$$

where the potential, V in eq. (5) is defined as the sum of a classical term and a quantum term i.e.

$$V(\theta, \bar{z}) = V_C + V_Q$$

where

$$V_C = -i(Ae^{i\theta} - c.c.) \quad (7)$$

is the classical component of the potential and

$$V_Q = -\frac{1}{2\bar{\rho}^2 R} \frac{\partial^2 R}{\partial \theta^2} \quad (8)$$

is the quantum component of the potential, which becomes negligible as $\bar{\rho} \rightarrow \infty$.

Defining fluid density and velocity variables

$$n = R^2 = |\Psi|^2, \quad u = \frac{\partial S}{\partial \theta}$$

we can also rewrite Eq. (4)-(6) in an alternative fluid form as

$$\frac{\partial n}{\partial \bar{z}} + \frac{\partial}{\partial \theta}(nu) = 0 \quad (9)$$

$$\frac{\partial u}{\partial \bar{z}} + u \frac{\partial u}{\partial \theta} = -\frac{\partial V}{\partial \theta} \quad (10)$$

$$\frac{dA}{d\bar{z}} = \int_0^{2\pi} n e^{-i\theta} d\theta + i\delta A. \quad (11)$$

It can be seen that Eq. (9) is a continuity equation and Eq. (10) is a Newton-like equation for a fluid. Note that integrating Eq. (9) with respect to θ , then the normalization condition becomes

$$\int_0^{2\pi} n(\theta, \bar{z}) d\theta = 1,$$

which is satisfied if n and u are periodic functions of θ between 0 and 2π .

A straightforward calculation shows that Eq.(9)-(11) admit two constants of motion,

$$\langle \bar{p} \rangle + |A|^2 = \mathcal{C}_1 \quad (12)$$

and

$$\frac{\langle \bar{p}^2 \rangle}{2} - i(Ab^* - c.c.) - \delta|A|^2 = \mathcal{C}_2 \quad (13)$$

where $\langle \bar{p} \rangle = \langle u \rangle = \int_0^{2\pi} d\theta nu$ is the average momentum, $\langle \bar{p}^2 \rangle = \langle u^2 + 2V_Q \rangle = \int_0^{2\pi} d\theta n(u^2 + 2V_Q)$ is the momentum variance and

$$b = \int_0^{2\pi} n e^{-i\theta} d\theta$$

is the bunching. These constants of motion are well-known in the classical FEL model [8] and describe energy conservation and a gain-spread relation. Notice the quantum contribution to the momentum variance proportional to the average quantum potential.

FOURIER EXPANSION AND LINEAR ANALYSIS

If R and S are periodic functions of θ , they can be expanded in a Fourier series:

$$R(\theta, \bar{z}) = \sum_m r_m(\bar{z}) e^{im\theta}, \quad (14)$$

$$S(\theta, \bar{z}) = \sum_n s_n(\bar{z}) e^{in\theta} \quad (15)$$

with $r_{-m} = s_m^*$ and $s_{-m} = r_m^*$, since R and S are real variables. Multiplying Eq.(5) by R and using (14) and (15) in Eqs.(4)-(6), we obtain:

$$\sum_m r_{k-m} \frac{ds_m}{d\bar{z}} = -\frac{1}{2} \sum_{m,n} n(n-m) r_{k-m} s_n s_{n-m}^* + i(Ar_{k-1} - A^*r_{k+1}) - \frac{k^2}{2\bar{\rho}^2} r_k \quad (16)$$

$$\frac{dr_k}{d\bar{z}} = \frac{1}{2} \sum_m (k^2 - m^2) r_m s_{m-k}^* \quad (17)$$

$$\frac{dA}{d\bar{z}} = \sum_m r_m r_{m-1}^* + i\delta A \quad (18)$$

Eqs.(16)- (18) are our working equations which can be numerically solved as it will be shown elsewhere.

Eqs.(16)-(18) admit an equilibrium solution with no field ($A = 0$) and unbunched electron beam ($n = 1/2\pi$, i.e. $r_n = \delta_{n0}$ and $s_n = 0$). Linearizing Eqs.(16)-(18) around this

equilibrium in the first order of the variables A , r_1 and s_1 , we obtain:

$$\frac{dA}{d\bar{z}} = 2r_1 + i\delta A \quad (19)$$

$$\frac{dr_1}{d\bar{z}} = \frac{s_1}{2} \quad (20)$$

$$\frac{ds_1}{d\bar{z}} = iA - \frac{1}{2\bar{\rho}^2}r_1 \quad (21)$$

Looking for solutions proportional to $\exp(i\lambda\bar{z})$, we obtain the well-known cubic equation of the quantum FEL [1]

$$(\lambda - \delta) \left(\lambda^2 - \frac{1}{4\bar{\rho}^2} \right) + 1 = 0. \quad (22)$$

which reduces to the classical dispersion relation in the limit $\bar{\rho} \gg 1$.

CONCLUSIONS

It has been shown that the quantum FEL model can be rewritten in a form where the electron beam is described a quantum fluid coupled to the electromagnetic field. The evolution of the quantum fluid is determined by a self-consistent potential which consists of a classical and quantum contribution. In the limit where $\bar{\rho} \gg 1$ the quantum contribution to the potential becomes negligible and the force equation reduces to that of a Newtonian fluid. Using a Fourier expansion, linear stability analysis of these quantum fluid equations produced a dispersion relation identical to that derived from the Schrodinger equation. These results show that there are interesting connections between the quantum FEL and quantum plasma instabilities.

-
- [1] R. Bonifacio, M.M. Cola, N. Piovella, and G.R.M. Robb, *Europhys. Lett.* **69**, 55 (2005) .
 - [2] R. Bonifacio, N. Piovella, G.R.M. Robb, *Nucl. Instrum. and Meth. in Phys. Res. A* **543**, 645 (2005).
 - [3] R. Bonifacio, N. Piovella, G.R.M. Robb, and A. Schiavi, *Phys. Rev. ST Accel. Beams* **9**, 090701 (2006).
 - [4] A. Serbeto, J. T. Mendonça, K. H Tsui, R. Bonifacio, *Physics of Plasmas* **15**, 013110 (2008).
 - [5] G. Preparata, *Phys. Rev.A* **38**, 233 (1988).
 - [6] R. Bonifacio, C. Pellegrini and L. Narducci, *Opt. Commun.* **50**, 373 (1984).

- [7] R. Bonifacio, L. De Salvo, P. Pierini, N. Piovella, and C. Pellegrini, *Phys. Rev. Lett.* **73**, 70 (1994).
- [8] R. Bonifacio, F. Casagrande, G. Cerchioni, L. De Salvo Souza, P. Pierini and N. Piovella, *Rivista del Nuovo Cimento* **13**, No. 9 (1990).
- [9] A. Messiah, *Quantum Mechanics*, Wiley (1966).