Quantum hypothesis testing and sufficient subalgebras

Anna Jenčová*

Mathematical Institute, Slovak Academy of Sciences Štefánikova 49, 814 73 Bratislava, Slovakia

We introduce a new notion of a sufficient subalgebra for quantum states: a subalgebra is 2- sufficient for a pair of states $\{\rho_0, \rho_1\}$ if it contains all Bayes optimal tests of ρ_0 against ρ_1 . In classical statistics, this corresponds to the usual definition of sufficiency. We show this correspondence in the quantum setting for some special cases. Furthermore, we show that sufficiency is equivalent to 2 - sufficiency, if the latter is required for $\{\rho_0^{\otimes n}, \rho_1^{\otimes n}\}$, for all n.

MSC: 46L53, 81R15, 62B05.

Key words: quantum hypothesis testing, sufficient subalgebras, 2sufficiency, quantum Chernoff bound

1 Introduction

In order to motivate our results, let us consider the following problem of classical statistics. Suppose that P_0 and P_1 are two probability distributions and the task is to discriminate between them by an *n*-dimensional observation vector X. The problem is, if there is a function (statistic) $T : X \to Y$, such that the vector Y = T(X) (usually of lower dimension) contains all information needed for the discrimination.

^{*}Email: jenca@mat.savba.sk. Supported by the Slovak Research and Development Agency under the contract No. APVV 0071-06, grant VEGA 2/0032/09, Center of Excellence SAS - Quantum Technologies and ERDF OP R&D Project CE QUTE ITMS 26240120009

In the setting of hypothesis testing, the null hypothesis $H_0 = P_0$ is tested against the alternative $H_1 = P_1$. In the most general formulation, a test is a measurable function $\varphi : X \to [0, 1]$, which can be interpreted as the probability of rejecting the hypothesis if $x \in X$ occurs. There are two kinds of errors appearing in hypothesis testing: it may happen that H_0 is rejected, although it is true (error of the first kind), or that it is not rejected when H_1 is true (error of the second kind). For a given test φ , the error probabilities are

$$\alpha(\varphi) = \int \varphi(x) P_0(dx) \text{ first kind}$$

$$\beta(\varphi) = \int (1 - \varphi(x)) P_1(dx) \text{ second kind}$$

The two kinds of errors are in some sense complementary and it is usually not possible to minimize both error probabilities simultaneously. In the Bayesian approach, we choose a prior probability distribution $\{\lambda, 1 - \lambda\}$, $\lambda \in [0, 1]$ on the two hypotheses and then minimize the average (Bayes) error probability

$$\int \varphi(x)\lambda P_0(dx) + \int (1-\varphi(x))(1-\lambda)P_1(dx) = \lambda\alpha(\varphi) + (1-\lambda)\beta(\varphi).$$

Suppose now that T is a sufficient statistic for $\{P_0, P_1\}$. Roughly speaking, this means that there exists a common version of the conditional expectation $E[\cdot|T] = E_{P_0}[\cdot|T]$, P_0 - a.s. and $E[\cdot|T] = E_{P_1}[\cdot|T]$, P_1 - a.s. If φ is any test, then $E[\varphi|T]$ is another test having the same error probabilities. It follows that we can always have an optimal test that is a function of T, so that only values of T(X) are needed for optimal discrimination between P_0 and P_1 .

The following theorem states that this can happen if and only if T is sufficient, so that the above property characterizes sufficient statistics. The theorem was proved by Pfanzagl, see also [16].

Theorem 1 [15] Let $T: X \to Y$ be a statistic. The following are equivalent.

1. For any $\lambda \in (0,1)$ and any test $\varphi : X \to [0,1]$, there exists a test $\psi : Y \to [0,1]$, such that

$$\lambda \alpha(\psi \circ T) + (1 - \lambda)\beta(\psi \circ T) \le \lambda \alpha(\varphi) + (1 - \lambda)\beta(\varphi)$$

2. T is a sufficient statistic for $\{P_0, P_1\}$.

The problem of hypothesis testing can be considered also in the quantum setting. Here we deal with a pair of density operators $\rho_0, \rho_1 \in B(\mathcal{H})$, where \mathcal{H} is a finite dimensional Hilbert space and all tests are given by operators $0 \leq M \leq 1, M \in B(\mathcal{H})$. The problem of finding the optimal tests (the quantum Neyman-Pearson tests) and average error probabilities was solved by Helstrom and Holevo [6, 8].

Here a question arises, if it is possible to discriminate the states optimally by measuring on a given subsystem. Then we can gain some information only on the restricted densities, which, in general, can be distinguished with less precision.

Let $M_0 \subseteq B(\mathcal{H})$ be the subalgebra describing the subsystem we have access to. The average error probabilities for tests in M_0 are usually higher than the optimal ones. We will consider the situation that this does not happen and M_0 contains some optimal tests for all prior probabilities. In agreement with classical terminology (see [16]), such a subalgebra will be called sufficient with respect to testing problems, or 2-sufficient, for $\{\rho_0, \rho_1\}$.

The quantum counterpart of sufficiency was introduced and studied by Petz, see Chap. 9. in [13], in a more general context. According to this definition, the subalgebra M_0 is sufficient for $\{\rho_0, \rho_1\}$, if there exists a completely positive, trace preserving map $M_0 \to B(\mathcal{H})$, that maps both restricted densities to the original ones. Then the restriction to M_0 preserves all information needed for discrimination between the states and it is quite easy to see that a sufficient subalgebra must be 2-sufficient.

The conditions for sufficiency seem to be quite restrictive (see for example the factorization conditions in [9]) and might be too strong, if only hypothesis testing is considered. It is therefore natural to ask if there is a quantum version of Theorem 1, that is, if every 2-sufficient subalgebra must be sufficient.

In this paper, we give a partial answer to this question. We show that 2-sufficiency and sufficiency are equivalent under each of the following conditions: 1) the subalgebra M_0 is invariant under the modular group of one of the states, 2) M_0 is commutative, 3) ρ_0 and ρ_1 commute. Moreover, we show that if the 2-sufficiency condition is strengthened to hold for n independent copies of the densities for all n, then the two notions become equivalent.

The organization of the paper is as follows. In Section 2, some basic notions are introduced and several characterizations of a sufficient subalgebra are given. A new characterization, based on a version of the Radon-Nikodym derivative, is found, this will be needed for the main results. Section 3 gives the quantum Neyman-Pearson lemma and quantum Chernoff bound. Section 4 contains the main results: a convenient necessary condition for 2-sufficiency is found and it is shown that it implies sufficiency in the three above described cases. Finally, the quantum Chernoff bound is utilized to treat the case when 2-sufficiency holds for n independent copies of the states, for all n.

2 Some basic definitions and facts

2.1 Generalized conditional expectation

Let \mathcal{H} be a finite dimensional Hilbert space and let ρ be an invertible density matrix. Let $M_0 \subseteq B(\mathcal{H})$ be a subalgebra and let $E : B(\mathcal{H}) \to M_0$ be the trace preserving conditional expectation. Then $E(\rho)$ is the restricted density of the state ρ .

As we have seen, the classical sufficient statistic is defined by certain property of the conditional expectations. It is well known that in the quantum case, a state preserving conditional expectation does not always exist. Therefore we need the generalized conditional expectation, defined by Accardi and Cecchini [1]. In our setting, it can be given as follows.

Let us introduce the inner product $\langle X, Y \rangle_{\rho} = \operatorname{Tr} X^* \rho^{1/2} Y \rho^{1/2}$ in $B(\mathcal{H})$. Then the generalized conditional expectation E_{ρ} is a map $B(\mathcal{H}) \to M_0$, defined by

$$\langle X_0, Y \rangle_{\rho} = \langle X_0, E_{\rho}(Y) \rangle_{E(\rho)}, \quad X_0 \in M_0, \ Y \in B(\mathcal{H})$$

It is easy to see that we have

$$E_{\rho}(X) = E(\rho)^{-1/2} E(\rho^{1/2} X \rho^{1/2}) E(\rho)^{-1/2}$$
(1)

It is known that E_{ρ} is completely positive and unital and that it is a conditional expectation if and only if $\rho^{it}M_0\rho^{-it} \subseteq M_0$, for all $t \in \mathbb{R}$. It is also easy to see that E_{ρ} preserves the state ρ , that is, $E_{\rho}^* \circ E(\rho) = \rho$.

Next we introduce two subalgebras, related to E_{ρ} . Let F_{ρ} be the set of fixed points of E_{ρ} and let $N_{\rho} \subseteq B(\mathcal{H})$ be the multiplicative domain of E_{ρ} ,

$$N_{\rho} = \{ X \in B(\mathcal{H}), E_{\rho}(X^*X) = E_{\rho}(X)^* E_{\rho}(X), E_{\rho}(XX^*) = E_{\rho}(X) E_{\rho}(X)^* \}$$

Then both F_{ρ} and N_{ρ} are subalgebras in $B(\mathcal{H})$. It is clear that $F_{\rho} \subseteq M_0 \cap N_{\rho}$, moreover, $X \in F_{\rho}$ if and only if $\rho^{it} X \rho^{-it} \in M_0$ for all $t \in \mathbb{R}$. As for N_{ρ} , we have the following result.

Lemma 1 $N_{
ho} = \rho^{1/2} M_0 \rho^{-1/2} \cap \rho^{-1/2} M_0 \rho^{1/2}$

Proof. It is clear from (1) that $X \in N_{\rho}$ if and only if

$$E(\rho^{1/2}X^*X\rho^{1/2}) = E(\rho^{1/2}X^*\rho^{1/2})E(\rho)^{-1}E(\rho^{1/2}X\rho^{1/2})$$
$$E(\rho^{1/2}XX^*\rho^{1/2}) = E(\rho^{1/2}X\rho^{1/2})E(\rho)^{-1}E(\rho^{1/2}X^*\rho^{1/2})$$

Let $A = X\rho^{1/2}$, $B = \rho^{1/2}$. Similarly as in [11], we put $M = A - B\Lambda$, with $\Lambda = E(\rho)^{-1}E(\rho^{1/2}X\rho^{1/2})$. Then from $E(M^*M) \ge 0$, we obtain

$$E(A^*A) \ge E(A^*B)E(\rho)^{-1}E(B^*A),$$

with equality if and only if M = 0, this implies

$$\rho^{-1/2} X \rho^{1/2} = E(\rho)^{-1} E(\rho^{1/2} X \rho^{1/2}) \in M_0.$$

Conversely, let $X_0 = \rho^{-1/2} X \rho^{1/2} \in M_0$, then $E(\rho^{1/2} X \rho^{1/2}) = E(\rho) X_0$, this implies that M = 0.

Similarly, we get that $\rho^{-1/2} X^* \rho^{1/2} \in M_0$ is equivalent with the second equality.

It is also known that $E_{\rho}(XY) = E_{\rho}(X)E_{\rho}(Y), E_{\rho}(YX) = E_{\rho}(Y)E_{\rho}(X)$ for all $X \in N_{\rho}, Y \in B(\mathcal{H})$, this can be also shown from the above Lemma. Note that in the case that E_{ρ} is a conditional expectation, $F_{\rho} = N_{\rho} = M_0$.

2.2 A Radon-Nikodym derivative and relative entropies

Let ρ_0, ρ_1 be invertible density matrices in $B(\mathcal{H})$. We will use the quantum version of the Radon-Nikodym derivative introduced in [5]. In our setting, the derivative d_{ρ_0,ρ_1} of ρ_1 with respect to ρ_0 is defined as the unique element in $B(\mathcal{H})$, such that $\operatorname{Tr} \rho_1 X = \langle X^*, d_{\rho_0,\rho_1} \rangle_{\rho_0}$. Then clearly

$$d_{\rho_0,\rho_1} = \rho_0^{-1/2} \rho_1 \rho_0^{-1/2}$$

so that d_{ρ_0,ρ_1} is positive, and $||d_{\rho_0,\rho_1}|| \leq \lambda$ for any $\lambda > 0$, such that $\rho_1 \leq \lambda \rho_0$. It is also easy to see that

$$E_{\rho_0}(d_{\rho_0,\rho_1}) = d_{E(\rho_0),E(\rho_1)}$$

Let us recall that the Belavkin - Staszewski relative entropy is defined as [5]

$$S_{BS}(\rho_1, \rho_0) = -\operatorname{Tr} \rho_0 \eta(\rho_0^{-1/2} \rho_1 \rho_0^{-1/2}) = -\operatorname{Tr} \rho_0 \eta(d_{\rho_0, \rho_1})$$

where $\eta(x) = -x \log(x)$. Let S be the Umegaki relative entropy

$$S(\rho_1, \rho_0) = \operatorname{Tr} \rho_1(\log \rho_1 - \log \rho_0)$$

then $S(\rho_1, \rho_0) \leq S_{BS}(\rho_1, \rho_0)$, [7] and $S(\rho_1, \rho_0) = S_{BS}(\rho_1, \rho_0)$ if ρ_0 and ρ_1 commute. Both relative entropies are monotone in the sense that

$$S(\rho_1, \rho_0) \ge S(E(\rho_1), E(\rho_0)), \quad S_{BS}(\rho_1, \rho_0) \ge S_{BS}(E(\rho_1), E(\rho_0))$$

holds for any subalgebra M_0 . As we will see in the next section, equality in the monotonicity for S is equivalent with sufficiency of the subalgebra M_0 with respect to $\{\rho_0, \rho_1\}$. For S_{SB} , we have the following result. Lemma 2 The following are equivalent.

- (i) $S_{BS}(\rho_1, \rho_0) = S_{BS}(E(\rho_1), E(\rho_0))$
- (*ii*) $d_{\rho_0,\rho_1} \in N_{\rho_0}$
- (*iii*) $\rho_1 \rho_0^{-1} \in M_0$
- (*iv*) $\rho_1 \rho_0^{-1} = E(\rho_1) E(\rho_0)^{-1}$

Proof. Since the function $-\eta(x) = x \log(x)$ is operator convex,

$$\eta(d_{E(\rho_0),E(\rho_1)}) = \eta(E_{\rho_0}(d_{\rho_0,\rho_1})) \le E_{\rho_0}(\eta(d_{\rho_0,\rho_1}))$$
(2)

by Jensen's inequality. We have

$$\operatorname{Tr} \rho_0(E_{\rho_0}(\eta(d_{\rho_0,\rho_1})) - \eta(E_{\rho_0}(d_{\rho_0,\rho_1}))) = S_{BS}(\rho_1,\rho_0) - S_{BS}(E(\rho_1),E(\rho_0))$$

and since ρ_0 is invertible, equality in the monotonicity of S_{BS} is equivalent with equality in (2). As it was proved in [14], this happens if and only if $d_{\rho_0,\rho_1} \in N_{\rho_0}$. This shows the equivalence (i) \leftrightarrow (ii). The equivalence of (ii) and (iii) follows by Lemma 1, (iii) \iff (iv) is rather obvious.

2.3 Sufficient subalgebras

We say that the subalgebra $M_0 \subseteq B(\mathcal{H})$ is sufficient for $\{\rho_0, \rho_1\}$ if there is a completely positive trace preserving map $T : M_0 \to B(\mathcal{H})$, such that $T \circ E(\rho_0) = \rho_0$ and $T \circ E(\rho_1) = \rho_1$. The following characterizations of sufficiency were obtained by Petz.

Theorem 2 [10, 13] The following are equivalent.

- (i) $M_0 \subseteq B(\mathcal{H})$ is sufficient for $\{\rho_0, \rho_1\}$
- (*ii*) $S(\rho_1, \rho_0) = S(E(\rho_1), E(\rho_0))$
- (*iii*) Tr $\rho_0^s \rho_1^{1-s} = \text{Tr } E(\rho_0)^s E(\rho_1)^{1-s}$ for some $s \in (0,1)$
- (iv) $\operatorname{Tr} E_{\rho_0}(X)\rho_1 = \operatorname{Tr} X\rho_1 \text{ for all } X \in B(\mathcal{H})$
- $(v) E_{\rho_0} = E_{\rho_1}.$

The next characterization is based on the Radon-Nikodym derivative.

Theorem 3 The subalgebra $M_0 \subseteq B(\mathcal{H})$ is sufficient for $\{\rho_0, \rho_1\}$ if and only if $d_{\rho_0,\rho_1} \in F_{\rho_0}$.

Proof. Let us denote $d = d_{\rho_0,\rho_1}$ and $d_0 = d_{E(\rho_0),E(\rho_1)}$. Since $d_0 \in M_0$, we have by definition that

Tr
$$\rho_1 E_{\rho_0}(X) = \langle d_0, E_{\rho_0}(X) \rangle_{E(\rho_0)} = \langle d_0, X \rangle_{\rho_0}$$

so that $\operatorname{Tr} \rho_1 E_{\rho_0}(X) = \operatorname{Tr} \rho_1 X$ if and only if $\langle d_0, X \rangle_{\rho_0} = \langle d, X \rangle_{\rho_0}$. It follows that $d = d_0$ is equivalent with sufficiency of M_0 , by Theorem 2 (iv). Since $E_{\rho_0}(d) = d_0$, this is equivalent with $d_{\rho_0,\rho_1} \in F_{\rho_0}$.

3 Quantum hypothesis testing

Let us now turn to the problem of hypothesis testing. Any test of the hypothesis $H_0 = \rho_0$ against the alternative $H_1 = \rho_1$ is represented by an operator $0 \le M \le 1$, which corresponds to rejecting the hypothesis. Then we have the error probabilities

$$\alpha(M) = \operatorname{Tr} \rho_0 M$$
 first kind
 $\beta(M) = \operatorname{Tr} \rho_1 (1 - M)$ second kind

For $\lambda \in (0, 1)$, we define the Bayes optimal test to be a minimizer of the expression

$$\lambda \alpha(M) + (1 - \lambda)\beta(M) \tag{3}$$

It is clear that minimizing (3) is the same as maximizing

$$\operatorname{Tr}(\rho_1 - t\rho_0)M, \qquad t = \frac{\lambda}{1-\lambda}$$

3.1 The quantum Neyman-Pearson lemma

The following is the quantum version of the Neyman-Pearson lemma. The obtained optimal tests are called the (quantum) Neyman-Pearson tests. We give a simple proof for completeness.

Lemma 3 Let $t \ge 0$ and let us denote $P_{t,+} := \operatorname{supp} (\rho_1 - t\rho_0)_+$, $P_{t,-} := \operatorname{supp} (\rho_1 - t\rho_0)_-$ and $P_{t,0} := 1 - P_{t,+} - P_{t,-}$. Then the operator $0 \le M_t \le 1$ is a Bayes optimal test of ρ_0 against ρ_1 if and only if

$$M_t = P_{t,+} + X_t$$

where $0 \leq X_t \leq P_{t,0}$.

Proof. Let $0 \le M \le 1$, then

$$\operatorname{Tr} (\rho_{1} - t\rho_{0})M = \operatorname{Tr} (\rho_{1} - t\rho_{0})_{+}M - \operatorname{Tr} (\rho_{1} - t\rho_{0})_{-}M \leq \operatorname{Tr} (\rho_{1} - t\rho_{0})_{+}M$$

$$\leq \operatorname{Tr} (\rho_{1} - t\rho_{0})_{+} = \operatorname{Tr} (\rho_{1} - t\rho_{0})P_{t,+}$$
(4)

It follows that $M_t = P_{t,+} + X_t$, $X_t \leq P_{t,0}$ is a Bayes optimal test. Conversely, let M_t be some Bayes optimal test, then we must have

$$Tr (\rho_1 - t\rho_0)M_t = Tr (\rho_1 - t\rho_0)_+ M_t = Tr (\rho_1 - t\rho_0)P_{t,+}$$

so that $\text{Tr}(\rho_1 - t\rho_0)_- M_t = 0$. By positivity, this implies that $P_{t,-}M_t = M_t P_{t,-} = 0$, so that

$$M_t(P_{t,+} + P_{t,0}) = (P_{t,+} + P_{t,0})M_t = M_t$$

which is equivalent with $M_t \leq P_{t,+} + P_{t,0}$. Furthermore, from

$$\operatorname{Tr} \left(\rho_1 - t\rho_0\right)_+ \left(P_{t,+} + P_{t,0} - M_t\right) = 0$$

we obtain $P_{t,+} - P_{t,+}M_tP_{t,+} = P_{t,+}(1 - M_t)P_{t,+} = 0$, hence $(1 - M_t)P_{t,+} = 0$. We obtain $P_{t,+} \le M_t$ and by putting $X_t := M_t - P_{t,+}$, we get the result.

Let us denote by $\Pi_{e,\lambda}$ the minimum Bayes error probability. Then

$$\Pi_{e,\lambda} = \lambda \alpha(M_{\lambda/(1-\lambda)}) + (1-\lambda)\beta(M_{\lambda/(1-\lambda)}) =$$

= $\frac{1}{2}(1 - \|(1-\lambda)\rho_1 - \lambda\rho_0\|_1)$ (5)

where the last equality follows from

$$1 - t = \text{Tr} (\rho_1 - t\rho_0) = \text{Tr} (\rho_1 - t\rho_0)_+ - \text{Tr} (\rho_2 - t\rho_0)_-$$

and

$$\|\rho_1 - t\rho_0\|_1 = \operatorname{Tr} |\rho_1 - t\rho_0| = \operatorname{Tr} (\rho_1 - t\rho_0)_+ + \operatorname{Tr} (\rho_2 - t\rho_0)_-$$

3.2 The quantum Chernoff bound

Suppose now that we have *n* copies of the states ρ_0 and ρ_1 , so that we test the hypothesis $\rho_0^{\otimes n}$ against $\rho_1^{\otimes n}$ by means of an operator $0 \leq M_n \leq 1$, $M_n \in \mathcal{B}(\mathcal{H}^{\otimes n})$. Again, we may use the Neyman-Pearson lemma to find the minimum Bayes error probability

$$\Pi_{e,\lambda,n} = \frac{1}{2} (1 - \|(1 - \lambda)\rho_1^{\otimes n} - \lambda\rho_0^{\otimes n}\|_1)$$

The following important result, obtained in [3] and [12] (see also [4]), is the quantum version of the classical Chernoff bound:

$$\lim_{n} \left(-\frac{1}{n} \log \Pi_{e,\lambda,n} \right) = -\log(\inf_{0 \le s \le 1} \operatorname{Tr} \rho_0^{1-s} \rho_1^s) =: \xi_{QCB}(\rho_0, \rho_1)$$
(6)

The expression ξ_{QCB} has a number of interesting properties. For example, it was proved that it is always nonnegative and equal to 0 if and only if $\rho_0 = \rho_1$, moreover, it is monotone in the sense that

$$\xi_{QCB}(\rho_0,\rho_1) \ge \xi_{QCB}(E(\rho_0),E(\rho_1))$$

Therefore, although it is not symmetric, ξ_{QCB} provides a reasonable distance measure on density matrices, called the quantum Chernoff distance. Note also that in the case that the matrices are invertible, the infimum is always attained in some $s^* \in [0, 1]$.

4 2-sufficiency

We say that M_0 is sufficient with respect to testing problems, or 2-sufficient, for $\{\rho_0, \rho_1\}$ if for any test M and any $\lambda \in (0, 1)$, there is some test $N_{\lambda} \in M_0$, such that

$$\lambda \alpha(N_{\lambda}) + (1 - \lambda)\beta(N_{\lambda}) \le \lambda \alpha(M) + (1 - \lambda)\beta(M)$$

It is quite clear that M_0 is 2-sufficient if and only if for all $t \ge 0$, we can find a Neyman-Pearson test $M_t \in M_0$. Moreover, suppose that M_0 is a sufficient subalgebra for $\{\rho_0, \rho_1\}$ and let $T = E_{\rho_0} = E_{\rho_1}$. Then, if M_t is a Neyman-Pearson test, then $T(M_t) \in M_0$ is a Neyman-Pearson test as well. Hence, a sufficient subalgebra is always 2-sufficient. In this section, we find the opposite implication in some special cases.

Lemma 4 $P_{t,0} \neq 0$ if and only if t is an eigenvalue of $d := d_{\rho_0,\rho_1}$. Moreover, the rank of $P_{t,0}$ is equal to multiplicity of t.

Proof. By definition,

$$(\rho_1 - t\rho_0)P_{t,0} = \rho_0^{1/2}(d-t)\rho_0^{1/2}P_{t,0} = 0$$

so that $(d-t)\rho_0^{1/2}P_{t,0}\rho_0^{1/2} = 0$. Suppose $P_{t,0} \neq 0$, then t is an eigenvalue of d and any vector in the range of $\rho_0^{1/2}P_{t,0}\rho_0^{1/2}$ is an eigenvector. This implies that $r(P_{t,0}) = r(\rho^{1/2}P_{t,0}\rho^{1/2}) \leq r(F)$, where F is the eigenprojection of t. Conversely, let t be an eigenvalue of d with the eigenprojection F, then

$$(\rho_1 - t\rho_0)\rho_0^{-1/2}F\rho_0^{-1/2} = \rho_0^{1/2}(d-t)F\rho_0^{-1/2} = 0,$$

so that the range of $\rho^{-1/2} F \rho^{-1/2}$ is in the kernel of $\rho_1 - t \rho_0$, this implies $r(F) \leq r(P_{t,0})$.

Let us denote $Q_{t,+} = \operatorname{supp} (E(\rho_1) - tE(\rho_0))_+, Q_{t,0} = \ker (E(\rho_1) - tE(\rho_0))$ and let $\Pi_{e,\lambda}^0$ be the minimal Bayes error probability for the restricted densities

$$\Pi_{e,\lambda}^{0} := \inf_{M \in M_{0}} \lambda \alpha(M) + (1-\lambda)\beta(M) = \frac{1}{2}(1 - \|(1-\lambda)E(\rho_{1}) - \lambda E(\rho_{0})\|_{1})$$

Lemma 5 The following are equivalent.

- (i) The subalgebra M_0 is 2-sufficient for $\{\rho_0, \rho_1\}$.
- (*ii*) $\Pi^0_{e,\lambda} = \Pi_{e,\lambda}$ for all $\lambda \in (0,1)$.
- (iii) $Q_{t,0} = P_{t,0}$ and $Q_{t,+} = P_{t,+}$ for all $t \ge 0$.

Proof. It is obvious that (i) implies (ii). Suppose (ii) and let us denote $f(t) := \max_{0 \le M \le 1} \operatorname{Tr} (\rho_1 - t\rho_0) M$. If N_t is any Neyman-Pearson test for $\{E(\rho_0), E(\rho_1)\}$, then

$$Tr (\rho_1 - t\rho_0)N_t = Tr (E(\rho_1) - tE(\rho_0))N_t = f(t),$$

so that N_t is a Neyman-Pearson test for $\{\rho_0, \rho_1\}$ as well. Putting $N_t = Q_{t,+}$ and $N_t = Q_{t,+} + Q_{t,0}$, we get by Lemma 3 that

$$Q_{t,+} = P_{t,+} + X_t, \qquad Q_{t,+} + Q_{t,0} = P_{t,+} + Y_t,$$

with $X_t, Y_t \leq P_{t,0}$. This implies that $Q_{t,0} \leq P_{t,0}$ and $Q_{t,+} = P_{t,+}$ if $P_{t,0} = 0$.

Let t be an eigenvalue of d_0 , then $P_{t,0} \ge Q_{t,0} \ne 0$, hence t is also an eigenvalue of d, and its multiplicity in d_0 is not greater that its multiplicity in d. Since the sum of multiplicities must equal to $m = \dim(\mathcal{H})$, we must have $r(Q_{t,0}) = r(P_{t,0})$, so that $Q_{t,0} = P_{t,0}$. This implies that $X_t \le Q_{t,0}$, hence $X_t = 0$ and $P_{t,+} = Q_{t,+}$ for all t.

The implication (iii) \rightarrow (i) is again obvious.

Note that the condition (ii) is equivalent with

$$||E(\rho_1) - tE(\rho_0)||_1 \ge ||\rho_1 - t\rho_0||_1$$
, for all $t \ge 0$

This condition, with $E(\rho_0)$ and $E(\rho_1)$ replaced by arbitrary densities σ_0 and σ_1 was studied in [2]. It was shown that for 2×2 matrices, this is equivalent with the existence of a completely positive trace preserving map T, such that $T(\rho_0) = \sigma_0$ and $T(\rho_1) = \sigma_1$. In our case, this means that 2-sufficiency implies sufficiency for 2×2 matrices. Since any nontrivial subalgebra in $\mathcal{M}(\mathbb{C}^2)$ is commutative, this agrees with our results below.

The above Lemma gives characterizations of 2-sufficiency, but the conditions are not easy to check. The next Theorem gives a simple necessary condition.

Theorem 4 Let M_0 be 2-sufficient for $\{\rho_1, \rho_0\}$. Then $d_{\rho_1,\rho_0} \in N_{\rho_0}$.

Proof. By the previous Lemma, we have $P_{t,0} = Q_{t,0} \in M_0$ for all t. Let t_1, \ldots, t_k be the eigenvalues of d and denote $P_i = P_{t_i,0}$. Then from $(d-t_i)\rho_0^{1/2}P_i = 0$ we get

$$d\rho_0^{1/2} \sum_i P_i = \rho_0^{1/2} \sum_i t_i P_i$$

By Lemma 4 and its proof, $\operatorname{supp}(\rho_0^{1/2}P_i\rho_0^{1/2}) \leq F_i$ and $r(P_i) = r(F_i)$, with F_i the eigenprojection of t_i . It follows that $\sum_i \rho_0^{1/2}P_i\rho_0^{1/2}$, and hence also $\sum_i P_i$, is invertible. Therefore,

$$d\rho_0^{1/2} = \rho_0^{1/2} c, \qquad c := \sum_i t_i P_i (\sum_j P_j)^{-1}$$

that is, $d = \rho_0^{1/2} c \rho_0^{-1/2}$, with $c \in M_0$. Moreover, $d = d^* = \rho_0^{-1/2} c^* \rho_0^{1/2}$, so that $d \in \rho_0^{1/2} M_0 \rho_0^{-1/2} \cap \rho_0^{-1/2} M_0 \rho_0^{1/2}$. By Lemma 1, this entails that $d \in N_{\rho_0}$.

Theorem 5 Let the subalgebra M_0 be 2-sufficient for $\{\rho_0, \rho_1\}$. Then M_0 is sufficient for $\{\rho_0, \rho_1\}$ in each of the following cases.

- (1) $\rho_0^{it} M_0 \rho_0^{-it} \subseteq M_0 \text{ for all } t \in \mathbb{R}$
- (2) M_0 is commutative
- (3) ρ_0 and ρ_1 commute

Proof. (1) By Theorem 4, we have $d \in N_{\rho_0}$. Since $\rho_0^{it} M_0 \rho_0^{-it} \subseteq M_0$, we have $d \in N_{\rho_0} = F_{\rho_0}$. By Theorem 3, this implies that M_0 is sufficient.

(2) Since $d \in N_{\rho_0}$, we have $S_{BS}(\rho_1, \rho_0) = S_{BS}(E(\rho_1), E(\rho_0))$, by Lemma 2. Since M_0 is commutative,

$$S(E(\rho_1), E(\rho_0)) = S_{BS}(E(\rho_1), E(\rho_0)) = S_{BS}(\rho_1, \rho_0) \ge S(\rho_1, \rho_0)$$

By monotonicity of the relative entropy, this implies $S(\rho_1, \rho_0) = S(E(\rho_1), E(\rho_0))$, so that M_0 is sufficient for $\{\rho_0, \rho_1\}$, by Theorem 2 (ii).

(3) Let M_1 be the subalgebra generated by all $P_{t,+}$, $t \in \mathbb{R}$. Then M_1 is commutative and 2-sufficient for $\{\rho_0, \rho_1\}$, hence sufficient by (2). If M_0 is 2-sufficient, we must have $M_1 \subseteq M_0$ by Lemma 5, so that M_0 must be sufficient for $\{\rho_0, \rho_1\}$ as well.

It is clear from the proof of (1) that 2-sufficiency implies sufficiency whenever $N_{\rho_0} = F_{\rho_0}$ (or, equivalently, $N_{\rho_1} = F_{\rho_1}$). In fact, it can be shown that $N_{\rho_0} = F_{\rho_0}$ whenever M_0 is commutative, which gives an alternative proof of (2). Next we give a further example of this situation.

Example 1 Let $\mathcal{H} = \mathbb{C}^4$ and let $M_0 = \mathcal{M}(\mathbb{C}^2) \otimes I \subset B(\mathcal{H})$. Let ρ be a block-diagonal density matrix $\rho = \begin{pmatrix} \rho_1 & 0 \\ 0 & \rho_2 \end{pmatrix}$, where ρ_1, ρ_2 are positive invertible matrices in $\mathcal{M}(\mathbb{C}^2)$, and let σ be any density matrix. Suppose that M_0 is 2-sufficient for $\{\rho, \sigma\}$.

By Theorem 4, $d_{\sigma,\rho} \in N_{\rho}$, which by Lemma 2 is equivalent with $\sigma \rho^{-1} \in M_0$. This implies that σ must be block-diagonal as well, $\sigma = \begin{pmatrix} \sigma_1 & 0 \\ 0 & \sigma_2 \end{pmatrix}$.

By Lemma 5, $P_{t,+} \in M_0$ for all $t \ge 0$, so that $P_{t,+} = \begin{pmatrix} p_t & 0 \\ 0 & p_t \end{pmatrix}$, where $p_t = \operatorname{supp} (\sigma_1 - t\rho_1)_+ = \operatorname{supp} (\sigma_2 - t\rho_2)_+$. Since p_t is a projection in $\mathcal{M}(\mathbb{C}^2)$, we have he following two possibilities: either $p_t = I$ for $t < t_0$ and $p_t = 0$ for $t \ge t_0$, or p_t is one-dimensional for t in some interval (t_0, t_1) . Since $\rho = \sigma$ in the first case, we may suppose that the latter is true, so that p_t is a common eigenprojection of $\sigma_1 - t\rho_1$ and $\sigma_2 - t\rho_2$ for $t \in (t_0, t_1)$. It follows that $\sigma_1 - t\rho_1$ commutes with $\sigma_2 - t\rho_2$ for $t \in (t_0, t_1)$, which implies that ρ_1 commutes with ρ_2 .

Let $X \in N_{\rho}$, then $X = \rho^{1/2} X_0 \rho^{-1/2}$, where both X_0 , $\rho X_0 \rho^{-1} \in M_0$. Let $X_0 = Y \otimes I \in M_0$, then $\rho X_0 \rho^{-1} \in M_0$ if and only if $\rho_1 Y \rho_1^{-1} = \rho_2 Y \rho_2^{-1}$, that is, Y commutes with $\rho_2^{-1} \rho_1$. If $\rho_2^{-1} \rho_1$ is a constant, then $\rho^{it} M_0 \rho^{-it} \subseteq M_0$, so that $F_{\rho} = M_0 = N_{\rho}$. Otherwise, Y must commute with both ρ_1 and ρ_2 and in this case, $X = \rho^{1/2} X_0 \rho^{-1/2} = X_0 \in F_{\rho}$.

In conclusion, if M_0 is 2-sufficient for $\{\rho, \sigma\}$, we must have $N_{\rho} = F_{\rho}$, so that M_0 must be a sufficient subalgebra.

Let us now suppose that we have *n* independent copies of the states, $\rho_0^{\otimes n}$ and $\rho_1^{\otimes n}$. An optimal test for $H_1 : \rho_0^{\otimes n}$ against $H_1 : \rho_1^{\otimes n}$ usually cannot be obtained as the product of optimal tests, but we may ask if there is some

optimal test in $M_0^{\otimes n}$. If this is the case for all λ , we say that M_0 is (2, n)-sufficient for $\{\rho_0, \rho_1\}$.

Theorem 6 The following conditions are equivalent.

- (i) M_0 is (2, n)-sufficient for $\{\rho_0, \rho_1\}$, for all n.
- (ii) M_0 is a sufficient subalgebra for $\{\rho_0, \rho_1\}$.

Proof. Let us denote

$$\Pi^{0}_{e,\lambda,n} := \frac{1}{2} (1 - \| (1 - \lambda) E(\rho_1)^{\otimes n} - \lambda E(\rho_0)^{\otimes n} \|_1)$$

By Lemma 5 (ii), the condition (i) implies that $\Pi_{e,\lambda,n} = \Pi^0_{e,\lambda,n}$ for all n, hence also

$$\lim_{n} \left(-\frac{1}{n} \log \Pi_{e,\lambda,n}\right) = \lim_{n} \left(-\frac{1}{n} \log \Pi_{e,\lambda,n}^{0}\right)$$

By (6), this entails that

$$\inf_{0 \le s \le 1} \operatorname{Tr} \rho_0^{1-s} \rho_1^s = \inf_{0 \le s \le 1} \operatorname{Tr} E(\rho_0)^{1-s} E(\rho_1)^s$$

By monotonicity, we have $\operatorname{Tr} \rho_0^{1-s} \rho_1^s \leq \operatorname{Tr} E(\rho_0)^{1-s} E(\rho_1)^s$ for all $s \in [0, 1]$. Suppose that the infimum on the RHS is attained in some $s_0 \in [0, 1]$. Then

$$\operatorname{Tr} E(\rho_0)^{1-s_0} E(\rho_1)^{s_0} = \inf_{0 \le s \le 1} \operatorname{Tr} \rho_0^{1-s} \rho_1^s \le \operatorname{Tr} \rho_0^{1-s_0} \rho_1^{s_0}.$$

If $s_0 = 0$ or 1, then the quantum Chernoff distance is equal to 0, so that $\rho_0 = \rho_1$ and the subalgebra M_0 is trivially sufficient. Otherwise, we must have $\operatorname{Tr} E(\rho_0)^{1-s_0} E(\rho_1)^{s_0} = \operatorname{Tr} \rho_0^{1-s_0} \rho_1^{s_0}$ for $s_0 \in (0, 1)$, which implies that M_0 is sufficient for $\{\rho_0, \rho_1\}$, by Theorem 2 (iii).

Conversely, let $E_{\rho^{\otimes n}}$ be the generalized conditional expectation $B(\mathcal{H}^{\otimes n}) \rightarrow M_0^{\otimes n}$. It is easy to see that for any invertible density matrix ρ , $E_{\rho^{\otimes n}} = E_{\rho^{\otimes n}}^{\otimes n}$, so that if $E_{\rho_0} = E_{\rho_1}$, then $E_{\rho_0^{\otimes n}} = E_{\rho_1^{\otimes n}}^{\otimes n}$ for all n. Hence if M_0 is sufficient for $\{\rho_0, \rho_1\}$, then $M_0^{\otimes n}$ is sufficient for $\{\rho_0^{\otimes n}, \rho_1^{\otimes n}\}$ for all n, this implies (i).

References

 L. Accardi, C. Cecchini, Conditional expectations in von Neumann algebras and a theorem of Takesaki, J. Functional. Anal. 45(1982), 245–273.

- [2] P. M. Alberti, A. Uhlmann, A problem relating to the positive linear maps on a matrix algebra, Rep. Math. Phys. 18 (1980), 163–176
- [3] K.M.R. Audenaert, J. Calsamiglia, L. Masanes, R. Munoz-Tapia, A. Acin, E. Bagan, F. Verstraete, Discriminating states: The quantum Chernoff bound, Phys. Rev. Lett. 98, 160501 (2007)
- [4] K.M.R. Audenaert, M. Nussbaum, A. Szkola, F. Verstraete, Asymptotic error rates in quantum hypothesis testing, Comm. Math. Phys. 279, 251-283 (2008)
- [5] V.P. Belavkin, P. Staszewski, C^{*}- algebraic generalizations of relative entropy and entropy, Ann. Ins. Henri Poincaré Sec. A 73 (1982), 51–58
- [6] C.W. Helstrom, Quantum Detection and Estimation Theory, Academic Press, New York, 1976
- [7] F. Hiai, D. Petz, The proper formula for relative entropy and its asymptotics in quantum probability, Comm. Math. Phys. **143** (1991), 99–114
- [8] A.S. Holevo, On asymptotically optimal hypothesis testing in quantum statistics, Theor. Prob. Appl. 23 (1978), 411–415
- [9] A. Jenčová, D. Petz, Sufficiency in quantum statistical inference. Commun. Math. Phys. 263, 259276 (2006).
- [10] A. Jenčová, D. Petz, Sufficiency in quantum statistical inference. A survey with examples, IDAQP 9 (2006), 331-351
- [11] E.H. Lieb, M.B. Ruskai, Some operator inequalities of the Schwarz type, Adv. Math. 12 (1974), 269–273
- [12] M. Nussbaum, A. Szkola, The Chernoff lower bound for symmetric quantum hypothesis testing, Annals of Statistics 37, No. 2, 1040-1057 (2009)
- [13] M. Ohya, D. Petz, Quantum Entropy and Its Use, Springer-Verlag, Heidelberg, 1993, 2nd edition 2004.
- [14] D. Petz, On the equality in Jensen's inequality for operator convex functions, Integral Equations and Operator Theory, 9 (1986), 744–747
- [15] J. Pfanzagl, A characterization of sufficiency by power functions, Metrika 21 (1974), 197–199
- [16] H. Strasser, Mathematical theory of statistics. Statistical experiments and asymptotic decision theory, Walter de Gruyter, Berlin, 1985.