

**Stabilisation of Seven (Toroidal) Directions
and Expansion of the remaining Three
in an M theoretic Early Universe Model**

Samrat Bhowmick, Sanatan Digal, and S. Kalyana Rama

Institute of Mathematical Sciences, C. I. T. Campus,
Tharamani, CHENNAI 600 113, India.

email: samrat, digal, krama@imsc.res.in

ABSTRACT

Our model consists of intersecting $22'55'$ brane configuration in M theory, distributed uniformly in the common transverse space and assumed to describe early universe. Equations of state are those following from U duality symmetries and a few standard assumptions. In this model, three spatial directions expand, and seven directions stabilise to constant sizes which depend on certain imbalance among initial values. One naturally obtains $M_{11} \simeq M_s \simeq M_4$ and $g_s \simeq 1$ within a few orders of magnitude. Smaller numbers, for example $M_s \simeq 10^{-16} M_4$, are also possible but require fine tuning.

1. In early universe, the temperature and energy densities are high. When they are of the order of Planck scale $M_4 \simeq 10^{19} GeV$, the dynamics of the early universe is expected to be described by a more fundamental theory such as string theory or M theory [1, 2].

If this is the case then the problem of spacetime dimensions needs to be resolved – spacetime is eleven dimensional in M theory whereas it is four dimensional in our observed universe.

A canonical resolution is that the early universe starts out being eleven dimensional. During its evolution, by some dynamics, seven of the spatial directions cease to expand and their sizes become stabilised. The remaining three spatial directions continue to expand and become the observed universe.

The stabilised sizes then relate the M theory scale M_{11} and the four dimensional Planck scale M_4 . Likewise, since string theory can be obtained by dimensionally reducing M theory, the sizes also relate M_{11} and the string scale M_s and the string coupling constant g_s . One may then enquire, for example, whether it is possible to have string/M theory scale in the TeV range as required in Large Volume compactification scenarios.

Various proposals have been made for obtaining a four dimensional universe from string/M theory [3, 4, 5]. Typically, one assumes that the spatial directions are all toroidal, and are wrapped by a gas of winding and anti winding strings or p -branes; and that the cosmological evolution is governed by a ten/eleven dimensional effective action. The earliest proposal [3], in the context of string theory, is based on the observation that winding and anti winding strings oppose the expansion, and are annihilated efficiently in four dimensional spacetime. Others [4, 5] are variants of this, or based on its generalisations to winding and anti winding p -branes in string/M theory. These proposals are quite appealing and have been used in a variety of ‘brane gas’ models [4, 5], but some important issues yet remain to be resolved [6].

In this letter, based on the ideas in [7, 8], we present an M theoretic early universe model where seven of the spatial directions cease to expand and their sizes become stabilised. The remaining three spatial directions continue to expand, thus leading to a four dimensional universe. The stabilised sizes, and thus the explicit relations among (M_{11}, M_4, M_s, g_s) , depend on certain imbalance among initial values. The exact values are obtained numerically, but can also be estimated analytically under certain approximations. In this model, one may obtain any value for M_{11} or M_s , including in the TeV range, by a corresponding fine tuning of initial values.

2. Our model is as follows. Let all the spatial directions be toroidal. Consider N – sets of branes in M theory, intersecting as per the rules given in [9]. Consider the $N = 4$ intersecting brane configuration [10], denoted by $22'55'$, which has vanishing net charges and consists of two sets each of 2 branes and 5 branes along (x^1, x^2) , (x^3, x^4) , $(x^1, x^3, x^5, x^6, x^7)$, and $(x^2, x^4, x^5, x^6, x^7)$ directions. Let this configuration be uniformly distributed in the common transverse space along (x^8, x^9, x^{10}) directions, which then is assumed to describe a homogeneous anisotropic universe whose evolution is governed by an eleven dimensional effective action [7, 8].

Let $I = 1, 2, 3, 4$ denote the branes $2, 2', 5, 5'$ respectively. We assume that, as in the case of black holes, the energy momentum tensors $T^A_{B(I)}$ of the I^{th} set of branes are mutually non interacting and seperately conserved [7, 12]. Then

$$T^A_B = \sum_I T^A_{B(I)} \quad , \quad \sum_A \nabla_A T^A_{B(I)} = 0 \quad (1)$$

where T^A_B is the total energy momentum tensor of the configuration [13]. Homogeneity implies that $T^A_B = \text{diag}(-\rho, p_i)$ and $T^A_{B(I)} = \text{diag}(-\rho_I, p_{iI})$. We take $\rho_I > 0$.

To obtain the equations of state $p_{iI}(\rho_I)$, let $p_{\parallel I}$ and $p_{\perp I}$ denote parallel and perpendicular components of pressure due to I^{th} set of branes. It is shown in [8] that U duality symmetries of M theory imply that the functions $p_{\perp I}(\rho_I)$ must be same for all I and that $p_{\parallel I} = 2p_{\perp I} - \rho_I$. The function $p_{\perp}(\rho)$ may be determined, in principle, by brane antibrane dynamics. In our model, we set $p_{\perp} \propto \rho$; more specifically, $p_{\perp} = (1 - u) \rho$ where u is a constant and $0 < u < 2$. (The corresponding result in [7] follows if $u = 1$.) Then, for the $22'55'$ configuration, we have $p_{iI} = (1 - u_i^I) \rho_I$ where

$$\begin{aligned} u_i^1 &= u (2, 2, 1, 1, 1, 1, 1, 1, 1, 1) \\ u_i^2 &= u (1, 1, 2, 2, 1, 1, 1, 1, 1, 1) \\ u_i^3 &= u (2, 1, 2, 1, 2, 2, 2, 1, 1, 1) \\ u_i^4 &= u (1, 2, 1, 2, 2, 2, 2, 1, 1, 1) \quad . \end{aligned} \quad (2)$$

3. Consider now the evolution of the $D = (10 + 1)$ – dimensional homogeneous anisotropic universe in the model described above. Let the line

element ds be given by

$$ds^2 = \sum_{AB} g_{AB} dx^A dx^B = -dt^2 + \sum_i e^{2\lambda^i} (dx^i)^2 \quad (3)$$

where λ^i are functions of t only. Einstein equations $R_{AB} - \frac{1}{2}g_{AB}R = T_{AB}$, with $8\pi G = 1$, and equations (1) lead to $\rho_I = e^{l^I - 2\Lambda}$ and

$$\sum_{ij} G_{ij} \lambda_t^i \lambda_t^j = 2 \sum_I e^{l^I - 2\Lambda} \quad (4)$$

$$\lambda_{tt}^i + \Lambda_t \lambda_t^i = \sum_I u^{iI} e^{l^I - 2\Lambda} \quad (5)$$

where $l^I = \sum_i u_i^I \lambda^i + l_0^I$, $\Lambda = \sum_i \lambda^i$, the subscripts t denote time derivatives, and

$$G_{ij} = 1 - \delta_{ij} \quad , \quad G^{ij} = \frac{1}{D-2} - \delta^{ij} \quad , \quad u^{iI} = \sum_j G^{ij} u_j^I \quad . \quad (6)$$

Let

$$d\tau = e^{-\Lambda} dt \quad , \quad \mathcal{G}^{IJ} = \sum_{ij} G^{ij} u_i^I u_j^J \quad , \quad u_I^i = \sum_J \mathcal{G}_{IJ} u^{iJ} \quad (7)$$

where \mathcal{G}_{IJ} is defined by $\sum_J \mathcal{G}^{IJ} \mathcal{G}_{JK} = \delta_K^I$. Then, manipulating equations (4) and (5), one obtains

$$\lambda^i = \sum_I u_I^i (l^I - l_0^I) + L^i \tau \quad (8)$$

$$l_{\tau\tau}^I = \sum_J \mathcal{G}^{IJ} e^{l^J} \quad (9)$$

$$\sum_{IJ} \mathcal{G}_{IJ} l_{\tau}^I l_{\tau}^J = 2 (E + \sum_I e^{l^I}) \quad (10)$$

where the subscripts τ denote τ -derivatives, L^i are integration constants satisfying $\sum_i u_i^I L^i = 0$, and $2E = -\sum_{ij} G_{ij} L^i L^j$. Also, with no loss of generality, we have taken the initial values to be

$$\left(\lambda^i, \lambda_t^i, l^I, l_t^I, \rho_I, \tau \right)_{t=0} = \left(0, k^i, l_0^I, K^I, \rho_{I0}, 0 \right) \quad (11)$$

where $\rho_{I0} = e^{l_0^I}$, $k^i = \sum_I u_I^i K^I + L^i$ since $f_t(0) = f_{\tau}(0)$ for any f , and

$$\sum_{IJ} \mathcal{G}_{IJ} K^I K^J = 2 (E + \sum_I e^{l_0^I}) \quad . \quad (12)$$

For the 22'55' configuration, \mathcal{G}^{IJ} , \mathcal{G}_{IJ} , u^{iI} , and u_I^i are as given below:

$$\mathcal{G}^{IJ} = 2u^2 (1 - \delta^{IJ}) \quad , \quad \mathcal{G}_{IJ} = \frac{1}{6u^2} (1 - 3\delta_{IJ}) \quad . \quad (13)$$

I	$\frac{u^{iI}}{u/3}$	$\frac{u_I^i}{1/6u}$
1	-2, -2, 1, 1, 1, 1, 1, 1, 1, 1	2, 2, -1, -1, -1, -1, -1, 1, 1, 1
2	1, 1, -2, -2, 1, 1, 1, 1, 1, 1	-1, -1, 2, 2, -1, -1, -1, 1, 1, 1
3	-1, 2, -1, 2, -1, -1, -1, 2, 2, 2	1, -2, 1, -2, 1, 1, 1, 0, 0, 0
4	2, -1, 2, -1, -1, -1, -1, 2, 2, 2	-2, 1, -2, 1, 1, 1, 1, 0, 0, 0

Table I : u^{iI} and u_I^i for the 22'55' configuration.

There is an interesting similarity with black holes: When L^i all vanish, e^{λ^i} here have the same form as those for extremal 22'55' black holes and e^{2uh_I} , where $h_I = \sum_J \mathcal{G}_{IJ} (l^J - l_0^J)$, play the role of harmonic functions $H_I = 1 + \frac{Q_I}{r}$. See equation (8), Table I, and equation (18) in [11]. Also, the asymptotic limit $t \rightarrow \infty$ here, see below, corresponds to the near horizon limit $r \rightarrow 0$ and (certain combination of) ρ_{I0} play the role of Q_I .

4. To obtain $\lambda^i(t)$ for the 22'55' configuration, and thus the evolution of the universe, one may solve equations (9) – (13) for $l^I(\tau)$ and obtain $\lambda^i(\tau)$ from equation (8). Then $t(\tau)$ follows from $dt = e^\Lambda d\tau$, and thus $\tau(t)$ and $\lambda^i(t)$. Note that l^I depend on (l_0^I, K^I) and λ^i depend further on L^i .

We are unable to solve equations (9) – (13) analytically. Nevertheless, the important features of the evolution can be obtained as follows. For the 22'55' configuration, the following two results can be proved [14]:

- The constraints $\sum_i u_i^I L^i = 0$ imply that $0 \leq c_i (L^i)^2 \leq E$ where c_i are constants of $\mathcal{O}(1)$. Hence $E = 0$ if and only if $L^i = 0$ for all i .
- If $E \geq 0$ then equations (8) and (10) imply that none of (Λ_τ, l_τ^I) may vanish, and that they must be *all positive* or *all negative*.

Let $K^I = l_\tau^I(0) > 0$ for all I . The above results then imply that $l^I(\tau)$ all increase as τ increases. Equations (9) and (13) imply that $l^I(\tau)$ all diverge at finite $\tau = \tau_\infty$. The asymptotic solutions can be easily obtained in the limit $\tau \rightarrow \tau_\infty$. Using $dt = e^\Lambda d\tau$ and the values of u_I^i given in Table I, the asymptotic solutions are given to the leading order by

$$e^{l^I} = \frac{1}{3u^2} \frac{1}{(\tau_\infty - \tau)^2}, \quad t = t_* + A (\tau_\infty - \tau)^{-\frac{2-u}{u}}$$

$$e^{\lambda^i} = e^{v^i} \left(\frac{1}{3u^2} \frac{1}{(\tau_\infty - \tau)^2} \right)^{\sum_I u_I^i} = e^{v^i} \{B (t - t_*)\}^{\beta^i} \quad (14)$$

where t_* and τ_∞ are finite constants and depend on the details of evolution, A and B are u -dependent constants, $v^i = -\sum_I u_I^i l_0^I + L^i \tau_\infty$, and $\beta^i = \frac{2u}{2-u} \sum_I u_I^i$. Explicitly, we have

$$\beta^i = \frac{2}{3(2-u)} (0, 0, 0, 0, 0, 0, 0, 1, 1, 1). \quad (15)$$

Thus, asymptotically, $t \rightarrow \infty$ since $0 < u < 2$ in our model. And, $e^{\lambda^i} \rightarrow t^{\frac{2}{3(2-u)}}$ for the common transverse directions $i = 8, 9, 10$. Hence, these directions continue to expand, their expansion being precisely that of a $(3+1)$ -dimensional homogeneous, isotropic universe containing a perfect fluid whose equation of state is $p = (1-u)\rho$. Also, $e^{\lambda^i} \rightarrow e^{v^i}$ for the brane directions $i = 1, \dots, 7$. Hence, these directions cease to expand and their final sizes are given by e^{v^i} .

In our model, irrespective of initial values, three spatial directions will always expand and seven brane directions will always be stabilised and reach constant sizes. The underlying dynamics is distinct from those in [3, 4, 5] and can be described as follows. From equation (5) and u^I in Table I, one sees that parallel brane directions contract and transverse ones expand, at opposite rates for 2 branes and 5 branes. If the brane energy densities ρ_I are all different then, generically, so will be the corresponding expansion and

contraction rates, and the brane directions will have net expansion or contraction. Only if the expansion rates equal contraction rates, will the brane directions cease to expand or contract and their sizes stabilise to constant values.

Such an equality ensues eventually in our model as a result of two crucial features : **(i)** The dynamics of the evolution, given by u_i^I which in turn follow from U duality symmetries [8], is such that ρ_I , even if different initially, evolve to become all equal. This equality is due to each $\rho_I \sim e^{l^I}$ being ‘sourced’ by the sum of other three, see equations (9) and (13). **(ii)** The 22’55’ configuration is such that each brane direction is parallel to two sets of branes, and transverse to other two in just the right way. Hence, its expansion and contraction rates become equal once ρ_I become all equal.

The stabilised sizes of the brane directions should then depend, for example, on the imbalance among ρ_{I0} [15]. That this is the case can be seen from explicit expressions for e^{v^i} ; for example, we have

$$e^{v^1} = e^{L^1} \tau_\infty \left(\frac{\rho_{20} \rho_{40}^2}{\rho_{30} \rho_{10}^2} \right)^{\frac{1}{6u}} , \quad e^{v^c} = e^{L^c} \tau_\infty \left(\frac{\rho_{10} \rho_{20}}{\rho_{30} \rho_{40}} \right)^{\frac{1}{6u}} , \quad (16)$$

where we also define $v^c = \sum_{i=1}^7 v^i$ and $L^c = \sum_{i=1}^7 L^i$, needed below. The dependence of e^{v^i} on L^i is due to the imbalance among $\lambda_t^i(0)$.

Thus, asymptotically as $t \rightarrow \infty$, the $(10 + 1)$ – dimensional universe effectively becomes $(3 + 1)$ – dimensional. Also, dimensional reduction of M theory along, for example, x^1 direction gives string theory. Therefore, upto numerical factors of $\mathcal{O}(1)$, the corresponding scales and coupling constant (M_{11}, M_4, M_s, g_s) are related asymptotically by

$$M_4^2 \simeq e^{v^c} M_{11}^2 \simeq e^{v^c - v^1} M_s^2 , \quad g_s^2 \simeq e^{3v^1} \quad (17)$$

where coordinate sizes are assumed to be of order $\frac{1}{M_{11}}$, and e^{v^i} measure physical sizes in units of $\frac{1}{M_{11}}$ [16].

5. To determine the sizes of brane directions and the relations in equation (17) explicitly for a given set of initial values (l_0^I, K^I, L^i) , we need τ_∞ if $L^i \neq 0$. We will obtain τ_∞ numerically since it depends on the details of evolution and we do not have explicit solutions.

We assume that L^i do not all vanish and set $E = 1$ [15]. Note that if $e^{l_0^I} \ll 1$ for all I then equations (9) and (10) imply that $l^I(\tau)$ may be taken

as evolving ‘freely’, *i.e.* $l^I(\tau) = l_0^I + K^I \tau$ where $K^I = l_\tau^I(0) > 0$, until one of the $e^{l^I} = 1$; from then on, all e^{l^I} will evolve quickly and diverge soon after. Consequently, τ_∞ may be given approximately by

$$\tau_\infty \simeq \tau_a = \min \left\{ -\frac{l_0^I}{K^I} \right\} . \quad (18)$$

Also, τ_a is maximum, and $\tau_{a, max} = \frac{1}{K}$, when $K^1 = x^1$, $K^2 = x^2$, $K^3 = \min \{x^1 + x^2, x^3\}$ and $K^4 = \min \{x^1 + x^2, \frac{1}{2}(x^1 + x^2 + x^3), x^4\}$ where $x^I = -l_0^I K$, equation (12) determines $K > 0$, and we assume with no loss of generality that $0 < x^1 \leq \dots \leq x^4$. No explicit solution is needed to evaluate τ_a and $\tau_{a, max}$.

We studied several sets of (l_0^I, K^I) numerically and obtained $\tau_{\infty, max}$, the maximum of τ_∞ , by sampling 25000 random sets of K^I for each set of l_0^I . We find that l^I all diverge at finite $\tau = \tau_\infty$ and that, when $e^{l_0^I} \ll 1$ for all I , the approximations given above are quite good: $l^I > l_0^I + K^I \tau$ discernibly only for $\tau \gtrsim \tau_\infty - 4$; $\tau_a \sim (0.5 - 1.1) \tau_\infty$ generically; and, for K^I which maximise τ_a , we get $\tau_a = \tau_{a, max} \sim (0.9 - 1.1) \tau_\infty \sim (0.9 - 1.1) \tau_{\infty, max}$.

To convey an idea of what values are possible in equation (17), and also an idea of how good the approximations given above are, we consider two illustrative sets of l_0^I , choose K^I which maximise τ_a , choose $L^i = \sqrt{\frac{1}{6}} (-1, 2, 2, -1, 0, 0, 0, -1, -1, -1)$ so that g_s can be small, and choose $u = \frac{2}{3}$ which corresponds to radiation filled universe in $(3 + 1)$ – dimensions. The corresponding numerical results are given in Table II, from which e^{v^1} and e^{v^c} can be read off easily using equation (17). Also, $(\tau_{a, max}, \tau_{\infty, max}) = (5.27, 5.82)$ for the first set, and $= (25.43, 25.69)$ for the second set of l_0^I in Table II.

$\{-l_0^I = -\ln \rho_{I0}\}$	τ_∞	$\frac{M_{11}}{M_4}$	$\frac{M_s}{M_4}$	g_s
5, 5, 12, 12	5.73	$1.67 * 10^{-2}$	$2.17 * 10^{-3}$	$2.17 * 10^{-3}$
20, 30, 40, 50	25.64	$1.92 * 10^{-7}$	$1.97 * 10^{-12}$	$1.09 * 10^{-15}$

Table II : *The numerical results for $(\tau_\infty; \frac{M_{11}}{M_4}, \frac{M_s}{M_4}, g_s)$ for two illustrative sets of l_0^I . Other parameters are chosen as explained in the text.*

For a given set of l_0^I , our choice of (K^I, L^i) in Table II results in near-minimum values for $(\frac{M_{11}}{M_4}, \frac{M_s}{M_4}, g_s)$ within about an order of magnitude. Our numerical studies confirm this. Also note that, since $E = 1$, $\lambda_t^i(0) = k^i \simeq K^I \simeq L^i \simeq \mathcal{O}(1)$ naturally whereas ensuring that $\rho_{I0} = e^{l_0^I} \ll 1$ for all I requires (fine) tuning. Thus, we conclude that our model naturally leads to $M_{11} \simeq M_s \simeq M_4$ and $g_s \simeq 1$ within a few orders of magnitude; and that smaller M_{11} and M_s , for example $M_s \simeq TeV \simeq 10^{-16} M_4$ as required in Large Volume compactification scenarios, are also possible but require a corresponding fine tuning of initial values.

6. We have shown that, in our model, three spatial directions expand and seven directions stabilise to constant sizes e^{v^i} , $i = 1, \dots, 7$. We have also given exact expressions for v^i , which depend on initial values and τ_∞ . τ_∞ can be evaluated explicitly if solutions are known, otherwise numerically. Also, we give approximate expressions for τ_∞ which are easy to evaluate and work well under certain conditions. Explicit relations among (M_{11}, M_4, M_s, g_s) then follow from which we see, for example, that obtaining $M_s \simeq TeV$ requires fine tuning.

We conclude by listing a few questions of obvious importance for further studies. **(i)** How to solve equations (9) – (13) analytically? **(ii)** Is there any way of obtaining $M_s \simeq TeV$ in the present model without fine tuning? **(iii)** Why 22'55' configuration and why not, for example, 22'2'' (which will lead [8] to four spatial directions expanding)? The likely answer is that 22'55' configuration is entropically favourable [2, 7, 8], but dynamical details are not clear. **(iv)** What is the evolution when topology of spatial directions is more general? **(v)** We pointed out an interesting similarity with black holes. Does it have any deeper significance?

Acknowledgement: We thank B. Sathiapalan and N. V. Suryanarayana for their comments.

References

- [1] M. J. Bowick and L. C. R. Wijewardhana, *Gen. Rel. Grav.* **18** (1986) 59; S. Kalyana Rama, *Phys. Lett. B* **638** (2006) 100 [arXiv: hep-th/0603216].
- [2] S. Kalyana Rama, *Phys. Lett. B* **645** (2007) 365 [arXiv: hep-th/0610071].
- [3] R. H. Brandenberger and C. Vafa, *Nucl. Phys. B* **316** (1989) 391; A. A. Tseytlin and C. Vafa, *Nucl. Phys. B* **372** (1992) 443 [arXiv: hep-th/9109048]. See also J. Kripfganz and H. Perlt, *Class. Quant. Grav.* **5** (1988) 453.
- [4] See, for example, S. Alexander, R. H. Brandenberger and D. Easson, *Phys. Rev. D* **62** (2000) 103509 [arXiv:hep-th/0005212]; R. Brandenberger, D. A. Easson and D. Kimberly, *Nucl. Phys. B* **623** (2002) 421 [arXiv:hep-th/0109165]; A. Kaya, *JCAP* **08** (2004) 014 [arXiv:hep-th/0405099]; S. Arapoglu and A. Kaya, *Phys. Lett. B* **603** (2004) 107 [arXiv:hep-th/0409094]; T. Rador, *Eur. Phys. J. C* **49** (2007) 1083 [arXiv:hep-th/0504047]; M. Sano and H. Suzuki, *Phys. Rev. D* **78** (2008) 064045, arXiv:0804.0176 [hep-th]. See also the recent reviews T. Battefeld and S. Watson, *Rev. Mod. Phys.* **78** (2006) 435 [arXiv:hep-th/0510022]; R. H. Brandenberger, arXiv:0808.0746 [hep-th]; and the references therein.
- [5] R. Easther, B. R. Greene, M. G. Jackson and D. Kabat, *Phys. Rev. D* **67** (2003) 123501 [arXiv: hep-th/0211124]; R. Durrer, M. Kunz and M. Sakellariadou, *Phys. Lett. B* **614** (2005) 125 [arXiv: hep-th/0501163]; A. Karch and L. Randall, *Phys. Rev. Lett.* **95** (2005) 161601 [arXiv: hep-th/0506053].
- [6] R. Easther, B. R. Greene, M. G. Jackson and D. Kabat, *JCAP* **01** (2004) 006 [arXiv: hep-th/0307233]; R. Easther, B. R. Greene, M. G. Jackson and D. Kabat, *JCAP* **02** (2005) 009 [arXiv: hep-th/0409121]; R. Danos, A. R. Frey and A. Mazumdar, *Phys. Rev. D* **70** (2004) 106010 [arXiv: hep-th/0409162].

- [7] B. D. Chowdhury and S. D. Mathur, *Class. Quant. Grav.* **24** (2007) 2689 [arXiv:hep-th/0611330]; S. D. Mathur, *What is the state of the Early Universe?*, arXiv:0803.3727 [hep-th].
- [8] S. Kalyana Rama, *Phys. Lett. B* **656** (2007) 226 [arXiv:0707.1421 [hep-th]]. See also S. Kalyana Rama, *Gen. Rel. Grav.* **39** (2007) 1773 [arXiv:hep-th/0702202].
- [9] A. A. Tseytlin, *Nucl. Phys. B* **475** (1996) 149 [arXiv: hep-th/9604035].
- [10] The $22'55'$ configuration, when localised in the common transverse space, describes a four charged black hole [11]. The configuration may be thought of as consisting of four sets of coincident branes and anti-branes. The branes and anti-branes are created and annihilated in pairs and are in thermal equilibrium. The differences in their numbers denote the charges of the black hole [12]. Similar picture is likely to be valid in our model also. For the sake of convenience, we simply refer to such a collection of branes and anti-branes as branes.
- [11] M. Cvetič and A. A. Tseytlin, *Nucl. Phys. B* **478** (1996) 181 [arXiv: hep-th/9606033].
- [12] G. T. Horowitz, J. M. Maldacena and A. Strominger, *Phys. Lett. B* **383** (1996) 151 [arXiv: hep-th/9603109]; G. T. Horowitz, D. A. Lowe and J. M. Maldacena, *Phys. Rev. Lett.* **77** (1996) 430 [arXiv: hep-th/9603195]; U. H. Danielsson, A. Guijosa and M. Kruczenski, *JHEP* **09** (2001) 011 [arXiv: hep-th/0106201].
- [13] We indicate the summation variables explicitly since the repeated indices are not always summed over. The variables $A, B, \dots = 0, 1, \dots, 10$, $I, J, \dots = 1, \dots, 4$, and $i, j, \dots = 1, \dots, 10$ unless indicated otherwise.
- [14] The constraints $\sum_i u_i^I L^i = 0$ imply that $L^1 - L^4 = L^2 - L^3 = \sum_{i=5}^7 L^i = 3(L^1 + L^2) + \sum_{i=8}^{10} L^i = 0$. The two results then follow upon judicious use of the relation $(\sum_{a=1}^n v_a)^2 = n \cos^2 \theta \sum_{a=1}^n (v_a)^2$ and some manipulation of equations (8), (10), and $2E = -\sum_{ij} G_{ij} L^i L^j$.
- [15] Scaling all $\rho_{I0} = e^{I\sigma}$ by an overall factor does not change the stabilised sizes since $e^{v^i} \rightarrow e^{v^i}$ for $i = 1, \dots, 7$ under $(t, \tau) \rightarrow \frac{1}{\sigma} (t, \tau)$;

$(K^I, L^i) \rightarrow \sigma (K^I, L^i) ; \quad (e^{l_0}, E) \rightarrow \sigma^2 (e^{l_0}, E) .$ Such a scaling changes only the rate of evolution but not the stabilised sizes. It can also be used, for example, to set $E = 1$ if $E > 0$, equivalently if L^i do not all vanish.

- [16] In string units $\frac{1}{M_s}$, the sizes are given by $e^{\tilde{v}^i} = \frac{M_s}{M_{11}} e^{v^i} = g_s^{\frac{1}{3}} e^{v^i}$ and the relation between M_4 and M_s becomes, in an obvious notation, the more familiar one : $G_4 \simeq \frac{g_s^2 l_s^8}{V_6} .$