Strong NP-Hardness of the Quantum Separability Problem

Sevag Gharibian *

March 13, 2019

Abstract

We show that the Weak Membership problem over the set of separable (equivalently, unentangled) bipartite quantum states is "strongly" NP-hard, implying it is NP-hard even when the error margin allowed is as large as inverse polynomial in the dimension, i.e. is "moderately large". This shows that it is NP-hard to determine whether an arbitrary quantum state located within an inverse polynomial distance from the border of the separable set is entangled. The result here extends the previous work of Gurvits, which shows NP-hardness for the case of inverse exponential distance. Based on our result, we observe an immediate lower bound on the maximum distance possible between a bound entangled state and the separable set (assuming $P \neq NP$). We also show that determining whether a completely positive trace-preserving linear map (i.e. a quantum channel) is entanglement-breaking is NP-hard.

1 Introduction

Given a bipartite quantum state ρ acting on Hilbert space $\mathcal{H}^M \otimes \mathcal{H}^N$, specified in terms of its density matrix with respect to some known orthonormal basis, and where M and N denote the dimensions of its respective subsystems, we consider the Quantum Separability problem (QUSEP) — is ρ separable, or equivalently, unentangled? With the multitude of applications known for entanglement within the quantum computing and information community, this problem has naturally been the focus of intense research over recent years (see [1, 2] for surveys). Let $\mathcal{S}_{M,N}$ denote the set of separable quantum states acting on $\mathcal{H}^M \otimes \mathcal{H}^N$. Then, informally, since $\mathcal{S}_{M,N}$ is convex, one way to think about QUSEP is in terms of membership in a convex set with some allowed margin of error $\beta > 0$. Briefly, β specifies the (Euclidean) distance from the border of $\mathcal{S}_{M,N}$ within which an algorithm for QUSEP is allowed to err. This margin of error is introduced to account for the fact that in encoding the entries of the density matrix of ρ using finite precision, it is possible that if ρ is very close to the border of $\mathcal{S}_{M,N}$, the encoding process may accidentally shift the resulting state into or out of $\mathcal{S}_{M,N}$. The above formulation, first considered by Gurvits [3], is dubbed the Weak Membership problem over $\mathcal{S}_{M,N}$, and is denoted WMEM_{β}($\mathcal{S}_{M,N}$).

In 2003, Gurvits showed that WMEM_{β}($S_{M,N}$) is NP-hard [3] via a polynomial time Turing reduction from the NP-complete problem PARTITION. PARTITION, however, is only weakly NPhard, since it can be solved efficiently if the values of its numerical parameters are polynomially bounded in the dimension (in this case, using a dynamic programming approach [4]). It follows that the reduction of [3] shows NP-hardness for WMEM_{β}($S_{M,N}$) only when $\beta \leq 1/\exp(M, N)$, i.e.

^{*}School of Computer Science and Institute for Quantum Computing, University of Waterloo, Waterloo, Canada

when the allowed margin of error is exponentially small (as observed by Aaronson and documented in [5]). This leaves open the possibility of an approximation algorithm when a larger margin of error is allowed.

In an attempt to strengthen this result, Gurvits suggested [5] the following reduction from the NP-complete problem CLIQUE, where RSDF and WVAL_{α}($S_{M,N}$) are the problems Robust Semidefinite Feasibility and Weak Validity over $S_{M,N}$, respectively (all necessary formal definitions will be given later in Section 2, and by \leq_K and \leq_C , we indicate a Karp and Cook reduction¹, respectively):

$$\operatorname{CLIQUE} \leq_K \operatorname{RSDF} \leq_K \operatorname{WVAL}_{\alpha}(\mathcal{S}_{M,N}) \leq_C \operatorname{WMEM}_{\beta}(\mathcal{S}_{M,N}).$$
(1)

This time, the link $WVAL_{\alpha}(\mathcal{S}_{M,N}) \leq_C WMEM_{\beta}(\mathcal{S}_{M,N})$ is problematic, as the only known² deterministic polynomial time Turing reduction from $WVAL_{\alpha}(K)$ to $WMEM_{\beta}(K)$ for arbitrary choice of $\alpha > 0$ is via the Yudin-Nemirovski theorem [7], which is based on the shallow-cut ellipsoid method. This also results in exponential scaling for β [5], leaving us with the same problem as before — $WMEM_{\beta}(\mathcal{S}_{M,N})$ is known to be NP-hard only for $\beta \leq 1/\exp(M, N)$. The main result we show in this note is hence as follows.

Theorem 1. WMEM_{β}($S_{M,N}$) is NP-hard for $\beta \leq 1/\operatorname{poly}(M,N)$, i.e. is strongly NP-hard.

Here, we define a problem as strongly NP-hard if it is NP-hard even if the magnitudes of its numerical parameters are polynomially bounded in the length of its input [8]. To show Theorem 1, we observe that there exists a recent non-ellipsoidal Turing reduction of Liu [9] from the problem³ Weak Optimization (WOPT_{ϵ}(K)) to WMEM_{β}(K). To use this reduction, we begin with Equation (1), and show a Karp reduction from RSDF to WOPT_{ϵ}($\mathcal{S}_{M,N}$) (based partly on work by Gurvits [5]). We then apply the reduction of Liu from WOPT_{ϵ}($\mathcal{S}_{M,N}$) to WMEM_{β}($\mathcal{S}_{M,N}$). Our reduction chain is hence (where \leq_T denotes a Turing reduction):

$$\operatorname{CLIQUE} \leq_K \operatorname{RSDF} \leq_K \operatorname{WOPT}_{\epsilon}(\mathcal{S}_{M,N}) \leq_T \operatorname{WMEM}_{\beta}(\mathcal{S}_{M,N}).$$
(2)

We remark that the reduction of Liu runs in polynomial time only if we demand at most "moderate" precision for WOPT_{ϵ}(K), i.e. $\epsilon \geq 1/\operatorname{poly}(M, N)$, for ϵ the error parameter for Weak Optimization. We show that to solve an instance of CLIQUE, it suffices to choose $\epsilon \geq 1/\operatorname{poly}(M, N)$ here.

We next discuss two applications of Theorem 1. By considering the Positive Partial Transpose entanglement detection criterion [11, 12], we observe immediate lower bounds on the maximum (Euclidean) distance possible between a bound entangled [13] state and $S_{M,N}$. Second, using the the Jamiołkowski isomorphism [14], we show that determining whether a completely positive tracepreserving linear map is entanglement-breaking [15] is NP-hard.

This note is organized as follows. We begin in Section 2 by introducing all necessary definitions and notation. In Section 3, we prove Theorem 1. Section 4 discusses applications of Theorem 1. We conclude in Section 5 with brief comments and an open problem.

¹A Cook reduction is a polynomial time Turing reduction from problem A to B, where, intuitively, a Turing reduction demonstrates how to solve A using an algorithm for B possibly multiple times. A Karp reduction, on the other hand, invokes the algorithm for B once, the output of which it returns as the solution for A.

 $^{^{2}}$ There is, however, a *randomized* reduction following from the random walk algorithm for convex optimization of Bertsimas and Vempala [6].

³What we call Weak Optimization here can be shown to be equivalent to the problem Weak Validity in [10]. We follow the naming convention in [9], however, in order to avoid confusion when applying the reduction therein.

2 Definitions and Notation

There are five problems we need to define in order to show Theorem 1: CLIQUE, Robust Semidefinite Feasibility (RSDF), Weak Optimization, Weak Membership, and QUSEP. We begin by presenting and discussing their definitions here. All norms are taken as the Euclidean norm (indicated $\| \|_2$ or $\| \|_F$, the latter denoting the Frobenius norm for matrices). We denote (column) vector vby \mathbf{v} , its conjugate transpose as \mathbf{v}^{\dagger} , and its *i*th entry as v_i . We use the notation $\langle \alpha \rangle$ to signify the number of bits necessary to encode an entity α . Specifically, if $\alpha = a/b$ is rational, we define $\langle \alpha \rangle = \langle a \rangle + \langle b \rangle$, and for matrix A, we let $\langle A \rangle = \sum_{ij} \langle A_{ij} \rangle$ (similarly for vectors).

First, the NP-complete problem CLIQUE is stated as follows.

Definition 1 (CLIQUE). Given a simple graph G on n vertices, and $c \leq n$, for $n, c \in \mathbb{Z}^+$, decide, with respect to the complexity measure $\langle G \rangle + \langle c \rangle$:

If the number of vertices in the largest complete subgraph of G is at least c, output "YES". Otherwise, output "NO".

Here, we take $\langle G \rangle = \langle A_G \rangle$, where A_G is the $n \times n$ adjacency matrix for G, such that $A_G[i, j] = 1$ if vertices i and j are connected by an edge, and $A_G[i, j] = 0$ otherwise. Next, we have the problem Robust Semidefinite Feasibility.

Definition 2 (Robust Semidefinite Feasibility (RSDF)). Given k rational, symmetric $l \times l$ matrices B_1, \ldots, B_k , and $\zeta, \eta \in \mathbb{Q}$, with $\zeta, \eta \geq 0$, define $g(B_1, \ldots, B_k) = \max_{\mathbf{x} \in \mathbb{R}^l, \|\mathbf{x}\|_2 = 1} \sum_{i=1}^k (\mathbf{x}^T B_i \mathbf{x})^2$. Then, decide, with respect to the complexity measure $lk + \sum_{i=1}^k \langle B_i \rangle + \langle \zeta \rangle + \langle \eta \rangle$:

If $g(B_1, \ldots, B_k) \ge \zeta + \eta$, output "YES". If $g(B_1, \ldots, B_k) \le \zeta - \eta$, output "NO".

We have assumed $\zeta \geq 0$ without loss of generality above, since $g(B_1, \ldots, B_k) \geq 0$. This will be necessary later in Lemma 4, when we need to take $\sqrt{g(B_1, \ldots, B_k)}$. We have also defined RSDF as a *promise* problem, meaning we are promised the input will fall into one of two disjoint cases which may separated by a non-zero gap, and we must distinguish between the two cases. One could equivalently lift the promise and allow input falling in the "gap" region (i.e. $\zeta - \eta < g(B_1, \ldots, B_k) < \zeta + \eta$) — for any such input, we would consider any output to be correct (i.e. "YES" or "NO").

Moving on, for Weak Optimization and Weak Membership, let $K \subseteq \mathbb{R}^m$ be a convex and compact set, and define related sets $S(K, \delta) := \{\mathbf{x} \in \mathbb{R}^m \mid \exists \mathbf{y} \in K \text{ s.t. } \|\mathbf{x} - \mathbf{y}\|_2 \leq \delta\}$, and $S(K, -\delta) := \{\mathbf{x} \in K \mid S(\mathbf{x}, \delta) \subseteq K\}$. Roughly, $S(K, \delta)$ can be thought of as extending the border of K by δ (in Euclidean distance), and $S(K, -\delta)$ can be thought of as taking the core of K, which is δ away from the border of K. As per [10], we shall require that K be a well-bounded p-centered set, meaning $K \subseteq S(\mathbf{0}, R)$ for $\mathbf{0}$ the origin and some rational R > 0, and $S(\mathbf{p}, r) \subseteq K$ for known point $\mathbf{p} \in K$ and some rational r > 0. This ensures K is full-dimensional and bounded. We will see shortly that $S_{M,N}$ is in fact such a set. Finally, we set the encoding size of K as $\langle K \rangle = m + \langle r \rangle + \langle R \rangle + \langle \mathbf{p} \rangle$ [3]. Weak Optimization and Weak Membership over the set K are defined as follows.

Definition 3 (Weak Optimization (WOPT_{ϵ}(K))[9]). Given $\mathbf{c} \in \mathbb{Q}^m$ with $\|\mathbf{c}\|_2 = 1$, and $\gamma, \epsilon \in \mathbb{Q}$, such that $\epsilon > 0$, decide, with respect to the complexity measure $\langle K \rangle + \langle \mathbf{c} \rangle + \langle \gamma \rangle + \langle \epsilon \rangle$:

If there exists $\mathbf{y} \in S(K, -\epsilon)$ with $\mathbf{c}^{\mathrm{T}}\mathbf{y} \geq \gamma + \epsilon$, then output "YES". If for all $\mathbf{x} \in S(K, \epsilon)$, $\mathbf{c}^{\mathrm{T}}\mathbf{x} \leq \gamma - \epsilon$, then output "NO". **Definition 4** (Weak Membership (WMEM_{β}(K))). Given $\mathbf{y} \in \mathbb{Q}^m$, and error parameter $\beta \in \mathbb{Q}$, such that $\beta > 0$, decide, with respect to the complexity measure $\langle K \rangle + \langle \mathbf{y} \rangle + \langle \beta \rangle$:

If $\mathbf{y} \in S(K, -\beta)$, then output "YES". If $\mathbf{y} \notin S(K, \beta)$, then output "NO".

Both of these problems are also stated as promise problems. Roughly speaking, the former asks whether there exists a point in the "core" of K that achieves a threshold value slightly higher than γ for a linear function defined by **c**, and the latter asks to distinguish whether a given point **y** is in the "core" of K or "far away" from K. We remark that unlike here, in Ref. [9] the inputs to WOPT_{ϵ}(K) and WMEM_{β}(K) are real (as opposed to rational), and specified using poly(m) bits of precision. The latter is because such precision suffices if one demands ϵ and β to be at least inverse polynomial in the input size [9], i.e. if one allows at least "moderate" error (which we also demand here). It is easy to see that we can exactly represent any such poly(m)-bit real numbers as rational numbers in poly-time using poly(m) bits as well, and hence the case of Ref. [9] is a special case of our definitions here. We remark that in our reduction, the choice for m is polynomial in the encoding size of the given CLIQUE instance.

Finally, let us formally define the Quantum Separability problem, and discuss how it relates to the problems over convex sets we have just seen.

Definition 5 (Quantum Separability Problem (QUSEP)). Let $H_{M,N}$ denote the set of Hermitian operators mapping $\mathbb{C}^M \otimes \mathbb{C}^N \mapsto \mathbb{C}^M \otimes \mathbb{C}^N$, for $M, N \geq 2$. Denote the set of separable quantum states in $H_{M,N}$ as $\mathcal{S}_{M,N} = \operatorname{conv}\{\mathbf{x}\mathbf{x}^{\dagger} \otimes \mathbf{y}\mathbf{y}^{\dagger} \mid \mathbf{x} \in \mathbb{C}^M, \mathbf{y} \in \mathbb{C}^N, \|\mathbf{x}\|_2 = \|\mathbf{y}\|_2 = 1\}$, where $\operatorname{conv}\{S\}$ denotes the convex hull generated by the set S. Then, given a quantum state $\rho \in H_{M,N}$, decide:

If $\rho \in \mathcal{S}_{M,N}$, output "YES". Otherwise, output "NO".

It is clear from Definition 5 that $S_{M,N}$ is a convex set. In fact, $S_{M,N}$ is p-centered and well-bounded, since it is contained in an origin-centered ball of radius $R = \sqrt{(MN-1)/MN}$ [2], and contains a ball of radius $r = \sqrt{1/MN(MN-1)}$ [16] centered at point p = I/MN [2], where I denotes the identity. Further, $S_{M,N}$ is compact, since the set of pure product states is closed and bounded, and the convex hull of a convex compact set is also compact [17]. Thus, in this paper we rephrase QUSEP as WMEM_{β}($S_{M,N}$), and investigate hardness of WMEM_{β}($S_{M,N}$) to draw conclusions about QUSEP (i.e. we never work with QUSEP directly).

Observe now that WOPT_{ϵ}(K) and WMEM_{β}(K) are phrased over $K \subseteq \mathbb{R}^m$, whereas $S_{M,N} \subseteq H_{M,N}$. To deal with this, recall that $H_{M,N}$ is isomorphic to $\mathbb{R}^{M^2N^2}$, and specifically one can write for any quantum state $\rho \in H_{M,N}$ [18]:

$$\rho = \frac{I}{MN} + \frac{1}{2} \sum_{i=1}^{M^2 N^2 - 1} r_i \lambda_i, \tag{3}$$

where we have chosen as a basis for $H_{M,N}$ the identity and the traceless Hermitian generators of SU(MN), the latter denoted by λ_i and such that $\text{Tr}(\lambda_i\lambda_j) = 2\delta_{ij}$ [18] (for δ_{ij} the Kronecker delta), and the terms $r_i = \text{Tr}(\rho\lambda_i)$ define the components of the real Bloch vector \mathbf{r} of ρ . By setting $m = M^2N^2 - 1$, we can hence let $K \subseteq \mathbb{R}^m$ be the convex set of Bloch vectors corresponding to valid quantum states in $H_{M,N}$. We remark that there is a scaling factor of $\sqrt{2}$ in Euclidean distance

between two states when switching between the two spaces (see Appendix A), which will not affect our analysis.

We can now formally state what is required to show Theorem 1. From the values of r, R, and p for $S_{M,N}$, it follows that $\langle S_{M,N} \rangle = m + \langle R \rangle + \langle r \rangle + \langle \mathbf{p} \rangle \leq \text{poly}(MN)$. Thus, by the complexity measure for Definition 4, our aim is to show NP-hardness of WMEM_{β}($S_{M,N}$) with respect to $MN + \langle \mathbf{y} \rangle + \lceil \frac{1}{\beta} \rceil$, where the $\lceil \frac{1}{\beta} \rceil$ term (as opposed to $\langle \beta \rangle$) follows from the fact that we wish to prove strong NP-hardness.

3 The Reduction

Let us show our main result, Theorem 1. To do so, we show the Turing reduction of Equation (2). We shall see that the third reduction in this chain runs in time polynomial in $1/\epsilon$, implying we must be able to choose $\epsilon \ge 1/\operatorname{poly}(M, N)$ in order for the entire chain to run in polynomial time. We return to this point later.

To begin, the first link in Equation (2) is given by the following theorem [5]. Unless otherwise stated, by a poly-time reduction, we mean with respect to the encoding size of the problem instance, as defined in Section 2. We use the notation $\Pi = (input \ parameters)$ to denote an instance Π of a given problem, with Π specified by the given input parameters.

Theorem 2 (Gurvits and Ioannou [5]). There exists a polynomial time Karp reduction which maps instance $\Pi_1 = (G, n, c)$ of CLIQUE to instance $\Pi_2 = (k, l, B_1, \ldots, B_k, \zeta, \eta)$ of RSDF, such that k = n(n-1)/2, l = n, $B_i \in \mathbb{Q}^{n \times n}$ and $\|B_i\|_{\mathrm{F}} \in \Theta(1)$ for all $1 \le i \le k$, $\zeta = \Theta(1)$, $\eta \in \Omega(n^{-2})$.

We refer the reader to Ref. [5] for the details of the proof of this theorem, but highlight here that it follows from direct application of the following theorem of Motzkin and Straus [19], which ties the maximum clique number of a graph to optimization of a square-free quadratic form over the unit simplex:

Theorem 3 (Motzkin and Straus [19]). Denote by $(i, j) \in G$ an edge in graph G between vertices i and j, and let k be the order of the maximal complete graph contained in G. Let Δ_n denote the simplex $\Delta_n := \{ \mathbf{x} \in \mathbb{R}^n \mid x_i \geq 0, \|\mathbf{x}\|_1 = 1 \}$. Then

$$\max_{\mathbf{x}\in\Delta_n}\sum_{(i,j)\in G} x_i x_j = \frac{1}{2}\left(1-\frac{1}{k}\right).$$
(4)

We remark that the matrices $B_i \in \mathbb{R}^{n \times n}$ from Theorem 2 will have the following structure — to each B_i , we uniquely assign an index (s, t) from the adjacency matrix A_G of G, $1 \le s < t \le n$, such that B_i has all entries zero, except for entries (s, t) and (t, s), which are set to the (s, t)th entry of A_G . We hence require k = n(n-1)/2 matrices B_i , as claimed by Theorem 2. Let us now state and prove the next link in Equation (2).

Lemma 4. There exists a poly-time Karp reduction which maps instance $\Pi_1 = (k, l, B_1, \ldots, B_k, \zeta, \eta)$ of RSDF to instance $\Pi_2 = (\mathbf{c}, \gamma)$ of WOPT_{ϵ}($\mathcal{S}_{M,N}$), where we define for convenience $\Delta := \sqrt{2\sum_{i=1}^k \|B_i\|_{\mathrm{F}}^2}$, and such that:

• M = k + 1

N = ^{l(l-1)}/₂ + 1
c = ĉ/ ||ĉ||₂ for some ĉ ∈ Q^m with ||ĉ||₂ ∈ O(m^{1/2}Δ) and m = M²N² - 1
γ = ¹/_{2||ĉ||₂} (√ζ + η + √ζ - η)
ε ≤ ^{√ζ+η-√ζ-η}/_{4||ĉ||₂(MN-1)+1}

Proof. Given instance Π_1 of RSDF, we map it to an instance Π_2 of WOPT_{ϵ}($\mathcal{S}_{M,N}$) as follows. We first rephrase the function $g(B_1, \ldots, B_k)$ from RSDF in terms of convex optimization over $\mathcal{S}_{M,N}$ via the following argument of Ref. [5]. Let M = k + 1, $N = \frac{l(l-1)}{2} + 1$, and consider the matrix $C \in \mathbb{R}^{MN \times MN}$, such that

$$C := \begin{pmatrix} 0 & A_1 & \dots & A_{M-1} \\ A_1 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ A_{M-1} & 0 & \dots & 0 \end{pmatrix},$$
 (5)

where each $A_i \in \mathbb{R}^{N \times N}$ is symmetric and all zeroes except for its upper-left $l \times l$ -dimensional submatrix, which we set to B_i . It is easy to see that $||C||_{\rm F} = \Delta$, as defined in the statement of our claim. Using C and Proposition 6.5 of [3], one can write:

$$\sqrt{g(B_1,\ldots,B_k)} = \max_{\mathbf{x}\in\mathbb{R}^N, \|\mathbf{x}\|_2=1} \sqrt{\sum_{i=1}^{M-1} (\mathbf{x}^{\mathrm{T}}A_i\mathbf{x})^2} = \max_{\rho\in\mathcal{S}_{M,N}} \mathrm{Tr}(C\rho).$$
(6)

Thus, Π_1 is reduced to optimizing the linear objective function $\operatorname{Tr}(C\rho)$ over all $\rho \in \mathcal{S}_{M,N}$.

Next, we use Equation (3) and the fact that $\operatorname{Tr}(C) = 0$ to rephrase $\operatorname{Tr}(C\rho)$ in terms of Bloch vectors, as required by WOPT_{ϵ}($\mathcal{S}_{M,N}$). We thus have

$$f(\rho) = \frac{1}{2} \sum_{i=1}^{M^2 N^2 - 1} r_i \cdot \operatorname{Tr}(C\sigma_i) = \hat{\mathbf{c}}^{\mathrm{T}} \mathbf{r},$$
(7)

for σ_i the generators of SU(MN), $\hat{c}_i = \frac{1}{2} \operatorname{Tr}(C\sigma_i)$, and **r** the Bloch vector of ρ . Set $m = M^2 N^2 - 1$, $\mathbf{c} = \hat{\mathbf{c}}/\|\hat{\mathbf{c}}\|_2$, and define objective functions $f(\mathbf{r}) = \mathbf{c}^{\mathrm{T}}\mathbf{r}$, and $f_{\max} = \max_{\mathbf{r}\in\mathcal{S}_{M,N}} f(\mathbf{r})$, where by $\mathbf{r} \in \mathcal{S}_{M,N}$, we denote a Bloch vector corresponding to some separable state $\rho \in \mathcal{S}_{M,N}$. Note that unless C is the zero matrix (i.e. the input graph to CLIQUE has no edges), we have $\|\hat{\mathbf{c}}\|_2 > 0$. Also, since $\operatorname{Tr}(\sigma_i \sigma_j) = 2\delta_{ij}$, it follows from Equation (7) and the Cauchy-Schwarz inequality that $\|\hat{\mathbf{c}}\|_2 \in O(m^{1/2}\Delta)$.

It hence remains to show the following (for γ and ϵ to be chosen): If $f_{\max} \geq \|\hat{\mathbf{c}}\|_2^{-1} \sqrt{\zeta + \eta}$, then there exists an $\mathbf{r} \in S(\mathcal{S}_{M,N}, -\epsilon)$ such that $f(\mathbf{r}) \geq \gamma + \epsilon$ (i.e. a "YES" instance of RSDF implies a "YES" instance of WOPT $_{\epsilon}(\mathcal{S}_{M,N})$). If $f_{\max} \leq \|\hat{\mathbf{c}}\|_2^{-1} \sqrt{\zeta - \eta}$, then for all $\mathbf{r} \in S(\mathcal{S}_{M,N}, \epsilon)$, $f(\mathbf{r}) \leq \gamma - \epsilon$ (i.e. a "NO" instance of RSDF implies a "NO" instance of WOPT $_{\epsilon}(\mathcal{S}_{M,N})$). The $\|\hat{\mathbf{c}}\|_2^{-1}$ term in these expressions follows from our definition of \mathbf{c} , and the square root in $\sqrt{\zeta + \eta}$ follows from Equation (6). We proceed case by case. Set $\gamma = \frac{1}{2\|\hat{\mathbf{c}}\|_2}(\sqrt{\zeta + \eta} + \sqrt{\zeta - \eta})$, and let us choose ϵ as needed. • Case 1: $f_{\max} \ge \frac{1}{\|\hat{\mathbf{c}}\|_2} \sqrt{\zeta + \eta}$.

Let $\mathbf{r}^* \in \mathcal{S}_{M,N}$ be such that $f(\mathbf{r}^*) = f_{\max}$. To find an $\mathbf{r}' \in S(\mathcal{S}_{M,N}, -\epsilon)$ such that $f(\mathbf{r}') \ge \gamma + \epsilon$, we first use the fact that for any well-bounded origin-centered convex set K, it holds that for all $\mathbf{x} \in K$, there exists a $\mathbf{y} \in S(K, -\epsilon)$ such that $\|\mathbf{x} - \mathbf{y}\|_2 \le 2\epsilon R/r$ [10] (where R and r are the radii of the ball containing K and the origin-centered ball contained within K, respectively). Plugging in the definitions of r and R for $\mathcal{S}_{M,N}$ from Section 2, it follows that there exists an $\mathbf{r}' \in S(\mathcal{S}_{M,N}, -\epsilon)$ such that $\|\mathbf{r}' - \mathbf{r}^*\|_2 \le 2(MN - 1)\epsilon$. Since f is linear, we can then write:

$$\left|f(\mathbf{r}') - f(\mathbf{r}^*)\right| = \left|\mathbf{c}^{\mathrm{T}}(\mathbf{r}' - \mathbf{r}^*)\right| \le \left\|\mathbf{c}\right\|_2 \left\|\mathbf{r}' - \mathbf{r}^*\right\|_2 \le 2(MN - 1)\epsilon,\tag{8}$$

where the first inequality follows from the Cauchy-Schwarz inequality. Thus, in order to have $f(\mathbf{r}') \geq \gamma + \epsilon$ as desired, it suffices to have

$$f(\mathbf{r}') \ge f_{\max} - 2(MN - 1)\epsilon \ge \gamma + \epsilon, \tag{9}$$

into which substitution of our values for γ and f_{max} gives that setting

$$\epsilon \le \frac{\sqrt{\zeta + \eta} - \sqrt{\zeta - \eta}}{4 \left\| \hat{\mathbf{c}} \right\|_2 (MN - 1) + 1} \tag{10}$$

suffices to conclude we have a "YES" instance of WOPT_{ϵ}($S_{M,N}$).

• CASE 2: $f_{\max} \leq \frac{1}{\|\hat{\mathbf{c}}\|_2} \sqrt{\zeta - \eta}$.

Let $\mathbf{r}^* \in \mathcal{S}_{M,N}$ be such that $f(\mathbf{r}^*) = f_{\max}$. To see that for all $\mathbf{r}' \in S(\mathcal{S}_{M,N}, \epsilon)$, we have $f(\mathbf{r}') \leq \gamma - \epsilon$, let $\mathbf{r}' \in S(\mathcal{S}_{M,N}, \epsilon)$. Then by the definition of $S(\mathcal{S}_{M,N}, \epsilon)$, there exists some $\mathbf{r} \in \mathcal{S}_{M,N}$ such that $\|\mathbf{r}' - \mathbf{r}\|_2 \leq \epsilon$. By Equation (8), it follows that

$$\left| f(\mathbf{r}') - f(\mathbf{r}) \right| \le \epsilon. \tag{11}$$

Thus, by considering $\mathbf{r} = \mathbf{r}^*$, we can conclude that $f(\mathbf{r}') \leq f_{\max} + \epsilon$ for any $\mathbf{r}' \in S(\mathcal{S}_{M,N}, \epsilon)$. To achieve $f(\mathbf{r}') \leq \gamma - \epsilon$ then, we set $f(\mathbf{r}') \leq f_{\max} + \epsilon \leq \gamma - \epsilon$, into which substitution of our values for γ and f_{\max} yields that setting

$$\epsilon \le \frac{\sqrt{\zeta + \eta} - \sqrt{\zeta - \eta}}{2 \left\| \hat{\mathbf{c}} \right\|_2 + 2} \tag{12}$$

suffices to conclude we have a "NO" instance of WOPT_{ϵ}($\mathcal{S}_{M,N}$).

Observe that combining Theorem 2 and Lemma 4 gives $M = N = \frac{n(n-1)}{2} + 1$. Following an argument of Ioannou [5], one can likewise show Lemma 4 for $M \ge N$ by padding the matrix C from its proof with extra $N \times N$ -dimensional zero matrices. Thus, the hardness result we show for WMEM_{β}($S_{M,N}$) by building on this link will be valid for $M \ge N$.

To show the last link of Equation (2), WOPT_{ϵ}($\mathcal{S}_{M,N}$) \leq_T WMEM_{β}($\mathcal{S}_{M,N}$), we finally apply the non-ellipsoidal Turing reduction of Liu [9], which holds for arbitrary p-centered well-bounded compact convex set $K \subseteq \mathbb{R}^m$. From Section 2, we know that $\mathcal{S}_{M,N}$ is such a set. **Theorem 5** (Liu [9]). Given instance $\Pi = (K, \mathbf{c}, \gamma, \epsilon)$ of WOPT_{ϵ}(K), for K with associated parameters (m, R, r, p), and $0 < \epsilon < 1$, there exists an algorithm which runs in time poly($\langle K \rangle, R, \lceil 1/\epsilon \rceil$), and solves Π using an oracle for WMEM_{β}(K) with $\beta = r^3 \epsilon^3 / [2^{13} 3^3 m^5 R^4 (R+r)]$.

Observe the dependence on $\lceil 1/\epsilon \rceil$ in the runtime of this reduction. We thus must be able to choose $\epsilon \geq 1/\operatorname{poly}(M, N)$ in order for our reduction chain of Equation (2) to run in polynomial time. Let us show that we can choose ϵ so. Specifically, by Lemma 4, we can set $\epsilon = (\sqrt{\zeta + \eta} - \sqrt{\zeta - \eta})/(4 \|\hat{\mathbf{c}}\|_2 (MN - 1) + 1)$. Piecing together Lemma 2 and Lemma 4, we have $\zeta \in \Theta(1)$ and $\eta \in \Omega(1/N)$. It follows that $\sqrt{\zeta + \eta} - \sqrt{\zeta - \eta} \in \Omega(1/N)$. To see this, recall that for given functions f(x) and g(x), we say $f(x) \in \Omega(g(x))$ if $f(x) \geq cg(x) \geq 0$ for all $x > x_0$ and positive constants c and x_0 [20]. Then one has, for positive constants c_1, c_2, N_1 , and N_2 :

$$\sqrt{\zeta + \eta} - \sqrt{\zeta - \eta} \geq \sqrt{c_1 + \frac{c_2}{N}} - \sqrt{c_1 - \frac{c_2}{N}} \qquad \forall N \geq \max(N_1, N_2)$$
(13)

$$= \frac{(c_1 + \frac{c_2}{N}) - (c_1 - \frac{c_2}{N})}{\sqrt{c_1 + \frac{c_2}{N}} + \sqrt{c_1 - \frac{c_2}{N}}}$$
(14)

$$\geq \frac{2c_2}{N(\sqrt{c_1 + c_2} + \sqrt{c_1})}.$$
(15)

The claim immediately follows. With a little thought, we further have that $\|\hat{\mathbf{c}}\|_2 \in O(\sqrt{N})$ (see Appendix A). Thus, we can always solve an instance of CLIQUE by choosing some $\epsilon \in \Omega(M^{-1}N^{-5/2})$ for WOPT_{ϵ}($\mathcal{S}_{M,N}$), as desired.

Substituting our values for r and R for $S_{M,N}$ and for m from Lemma 4 into the expression for β in Theorem 5 gives that $\beta = \text{poly}(M^{-1}, N^{-1}, \epsilon)$. Combining Theorem 2, Lemma 4, and Theorem 5 thus yields our desired result — a polynomial time Turing reduction from CLIQUE to WMEM_{β}($S_{M,N}$) with $\beta \ge 1/\text{poly}(M,N)$. Specifically, we have the following, which immediately implies Theorem 1.

Theorem 6. Given instance $\Pi = (G, n, c)$ of CLIQUE, there exists an algorithm which solves Π in time poly(n) using an oracle for WMEM_{β}($S_{M,N}$) with parameters $M = N = \frac{n(n-1)}{2} + 1$ and some $\beta \in \Omega(n^{-73})$. More generally, for $N = \frac{n(n-1)}{2} + 1$ and any choice of $M \ge N$, the result holds for some $\beta \in \Omega(M^{-16}N^{-20.5})$.

Theorem 1 (*revised*). WMEM_{β}($\mathcal{S}_{M,N}$) is NP-hard for $M \ge N$ and $\beta \le \text{poly}(M^{-16}N^{-20.5})$, or equivalently, is strongly NP-hard.

We stress the phrase "some $\beta \in \Omega(n^{-73})$ " in the statement of Theorem 6 above — specifically, we cannot have $\beta \in O(1)$ in our reduction, due, for example, to the expression for ϵ in Lemma 4. We remark that the major contributor to the large negative exponent on n in the estimate for β is the reduction of Theorem 5, which is based largely on well-known results by Grötschel *et al* [10] dating from the late 1980's. In Section 5, we briefly discuss a possible alternate reduction from WOPT_{ϵ}(K) to WMEM_{β}(K) based on random walks, as suggested by Liu [9], which may lead to a better estimate for β .

4 Applications

Let us now observe two applications of Theorem 6. First, one immediately has a lower bound on the maximum distance a bound entangled state can have from $S_{M,N}$. To see this, recall that bound entangled states are mixed entangled quantum states from which no pure (state) entanglement can be distilled [13], and are the only entangled states whose entanglement is capable of escaping detection by the Peres-Horodecki Positive Partial Transpose (PPT) [11, 12] criterion⁴. Theorem 1 implies that unless P = NP, any test of membership for $S_{M,N}$ must be unable to efficiently resolve $S_{M,N}$ within distance $\beta \in \Omega(M^{-16}N^{-20.5})$ of its border in the general case. It follows that unless P = NP, there must exist bound entangled state(s) $\rho_{be} \in H_{M,N}$ such that for any separable state $\rho_{sep} \in S_{M,N}$, $\|\rho_{be} - \rho_{sep}\|_{\rm F} \in \Omega(M^{-16}N^{-20.5})$ — if not, one could determine the separability of any quantum state within this region using the PPT test, contradicting Theorem 1. For this reason, it would be of interest to determine precisely how large one can make β before NP-hardness of WMEM_{β}($S_{M,N}$) ceases to hold.

Second, we use the Jamiołkowski isomorphism [14] to show NP-hardness of the problem of determining whether a completely positive (CP) trace-preserving (TP) linear map (also known as a *channel*) is entanglement-breaking (which we define shortly). Let $H_{MN,MN}$ denote the set of linear operators mapping ($\mathbb{C}^M \otimes \mathbb{C}^N$) \otimes ($\mathbb{C}^M \otimes \mathbb{C}^N$) to itself, where the second tensor product is between subsystems A and B, and let $S_{MN,MN} \subset H_{MN,MN}$ denote the subset of separable density operators. For any linear map $\Phi : H_{M,N} \mapsto H_{M,N}$, one can associate a unique operator in $H_{MN,MN}$ via the Jamiołkowski isomorphism [14], denoted $J(\Phi)$, such that:

$$J(\Phi) := [\Phi \otimes I] \left(|\phi^+\rangle \langle \phi^+| \right), \tag{16}$$

where $|\phi^+\rangle$ is the maximally entangled state $|\phi^+\rangle = \frac{1}{\sqrt{MN}} \sum_{k=0}^{MN-1} |k\rangle \otimes |k\rangle$, for $\{|k\rangle\}_{k=0}^{MN-1}$ an arbitrary orthonormal basis for $\mathbb{C}^M \otimes \mathbb{C}^N$. In particular, if and only if Φ is completely positive (CP), then $J(\Phi)$ is positive semidefinite [22], and if and only if Φ is trace-preserving (TP), then $\operatorname{Tr}_A(J(\Phi)) = I/MN$ [2]. Via this isomorphism, it is known that the set of *entanglement-breaking* (EB) channels [2] acting on $H_{M,N}$ is isomorphic to the set of separable states $\rho \in \mathcal{S}_{MN,MN}$ with $\operatorname{Tr}_A(\rho) = I/MN$. Intuitively, an EB channel Φ has the property that for any input quantum state $\rho \in H_{M,N}$, $(\Phi \otimes I)(\rho)$ is separable. Such channels can be defined formally as follows:

Definition 6 (Entanglement-Breaking (EB) Channels [15]). Given a CP and TP linear map Φ : $H_{M,N} \mapsto H_{M,N}$, the following are equivalent:

- 1. Φ is entanglement-breaking (EB).
- 2. $J(\Phi)$ is separable and $\operatorname{Tr}_A(J(\Phi)) = I/MN$.
- 3. Φ can be written⁵ $\Phi(\rho) = \sum_{i} \sigma_i \operatorname{Tr}(F_i \rho)$, for σ_i density matrices and $\{F_i\}_i$ a Positive Operator Valued Measure (POVM).
- 4. Φ can be written $\Phi(\rho) = \sum_i K_i \rho K_i^{\dagger}$ for Kraus operators [24] K_i strictly of rank 1, and such that $\sum_i K_i^{\dagger} K_i = I$.
- 5. $\Gamma \circ \Phi$ and $\Phi \circ \Gamma$ are completely positive for all positive TP maps Γ .

Note first that if Φ is not TP, an analogue of the above definition holds [15] for $\operatorname{Tr}_A(J(\Phi)) \neq I/MN$, $\{F_i\}_i$ positive semidefinite but not a POVM, $\sum_i K_i^{\dagger} K_i < I$, and non-TP maps Γ . In

 $^{^{4}}$ The converse question of whether *all* bound entangled states escape such detection by the PPT test is, however, a major open question [21].

⁵This is sometimes dubbed the Holevo form, as introduced by Holevo [23].

this case, the set of CP but not necessarily TP entanglement-breaking maps Φ acting on $H_{M,N}$ is isomorphic to the entire separable set $S_{MN,MN}$ — it thus trivially follows from Theorem 6 that if we denote the (convex) set of EB maps acting on $H_{M,N}$ as $\mathcal{B}_{M,N}$ (where we assume the elements of $\mathcal{B}_{M,N}$ are given in their Jamiołkowski form), then WMEM_{β}($\mathcal{B}_{M,N}$) is also NP-hard for $\beta \leq \text{poly}(M^{-16}N^{-20.5})$, i.e. is strongly NP-hard.

Let us show that the task of determining whether a given map is EB is still NP-hard when restricted to the case of quantum channels, which is an important case of interest, as channels correspond to physically realizable processes. Assume we are asked to determine the separability of an arbitrary quantum state $\rho \in H_{MN,MN}$, and that we have an oracle Q which, given $J(\Phi)$ for some CP and TP map Φ acting on $H_{M,N}$, efficiently determines whether Φ is EB. Let $\rho_B = \text{Tr}_A(\rho)$. If $\rho_B = I/MN$, by Definition 6, we can determine the separability of ρ by simply feeding ρ into Q and returning the oracle's answer. Suppose then that $\rho_B \neq I/MN$. If ρ_B is positive definite, we can apply local filters (as first considered in [25]) to obtain a state ρ' with a maximally mixed subsystem B. Specifically, define the locally acting CP map

$$\Psi(\rho) := \left(I \otimes \rho_B^{-1/2}\right) \rho \left(I \otimes \rho_B^{-1/2}\right).$$
(17)

Note that Ψ corresponds to a probabilistic local operation⁶, since it is not TP (i.e. if $K_1 := I \otimes \rho_B^{-1/2}$ is a Kraus operator, then $K_1^{\dagger}K_1 < I$). It is easy to see that $\operatorname{Tr}_A(\Psi(\rho)) = I/MN$. Further, Ψ is an invertible map, and so with non-zero probability, one can recover ρ from $\Psi(\rho)$. Thus, since entanglement cannot be created via local operations and classical communication (LOCC), $\Psi(\rho)$ is entangled if and only if ρ is. We can hence feed $\Psi(\rho)$ into Q and return its response, as before. Finally, if ρ_B is not positive definite, then before applying the local filter, we process ρ via the following channel Υ [26]:

$$\Upsilon(\rho) := (1-p)\rho + p\frac{1}{\alpha}I \otimes P_{\operatorname{supp}(\rho_B)}^{\perp}.$$
(18)

Here, $p \in (0,1)$, $P_{\text{supp}(\rho_B)}^{\perp}$ is the projector onto the orthogonal complement of the support of ρ_B , and $\alpha = \text{Tr}(I \otimes P_{\text{supp}(\rho_B)}^{\perp}) = MN(MN - \text{rank}(\rho_B))$. Observe that

$$\operatorname{Tr}_{A}(\Upsilon(\rho)) = (1-p)\rho_{B} + p\frac{1}{\alpha_{B}}P_{\operatorname{supp}(\rho_{B})}^{\perp}$$
(19)

is full rank and hence positive definite, where $\alpha_B = MN - \operatorname{rank}(\rho_B)$. Further, given $\Upsilon(\rho)$, one can recover the initial state ρ with non-zero probability by performing a von Neumann measurement on $\Upsilon(\rho)$ consisting of projectors $\left\{ I \otimes P_{\operatorname{supp}}(\rho_B), I \otimes P_{\operatorname{supp}}^{\perp}(\rho_B) \right\}$. Since this channel can be implemented using LOCC (i.e. with probability p two parties, Alice and Bob, agree to act invariantly on their halves of ρ , and with probability 1 - p, they agree to replace the state completely by locally preparing the states I/MN and $P_{\operatorname{supp}}^{\perp}(\rho_B)/\alpha_B$, respectively), it follows that $\Upsilon(\rho)$ is entangled if and only if ρ is. We thus apply the local filter of Equation 17 to $\Upsilon(\rho)$ and proceed as in the previous case. Therefore, if one can determine whether a channel $\Phi : H_{M,N} \mapsto H_{M,N}$ is EB, then one could determine the separability of $\rho \in H_{MN,MN}$. The former task must hence be NP-hard.

Finally, we remark that although we have assumed here that the input map Φ is specified in Jamiołkowski form, it is typically straightforward to move to another representation, such as the

 $^{^{6}}$ Such maps are sometimes classified under stochastic local operations and classical communication (SLOCC) [1]

Kraus representation. For example, the canonical⁷ way to determine a Kraus representation for Φ given the Jamiołkowski form $J(\Phi)$ is to reshuffle the eigenvectors of $J(\Phi)$ into matrices (i.e. Kraus operators) K_i . We refer the reader to [2] for further details.

5 Concluding Comments

We have shown that the problem of Weak Membership over the set of separable quantum states $S_{M,N}$ is strongly NP-hard, implying it is NP-hard even if one allows "moderate" error, i.e. $\beta \leq 1/\operatorname{poly}(M,N)$. This shows that it is NP-hard to determine whether an arbitrary quantum state within an inverse polynomial distance from the separable set is entangled. It would be interesting to know whether our estimate for β here can be improved to O(1), or perhaps more ambitiously, if one could determine precisely how large β can be before NP-hardness of WMEM_{β}($S_{M,N}$) ceases to hold. As discussed in Section 4, such a result would have immediate implications to the study of bound entanglement.

One possible approach to improving our estimate for β , following a suggestion of Liu [9], may be to replace the reduction of Theorem 5 with the Bertsimas-Vempala random walk algorithm for convex optimization [6]. The latter solves optimization problems over a convex set K given a membership oracle for K. This alone, however, would not suffice to improve our estimate to $\beta \in O(1)$, as an inverse dependence on the dimension is already introduced in reducing CLIQUE to RSDF (i.e. the first link of the reduction chain). We refer the interested reader to [9] for further details regarding this alternate approach.

6 Acknowledgements

We thank the following: Ben Toner for ideas which spurred this line of thought, Richard Cleve and Stephen Vavasis for helpful discussions, and John Watrous and Marco Piani for bringing local filters to our attention. This work was partially supported by Canada's NSERC, CIAR and MITACS.

A Appendix

Lemma 7. Given density operators ρ_1 and ρ_2 of quantum states of dimension N, with corresponding Bloch vectors $\vec{\alpha}$ and $\vec{\beta}$ given by equations $\rho_1 = \frac{I}{N} + \frac{1}{2}\vec{\alpha} \cdot \vec{\sigma}$ and $\rho_2 = \frac{I}{N} + \frac{1}{2}\vec{\beta} \cdot \vec{\sigma}$ (as per Equation (3)), respectively, we have $\sqrt{2} \|\rho_1 - \rho_2\|_{\rm F} = \|\vec{\alpha} - \vec{\beta}\|_2$.

⁷Unlike the Jamiołkowski representation, the operator sum representation is not unique.

Proof. Via straightforward manipulation and the fact that $Tr(\sigma_i \sigma_j) = 2\delta_{ij}$, we have:

$$\begin{aligned} \|\rho_{1} - \rho_{2}\|_{\mathrm{F}} &= \frac{1}{2} \left\| \sum_{i=1}^{N^{2}-1} (\alpha_{i} - \beta_{i})\sigma_{i} \right\|_{\mathrm{F}} \\ &= \frac{1}{2} \sqrt{\mathrm{Tr} \left((\sum_{i=1}^{N^{2}-1} (\alpha_{i} - \beta_{i})\sigma_{i})^{\dagger} (\sum_{j=1}^{N^{2}-1} (\alpha_{j} - \beta_{j})\sigma_{j}) \right)} \\ &= \frac{1}{2} \sqrt{\sum_{i=1,j=1}^{N^{2}-1} (\alpha_{i} - \beta_{i})(\alpha_{j} - \beta_{j})2\delta_{ij}} \\ &= \frac{1}{\sqrt{2}} \|\vec{\alpha} - \vec{\beta}\|_{2} \end{aligned}$$

Lemma 8. Combining Theorem 2 and Lemma 4 gives $\|\hat{\mathbf{c}}\|_2 \in O(\sqrt{N})$.

Proof. By definition, we have:

$$\|\hat{\mathbf{c}}\|_{2} = \sqrt{\sum_{i=1}^{m} \left[\frac{1}{2} \operatorname{Tr}(C\sigma_{i})\right]^{2}},\tag{20}$$

where $m = M^2 N^2 - 1$. Recall now the definition of C from Equation (5), where in our case, each $A_i \in \mathbb{R}^{N \times N}$ is all zeroes except for its upper left corner, which is set to submatrix $B_i \in \mathbb{R}^{l \times l}$ from Theorem 2. Each B_i in turn is all zeroes, except for some index (k, l) (and hence (l, k), by symmetry), $1 \leq k < l \leq n$, which is set to the (k, l)th entry of the adjacency matrix A_G of graph G (see Theorem 2 and ensuing discussion). We also require an explicit construction for the generators σ_i of SU(MN), given for example in [18], where $\{\sigma_i\}_{i=1}^{M^2N^2-1} = \{U_{pq}, V_{pq}, W_r\}$, such that for $1 \leq p < q \leq MN$ and $1 \leq r \leq MN - 1$, and $\{\mathbf{x}_i\}_{i=1}^{MN}$ an orthonormal basis for Hilbert space \mathcal{H}^{MN} :

$$U_{pq} = \mathbf{x}_p \mathbf{x}_q^{\dagger} + \mathbf{x}_q \mathbf{x}_p^{\dagger} \tag{21}$$

$$V_{pq} = -i\mathbf{x}_p \mathbf{x}_q^{\dagger} + i\mathbf{x}_q \mathbf{x}_p^{\dagger} \tag{22}$$

$$W_r = \sqrt{\frac{2}{r(r+1)}} \left[\left(\sum_{k=1}^r \mathbf{x}_k \mathbf{x}_k^{\dagger} \right) - r \mathbf{x}_{r+1} \mathbf{x}_{r+1}^{\dagger} \right].$$
(23)

Due to the symmetry of C and the fact that $\operatorname{Tr}(C) = 0$, it is clear that only the generators of the form U_{pq} will contribute to the sum in Equation (20). Further, for each edge in G, $\operatorname{Tr}(CU_{pq}) = 2$ for each U_{pq} whose non-zero indices match those of the entries in C corresponding to that edge. Since each edge contributes four (symmetrically placed) entries of 1 to C, we hence have $\|\hat{\mathbf{c}}\|_2 = \frac{1}{2}\sqrt{(2\hat{c})2^2} = \sqrt{2\hat{e}}$, where \hat{e} denotes the number of edges in G. Since $\hat{e} \in O(n^2)$ (n the number of vertices in G), and $N \in \Theta(n^2)$, we have $\|\hat{\mathbf{c}}\|_2 \in O(\sqrt{N})$, as required.

References

- R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki. Quantum entanglement. arXiv:quant-ph/0702225v2, 2007.
- [2] I. Bengtsson and K. Życzkowski. *Geometry of Quantum States: An Introduction to Quantum Entanglement.* Cambridge University Press, 2006.
- [3] L. Gurvits. Classical deterministic complexity of edmonds problem and quantum entanglement. In Proceedings of the thirty-fifth ACM symposium on Theory of computing, pages 10–19, New York, 2003. ACM Press.
- [4] M. Garey and D. Johnson. Computers and Intractability: A Guide to the Theory of NPcompleteness. W.H. Freeman and Company, New York, 1979.
- [5] L. Ioannou. Computational complexity of the quantum separability problem. Quantum Information and Computation, 7(4):335, 2007.
- [6] D. Bertsimas and S. Vempala. Solving convex programs by random walks. Journal of the ACM, 51(4), 2004.
- [7] D. B. Yudin and A. S. Nemirovskii. Informational complexity and efficient methods for the solution of convex extremal problems (in Russian). *Ekonomica i Matematicheskie Metody*, 12:357–369, 1976.
- [8] M. R. Garey and D. S.Johnson. Computers and Intractability: A Guide to the theory of NP-completeness. W. H. Freeman and Company, New York, 1979.
- [9] Y.K. Liu. The complexity of the consistency and N-representability problems for quantum states. http://arxiv.org/abs/0712.3041, 2007. PhD Thesis.
- [10] M. Grötschel, L. Lovász, and A. Schrijver. Geometric Algorithms and Combinatorial Optimization. Springer-Verlag, Berlin, 1988.
- [11] A. Peres. Separability criterion for density matrices. *Physical Review Letters*, 77:1413, 1996.
- [12] M. Horodecki, P. Horodecki, and R. Horodecki. Separability of mixed states: necessary and sufficient conditions. *Physics Letters A*, 223(1):1–8, 1996.
- [13] M. Horodecki, P. Horodecki, and R. Horodecki. Mixed-state entanglement and distillation: Is there a "bound" entanglement in nature? *Phys. Rev. Lett.*, 80(24):5239–5242, 1998.
- [14] A. Jamiołkowski. Linear transformations which preserve trace and positive semi-definiteness of operators. *Rep. Math. Phys.*, 3:275, 1972.
- [15] M. Horodecki, P. W. Shor, and M. B. Ruskai. Entanglement breaking channels. *Rev. Math. Phys.*, 15:629, 2003.
- [16] L. Gurvits and H. Barnum. Largest separable balls around the maximally mixed bipartite quantum state. *Phys. Rev. A*, 66(6):062311, 2002.
- [17] E. B. Vinberg. Linear Representations of Groups. Birkhäuser Basel, 1989.

- [18] G. Kimura. The bloch vector for N-level systems. *Physics Letters A*, 314(5):339–349, 2003.
- [19] T. S. Motzkin and E. G. Straus. Maxima for graphs and a new proof of a theorem of Túran. Canadian J. Math., 17:533–540, 1965.
- [20] T. H. Cormen, C. E. Leiserson, and R. L. Rivest. Introduction to Algorithms. The MIT Press, 1990.
- [21] D. Bruß. Characterizing entanglement. Journal of Mathematical Physics, 43(9), 2002.
- [22] M.-D. Choi. Completely positive linear maps on complex matrices. Linear Alg. Appl., 10:285, 1975.
- [23] A. S. Holevo. Coding theorems for quantum channels. Russian Math. Surveys, 53:1295–1331, 1999.
- [24] K. Kraus. General state changes in quantum theory. Ann. Phys., 64:311, 1971.
- [25] N. Gisin. Hidden quantum nonlocality revealed by local filters. Phys. Lett. A, 210(3):151– 156(6), 1996.
- [26] M. Piani. Private communication, 2008.